

interleukin-6 in cerebrospinal fluid: association with clinical outcome for children with bacterial meningitis. Clinical Infectious Diseases 25: 328-329.

Ichiyama T, Nishikawa M, Yoshitomi T, Hayashi T, Furukawa S (1998): Tumor necrosis factor- α , interleukin-1 β , and interleukin-6 in cerebrospinal fluid from children with prolonged febrile seizures. Neurology 50: 407-411.

Kimura H, Futamura M, Kito H, Ando T, Goto M, Kuzushima K, Shibata M, Morishima T (1991): Detection of viral DNA in neonatal herpes simplex virus infections: frequent and prolonged presence in serum and cerebrospinal fluid. Journal of Infectious Diseases 164: 289-293.

Kimura H, Aso K, Kuzushima K, Hanada N, Shibata M, Morishima T (1992): Relapse of herpes simplex encephalitis in children. Pediatrics 89: 891-894.

Lehmann NI, Gust ID (1971): Viremia in influenza. A report of two cases. Medical Journal of Australia 2: 1166-1169.

López-Cortés LF, Cruz-Ruiz M, Gómez-Mateos J, Jiménez-Hernández D, Polomino J, Jiménez E (1993): Measurement of levels of tumor necrosis factor- α and interleukin-1 β in the CSF of patients with meningitis of different etiologies: utility in the differential diagnosis. Clinical Infectious Diseases 16: 534-539.

Martín-Ancel A, García-Alix A, Pascual-Salcedo D, Cabañas F, Valcarce M, Quero J (1997): Interleukin-6 in the cerebrospinal fluid after perinatal asphyxia is related to early and late neurological manifestations. *Pediatrics* 100: 789-794.

McConkey B, Oxon BM, Daws RA (1958): Neurological disorders associated with Asian influenza. *Lancet* ii: 15-17.

Messer J, Eyer D, Donato L, Gallati H, Matis J, Simeoni U (1996): Evaluation of interleukin-6 and soluble receptors of tumor necrosis factor for early diagnosis of neonatal infection. *Journal of Pediatrics* 129: 574-580.

Mori T, Nagafuji H, Matsumoto K, Kimura Y (1997): Use of polymerase chain reaction for demonstration of influenza virus dissemination in children. *Clinical Infectious Diseases* 24: 736-737.

Murphy BR, Webster RG (1996): Orthomyxoviruses. In Fields BN, Knipe DM, Howley PM (eds.): "Fields Virology. 3rd ed." Philadelphia: Lippincott-Raven Publishers, pp.1397-1445.

Mustafa MM, Lebel MH, Ramilo O, Olsen KD, Reisch JS, Beutler B, McCracken Jr GH (1989): Correlation of interleukin-1 β and cachectin concentrations in cerebrospinal fluid and outcome from bacterial meningitis. *Journal of Pediatrics* 115: 208-213.

Nakajima S, Sugiura A (1980): Neurovirulence of influenza virus in mice II. Mechanism of virulence as studied in a neuroblastoma

cell line. Virology 101: 450-457.

Panero A, Pacifico L, Rossi N, Mancuso G, Stegagno M, Chiesa C (1997): Interleukin 6 in neonates with early and late onset infection. Pediatric Infectious Disease Journal 16: 370-375.

Protheroe SM, Mellor DH (1991): Imaging in influenza A encephalitis. Archives of Disease in Childhood 66: 702-705.

Reilly PL, Simpson DA, Sprod R, Thomas L (1988): Assessing the conscious level in infants and young children: a paediatric version of the Glasgow Coma Scale. Childs Nervous System 4: 30-33.

Saija A, Princi P, Lanza M, Scalese M, Aramnejad E, De Sarro A. (1995): Systemic cytokine administration can affect blood-brain barrier permeability in the rat. Life Sciences 56: 775-784.

Stanley ED, Jackson GG (1969): Viremia in Asian influenza. Transactions of the Association of American Physicians 79: 376-387.

Sugiura A, Ueda M (1980): Neurovirulence of influenza virus in mice I. Neurovirulence of recombinants between virulent and avirulent virus strains. Virology 101: 440-449.

Takahashi M, Yamada T (1995): Nakajima S, Nakajima K, Yamamoto T, Okada H. The substantia nigra is a major target for

neurovirulent influenza A virus. Journal of Experimental Medicine 181: 2161-2169.

Teasdale G, Jennett B (1974): Assessment of coma and impaired consciousness. A practical scale. Lancet ii: 81-84.

Tsuruoka H, Xu H, Kuroda K, Hayashi K, Yasui O, Yamada A, Ishizaki T, Yamada Y, Watanabe T, Hosaka Y (1997): Detection of influenza virus RNA in peripheral blood mononuclear cells of influenza patients. Japanese Journal of Medical Science & Biology 50: 27-34.

Ueda T, Miyake Y, Imoto K, Hattori S, Miyake S, Ishizaki T, Yamada A, Kurata T, Nagai T, Suga S, Asano Y (1996): Distribution of human herpesvirus 6 and varicella-zoster virus in organs of fatal case with exanthem subitum and varicella. Acta Paediatrica Japonica 38: 590-595.

TABLE 1. Clinical features of patients with influenza-virus-associated encephalopathy

Patient no.	Age/Sex	GCS ^a	Convulsion	Cerebrospinal fluid		Serum			Mortality and morbidity
				Cell count (/μL)	Protein (mg/dL)	AST ^b (IU/L)	ALT ^c (IU/L)	NH ₃ (μg/dL)	
1	2y/F	11	yes	2	24	92	24	18	recovery
2	2y/M	12	yes	15	27	49	15	34	recovery
3	2y/M	13	yes	0	13	26586	13879	74	recovery
4	2y/M	12	yes	6	21	56	41	53	recovery
5	3y/M	3	yes	0	57	18088	10472	50	sequelae
6	5y/M	3	yes	6	20	1276	1667	37	sequelae
7	6y/F	3	yes	NA ^d	NA	32	14	NA	recovery
8	6y/F	11	no	0	10	39	17	NA	sequelae
9	11y/M	11	no	NA	NA	200	72	NA	sequelae
10	11y/M	11	yes	3	23	35	13	21	recovery
11	13y/F	3	yes	NA	NA	10510	3160	NA	death

^aGCS, Glasgow Coma Scale.^bAST, aspartate aminotransferase.^cALT, alanine aminotransferase.^dNA, not applicable.

TABLE 2. Results of RT-PCR in patients with influenza-virus-associated encephalopathy

Samples	Patients	Controls
Throat swab	9/9	29/29
Plasma	0/11	0/29
PBMC ^a	1/9	0/29
Erythrocytes	0/9	0/29
CSF ^b	1/11	ND ^c

^aPBMC, peripheral blood mononuclear cells.

^bCSF, cerebrospinal fluid.

^cND, not done.

TABLE 3. Cerebrospinal fluid concentrations of TNF- α , sTNF-R1, IL-1 β , and IL-6 in patients with influenza-virus-associated encephalopathy

Patient no.	TNF- α ^a (pg/mL)	sTNF-R1 ^b (pg/mL)	IL-1 β (pg/mL)	IL-6 (pg/mL)
1	NA ^d	NA	NA	NA
2	NA	NA	NA	NA
3	<15	1196	<4	324 ^f
4	<15	2934	NA	<31.2
5	<15	1848	<4	<31.2
6	NA	NA	NA	NA
7	<15	555	<4	<31.2
8	<15	433	<4	<31.2
9	<15	553	<4	<31.2
10	<15	635	<4	<31.2
11	NA	NA	NA	NA
Normal range	<15	836 ± 402 ^e	<4	<31.2

^aTNF- α , tumor necrosis factor- α .

^bsTNF-R1, soluble tumor necrosis factor receptor 1.

^cIL, interleukin.

^dNA, not applicable.

^eMean ± SD.

^fUnderscores represent the level considered abnormal.

TABLE 4. Plasma concentrations of TNF- α , sTNF-R1, IL-1 β , and IL-6 in patients with influenza-virus-associated encephalopathy

Patient no.	TNF- α ^a (pg/mL)	sTNF-R1 ^b (pg/mL)	IL-1 β (pg/mL)	IL-6 (pg/mL)
1	NA ^d	NA	NA	NA
2	NA	NA	NA	NA
3	<31.2	2232	<8	860 ^f
4	<31.2	810	30.2	18.2
5	<31.2	702	<8	<12.5
6	NA	426	<8	<12.5
7	<31.2	760	<8	<12.5
8	<31.2	869	<8	100
9	<31.2	\geq 5000	<8	1295
10	<31.2	745	<8	<12.5
11	<31.2	270	21.1	<12.5
Normal range	<15.6	1020 \pm 495 ^e	<4	<12.5

^aTNF- α , tumor necrosis factor- α .

^bsTNF-R1, soluble tumor necrosis factor receptor 1.

^cIL, interleukin.

^dNA, not applicable.

^eMean \pm SD.

^fUnderscores represent the level considered abnormal.

TABLE 5. Comparison of the percentage of patients exhibiting plasma cytokines

Cytokines (pg/mL)	Patients (%) (n=9)	Controls (%) (n=29)	P
TNF- α ^a	0 (0)	0 (0)	1.00
sTNF-R1 ^b	2 (22)	1 (3)	0.13
IL-1 β ^c	2 (22)	2 (7)	0.23
IL-6	4 (44)	12 (41)	0.58
IL-6 (≥ 100)	3 (30)	0 (0)	0.01

^aTNF- α , tumor necrosis factor- α .

^bsTNF-R1, soluble tumor necrosis factor receptor 1.

^cIL, interleukin.




Fig. 1. Sensitivity of the RT-PCR in detecting influenza virus NS gene. Lane 1, 200bp DNA marker ladder; lanes 2 to 4, 3×10^2 , 3×10^1 , 3 copies of NS gene, respectively; lane 5, no template control.