

Antisense & nucleic acid drug development 8:441-450, Phosphorothioate hammerhead ribozymes targeting a conserved sequence in V3 loop region inhibit HIV-1 entry.	1998		Zhang X., Iwatani Y, Shimayama T., Yamada R., Koito A., Xu Y., Sakai, H., Uchiyama T., and Hattori T.
J. Virol., 72: 5414-5424 Threshold number of provirus copies required per cell for efficient virus production and interference in Moloney murine leukemia virus-infected NIH 3T3 cells.	1998		Odawara T., Oshima M., Doi K., Iwamoto A., and Yoshikura H.
J. Virol., 72: 6414-6420 cis Elements required for high-level expression of unspliced gag-containing message in Moloney murine leukemia virus.	1998		Oshima M., Odawara T., Hanaki K., Igarashi H., and Yoshikura H.
Microbiol. Immunol., 42, 305-311 Detection of HIV-Gag p24-specific antibodies in sera and saliva of HIV-1-infected adults and in sera of infants born to HIV-1-infected mothers.	1998		Yasuda, S., Iwasaki, M., Oka, S., Naganawa, S., Nakasone, T., Honda, M., Sata, T., Kojima, A., Matsuda, S., Takemori, T., and Tsunetsugu-Yokota, Y.
J. Virol., 72, 6257-6259 Inhibition of human Immunodeficiency virus type 1 virion entry by dominant-negative Hck.	1998		Tokunaga, K., Kiyokawa, E., Nakaya, M., Otsuka, N., Kojima, A., Kurata, T. and Matsuda, M
J. Gen. Virol., 78, 2447-2453 Enhancement of human immunodeficiency virus type 1 infectivity by Nef is producer cell-dependent.	1998		Tokunaga, K., Kojima, A., Kurata, T., Ikuta, K., Akari, H., Koyama, A H., Kawamura, M., Inubushi, R., Shimano, R. and Adachi, A.
Biochem. Biophys. Res. Commun., 250, 565-568 Producer cell-dependent requirement of the Nef protein for efficient entry of HIV-1 into cells.	1998		Tokunaga, K., Kojima, A., Kurata, T., Ikuta, K., Inubushi, R., Shimano, R. Kawamura, M., Akari, H., Koyama, H. and Adachi, A

5. 研究成果による特許権等の知的財産権の取得状況