

199900397A

以降は雑誌/図書等に掲載された論文となりますので「研究成果の刊行に関する一覧」をご参照ください。

「研究成果の刊行に関する一覧」

YAC and cosmid contigs encompassing the Fukuyama-type congenital muscular dystrophy (FCMD) candidate region on 9q31.

Miyake M, Nakahori Y, Matsushita I, Kobayashi K, Mizuno K, Hirai M, Kanazawa I, Nakagome Y, Tokunaga K, Toda T.
Genomics. 1997 Mar 1;40(2):284–93.

Polymorphism analysis of Fukuyama type congenital muscular dystrophy (FCMD) siblings with different phenotypes.

Kondo-Iida E, Saito K, Osawa M, Ishihara T, Toda T, Fukuyama Y.
Brain Dev. 1997 Apr;19(3):181–6.

Molecular genetic evidence of clinical heterogeneity in Fukuyama-type congenital muscular dystrophy.

Kondo-Iida E, Saito K, Tanaka H, Tsuji S, Ishihara T, Osawa M, Fukuyama Y, Toda T.
Hum Genet. 1997 Apr;99(4):427–32.

The clinical spectrum and genetic studies of Fukuyama congenital muscular dystrophy

Mieko Yoshioka, Tatsushi Toda, and Shigekazu Kuroki
Congenital muscular dystrophies Elsevier, 1997

Toward identification of the Fukuyama type congenital muscular dystrophy gene

Tatsushi Toda, Masashi Miyake, Yutaka Nakahori, Masaya Segawa, Yoshiko Nomura, Ikuya Nonaka, Shiro Ikegawa, Eri Kondo, Kayoko Saito,

Makiko Osawa, Yukiko Fukuyama, Mieko Yoshioka, Teruo Shimizu, Ichiro Kanazawa, Yusuke Nakamura and Yasuo Nakagome
Congenital muscular dystrophies Elsevier, 1997

Reconfirmation of the Fukuyama congenital muscular dystrophy (FCMD) gene locus at chromosome 9q31, and a successful prenatal diagnosis of FCMD in two families

Eri Kondo, Kayoko Saito, Tatsushi Toda, Makiko Osawa, Hajime Tanaka, Shoji Tsuji, Tomoko Yamamoto, Makio Kobayashi, Yusuke Nakamura and Yukiko Fukuyama

Congenital muscular dystrophies Elsevier, 1997

Prenatal diagnosis of Fukuyama type congenital muscular dystrophy in eight Japanese families by haplotype analysis using new markers closest to the gene.

Saito K, Kondo-Iida E, Kawakita Y, Juan D, Ikeya K, Osawa M, Fukuyama Y, Toda T, Nakabayashi M, Yamamoto T, Kobayashi M.
Am J Med Genet. 1998 May 26;77(4):310-6.

Prenatal diagnosis of Fukuyama-type congenital muscular dystrophy by microsatellite analysis.

Takai Y, Tsutsumi O, Harada I, Fujii T, Kashima T, Kobayashi K, Toda T, Taketani Y.

Hum Reprod. 1998 Feb;13(2):320-3.

An ancient retrotransposal insertion causes Fukuyama-type congenital muscular dystrophy.

Kobayashi K, Nakahori Y, Miyake M, Matsumura K, Kondo-Iida E, Nomura Y, Segawa M, Yoshioka M, Saito K, Osawa M, Hamano K, Sakakihara Y, Nonaka I, Nakagome Y, Kanazawa I, Nakamura Y, Tokunaga K, Toda T.
Nature. 1998 Jul 23;394(6691):388-92.

Founder-haplotype analysis in Fukuyama-type congenital muscular dystrophy (FCMD).

Kobayashi K, Nakahori Y, Mizuno K, Miyake M, Kumagai T, Honma A, Nonaka I, Nakamura Y, Tokunaga K, Toda T.
Hum Genet. 1998 Sep;103(3):323–7.

Merosin-positive congenital muscular dystrophy with transient brain dysmyelination, pontocerebellar hypoplasia and mental retardation.

Voit T, Cohn RD, Sperner J, Leube B, Sorokin L, Toda T, Herrmann R.
Neuromuscul Disord. 1999 Mar;9(2):95–101.

Congenital muscular dystrophy with central and peripheral nervous system involvement in a Belgian patient.

Belpaire-Dethiou MC, Saito K, Fukuyama Y, Kondo-Iida E, Toda T, Duprez T, Verellen-Dumoulin C, Van den Bergh PY.
Neuromuscul Disord. 1999 Jun;9(4):251–6.

Broader clinical spectrum of Fukuyama-type congenital muscular dystrophy manifested by haplotype analysis.

Yoshioka M, Toda T, Kuroki S, Hamano K.
J Child Neurol. 1999 Nov;14(11):711–5.

Novel mutations and genotype-phenotype relationships in 107 families with Fukuyama-type congenital muscular dystrophy (FCMD).

Kondo-Iida E, Kobayashi K, Watanabe M, Sasaki J, Kumagai T, Koide H, Saito K, Osawa M, Nakamura Y, Toda T.
Hum Mol Genet. 1999 Nov;8(12):2303–9.

Fukuyama-type congenital muscular dystrophy: the first human disease to be caused by an ancient retrotransposal integration.

Toda T, Kobayashi K.
J Mol Med (Berl). 1999 Dec;77(12):816–23.

Differential expression of the parkin gene in the human brain and peripheral leukocytes.

Sunada Y, Saito F, Matsumura K, Shimizu T.

Neurosci Lett. 1998 Oct 2;254(3):180–2.

Detection of O-mannosyl glycans in rabbit skeletal muscle alpha-dystroglycan.

Sasaki T, Yamada H, Matsumura K, Shimizu T, Kobata A, Endo T.

Biochim Biophys Acta. 1998 Nov 27;1425(3):599–606.

Characterization of the transmembrane molecular architecture of the dystroglycan complex in schwann cells.

Saito F, Masaki T, Kamakura K, Anderson LV, Fujita S, Fukuta-Ohi H, Sunada Y, Shimizu T, Matsumura K.

J Biol Chem. 1999 Mar 19;274(12):8240–6.

A 21-kDa surface protein of *Mycobacterium leprae* binds peripheral nerve laminin-2 and mediates Schwann cell invasion.

Shimoji Y, Ng V, Matsumura K, Fischetti VA, Rambukkana A.

Proc Natl Acad Sci U S A. 1999 Aug 17;96(17):9857–62.

Sarcoglycan complex: a muscular supporter of dystroglycan-dystrophin interplay?

Matsumura K, Saito F, Yamada H, Hase A, Sunada Y, Shimizu T.

Cell Mol Biol (Noisy-le-grand). 1999 Sep;45(6):751–62.

Quick-freeze deep-etch electron microscopy of the actin-heavy meromyosin complex during the in vitro motility assay.

Katayama E.

J Mol Biol. 1998 May 1;278(2):349–67.

11 Electron microscopy coupled with quick freezing

Eisaku Katayama

287–301

Three-dimensional image analysis of myosin head in function as captured by quick-freeze deep-etch replica electron microscopy.

Katayama E, Ohmori G, Baba N

Adv Exp Med Biol. 1998;453:37–45.

Further characterization of the type 3 ryanodine receptor (RyR3) purified from rabbit diaphragm.

Murayama T, Oba T, Katayama E, Oyamada H, Oguchi K, Kobayashi M, Otsuka K, Ogawa Y.

J Biol Chem. 1999 Jun 11;274(24):17297–308.

Inner-arm dynein c of Chlamydomonas flagella is a single-headed processive motor.

Sakakibara H, Kojima H, Sakai Y, Katayama E, Oiwa K.

Nature. 1999 Aug 5;400(6744):586–90.

Electron microscopic examination of basal lamina in Fukuyama congenital muscular dystrophy.

Ishii H, Hayashi YK, Nonaka I, Arahata K.

Neuromuscul Disord. 1997 May;7(3):191–7.

Laminin alpha2 chain-null mutant mice by targeted disruption of the Lama2 gene: a new model of merosin (laminin 2)-deficient congenital muscular dystrophy.

Miyagoe Y, Hanaoka K, Nonaka I, Hayasaka M, Nabeshima Y, Arahata K, Nabeshima Y, Takeda S.

FEBS Lett. 1997 Sep 22;415(1):33–9.

Distal myopathies.

Nonaka I.

Curr Opin Neurol. 1999 Oct;12(5):493–9.

Calcineurin-dependent nuclear translocation of a murine transcription factor NFATx: molecular cloning and functional characterization.

Liu J, Koyano-Nakagawa N, Amasaki Y, Saito-Ohara F, Ikeuchi T, Imai S, Takano T, Arai N, Yokota T, Arai K.
Mol Biol Cell. 1997 Jan;8(1):157-70.

Assignment of the mouse heme oxygenase genes: heme oxygenase-1 (Hmox1) to chromosome 10 band C1 and heme oxygenase-2 (Hmox2) to chromosome 16 band B1.

Saito-Ohara F, Ikeuchi T, Matsumoto M, Kurata S.
Cytogenet Cell Genet. 1997;77(3-4):180-1.

Assignment of the gene for very-long-chain acyl-CoA dehydrogenase (Acadvl) to mouse chromosome band 11B2-B5 by in situ hybridization.

Orii KO, Saito-Ohara F, Ikeuchi T, Orii T, Kondo N, Aoyama T, Hashimoto T.

Cytogenet Cell Genet. 1997;78(1):25-6.

Assignment of the BMPR1A and BMPR1B genes to human chromosome 10q22.3 and 4q23-->q24 by in situ hybridization and radiation hybrid mapping.

Ide H, Saito-Ohara F, Ohnami S, Osada Y, Ikeuchi T, Yoshida T, Terada M.
Cytogenet Cell Genet. 1998;81(3-4):285-6.

The human caspase-activated DNase gene (hCAD): genomic structure, exonic single-nucleotide polymorphisms, and a highly polymorphic dinucleotide repeat at the hCAD locus.

Sugimoto N, Fukuda Y, Saito-Ohara F, Kamiyama R, Nakagawara A, Mukae N, Nagata S, Inazawa J.
J Hum Genet. 1999;44(6):408-11.

Cloning, expression and mapping of a novel RING-finger gene (RNF5), a

human homologue of a putative zinc-finger gene from *Caenorhabditis elegans*.

Kyushiki H, Kuga Y, Suzuki M, Takahashi E, Horie M.

Cytogenet Cell Genet. 1997;79(1-2):114-7.

Isolation and characterization of a new member of the human Ly6 gene family (LY6H).

Horie M, Okutomi K, Taniguchi Y, Ohbuchi Y, Suzuki M, Takahashi E.

Genomics. 1998 Nov 1;53(3):365-8.

Cloning of a novel gene (ING1L) homologous to ING1, a candidate tumor suppressor.

Shimada Y, Saito A, Suzuki M, Takahashi E, Horie M.

Cytogenet Cell Genet. 1998;83(3-4):232-5.

Cloning, expression, and chromosomal mapping of the human 14-3-3gamma gene (YWHAG) to 7q11.23.

Horie M, Suzuki M, Takahashi E, Tanigami A.

Genomics. 1999 Sep 1;60(2):241-3.