





(图3)





(図4)





(**図**5)

C



B

	Wild				<i>Chk2</i> ^{-/-}				<i>Chk2</i> ^{-/-} + <i>hChk2</i>			
	0	20	60	120	0	20	60	120	0	20	60	120





E

	Wild				<i>Chk2</i> ^{-/-}			
Caffeine	-	+	-	+	-	+	-	+
X-ray	-	-	+	+	-	-	+	+




(图6)





(图7)

0h



12 hr after 10Gy X-ray



20000154

以降のページは雑誌／図書等に掲載された論文となりますので
「研究成果の刊行に関する一覧表」をご参照ください。

「研究成果の刊行に関する一覧表」

- Hiroyuki Takai, Kaoru Tominaga, Noboru Motoyama, Yohji A. Minamishima, Hiroyasu Nagahama, Tadasuke Tsukiyama, Kyoji Ikeda, Keiko Nakayama, Makoto Nakanishi, and Kei-ichi Nakayama. Aberrant Cell Cycle Checkpoint Function and Early Embryonic Death in *Chk1^{-/-}* Mice. *Genes Dev.* 14: 1439-1447, 2000.
- Nakayama, K., Nagahama, H., Minamishima, Y. A., Matsumoto, M., Nakamichi, I., Kitagawa, K., Shirane, M., Tsunematsu, R., Tsukiyama, T., Ishida, N., Kitagawa, M., Nakayama, K.i., Hatakeyama, S.: Targeted disruption of *Skp2* results in accumulation of cyclin E and *p27^{Kip1}*, polyploidy and centrosome overduplication. *EMBO J.* 19: 2069-2081. 2000.
- Takahashi, K., Nakayama, K.i., Nakayama, K.: Mice lacking a CDK inhibitor, *p57^{Kip2}*, exhibit skeletal abnormalities and growth retardation. *J. Biochem.*, 127: 73-83. 2000.
- Tsukiyama, T., Ishida, N., Shirane, M., Minamishima, Y. A., Hatakeyama, S., Kitagawa, M., Nakayama, K., Nakayama, K.i.: Down-regulation of *p27^{Kip1}* expression is required for development and function of T cells. *J. Immunol.*, 166: 304-312. 2001.