

クログリアの神経細胞に対する保
護作用、第 43 回日本神経化学会、
金沢、Oct 18- 20.

研究成果一覽表

Oh-hashi K, Maruyama W, Isobe K. Peroxynitrite induces gadd34, 45 and 153 via p38 MAPK in human neuroblastoma SH-SY5Y cells. Free Rad. Biol. Med. 30: 213-221, 2001.

Hasegawa T, Xiao H, Hamajima F, Isobe K ;Interaction between DNA-damage protein GADD34 and a new member of the hsp40 family of heat shock proteins that is induced by a DNA-damaging reagent. Biochem J 2000 15; 795-800. 2000.

Maebara, K., Hasegawa T. and Isobe K. A NF-kB p65 subunit is indispensable for activating manganese superoxide Dismutase Gene Transcription mediated by tumor necrosis factor-a. J. Cellular Biochem. 77: 474-486, 2000.

Takeuchi T., Miyaishi O., Kiuchi K. and Isobe K. Cu/Zn- and Mn-Superoxide Dismutase Are Specifically Up-regulated in Neuron After Focal Brain Injury. J. Neurobiology 45:39-46, 2000.

.Isobe K. Stress and aging. Current Genomics 1;1-10;2000

Hasegawa T., Takeuchi A., Miyaishi O., Xiao H., Mao J, and Isobe K. PTRF is tissue-specific and interacts with the BFCOL1 zinc-finger transcription factor which binds to the two mouse type I collagen gene promoters. Biochem. J.347,55-59, 2000.

Xiao, H., Hasegawa, T.. and Isobe, K. p300 Collaborates with Sp1 and Sp3 in p21waf1/cip1 promoter activation Induced by Histone Deacetylase Inhibitor. J. Biol. Chem. 275; 1371-1376,2000.

Hasegawa T., Yagi A. and Isobe K.; Interaction between GADD34 and kinesin superfamily,KIF3A. Biochem Biophys Res Commun 267; 593-596,2000.

Takeda, K., Kato, M., Wu, J., Iwashita, T., Suzuki, H., Takahashi, M.and Nakashima, I.: Osmotic stress-mediated activation of RET kinases involves intracellular disulfide-bonded dimer formation. Antioxidants & Redox Signaling, in press

Kato, M., Iwashita, T., Akhand, A.A., Liu, W., Takeda, K., Takeuchi, K., Yoshihara, M., Hossain, K., Wu, J., Du, J., Oh, C., Kawamoto, Y., Suzuki,K., Takahashi, M. and Nakashima, I. Molecular mechanism of activation and superactivation of Ret tyrosine

kinases by ultraviolet light irradiation. *Antiox. Redox. Signal.*, in press.

Takeuchi, K., Kato, N., Suzuki, H., Akhand, A.A., Wu, J., Hossain, K., Miyata, T., Matsumoto, Y., Nimura, Y. and Nakashima, I.: Acrolein induces activation of the epidermal growth factor receptor of human keratinocytes for cell death. *J. Cell. Biochem.*, in press.

Wu, J., Suzuki, H., Zhou, Y.W., Liu, W., Yoshihara, M., Kato, M., Akhand, A.A., Hayakawa, A., Takeuchi, K., Hossain, K., Kurosawa, M. and Nakashima, I.: Cepharantine activates caspases and induces apoptosis in Jurkat and K562 human leukeimia cell lines. *J. Cell. Biochem.*, in press.

Kato, M., Liu, W., Akhand, A.A., Hossain, K., Takeda, K., Takahashi, M. and Nakashima, I.: Ultraviolet radiation induces both full activation of Ret kinase and malignant melanocytic tumor promotion in RFP-RET-transgenic mice. *J. Invest. Dermatol.* 115:1157-1158, 2000.

Nakashima, I., Kato, M., Akhand, A.A., Du, J., Liu, W., Dai, Y., Suzuki, H., Senga, T., Hamaguchi, M., Iwashita, T., Takahashi, M., Miyata, T., Hossain, K., Takeda, K., Wu, J., Takeuchi, K., Yoshihara, M. and Kawamoto, Y.: Chemical reaction-mediated alternative signalling pathway in cells of the immune system. *Current Trends in Immunology* 3:45-58, 2000.

Hossain, K., Akhand, A.A., Kato, M., Du, J., Takeda, K., Wu, J., Takeuchi, K., Liu, W., Suzuki, H. and Nakashima, I.: Arsenic induces apoptosis of murine T lymphocytes through membrane rafts-linked signaling for activation of c-Jun amino terminal kinase (JNK). *J. Immunol.* 165:4290-4297, 2000.

Du, J., Suzuki, H., Nagase, F., Akhand, A. A., Yokoyama, T. and Nakashima, I.: Evidence of a mercury-stimulated but interleukin2-independent signal transduction pathway for the proliferation of acytotoxic T cell line, CTLL-2. *J. Cell. Biochem.* 78:500-508, 2000.

Kato, M., Isobe, K., Dai, Y., Takahashi, M. and Nakashima, I.: Furthercharacterization of the Sho-saiko-to-mediated anti-tumor effect on melanoma developed in RET-transgenic mice. *J. Invest. Dermatol.* 114:599-601, 2000.

Du, J., Suzuki, H., Nagase, F., Akhand, A.A., Yokoyama, T., Miyata, T., Kurokawa, K. and Nakashima, I.: Methylglyoxal induces apoptosis in Jurkatleukemia T-cells by activating c-Jun N-terminal kinase. *J. Cell. Biochem.* 77:333-344, 2000.

Liu, W., Kato, M., Akhand, A.A., Hayakawa, A., Suzuki, H., Miyata, T., Kurokawa, K., Hotta, Y., Ishikawa, N. and Nakashima, I.: 4-Hydroxynonenalinduces a cellular redox status-related activation of the caspase cascade for apoptotic cell death. *J. Cell. Sci.* 113:635-641, 2000.

Kato, M., Iwashita, T., Takeda, K., Akhand, A.A., Liu, W., Yoshihara, M., Asai, N., Suzuki, H., Takahashi, M., and Nakashima, I.: Ultraviolet inudces redox reaction-mediated dimerization and superactivation of oncogenic Ret tyrosine kinases. *Mol. Biol. Cell* 11:93-101, 2000.

Senga, T., Miyazaki, K., Machida, K., Iwata, H., Matsuda, S., Nakashima, I. and Hamaguchi, M. Clustered cysteine residues in the kinase domain of v-Src: critical role for protein stability, cell transformation and sensitivity to herbimycin A. *Oncogene* 19:273-279, 2000.

Oda, E., Ohki, R., Murasawa, H., Nemoto, J., Shibue, T., Yamashita, T., Tokino, T., Taniguchi, T. and Tanaka, N.; Noxa, a BH3-only member of the Bcl-2 family, and candidate mediator of p53-induced apoptosis. *Science*, 288, 1053-1058. 2000.

Takaoka, A., Mitani, Y., Suemori, H., Sato, S., Yokochi, T., Noguchi, S., Tanaka, N., and Taniguchi, T.; Crosstalk between interferon-g and -a/b signaling components at caveolar membrane domain.. *Science*, 288, 2357-2360. 2000.

Ohki, R., Nemoto, J., Murasawa, H, Oda, E., Inazawa, J., Tanaka, N., and Taniguchi, T.; Reprimo: A new candidate mediator of the p53-mediated cell cycle arrest at the G2 phase. *J. Biol. Chem.*, 275, 22627-22630. 2000.

Sato, M., Suemori, H., Hata, N., Asagiri, M., Ogasawara, K., Nakao, K., Nakaya, T., Katsuki, M., Noguchi, S., Tanaka, N., and Taniguchi, T.; Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to viruses for IFN-a/b gene induction. *Immunity*, 13, 539-548. 2000.

Hida, S., Ogasawara, K., Sato, K., Abe, M., Takayanagi, H., Yokochi, T., Sato, T., Hirose, S., Shirai, T., Taki, S., and Taniguchi, T.; CD8+ T cell-mediated skin disease in mice lacking IRF-2, the transcriptional attenuator of interferon a/b signaling. *Immunity*, 13, 643-655. 2000.

Takayanagi, H., Ogasawara, K., Hida, H., Chiba, T., Murata, S., Sato, K., Takaoka, A., Yokochi, T., Oda, H., Tanaka, K., Nakamura, K., and Taniguchi, T. ; T-cell-mediated regulation of osteoclastogenesis by signalling cross-talk between RANKL and IFN-g.

Nature, 408, 600-605. 2000.

Okado-Matsumoto A., Matusmoto Fujii J. and Taniguchi N.: Peroxiredoxin IV is a secretable protein with heparin-binding properties under reduced conditions., J. Biochem., 127, 493-501, 2000.

Fujii T., Hamaoka R., Fujii J. and Taniguchi N.: Redox capacity of cells affects inactivation of glutathione reductase by nitrosative stress., Archives of Biochemistry and Biophysics, 378, 123-130, 2000.

Endo T., Fujii T., Sato K., Taniguchi N. and Fujii J.: A pivotal role of Zn-binding residues in the function of the copper chaperone for SOD1., Biochem. Biophys. Res. Commun., 276, 999-1004, 2000.

Nakao C., Ookawara T., Sato Y., Kizaki T., Imazeki N., Matsubara O., Haga S., Suzuki K., Taniguchi N. and Ohno H.: Extracellular Superoxide dismutase in tissues from obese (ob/ob) mice., Free. Rad. Res., 33, 229-241, 2000.

Kim Y. H., Takahashi M., Noguchi N., Suzuki E., Suzuki K., Taniguchi N. and Niki E.: Inhibition of c-Jun expression induces antioxidant enzymes under serum deprivation., Arch. Biochem. Biophys., 374, 339-346, 2000.

Hisamoto K., Ohmichi M., Kurachi H., Hayakawa J., Kanda Y., Nishio Y., Adachi K., Tasaka K., Miyoshi E., Fujiwara N., Taniguchi N., Murata Y.: Estrogen Induces the Akt-dependent Activation of Endothelial Nitric-oxide Synthase in Vascular Endothelial Cells. J Biol Chem. 2001 Feb 2;276(5):3459-3467. J Biol Chem. 276:3459-3467. 2001.

Koh YH, Park YS, Takahashi M, Suzuki K, Taniguchi N. Aldehyde reductase gene expression by lipid peroxidation end products, MDA and HNE. Free Radic Res.;33(6):739-46. 2000.

Nishida M., Miyagawa J., Yamashita S., Higashiyama S., Nakata A., Ouchi N., Tamura R., Yamamori K., Kihara S., Taniguchi N. and Matsuzawa Y.: Localization of CD9, an enhancer protein for proheparin-binding EGF-like growth factor, in human atherosclerotic plaques - Possible involvement of juxtacrine growth mechanism on smooth muscle cell proliferation., Arterioscler Thromb Vasc Biol., 20, 1236-1243,

2000.

Naka M., Nanbu T., Kobayashi K., Kamanaka Y., Komeno M., Yanase R., Fukutomi T., Fujimura S., Seo H. G., Fujiwara N., Ohuchida S., Suzuki K., Kondo K. and Taniguchi N.: A potent inhibitor of inducible nitric oxide synthase, ONO-1714, a cyclic amidine derivative., *Biochem. Biophys. Res.Commun.*, 270, 663-667, 2000.

Park Y. S., Suzuki K., Mumby S., Taniguchi N. and Gutteridge J. M. C.: Antioxidant binding of caeruloplasmin to myeloperoxidase: myeloperoxidase is inhibited, but oxidase, peroxidase and immunoreactive properties of caeruloplasmin remain intact., *Free. Rad. Res.*, 33, 261-265, 2000.

Niwa J, Ishigaki S, Doyu M, Suzuki T, Tanaka K, Sobue G: A novel centromosomal RING-finger protein, Dorfin, mediates ubiquitin ligase activity.
Biochem Biophys Res Com, in press, 2001

Adachi H, Sobue G, et al: Transgenic mice with an expanded CAG repeat controlled by the human AR promoter show polyglutamine nuclear inclusions and neuronal dysfunction without neuronal cell death. *Hum Mol Genet*, in press, 2001

Shanlou Q, Iwashita T, Furukawa T, Yamamoto M, Sobue G, Takahashi M: Differential effects of LAR on biochemical and biological activities of RET-MEN2A and RET-MEN2B mutant protein. *J Biol Chem*, in press, 2001

McCormick A, Taylor JP, Taye AA, Robitschek J, Li M, Walcott J, Merry D, Chai Y, Paulson H, Sobue G, Fischbeck KH: CREB-binding protein sequestration by expanded polyglutamine. *Hum. Mol. Genet*, 9: 2197-2202, 2000

Kobayashi Y, Kume A, Li M, Doyu M, Hata M, Ohtsuka K, Sobue G: Chaperones, Hsp70 and Hsp40, suppress aggregate formation and apoptosis in cultured neuronal cells expressing truncated androgen receptor protein with expanded polyglutamine tract. *J Biol Chem*, 275(12): 8772-8778, 2000

Watanabe H, Tanaka F, Doyu M, Riku S, Yoshida M, Hashizume Y, Sobue G: Differential somatic CAG repeat instability in variable brain cell lineage in dentatorubral pallidoluysian atrophy (DRPLA): a laser-captured microdissection (LCM)-based analysis. *Hum Genet*, 107: 452-457, 2000

Ishigaki S, Niwa J, Yoshihara T, Mitsuma N, Doyu M, Sobue G: Two novel genes, human neugrin and mouse m-neugrin, are upregulated with neuronal differentiation in neuroblastoma cells. *Biochem Biophys Res Com*, 279: 526-533, 2000

Hasegawa, Y., T. Inagaki, M. Sawada and A. Suzumura (2000). Impaired cytokine production by peripheral blood mononuclear cells and monocytes/macrophages in Parkinson's disease. *Acta Neurol Scand* 101(3): 159-64.

Hasegawa, Y., M. Sawada, N. Ozaki, T. Inagaki and A. Suzumura (2000). Increased soluble tumor necrosis factor receptor levels in the serum of elderly people. *Gerontology* 46(4): 185-8.

Kanzawa, T., M. Sawada, K. Kato, K. Yamamoto, H. Mori and R. Tanaka (2000). Differentiated regulation of allo-antigen presentation by different types of murine microglial cell lines [In Process Citation]. *J Neurosci Res* 62(3): 383-388.

Morihata, H., J. Kawasaki, H. Sakai, M. Sawada, T. Tsutada and M. Kuno (2000). Temporal fluctuations of voltage-gated proton currents in rat spinal microglia via pH-dependent and -independent mechanisms [In Process Citation]. *Neurosci Res* 38(3): 265-271.

澤田誠：細胞を使って脳の疾患を治療する試み 治療、82, 128-130, 2000.

澤田誠：細胞周期とアポトーシス Clinical Neuroscience 18(4) 407-409, 2000.