

平成12年度事業報告

強皮症調査研究 第1回 班会議

平成13年1月19日 於：東京医科歯科大学

研究成果の刊行一覧

Ihn H, Sato S, Fujimoto M, Igarashi A, Yazawa N, Kubo M, Kikuchi K, Takehara K, Tamaki K. Characterization of autoantibodies to endothelial cells in systemic sclerosis' association with pulmonary fibrosis. *Clin Exp Immunol* 119; 203-209, 2000.

Ihn H, Yazawa N, Kubo M, Yamane K, Sato S, Fujimoto M, Kikuchi K, Soma Y, Tamaki K. Circulating levels of soluble CD30 are increased in patients with localized scleroderma and correlated with serological and clinical features of the disease. *J Rheumatol* 27; 698-702, 2000.

Ihn H, Tamaki K. Competition analysis of the human $\alpha 2(I)$ collagen promoter using synthetic oligonucleotides. *J Invest Dermatol* 114; 1011-1016, 2000.

Ihn H, Tamaki K. Oncostatin M stimulates the growth of dermal fibroblasts via a mitogen-activated protein kinase-dependent pathway. *J Immunol* 165; 2149-2155, 2000.

Ihn H, Tamaki K. Increased phosphorylation of transcription factor Sp1 in scleroderma fibroblasts'; association with increased expression of collagen gene. *Arthritis Rheum* 43; 2240-2247, 2000.

Yamane K, Ihn H, Kubo M, Yazawa N, Kikuchi K, Soma Y, Tamaki K. Increased serum levels of soluble VCAM-1 and E-selectin in patients with localized scleroderma. *J Am Acad Dermatol* 42; 64-66, 2000.

Yazawa N, Kikuchi K, Ihn H, Fujimoto M, Kubo M, Tamaki T, Tamaki K. Serum levels of tissue inhibitor of metalloproteinase-2 (TIMP-2) in patients with systemic sclerosis. *J Am Acad Dermatol* 42; 70-75, 2000.

Sato S, Hasegawa M, Ihn H, Kikuchi K, Takehara K. Clinical significance of anti-nuclear matrix antibody in sera from patients with anti-U1RNP antibody. *Arch Dermatol Res* 292; 55-59, 2000.

Yamane K, Ihn H, Kubo M, Yazawa N, Kikuchi K, Soma Y, Tamaki K. Serum levels of KL-6 as a useful marker in sera from patients with systemic sclerosis. *J Rheumatol* 27; 930-934, 2000.

Kubo M, Ihn H, Yamane K, Kikuchi K, Yazawa N, Kikuchi K, Yazawa N, Soma Y, Tamaki K. Serum KL-6 in adult patients with polymyositis and dermatomyositis. *Rheumatol* 39; 632-636, 2000.

Nagaoka T, Sato S, Hasegawa M, Ihn H, Takehara K. Serum levels of soluble interleukin-6 receptor and soluble gp130 in patients with localized scleroderma. *J Rheumatol* 27; 1917-1921, 2000.

Kubo M, Ihn H, Yamane K, Yazawa N, Kikuchi K, Soma Y, Tamaki K. Increased serum levels of soluble E-selectin in patients with polymyositis and dermatomyositis. *Br J Dermatol* 143; 392-398, 2000.

Yamane K, Ihn H, Asano Y, Yazawa N, Kubo M, Kikuchi K, Soma Y, Tamaki K. Clinical and laboratory features of scleroderma patients with pulmonary hypertension. *Rheumatol* 39; 1269-1271, 2000.

Yamamoto T, Eckes B, Mauch C, Hartmann K, Krieg T: Monocyte chemoattractant protein-1 enhances gene expression and synthesis of matrix metalloproteinase-1 in human fibroblasts via an autocrine IL-1_α loop. *J Immunol* 2000; 164: 6174-6179.

Yamamoto T, Hartmann K, Eckes B, Krieg T: Mast cells enhance gel contraction of three-

- dimensional collagen lattices by fibroblasts by cell-cell interaction: role of stem cell factor/c-kit. *Immunology* 2000; 99: 435-439.
- Yamamoto T, Takagawa S, Nishioka K: Effect of interferon- γ on experimental scleroderma induced by bleomycin. *Arch Derm Res* 2000; 292: 362-365.
- Yamamoto T, Kuroda M, Nishioka K: Animal model of sclerotic skin. III: Histopathological comparison of bleomycin-induced scleroderma in various mice strains. *Arch Dermatol Res* 2000; 292: 535-541.
- Yamamoto T, Eckes B, Krieg T: Bleomycin increases gene expression of $\alpha 1(I)$ collagen, fibronectin and decorin in human fibroblasts. *Arch Dermatol Res* 2000; 292: 556-561.
- Shinohara M, Hamasaki Y, Katayama I: Refractory pretibial myxoedema with response to intralesional insulin-like growth factor 1 antagonist (octreotide): downregulation of hyaluronic acid production by the lesional fibroblasts. *Br J Dermatol*, 2000, 143(5):1083-6
- Yamamoto T, Katayama I, Nishioka K: A rare association of systemic sclerosis with psoriasis vulgaris. *J Dermatol*, 2000, 27(5):346-9
- Miyazaki Y, Yokozeki H, Awad S, Igawa K, Minatohara K, Satoh T, Katayama I, Nishioka K: Glucocorticoids augment the chemically induced production and gene expression of interleukin-1 α through NF- κ B and AP-1 activation in murine epidermal cells. *J Invest Dermatol*, 2000, 115(4): 746-753
- Naito S, Shimizu K, Nakashima M, Nakayama T, Ito T, Ito M, Yamashita S, Sekine I: Overexpression of Ets-1 transcription factor in angiosarcoma of the skin. *Pathol Res Prac*, 2000, 96: 103-109
- Katayama I, Umeda T, Nishioka K: Adult Still's disease like-illness In a patient with silicone breast implants. *Review Series Rheumatology*. 1999, 1: 12-13
- Maeda A, Yamanouchi H, Lee JB, Katayama I: Letter: Oral prednisolone improved acetylcholine-induced sweating in Sjogren's syndrome-related anhidrosis. *Clin Rheumatol*, 2000, 19: 396-397
- Hata R., Akai J., Kimura A., Ishikawa O., Kuwana M, and Shinkai H.: Association of functional microsatellites in the human type I collagen $\alpha 2$ chain (COL1A2) gene with systemic sclerosis. *Biochem. Biophys. Res. Commun.* (2000) 272: 36-40.
- Akai J., Kimura A. and Hata R. : Analysis of the higher-order structure of the two dinucleotide repeat regions that enhance the transcriptional activity of the human type I collagen $\alpha 2$ (COL1A2) gene promoter. *Connective Tissue* (2000) 32:259-265.
- Hata R., Izukuri K., Furue M., Kato Y., and Koshika S. Complex gene diseases of the connective tissue. *Connective Tissue* in press.
- Takemoto, M., Yokote, K., Nishimura, M., Shigematsu, T., Hasegawa, T., Kon, S., Uede, T., Matsumoto, T., Saito, Y., Mori, S. : Enhanced expression of osteopontin in human diabetic artery and analysis of its functional role in accelerated atherogenesis. *Arterioscler. Thromb. Vasc.*

Biol. 20: 624-628, 2000.

Kobayashi, J., Murano, S., Yokote, K., Mori, S., Matsunaga, A., Sasaki, J., Takahashi, K., Bujo, H., Saito, Y.: Marked decrease in plasma apolipoprotein A-I and high density lipoprotein-cholesterol in a case with Werner syndrome. *Clinica. Chimica. Acta.* 293: 63-73, 2000.

Matsumoto, T., Yokote, K., Take, A., Takemoto, M., Asaumi, S., Hashimoto, Y., Matsuda, M., Saito, Y., Mori, S.: Differential interaction of CrkII adaptor protein with platelet-derived growth factor alpha- and beta-receptors is determined by its internal tyrosine phosphorylation. *Biochem. Biophys. Res. Commun.* 270: 28-33, 2000.

Tamura, K., Kanzaki, T., Tashiro, J., Yokote, K., Mori, S., Ueda, S., Saito, Y., Morisaki, N.: Increased atherogenesis in Otsuka Long-Evans Tokushima fatty rats before the onset of diabetes mellitus: association with overexpression of PDGF beta-receptors in aortic smooth muscle cells. *Atherosclerosis* 149: 351-358, 2000.

Takemoto, M., Yokote, K., Yamazaki, M., Ridall, A. L., Butler, W. T., Matsumoto, T., Tamura, K., Saito, Y., Mori, S.: Enhanced expression of osteopontin by high glucose: involvement of osteopontin in diabetic macroangiopathy. *Ann. N. Y. Acad. Sci.* 902: 357-363, 2000.

Wen Zhang, Jiongwen Ou, Yutaka Inagaki, Patricia Greenwel and Francesco Ramirez: Synergistic cooperation between Sp1 and Smad3/Smad4 mediates TGF- β 1 stimulation of α 2(I) collagen (COL1A2) transcription. 2000. *J Biol Chem* 275: 39237-39245

Yutaka Inagaki, Mizuko Mamura, Yutaka Kanamaru, Patricia Greenwel, Tomoyuki Nemoto, Kazuhiko Takehara, Peter ten Dijke and Atsuhito Nakao: Constitutive phosphorylation and nuclear localization of Smad3 are correlated with increased collagen gene transcription in activated hepatic stellate cells. *J Cell Physiol* (in press)

Kobayashi T, Munakata S, Sugiura H, Koizumi M, Sumida M, Murata K, Shinkai H. Angiotropic lymphoma: proliferation of B cells in the capillaries of cutaneous angiomas. *Br J Dermatol*, 143: 162-4, 2000

Kakinuma Y, Endo H, Tsukahara T, Futoeda T, Saito Y, Shinkai H. Collagenoma with pseudohypoparathyroidism. *Br J Dermatol*, 143: 1122-4, 2000

Kurasawa K, Hirose K, Sano H, Endo H, Shinkai H, Nawata Y, Takabayashi K, Iwamoto I. Increased interleukin-17 production in patients with systemic sclerosis. *Arthritis Rheum* 43:2455-63, 2000.

Hatamochi A, Nagayama H, Kuroda K, Shinkai H, Ishikiriyama S, Kobayashi M, Kobayashi K. Costello Syndrome with Decreased Gene Expression of Elastin in Cultured Dermal Fibroblasts. *Dermatology*. 201(4):366-369. 2000

Yokoyama Y, Akimoto S, Ishikawa O. Atrophoderma of Pasisi and Pierini: case reports and disaccharide analysis of skin glycosaminoglycans. *Clin Exp Dermatol* 25: 436-440, 2000

Amano H, Kurosawa M, Ishikawa O, Miyachi Y. Cultured human mast cells derived from umbilical cord blood cells in the presence of stem cell factor and interleukin-6 cannot be a model of

human skin mast cells: fluorescence microscopic analysis of intercellular calcium ion mobilization. *J Dermatol Sci* 24: 146-152, 2000

Yamanaka M, Ishikawa O. Hypoxic conditions decrease the mRNA expression of pro $\alpha 1$ (I) and $\alpha 1$ (III) collagens and increase matrix metalloproteinase-1 of dermal fibroblasts in three-dimensional culture. *J Dermatol Sci* 24:99-104, 2000

Abe M, Kurosawa M, Ishikawa O, Miyachi Y. Effect of mast cell-derived mediators and mast cell-related neutral proteases on human dermal fibroblast proliferation and type I collagen production. *J Allergy Clin Immun* 106: 78-84, 2000

Matsushima Y, Ishikawa O, Kurosawa M, Miyachi Y. Stem cell factor and interleukin-6 do not promote complete maturation of human cultured mast cells from umbilical cord blood cells: an ultrastructural study. *J Dermatol Sci* 24: 4-13, 2000

Amano H, Kurosawa M, Ishikawa O, Miyachi Y. Mast cells in cutaneous lesions of sarcoidosis: their subtypes and the relationship to systemic manifestation. *J Dermatol Sci* 24: 60-66, 2000

Sakai S, Yasuda R, Sayo T, Ishikawa O, Inoue O: Hyaluronic acid exists in the normal stratum corneum. *J Invest Dermatol* 114: 1184-1187, 2000

山本俊幸：強皮症モデルの応用。皮膚病診療、2000、22：1014-1017.

西村香織、松永義孝、竹中基、片山一朗： 染色体異常（トリソミー1）を認める骨髄異型性症候群に伴った壊疽性膿皮症の1例。日皮会誌、1999、109(10)：1479-1484

片山一朗、濱崎洋一郎、有馬優子、天満美輪、前田亜紀、野村昌代、武石恵美子、山本雅一： シェーグレン症候群における更年期症状様病態の臨床的検討。西日皮膚、2000、62（5）：648-654

濱崎洋一郎、片山一朗： 剣創状強皮症の酢酸オクトレオチド局所投与による治療。皮膚病診療、2000、22(11)：1059-1062

片山一朗： サルコイドーシスと膠原病。皮膚病診療、2000、22(8)：716-722

片山一朗： 抗リン脂質抗体症候群。皮膚臨床、2000、42(10)：1542-1544

片山一朗、西岡清： 全身性強皮症。year note 2001, Selected Articles, 2000, 823-835

片山一朗： シェーグレン症候群・腺外症状および合併症の検索 VI. 皮膚病変。シェーグレン症候群診断の手技・手法マニュアル、江口勝美編、日本シェーグレン研究会、2000、pp.147-153

稲垣 豊、根本朋幸：コラーゲン遺伝子の発現動態からみた肝壊死、再生、線維化の病態連繋と細胞種特異的転写調節機構。2000。肝臓 41：69-72.

稲垣 豊：コラーゲン産生の遺伝子制御。2000。最新医学 55：1771-1775.

石川 治、秋元幸子。皮膚科医から見た全身性強皮症。リウマチ 40：39-47, 2000

遠藤雪恵、石川 治。強皮症とアフェレーシス。アフェレーシス学会誌 19：204-208, 2000

根岸 泉, 石川 治. 強皮症・シェーグレン症候群とマイクロキメリズム. 内科 86: 259-261, 2000

根岸 泉, 遠藤雪恵, 石川 治. 強皮症・Sj 喩 ren 症候群の病因としてのキメリズム. 「臨床皮膚科_最近のトピックス_Clinical Dermatology_」. 臨皮 54 (5増): 180-181, 2000

関 姿恵, 秋元幸子, 石川 治. ニトログリセリンテープによる強皮症患者末梢循環不全の治療. 「臨床皮膚科_最近のトピックス_Clinical Dermatology_」. 臨皮 54 (5増): 133-137, 2000

厚生科学研究費補助金
(特定疾患対策事業)
強皮症調査研究
平成12年度研究報告書

発行 平成13年3月31日
発行所 厚生科学研究
強皮症調査研究班
〒260 千葉市中央区亥鼻1-8-1
千葉大学医学部皮膚科
主任研究者 新海 滋
事務局 ☎043-222-7171 (5332)