

2. 多発性硬化症

表1-5-1-2 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別


性・年齢階級		1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
			84889297	848892×	8488×	84×	8488×97	84×9297	84×92×	84××97		
男	0-4	6	1 (16.7)	1 (16.7)	1 (16.7)	3 (50.0)
	5-9	13	1 (7.7)	1 (7.7)	3 (23.1)	6 (46.2)	2 (15.4)
	10-14	25	5 (20.0)	1 (4.0)	5 (20.0)	13 (52.0)	1 (4.0)
	15-19	22	4 (18.2)	2 (9.1)	5 (22.7)	9 (40.9)	2 (9.1)
	20-24	26	11 (42.3)	3 (11.5)	5 (19.2)	6 (23.1)	1 (3.8)
	25-29	35	11 (31.4)	5 (14.3)	4 (11.4)	12 (34.3)	3 (8.6)
	30-34	45	17 (37.8)	6 (13.3)	4 (8.9)	13 (28.9)	5 (11.1)
	35-39	64	28 (43.8)	8 (12.5)	9 (14.1)	16 (25.0)	3 (4.7)
	40-44	59	26 (44.1)	7 (11.9)	9 (15.3)	14 (23.7)	3 (5.1)
	45-49	39	20 (51.3)	7 (17.9)	1 (2.6)	9 (23.1)	2 (5.1)
	50-54	44	26 (59.1)	3 (6.8)	9 (20.5)	6 (13.6)
	55-59	56	24 (42.9)	7 (12.5)	9 (16.1)	12 (21.4)	4 (7.1)
	60-64	21	4 (19.0)	4 (19.0)	5 (23.8)	8 (38.1)
	65-69	15	2 (13.3)	1 (6.7)	6 (40.0)	5 (33.3)	1 (6.7)
	70-74
	75-79
80-84	
85-	
全年齢		470	180 (38.3)	56 (11.9)	75 (16.0)	132 (28.1)	27 (5.7)	
女	0-4	4	2 (50.0)	.	.	2 (50.0)	
	5-9	13	3 (23.1)	2 (15.4)	3 (23.1)	5 (38.5)	
	10-14	33	10 (30.3)	1 (3.0)	6 (18.2)	15 (45.5)	1 (3.0)	
	15-19	64	18 (28.1)	5 (7.8)	8 (12.5)	27 (42.2)	6 (9.4)	
	20-24	63	32 (50.8)	6 (9.5)	6 (9.5)	14 (22.2)	5 (7.9)	
	25-29	86	43 (50.0)	7 (8.1)	9 (10.5)	17 (19.8)	10 (11.6)	
	30-34	134	74 (55.2)	11 (8.2)	10 (7.5)	28 (20.9)	11 (8.2)	
	35-39	147	77 (52.4)	22 (15.0)	15 (10.2)	23 (15.6)	10 (6.8)	
	40-44	164	78 (47.6)	21 (12.8)	18 (11.0)	38 (23.2)	9 (5.5)	
	45-49	125	57 (45.6)	13 (10.4)	13 (10.4)	40 (32.0)	2 (1.6)	
	50-54	166	59 (35.5)	28 (16.9)	25 (15.1)	48 (28.9)	6 (3.6)	
	55-59	136	50 (36.8)	26 (19.1)	25 (18.4)	31 (22.8)	4 (2.9)	
	60-64	97	24 (24.7)	14 (14.4)	24 (24.7)	31 (32.0)	4 (4.1)	
	65-69	31	5 (16.1)	4 (12.9)	7 (22.6)	14 (45.2)	1 (3.2)	
	70-74	6	.	.	1 (16.7)	5 (83.3)	
	75-79	
80-84		
85-		
全年齢		1,269	532 (41.9)	160 (12.6)	170 (13.4)	338 (26.6)	69 (5.4)	

2. 多発性硬化症

	1988年度 受給者数 (100%)		9年継続		4年継続		継続なし		中止後再開		1992年度 受給者数 (100%)		5年継続		継続なし	
			889297		8892×		88×		88×	97		9297		92×		
男 0-4											5	2	(40.0)	3	(60.0)	
5-9	18	11	(61.1)	4	(22.2)	3	(16.7)				27	12	(44.4)	15	(55.6)	
10-14	31	8	(25.8)	8	(25.8)	13	(41.9)	2	(6.5)		36	19	(52.8)	17	(47.2)	
15-19	30	12	(40.0)	4	(13.3)	14	(46.7)				45	22	(48.9)	23	(51.1)	
20-24	40	18	(45.0)	4	(10.0)	16	(40.0)	2	(5.0)		80	50	(62.5)	30	(37.5)	
25-29	74	31	(41.9)	18	(24.3)	22	(29.7)	3	(4.1)		91	58	(63.7)	33	(36.3)	
30-34	84	46	(54.8)	22	(26.2)	16	(19.0)				140	88	(62.9)	52	(37.1)	
35-39	128	75	(58.6)	22	(17.2)	30	(23.4)	1	(0.8)		147	92	(62.6)	55	(37.4)	
40-44	118	62	(52.5)	22	(18.6)	33	(28.0)	1	(0.8)		215	166	(77.2)	49	(22.8)	
45-49	101	58	(57.4)	12	(11.9)	29	(28.7)	2	(2.0)		150	105	(70.0)	45	(30.0)	
50-54	79	47	(59.5)	11	(13.9)	21	(26.6)				123	100	(81.3)	23	(18.7)	
55-59	79	46	(58.2)	14	(17.7)	19	(24.1)				102	80	(78.4)	22	(21.6)	
60-64	67	30	(44.8)	14	(20.9)	20	(29.9)	3	(4.5)		96	59	(61.5)	37	(38.5)	
65-69	21	3	(14.3)	4	(19.0)	13	(61.9)	1	(4.8)		51	33	(64.7)	18	(35.3)	
70-74	11	2	(18.2)	2	(18.2)	7	(63.6)				10	4	(40.0)	6	(60.0)	
75-79	2			1	(50.0)	1	(50.0)				3	1	(33.3)	2	(66.7)	
80-84											3			3	(100.0)	
85-																
全年齢	883	449	(50.8)	162	(18.3)	257	(29.1)	15	(1.7)		1,324	891	(67.3)	433	(32.7)	
女 0-4	2	2	(100.0)								7	4	(57.1)	3	(42.9)	
5-9	20	5	(25.0)	6	(30.0)	9	(45.0)				32	19	(59.4)	13	(40.6)	
10-14	45	15	(33.3)	10	(22.2)	17	(37.8)	3	(6.7)		56	33	(58.9)	23	(41.1)	
15-19	69	35	(50.7)	7	(10.1)	22	(31.9)	5	(7.2)		84	55	(65.5)	29	(34.5)	
20-24	124	70	(56.5)	21	(16.9)	29	(23.4)	4	(3.2)		153	118	(77.1)	35	(22.9)	
25-29	154	88	(57.1)	27	(17.5)	35	(22.7)	4	(2.6)		244	181	(74.2)	63	(25.8)	
30-34	172	102	(59.3)	27	(15.7)	37	(21.5)	6	(3.5)		268	194	(72.4)	74	(27.6)	
35-39	270	172	(63.7)	38	(14.1)	52	(19.3)	8	(3.0)		266	206	(77.4)	60	(22.6)	
40-44	241	151	(62.7)	39	(16.2)	47	(19.5)	4	(1.7)		411	324	(78.8)	87	(21.2)	
45-49	252	157	(62.3)	39	(15.5)	53	(21.0)	3	(1.2)		320	245	(76.6)	75	(23.4)	
50-54	220	145	(65.9)	27	(12.3)	38	(17.3)	10	(4.5)		319	254	(79.6)	65	(20.4)	
55-59	201	108	(53.7)	39	(19.4)	49	(24.4)	5	(2.5)		276	224	(81.2)	52	(18.8)	
60-64	160	71	(44.4)	33	(20.6)	54	(33.8)	2	(1.3)		223	146	(65.5)	77	(34.5)	
65-69	74	22	(29.7)	20	(27.0)	29	(39.2)	3	(4.1)		136	84	(61.8)	52	(38.2)	
70-74	18	5	(27.8)	6	(33.3)	7	(38.9)				49	26	(53.1)	23	(46.9)	
75-79	5	1	(20.0)			4	(80.0)				10	7	(70.0)	3	(30.0)	
80-84																
85-											1			1	(100.0)	
全年齢	2,027	1,149	(56.7)	339	(16.7)	482	(23.8)	57	(2.8)		2,855	2,120	(74.3)	735	(25.7)	


2. 多発性硬化症

図1-5-1-2 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




2. 多発性硬化症

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



▨ 5年継続 □ 継続なし

3 . 重症筋無力症

表 1 - 5 - 1 - 3 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別

性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
		84889297		848892×		8488×		84×		8488×97 84×9297 84×92×	84×97
男 0-4	60	21 (35.0)		8 (13.3)		6 (10.0)		22 (36.7)		3 (5.0)	
5-9	116	32 (27.6)		17 (14.7)		19 (16.4)		39 (33.6)		9 (7.8)	
10-14	126	37 (29.4)		19 (15.1)		28 (22.2)		37 (29.4)		5 (4.0)	
15-19	84	25 (29.8)		5 (6.0)		14 (16.7)		34 (40.5)		6 (7.1)	
20-24	63	28 (44.4)		8 (12.7)		7 (11.1)		18 (28.6)		2 (3.2)	
25-29	59	25 (42.4)		11 (18.6)		5 (8.5)		14 (23.7)		4 (6.8)	
30-34	73	38 (52.1)		7 (9.6)		6 (8.2)		18 (24.7)		4 (5.5)	
35-39	102	56 (54.9)		11 (10.8)		12 (11.8)		18 (17.6)		5 (4.9)	
40-44	86	45 (52.3)		9 (10.5)		7 (8.1)		20 (23.3)		5 (5.8)	
45-49	86	53 (61.6)		8 (9.3)		9 (10.5)		11 (12.8)		5 (5.8)	
50-54	102	53 (52.0)		9 (8.8)		15 (14.7)		18 (17.6)		7 (6.9)	
55-59	97	36 (37.1)		18 (18.6)		15 (15.5)		23 (23.7)		5 (5.2)	
60-64	98	37 (37.8)		21 (21.4)		16 (16.3)		19 (19.4)		5 (5.1)	
65-69	59	14 (23.7)		4 (6.8)		13 (22.0)		25 (42.4)		3 (5.1)	
70-74	14	1 (7.1)		1 (7.1)		4 (28.6)		8 (57.1)		.	.
75-79	4	.		.		1 (25.0)		3 (75.0)		.	.
80-84
85-	1	.		1 (100.0)	
全年齢	1,230	501 (40.7)		157 (12.8)		177 (14.4)		327 (26.6)		68 (5.5)	
女 0-4	106	39 (36.8)		13 (12.3)		17 (16.0)		34 (32.1)		3 (2.8)	
5-9	170	61 (35.9)		33 (19.4)		22 (12.9)		41 (24.1)		13 (7.6)	
10-14	225	83 (36.9)		36 (16.0)		33 (14.7)		58 (25.8)		15 (6.7)	
15-19	157	60 (38.2)		16 (10.2)		14 (8.9)		54 (34.4)		13 (8.3)	
20-24	120	56 (46.7)		7 (5.8)		16 (13.3)		26 (21.7)		15 (12.5)	
25-29	206	101 (49.0)		17 (8.3)		20 (9.7)		55 (26.7)		13 (6.3)	
30-34	299	185 (61.9)		25 (8.4)		29 (9.7)		46 (15.4)		14 (4.7)	
35-39	393	240 (61.1)		37 (9.4)		37 (9.4)		64 (16.3)		15 (3.8)	
40-44	342	222 (64.9)		41 (12.0)		20 (5.8)		46 (13.5)		13 (3.8)	
45-49	335	209 (62.4)		21 (6.3)		32 (9.6)		59 (17.6)		14 (4.2)	
50-54	287	166 (57.8)		32 (11.1)		22 (7.7)		55 (19.2)		12 (4.2)	
55-59	277	153 (55.2)		31 (11.2)		29 (10.5)		53 (19.1)		11 (4.0)	
60-64	209	95 (45.5)		28 (13.4)		27 (12.9)		51 (24.4)		8 (3.8)	
65-69	127	31 (24.4)		22 (17.3)		20 (15.7)		44 (34.6)		10 (7.9)	
70-74	33	8 (24.2)		5 (15.2)		5 (15.2)		13 (39.4)		2 (6.1)	
75-79	1	1 (100.0)	
80-84
85-	1	.		.		.		1 (100.0)		.	.
全年齢	3,288	1,710 (52.0)		364 (11.1)		343 (10.4)		700 (21.3)		171 (5.2)	


3. 重症筋無力症

	1988年度	9年継続			4年継続		継続なし		中止後再開		1992年度	5年継続			継続なし	
	受給者数 (100%)	889297			8892×		88×		88×97		受給者数 (100%)	9297			92×	
男 0-4	60	28	(46.7)	10	(16.7)	22	(36.7)	.	.		44	27	(61.4)	17	(38.6)	
5-9	87	47	(54.0)	16	(18.4)	24	(27.6)	.	.		87	55	(63.2)	32	(36.8)	
10-14	127	55	(43.3)	35	(27.6)	35	(27.6)	2	(1.6)		103	71	(68.9)	32	(31.1)	
15-19	125	50	(40.0)	24	(19.2)	49	(39.2)	2	(1.6)		132	79	(59.8)	53	(40.2)	
20-24	88	38	(43.2)	19	(21.6)	29	(33.0)	2	(2.3)		118	76	(64.4)	42	(35.6)	
25-29	93	55	(59.1)	10	(10.8)	26	(28.0)	2	(2.2)		112	74	(66.1)	38	(33.9)	
30-34	112	53	(47.3)	25	(22.3)	29	(25.9)	5	(4.5)		128	96	(75.0)	32	(25.0)	
35-39	161	100	(62.1)	26	(16.1)	32	(19.9)	3	(1.9)		173	124	(71.7)	49	(28.3)	
40-44	175	117	(66.9)	25	(14.3)	27	(15.4)	6	(3.4)		242	183	(75.6)	59	(24.4)	
45-49	149	100	(67.1)	24	(16.1)	25	(16.8)	.	.		246	193	(78.5)	53	(21.5)	
50-54	168	109	(64.9)	27	(16.1)	30	(17.9)	2	(1.2)		217	170	(78.3)	47	(21.7)	
55-59	189	125	(66.1)	26	(13.8)	37	(19.6)	1	(0.5)		265	205	(77.4)	60	(22.6)	
60-64	173	86	(49.7)	40	(23.1)	44	(25.4)	3	(1.7)		266	198	(74.4)	68	(25.6)	
65-69	123	55	(44.7)	24	(19.5)	42	(34.1)	2	(1.6)		218	143	(65.6)	75	(34.4)	
70-74	57	21	(36.8)	11	(19.3)	25	(43.9)	.	.		109	74	(67.9)	35	(32.1)	
75-79	16	4	(25.0)	4	(25.0)	8	(50.0)	.	.		43	25	(58.1)	18	(41.9)	
80-84	8	2	(25.0)	.	.	6	(75.0)	.	.		11	5	(45.5)	6	(54.5)	
85-	1	.	.	1	(100.0)		3	2	(66.7)	1	(33.3)	
全年齢	1,912	1,045	(54.7)	347	(18.1)	490	(25.6)	30	(1.6)		2,517	1,800	(71.5)	717	(28.5)	
女 0-4	75	44	(58.7)	12	(16.0)	18	(24.0)	1	(1.3)		69	45	(65.2)	24	(34.8)	
5-9	168	93	(55.4)	34	(20.2)	37	(22.0)	4	(2.4)		157	105	(66.9)	52	(33.1)	
10-14	204	98	(48.0)	58	(28.4)	45	(22.1)	3	(1.5)		186	131	(70.4)	55	(29.6)	
15-19	231	120	(51.9)	52	(22.5)	55	(23.8)	4	(1.7)		263	164	(62.4)	99	(37.6)	
20-24	197	116	(58.9)	28	(14.2)	47	(23.9)	6	(3.0)		278	183	(65.8)	95	(34.2)	
25-29	253	165	(65.2)	33	(13.0)	49	(19.4)	6	(2.4)		291	222	(76.3)	69	(23.7)	
30-34	357	228	(63.9)	56	(15.7)	65	(18.2)	8	(2.2)		332	257	(77.4)	75	(22.6)	
35-39	486	336	(69.1)	59	(12.1)	82	(16.9)	9	(1.9)		459	364	(79.3)	95	(20.7)	
40-44	545	405	(74.3)	58	(10.6)	74	(13.6)	8	(1.5)		652	551	(84.5)	101	(15.5)	
45-49	466	330	(70.8)	63	(13.5)	69	(14.8)	4	(0.9)		656	548	(83.5)	108	(16.5)	
50-54	440	326	(74.1)	44	(10.0)	61	(13.9)	9	(2.0)		586	504	(86.0)	82	(14.0)	
55-59	380	256	(67.4)	60	(15.8)	61	(16.1)	3	(0.8)		533	437	(82.0)	96	(18.0)	
60-64	348	215	(61.8)	65	(18.7)	60	(17.2)	8	(2.3)		519	413	(79.6)	106	(20.4)	
65-69	257	137	(53.3)	54	(21.0)	59	(23.0)	7	(2.7)		422	290	(68.7)	132	(31.3)	
70-74	124	58	(46.8)	28	(22.6)	35	(28.2)	3	(2.4)		251	173	(68.9)	78	(31.1)	
75-79	38	9	(23.7)	13	(34.2)	15	(39.5)	1	(2.6)		105	68	(64.8)	37	(35.2)	
80-84	3	.	.	1	(33.3)	2	(66.7)	.	.		31	14	(45.2)	17	(54.8)	
85-	2	1	(50.0)	.	.	1	(50.0)	.	.		6	1	(16.7)	5	(83.3)	
全年齢	4,574	2,937	(64.2)	718	(15.7)	835	(18.3)	84	(1.8)		5,796	4,470	(77.1)	1,326	(22.9)	

3. 重症筋無力症


図1-5-1-3 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別

1984年度受給者 継続状況




3. 重症筋無力症

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



■ 5年継続 □ 継続なし

4. 全身性エリテマトーデス

表1-5-1-4 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
		84889297		848892×		8488×		84×		8488×97	
男 0-4	1	.	.	1 (100.0)
5-9	12	3 (25.0)		3 (25.0)	1 (8.3)		5 (41.7)				
10-14	56	24 (42.9)		10 (17.9)	6 (10.7)		13 (23.2)		3 (5.4)		
15-19	127	58 (45.7)		11 (8.7)	19 (15.0)		29 (22.8)		10 (7.9)		
20-24	132	67 (50.8)		10 (7.6)	14 (10.6)		28 (21.2)		13 (9.8)		
25-29	120	73 (60.8)		8 (6.7)	16 (13.3)		18 (15.0)		5 (4.2)		
30-34	157	78 (49.7)		27 (17.2)	14 (8.9)		28 (17.8)		10 (6.4)		
35-39	151	84 (55.6)		14 (9.3)	14 (9.3)		35 (23.2)		4 (2.6)		
40-44	102	50 (49.0)		13 (12.7)	9 (8.8)		22 (21.6)		8 (7.8)		
45-49	76	39 (51.3)		11 (14.5)	9 (11.8)		14 (18.4)		3 (3.9)		
50-54	69	25 (36.2)		9 (13.0)	11 (15.9)		20 (29.0)		4 (5.8)		
55-59	90	27 (30.0)		18 (20.0)	18 (20.0)		25 (27.8)		2 (2.2)		
60-64	59	11 (18.6)		13 (22.0)	10 (16.9)		24 (40.7)		1 (1.7)		
65-69	42	7 (16.7)		6 (14.3)	11 (26.2)		17 (40.5)		1 (2.4)		
70-74	8	.	.	2 (25.0)	1 (12.5)		5 (62.5)		.	.	
75-79	1	.	.	1 (100.0)	
80-84	
85-	
全年齢	1,203	546 (45.4)		157 (13.1)	153 (12.7)		283 (23.5)		64 (5.3)		
女 0-4	8	1 (12.5)		1 (12.5)	1 (12.5)		5 (62.5)		.	.	
5-9	64	32 (50.0)		11 (17.2)	4 (6.3)		11 (17.2)		6 (9.4)		
10-14	318	195 (61.3)		23 (7.2)	39 (12.3)		43 (13.5)		18 (5.7)		
15-19	868	516 (59.4)		69 (7.9)	69 (7.9)		142 (16.4)		72 (8.3)		
20-24	1,231	728 (59.1)		108 (8.8)	107 (8.7)		186 (15.1)		102 (8.3)		
25-29	1,772	1,095 (61.8)		152 (8.6)	140 (7.9)		273 (15.4)		112 (6.3)		
30-34	2,552	1,618 (63.4)		213 (8.3)	198 (7.8)		351 (13.8)		172 (6.7)		
35-39	2,908	1,952 (67.1)		228 (7.8)	197 (6.8)		364 (12.5)		167 (5.7)		
40-44	2,442	1,589 (65.1)		262 (10.7)	167 (6.8)		313 (12.8)		111 (4.5)		
45-49	1,821	1,122 (61.6)		219 (12.0)	149 (8.2)		253 (13.9)		78 (4.3)		
50-54	1,438	789 (54.9)		186 (12.9)	144 (10.0)		256 (17.8)		63 (4.4)		
55-59	997	457 (45.8)		142 (14.2)	149 (14.9)		196 (19.7)		53 (5.3)		
60-64	600	183 (30.5)		108 (18.0)	142 (23.7)		143 (23.8)		24 (4.0)		
65-69	242	43 (17.8)		29 (12.0)	40 (16.5)		127 (52.5)		3 (1.2)		
70-74	42	5 (11.9)		5 (11.9)	6 (14.3)		25 (59.5)		1 (2.4)		
75-79	8	.	.	1 (12.5)	2 (25.0)		5 (62.5)		.	.	
80-84	
85-	1	1 (100.0)		.	.	
全年齢	17,312	10,325 (59.6)		1,757 (10.1)	1,554 (9.0)		2,694 (15.6)		982 (5.7)		

4. 全身性エリテマトーデス

	1988年度					1992年度				
	受給者数 (100%)	9年継続	4年継続	継続なし	中止後再開	受給者数 (100%)	5年継続	継続なし		
		889297	8892×	88××	88×97		9297	92×		
男 0-4	2		1 (50.0)	1 (50.0)		2	2 (100.0)			
5-9	12	6 (50.0)	3 (25.0)	3 (25.0)		12	8 (66.7)	4 (33.3)		
10-14	56	41 (73.2)	10 (17.9)	5 (8.9)		62	47 (75.8)	15 (24.2)		
15-19	179	108 (60.3)	32 (17.9)	33 (18.4)	6 (3.4)	195	144 (73.8)	51 (26.2)		
20-24	220	124 (56.4)	31 (14.1)	57 (25.9)	8 (3.6)	271	205 (75.6)	66 (24.4)		
25-29	208	138 (66.3)	21 (10.1)	44 (21.2)	5 (2.4)	299	232 (77.6)	67 (22.4)		
30-34	202	138 (68.3)	29 (14.4)	32 (15.8)	3 (1.5)	253	211 (83.4)	42 (16.6)		
35-39	247	163 (66.0)	40 (16.2)	42 (17.0)	2 (0.8)	287	221 (77.0)	66 (23.0)		
40-44	204	134 (65.7)	27 (13.2)	34 (16.7)	9 (4.4)	338	263 (77.8)	75 (22.2)		
45-49	148	102 (68.9)	19 (12.8)	25 (16.9)	2 (1.4)	265	212 (80.0)	53 (20.0)		
50-54	150	91 (60.7)	28 (18.7)	31 (20.7)		234	185 (79.1)	49 (20.9)		
55-59	155	92 (59.4)	32 (20.6)	31 (20.0)		226	168 (74.3)	58 (25.7)		
60-64	157	58 (36.9)	41 (26.1)	53 (33.8)	5 (3.2)	252	168 (66.7)	84 (33.3)		
65-69	93	27 (29.0)	23 (24.7)	42 (45.2)	1 (1.1)	196	119 (60.7)	77 (39.3)		
70-74	49	8 (16.3)	12 (24.5)	29 (59.2)		86	37 (43.0)	49 (57.0)		
75-79	10		3 (30.0)	7 (70.0)		35	13 (37.1)	22 (62.9)		
80-84	2			2 (100.0)		9	1 (11.1)	8 (88.9)		
85-						1		1 (100.0)		
全年齢	2,094	1,230 (58.7)	352 (16.8)	471 (22.5)	41 (2.0)	3,023	2,236 (74.0)	787 (26.0)		
女 0-4	15	5 (33.3)	5 (33.3)	5 (33.3)		5	3 (60.0)	2 (40.0)		
5-9	53	35 (66.0)	9 (17.0)	8 (15.1)	1 (1.9)	54	35 (64.8)	19 (35.2)		
10-14	406	290 (71.4)	58 (14.3)	50 (12.3)	8 (2.0)	343	293 (85.4)	50 (14.6)		
15-19	1,172	827 (70.6)	157 (13.4)	157 (13.4)	31 (2.6)	1,286	1,055 (82.0)	231 (18.0)		
20-24	1,685	1,193 (70.8)	197 (11.7)	242 (14.4)	53 (3.1)	2,151	1,732 (80.5)	419 (19.5)		
25-29	2,262	1,628 (72.0)	251 (11.1)	320 (14.1)	63 (2.8)	2,618	2,130 (81.4)	488 (18.6)		
30-34	2,830	2,085 (73.7)	328 (11.6)	370 (13.1)	47 (1.7)	3,141	2,595 (82.6)	546 (17.4)		
35-39	3,867	2,969 (76.8)	409 (10.6)	427 (11.0)	62 (1.6)	3,683	3,079 (83.6)	604 (16.4)		
40-44	3,770	2,913 (77.3)	405 (10.7)	399 (10.6)	53 (1.4)	5,165	4,417 (85.5)	748 (14.5)		
45-49	3,103	2,328 (75.0)	408 (13.1)	336 (10.8)	31 (1.0)	4,384	3,752 (85.6)	632 (14.4)		
50-54	2,395	1,695 (70.8)	360 (15.0)	304 (12.7)	36 (1.5)	3,537	2,934 (83.0)	603 (17.0)		
55-59	1,691	1,141 (67.5)	276 (16.3)	259 (15.3)	15 (0.9)	2,750	2,235 (81.3)	515 (18.7)		
60-64	1,205	651 (54.0)	253 (21.0)	267 (22.2)	34 (2.8)	1,807	1,402 (77.6)	405 (22.4)		
65-69	619	239 (38.6)	141 (22.8)	226 (36.5)	13 (2.1)	1,152	764 (66.3)	388 (33.7)		
70-74	149	58 (38.9)	28 (18.8)	60 (40.3)	3 (2.0)	444	259 (58.3)	185 (41.7)		
75-79	29	4 (13.8)	6 (20.7)	19 (65.5)		107	64 (59.8)	43 (40.2)		
80-84	13		2 (15.4)	11 (84.6)		17	6 (35.3)	11 (64.7)		
85-	4	1 (25.0)	1 (25.0)	2 (50.0)		4		4 (100.0)		
全年齢	25,268	18,062 (71.5)	3,294 (13.0)	3,462 (13.7)	450 (1.8)	32,648	26,755 (81.9)	5,893 (18.1)		


4. 全身性エリテマトーデス

図1-5-1-4 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




4. 全身性エリテマトーデス

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



■ 5年継続 □ 継続なし

5. スモン

表1-5-1-5 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開		
		84889297		848892×		8488×		84×		8488×97	84×9297	84×92×
男												
0-4
5-9
10-14
15-19
20-24	1	1 (100.0)
25-29	3	3 (100.0)
30-34	3	2 (66.7)	1 (33.3)
35-39	11	6 (54.5)	2 (18.2)	1 (9.1)	2 (18.2)
40-44	20	9 (45.0)	1 (5.0)	1 (5.0)	5 (25.0)	4 (20.0)	.	.	.	4 (20.0)	.	.
45-49	42	26 (61.9)	3 (7.1)	4 (9.5)	7 (16.7)	2 (4.8)	.	.	.	2 (4.8)	.	.
50-54	68	44 (64.7)	8 (11.8)	3 (4.4)	8 (11.8)	5 (7.4)	.	.	.	5 (7.4)	.	.
55-59	105	66 (62.9)	16 (15.2)	10 (9.5)	9 (8.6)	4 (3.8)	.	.	.	4 (3.8)	.	.
60-64	84	32 (38.1)	7 (8.3)	20 (23.8)	22 (26.2)	3 (3.6)	.	.	.	3 (3.6)	.	.
65-69	92	33 (35.9)	17 (18.5)	12 (13.0)	25 (27.2)	5 (5.4)	.	.	.	5 (5.4)	.	.
70-74	34	6 (17.6)	9 (26.5)	7 (20.6)	11 (32.4)	1 (2.9)	.	.	.	1 (2.9)	.	.
75-79	20	4 (20.0)	1 (5.0)	4 (20.0)	9 (45.0)	2 (10.0)	.	.	.	2 (10.0)	.	.
80-84	5	.	1 (20.0)	2 (40.0)	2 (40.0)
85-
全年齢	488	231 (47.3)	65 (13.3)	65 (13.3)	101 (20.7)	26 (5.3)	.	.	.	26 (5.3)	.	.
女												
0-4
5-9
10-14
15-19	4	3 (75.0)	.	.	1 (25.0)
20-24	3	2 (66.7)	.	.	1 (33.3)
25-29	1	1 (100.0)
30-34	10	6 (60.0)	.	.	2 (20.0)	2 (20.0)	.	.	.	2 (20.0)	.	.
35-39	30	20 (66.7)	2 (6.7)	4 (13.3)	3 (10.0)	1 (3.3)	.	.	.	1 (3.3)	.	.
40-44	73	50 (68.5)	4 (5.5)	2 (2.7)	12 (16.4)	5 (6.8)	.	.	.	5 (6.8)	.	.
45-49	144	103 (71.5)	9 (6.3)	7 (4.9)	13 (9.0)	12 (8.3)	.	.	.	12 (8.3)	.	.
50-54	228	173 (75.9)	20 (8.8)	6 (2.6)	23 (10.1)	6 (2.6)	.	.	.	6 (2.6)	.	.
55-59	329	202 (61.4)	36 (10.9)	45 (13.7)	30 (9.1)	16 (4.9)	.	.	.	16 (4.9)	.	.
60-64	384	193 (50.3)	40 (10.4)	55 (14.3)	54 (14.1)	42 (10.9)	.	.	.	42 (10.9)	.	.
65-69	292	110 (37.7)	30 (10.3)	38 (13.0)	82 (28.1)	32 (11.0)	.	.	.	32 (11.0)	.	.
70-74	157	53 (33.8)	29 (18.5)	31 (19.7)	33 (21.0)	11 (7.0)	.	.	.	11 (7.0)	.	.
75-79	49	11 (22.4)	7 (14.3)	12 (24.5)	16 (32.7)	3 (6.1)	.	.	.	3 (6.1)	.	.
80-84	15	.	5 (33.3)	2 (13.3)	7 (46.7)	1 (6.7)	.	.	.	1 (6.7)	.	.
85-	12	4 (33.3)	1 (8.3)	3 (25.0)	4 (33.3)
全年齢	1,731	931 (53.8)	183 (10.6)	206 (11.9)	280 (16.2)	131 (7.6)	.	.	.	131 (7.6)	.	.

5. スモン

	1988年度				1992年度			
	受給者数 (100%)	9年継続	4年継続	継続なし	中止後再開	受給者数 (100%)	5年継続	継続なし
		889297	8892×	88××	88×97		9297	92×
男 0-4
5-9
10-14
15-19
20-24	1	1 (100.0)
25-29	1	1 (100.0)	.	.	.	2	1 (50.0)	1 (50.0)
30-34	2	2 (100.0)	.	.	.	1	1 (100.0)	.
35-39	7	6 (85.7)	1 (14.3)	.	.	5	4 (80.0)	1 (20.0)
40-44	12	8 (66.7)	2 (16.7)	1 (8.3)	1 (8.3)	9	7 (77.8)	2 (22.2)
45-49	35	19 (54.3)	2 (5.7)	8 (22.9)	6 (17.1)	13	11 (84.6)	2 (15.4)
50-54	51	42 (82.4)	4 (7.8)	4 (7.8)	1 (2.0)	32	28 (87.5)	4 (12.5)
55-59	85	64 (75.3)	11 (12.9)	7 (8.2)	3 (3.5)	60	52 (86.7)	8 (13.3)
60-64	122	83 (68.0)	18 (14.8)	16 (13.1)	5 (4.1)	98	81 (82.7)	17 (17.3)
65-69	74	38 (51.4)	12 (16.2)	21 (28.4)	3 (4.1)	97	73 (75.3)	24 (24.7)
70-74	71	28 (39.4)	22 (31.0)	19 (26.8)	2 (2.8)	58	42 (72.4)	16 (27.6)
75-79	25	7 (28.0)	11 (44.0)	5 (20.0)	2 (8.0)	52	27 (51.9)	25 (48.1)
80-84	14	3 (21.4)	2 (14.3)	9 (64.3)	.	19	10 (52.6)	9 (47.4)
85-	4	.	.	4 (100.0)	.	8	4 (50.0)	4 (50.0)
全年齢	504	302 (59.9)	85 (16.9)	94 (18.7)	23 (4.6)	454	341 (75.1)	113 (24.9)
女 0-4
5-9
10-14
15-19
20-24	4	3 (75.0)	.	1 (25.0)
25-29	2	1 (50.0)	.	1 (50.0)	.	3	3 (100.0)	.
30-34	2	2 (100.0)	.	.	.	2	2 (100.0)	.
35-39	20	14 (70.0)	3 (15.0)	3 (15.0)	.	2	2 (100.0)	.
40-44	42	31 (73.8)	6 (14.3)	4 (9.5)	1 (2.4)	27	24 (88.9)	3 (11.1)
45-49	84	75 (89.3)	4 (4.8)	3 (3.6)	2 (2.4)	53	46 (86.8)	7 (13.2)
50-54	173	141 (81.5)	14 (8.1)	13 (7.5)	5 (2.9)	99	91 (91.9)	8 (8.1)
55-59	262	216 (82.4)	25 (9.5)	18 (6.9)	3 (1.1)	179	163 (91.1)	16 (8.9)
60-64	362	235 (64.9)	48 (13.3)	65 (18.0)	14 (3.9)	295	252 (85.4)	43 (14.6)
65-69	334	189 (56.6)	42 (12.6)	74 (22.2)	29 (8.7)	320	250 (78.1)	70 (21.9)
70-74	232	120 (51.7)	42 (18.1)	58 (25.0)	12 (5.2)	253	200 (79.1)	53 (20.9)
75-79	131	56 (42.7)	30 (22.9)	42 (32.1)	3 (2.3)	173	120 (69.4)	53 (30.6)
80-84	54	19 (35.2)	15 (27.8)	20 (37.0)	.	105	61 (58.1)	44 (41.9)
85-	20	3 (15.0)	4 (20.0)	13 (65.0)	.	44	21 (47.7)	23 (52.3)
全年齢	1,722	1,105 (64.2)	233 (13.5)	315 (18.3)	69 (4.0)	1,555	1,235 (79.4)	320 (20.6)


5. スモン

図1-5-1-5 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




5. スモン

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



■ 5年継続 □ 継続なし

6. 再生不良性貧血

表1-5-1-6 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
		84889297		848892×		8488×		84×××		8488×97 84×9297 84×92× 84××97	
男 0-4	69	20 (29.0)	8 (11.6)	12 (17.4)	27 (39.1)	2 (2.9)					
5-9	147	49 (33.3)	16 (10.9)	22 (15.0)	56 (38.1)	4 (2.7)					
10-14	215	53 (24.7)	36 (16.7)	53 (24.7)	67 (31.2)	6 (2.8)					
15-19	189	41 (21.7)	25 (13.2)	40 (21.2)	74 (39.2)	9 (4.8)					
20-24	152	34 (22.4)	20 (13.2)	25 (16.4)	69 (45.4)	4 (2.6)					
25-29	98	27 (27.6)	21 (21.4)	17 (17.3)	28 (28.6)	5 (5.1)					
30-34	122	38 (31.1)	19 (15.6)	21 (17.2)	38 (31.1)	6 (4.9)					
35-39	93	32 (34.4)	11 (11.8)	16 (17.2)	31 (33.3)	3 (3.2)					
40-44	95	27 (28.4)	10 (10.5)	23 (24.2)	34 (35.8)	1 (1.1)					
45-49	105	36 (34.3)	13 (12.4)	17 (16.2)	36 (34.3)	3 (2.9)					
50-54	102	35 (34.3)	12 (11.8)	17 (16.7)	31 (30.4)	7 (6.9)					
55-59	130	28 (21.5)	16 (12.3)	28 (21.5)	53 (40.8)	5 (3.8)					
60-64	131	28 (21.4)	14 (10.7)	27 (20.6)	60 (45.8)	2 (1.5)					
65-69	150	14 (9.3)	9 (6.0)	28 (18.7)	96 (64.0)	3 (2.0)					
70-74	31	3 (9.7)	3 (9.7)	2 (6.5)	21 (67.7)	2 (6.5)					
75-79	11	.	.	.	11 (100.0)	.					
80-84	4	.	.	.	4 (100.0)	.					
85-					
全年齢	1,844	465 (25.2)	233 (12.6)	348 (18.9)	736 (39.9)	62 (3.4)					
女 0-4	52	16 (30.8)	5 (9.6)	8 (15.4)	23 (44.2)	.					
5-9	117	37 (31.6)	15 (12.8)	28 (23.9)	34 (29.1)	3 (2.6)					
10-14	167	67 (40.1)	26 (15.6)	36 (21.6)	31 (18.6)	7 (4.2)					
15-19	154	48 (31.2)	14 (9.1)	24 (15.6)	47 (30.5)	21 (13.6)					
20-24	132	51 (38.6)	18 (13.6)	19 (14.4)	39 (29.5)	5 (3.8)					
25-29	175	80 (45.7)	19 (10.9)	29 (16.6)	42 (24.0)	5 (2.9)					
30-34	199	100 (50.3)	16 (8.0)	28 (14.1)	47 (23.6)	8 (4.0)					
35-39	187	96 (51.3)	15 (8.0)	20 (10.7)	48 (25.7)	8 (4.3)					
40-44	215	98 (45.6)	25 (11.6)	24 (11.2)	58 (27.0)	10 (4.7)					
45-49	209	79 (37.8)	36 (17.2)	22 (10.5)	64 (30.6)	8 (3.8)					
50-54	276	119 (43.1)	40 (14.5)	33 (12.0)	76 (27.5)	8 (2.9)					
55-59	306	113 (36.9)	32 (10.5)	57 (18.6)	89 (29.1)	15 (4.9)					
60-64	337	94 (27.9)	73 (21.7)	66 (19.6)	95 (28.2)	9 (2.7)					
65-69	281	49 (17.4)	28 (10.0)	52 (18.5)	144 (51.2)	8 (2.8)					
70-74	61	6 (9.8)	9 (14.8)	7 (11.5)	37 (60.7)	2 (3.3)					
75-79	8	.	1 (12.5)	1 (12.5)	6 (75.0)	.					
80-84	3	.	.	.	3 (100.0)	.					
85-	3	.	.	.	3 (100.0)	.					
全年齢	2,882	1,053 (36.5)	372 (12.9)	454 (15.8)	886 (30.7)	117 (4.1)					

6. 再生不良性貧血

	1988年度 受給者数 (100%)		9年継続		4年継続		継続なし		中止後再開		1992年度 受給者数 (100%)		5年継続		継続なし	
			889297		8892×		88××		88×97			9297		92×		
男 0-4	58	22 (37.9)	12 (20.7)	24 (41.4)	.	.	51	25 (49.0)	26 (51.0)							
5-9	137	64 (46.7)	31 (22.6)	41 (29.9)	1 (0.7)		123	79 (64.2)	44 (35.8)							
10-14	261	129 (49.4)	65 (24.9)	62 (23.8)	5 (1.9)		211	146 (69.2)	65 (30.8)							
15-19	285	102 (35.8)	68 (23.9)	106 (37.2)	9 (3.2)		335	205 (61.2)	130 (38.8)							
20-24	190	72 (37.9)	44 (23.2)	71 (37.4)	3 (1.6)		277	151 (54.5)	126 (45.5)							
25-29	170	74 (43.5)	39 (22.9)	53 (31.2)	4 (2.4)		216	138 (63.9)	78 (36.1)							
30-34	143	61 (42.7)	35 (24.5)	44 (30.8)	3 (2.1)		204	135 (66.2)	69 (33.8)							
35-39	167	71 (42.5)	45 (26.9)	51 (30.5)	.	.	164	106 (64.6)	58 (35.4)							
40-44	144	74 (51.4)	21 (14.6)	48 (33.3)	1 (0.7)		213	134 (62.9)	79 (37.1)							
45-49	145	67 (46.2)	31 (21.4)	46 (31.7)	1 (0.7)		179	115 (64.2)	64 (35.8)							
50-54	169	71 (42.0)	39 (23.1)	58 (34.3)	1 (0.6)		193	125 (64.8)	68 (35.2)							
55-59	184	79 (42.9)	37 (20.1)	65 (35.3)	3 (1.6)		245	147 (60.0)	98 (40.0)							
60-64	221	57 (25.8)	59 (26.7)	103 (46.6)	2 (0.9)		249	147 (59.0)	102 (41.0)							
65-69	202	52 (25.7)	29 (14.4)	118 (58.4)	3 (1.5)		275	117 (42.5)	158 (57.5)							
70-74	95	13 (13.7)	20 (21.1)	59 (62.1)	3 (3.2)		161	81 (50.3)	80 (49.7)							
75-79	47	7 (14.9)	11 (23.4)	29 (61.7)	.	.	70	25 (35.7)	45 (64.3)							
80-84	13	.	1 (7.7)	12 (92.3)	.	.	36	9 (25.0)	27 (75.0)							
85-	4	.	.	4 (100.0)	.	.	9	1 (11.1)	8 (88.9)							
全年齢	2,635	1,015 (38.5)	587 (22.3)	994 (37.7)	39 (1.5)		3,211	1,886 (58.7)	1,325 (41.3)							
女 0-4	47	11 (23.4)	12 (25.5)	24 (51.1)	.	.	37	25 (67.6)	12 (32.4)							
5-9	124	51 (41.1)	22 (17.7)	50 (40.3)	1 (0.8)		123	74 (60.2)	49 (39.8)							
10-14	216	108 (50.0)	49 (22.7)	58 (26.9)	1 (0.5)		184	122 (66.3)	62 (33.7)							
15-19	250	116 (46.4)	41 (16.4)	83 (33.2)	10 (4.0)		266	185 (69.5)	81 (30.5)							
20-24	195	105 (53.8)	38 (19.5)	47 (24.1)	5 (2.6)		242	170 (70.2)	72 (29.8)							
25-29	189	105 (55.6)	30 (15.9)	51 (27.0)	3 (1.6)		256	183 (71.5)	73 (28.5)							
30-34	240	142 (59.2)	36 (15.0)	59 (24.6)	3 (1.3)		240	176 (73.3)	64 (26.7)							
35-39	252	161 (63.9)	34 (13.5)	54 (21.4)	3 (1.2)		258	199 (77.1)	59 (22.9)							
40-44	225	135 (60.0)	34 (15.1)	53 (23.6)	3 (1.3)		323	246 (76.2)	77 (23.8)							
45-49	274	165 (60.2)	47 (17.2)	58 (21.2)	4 (1.5)		276	207 (75.0)	69 (25.0)							
50-54	278	147 (52.9)	59 (21.2)	69 (24.8)	3 (1.1)		334	240 (71.9)	94 (28.1)							
55-59	374	202 (54.0)	66 (17.6)	104 (27.8)	2 (0.5)		395	273 (69.1)	122 (30.9)							
60-64	413	192 (46.5)	92 (22.3)	120 (29.1)	9 (2.2)		461	318 (69.0)	143 (31.0)							
65-69	389	120 (30.8)	104 (26.7)	157 (40.4)	8 (2.1)		515	310 (60.2)	205 (39.8)							
70-74	187	66 (35.3)	39 (20.9)	78 (41.7)	4 (2.1)		318	161 (50.6)	157 (49.4)							
75-79	56	5 (8.9)	14 (25.0)	36 (64.3)	1 (1.8)		136	72 (52.9)	64 (47.1)							
80-84	20	3 (15.0)	6 (30.0)	11 (55.0)	.	.	47	15 (31.9)	32 (68.1)							
85-	5	.	.	5 (100.0)	.	.	20	4 (20.0)	16 (80.0)							
全年齢	3,734	1,834 (49.1)	723 (19.4)	1,117 (29.9)	60 (1.6)		4,431	2,980 (67.3)	1,451 (32.7)							


6. 再生不良性貧血

図1-5-1-5 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




6. 再生不良性貧血

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



■ 5年継続 □ 継続なし