

7. サルコイドーシス

表1-5-1-7 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
		84889297		848892×		8488×		84×		8488×97	84×9297
男										84×92×	84×97
0-4	1	1	(100.0)
5-9	2	.	.	1	(50.0)	.	.	1	(50.0)	.	.
10-14	15	2	(13.3)	5	(33.3)	3	(20.0)	5	(33.3)	.	.
15-19	58	10	(17.2)	2	(3.4)	15	(25.9)	29	(50.0)	2	(3.4)
20-24	137	17	(12.4)	9	(6.6)	18	(13.1)	86	(62.8)	7	(5.1)
25-29	145	33	(22.8)	19	(13.1)	17	(11.7)	67	(46.2)	9	(6.2)
30-34	105	47	(44.8)	12	(11.4)	8	(7.6)	37	(35.2)	1	(1.0)
35-39	73	34	(46.6)	6	(8.2)	9	(12.3)	23	(31.5)	1	(1.4)
40-44	76	29	(38.2)	7	(9.2)	12	(15.8)	23	(30.3)	5	(6.6)
45-49	61	28	(45.9)	10	(16.4)	4	(6.6)	17	(27.9)	2	(3.3)
50-54	66	30	(45.5)	12	(18.2)	5	(7.6)	15	(22.7)	4	(6.1)
55-59	54	25	(46.3)	5	(9.3)	9	(16.7)	15	(27.8)	.	.
60-64	53	13	(24.5)	10	(18.9)	13	(24.5)	16	(30.2)	1	(1.9)
65-69	30	7	(23.3)	6	(20.0)	4	(13.3)	12	(40.0)	1	(3.3)
70-74	5	2	(40.0)	3	(60.0)	.	.
75-79	2	2	(100.0)	.	.
80-84
85-
全年齢	883	276	(31.3)	104	(11.8)	119	(13.5)	351	(39.8)	33	(3.7)
女											
0-4	2	2	(100.0)	.	.
5-9	5	3	(60.0)	.	.	1	(20.0)	1	(20.0)	.	.
10-14	19	8	(42.1)	.	.	5	(26.3)	5	(26.3)	1	(5.3)
15-19	63	16	(25.4)	5	(7.9)	8	(12.7)	30	(47.6)	4	(6.3)
20-24	118	32	(27.1)	10	(8.5)	16	(13.6)	54	(45.8)	6	(5.1)
25-29	186	62	(33.3)	21	(11.3)	21	(11.3)	72	(38.7)	10	(5.4)
30-34	246	110	(44.7)	19	(7.7)	30	(12.2)	71	(28.9)	16	(6.5)
35-39	212	102	(48.1)	20	(9.4)	31	(14.6)	50	(23.6)	9	(4.2)
40-44	199	105	(52.8)	22	(11.1)	18	(9.0)	42	(21.1)	12	(6.0)
45-49	224	138	(61.6)	23	(10.3)	18	(8.0)	39	(17.4)	6	(2.7)
50-54	410	241	(58.8)	47	(11.5)	44	(10.7)	67	(16.3)	11	(2.7)
55-59	463	244	(52.7)	69	(14.9)	49	(10.6)	86	(18.6)	15	(3.2)
60-64	373	151	(40.5)	67	(18.0)	73	(19.6)	69	(18.5)	13	(3.5)
65-69	185	56	(30.3)	27	(14.6)	30	(16.2)	60	(32.4)	12	(6.5)
70-74	41	6	(14.6)	4	(9.8)	11	(26.8)	18	(43.9)	2	(4.9)
75-79	3	1	(33.3)	1	(33.3)	.	.	1	(33.3)	.	.
80-84	1	1	(100.0)	.	.
85-
全年齢	2,750	1,275	(46.4)	335	(12.2)	355	(12.9)	668	(24.3)	117	(4.3)

7. サルコイドーシス

	1988年度					1992年度				
	受給者数 (100%)	9年継続	4年継続	継続なし	中止後再開	受給者数 (100%)	5年継続	継続なし		
		889297	8892×	88×	88×97		9297	92×		
男 0-4	1	1 (100.0)	.	.	.	1	.	1 (100.0)		
5-9	2	1 (50.0)	.	1 (50.0)	.	3	1 (33.3)	2 (66.7)		
10-14	13	2 (15.4)	7 (53.8)	4 (30.8)	.	13	7 (53.8)	6 (46.2)		
15-19	41	14 (34.1)	9 (22.0)	18 (43.9)	.	36	10 (27.8)	26 (72.2)		
20-24	224	69 (30.8)	41 (18.3)	111 (49.6)	3 (1.3)	238	104 (43.7)	134 (56.3)		
25-29	335	113 (33.7)	67 (20.0)	148 (44.2)	7 (2.1)	532	288 (54.1)	244 (45.9)		
30-34	231	106 (45.9)	48 (20.8)	72 (31.2)	5 (2.2)	400	239 (59.8)	161 (40.3)		
35-39	181	104 (57.5)	25 (13.8)	50 (27.6)	2 (1.1)	277	186 (67.1)	91 (32.9)		
40-44	115	72 (62.6)	18 (15.7)	22 (19.1)	3 (2.6)	258	193 (74.8)	65 (25.2)		
45-49	138	77 (55.8)	26 (18.8)	34 (24.6)	1 (0.7)	200	146 (73.0)	54 (27.0)		
50-54	146	90 (61.6)	29 (19.9)	25 (17.1)	2 (1.4)	224	167 (74.6)	57 (25.4)		
55-59	151	86 (57.0)	22 (14.6)	39 (25.8)	4 (2.6)	279	204 (73.1)	75 (26.9)		
60-64	118	64 (54.2)	25 (21.2)	29 (24.6)	.	253	188 (74.3)	65 (25.7)		
65-69	88	35 (39.8)	16 (18.2)	35 (39.8)	2 (2.3)	159	98 (61.6)	61 (38.4)		
70-74	27	9 (33.3)	10 (37.0)	8 (29.6)	.	79	51 (64.6)	28 (35.4)		
75-79	4	.	1 (25.0)	3 (75.0)	.	20	8 (40.0)	12 (60.0)		
80-84	2	1 (50.0)	1 (50.0)		
85-		
全年齢	1,815	843 (46.4)	344 (19.0)	599 (33.0)	29 (1.6)	2,974	1,891 (63.6)	1,083 (36.4)		
女 0-4	1	1 (100.0)	.	.	.	2	2 (100.0)	.		
5-9	5	3 (60.0)	1 (20.0)	1 (20.0)	.	2	1 (50.0)	1 (50.0)		
10-14	22	10 (45.5)	4 (18.2)	7 (31.8)	1 (4.5)	19	13 (68.4)	6 (31.6)		
15-19	49	15 (30.6)	12 (24.5)	22 (44.9)	.	46	26 (56.5)	20 (43.5)		
20-24	143	60 (42.0)	28 (19.6)	49 (34.3)	6 (4.2)	156	94 (60.3)	62 (39.7)		
25-29	300	129 (43.0)	51 (17.0)	110 (36.7)	10 (3.3)	370	220 (59.5)	150 (40.5)		
30-34	329	180 (54.7)	65 (19.8)	75 (22.8)	9 (2.7)	433	286 (66.1)	147 (33.9)		
35-39	323	207 (64.1)	39 (12.1)	71 (22.0)	6 (1.9)	435	312 (71.7)	123 (28.3)		
40-44	306	189 (61.8)	46 (15.0)	66 (21.6)	5 (1.6)	467	362 (77.5)	105 (22.5)		
45-49	328	222 (67.7)	51 (15.5)	51 (15.5)	4 (1.2)	459	354 (77.1)	105 (22.9)		
50-54	578	403 (69.7)	98 (17.0)	72 (12.5)	5 (0.9)	717	592 (82.6)	125 (17.4)		
55-59	713	520 (72.9)	94 (13.2)	92 (12.9)	7 (1.0)	1,094	905 (82.7)	189 (17.3)		
60-64	686	426 (62.1)	137 (20.0)	112 (16.3)	11 (1.6)	1,131	951 (84.1)	180 (15.9)		
65-69	442	220 (49.8)	91 (20.6)	126 (28.5)	5 (1.1)	889	629 (70.8)	260 (29.2)		
70-74	158	58 (36.7)	38 (24.1)	60 (38.0)	2 (1.3)	405	263 (64.9)	142 (35.1)		
75-79	35	8 (22.9)	8 (22.9)	17 (48.6)	2 (5.7)	121	71 (58.7)	50 (41.3)		
80-84	7	.	.	7 (100.0)	.	23	13 (56.5)	10 (43.5)		
85-	2	.	.	2 (100.0)	.	3	1 (33.3)	2 (66.7)		
全年齢	4,427	2,651 (59.9)	763 (17.2)	940 (21.2)	73 (1.6)	6,772	5,095 (75.2)	1,677 (24.8)		

7. サルコイドーシス



図1-5-1-7 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別



7. サルコイドーシス

1988年度受給者 継続状況

1992年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

□ 5年継続 □ 継続なし

8. 筋萎縮性側索硬化症

表1-5-1-8 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
		84889297		848892×		8488×		84×××		8488×97 84×9297 84×92× 84××97	
男 0-4	3	.	.	1 (33.3)	.	1 (33.3)	.	1 (33.3)	.	.	.
5-9	2	1 (50.0)	1 (50.0)	.	.	.
10-14	4	1 (25.0)	3 (75.0)	.	.	.
15-19	7	1 (14.3)	.	.	.	1 (14.3)	.	5 (71.4)	.	.	.
20-24	10	2 (20.0)	2 (20.0)	6 (60.0)	.	.	.
25-29	14	3 (21.4)	1 (7.1)	2 (14.3)	8 (57.1)
30-34	21	4 (19.0)	.	5 (23.8)	12 (57.1)
35-39	31	5 (16.1)	4 (12.9)	8 (25.8)	13 (41.9)	1 (3.2)
40-44	53	12 (22.6)	3 (5.7)	8 (15.1)	30 (56.6)
45-49	96	24 (25.0)	9 (9.4)	10 (10.4)	49 (51.0)	4 (4.2)
50-54	148	16 (10.8)	18 (12.2)	16 (10.8)	96 (64.9)	2 (1.4)
55-59	183	22 (12.0)	17 (9.3)	30 (16.4)	109 (59.6)	5 (2.7)
60-64	180	12 (6.7)	17 (9.4)	26 (14.4)	122 (67.8)	3 (1.7)
65-69	122	5 (4.1)	4 (3.3)	18 (14.8)	93 (76.2)	2 (1.6)
70-74	28	1 (3.6)	.	1 (3.6)	26 (92.9)
75-79	4	.	.	.	4 (100.0)
80-84
85-
全年齢	906	109 (12.0)	76 (8.4)	126 (13.9)	578 (63.8)	17 (1.9)
女 0-4	2	.	.	.	2 (100.0)
5-9	1	.	.	.	1 (100.0)
10-14	2	1 (50.0)	.	1 (50.0)
15-19	3	2 (66.7)	.	1 (33.3)
20-24
25-29	6	2 (33.3)	1 (16.7)	1 (16.7)	2 (33.3)
30-34	13	5 (38.5)	3 (23.1)	.	4 (30.8)	1 (7.7)
35-39	28	6 (21.4)	2 (7.1)	4 (14.3)	16 (57.1)
40-44	41	9 (22.0)	4 (9.8)	6 (14.6)	21 (51.2)	1 (2.4)
45-49	56	6 (10.7)	10 (17.9)	12 (21.4)	28 (50.0)
50-54	97	13 (13.4)	5 (5.2)	13 (13.4)	66 (68.0)
55-59	108	11 (10.2)	7 (6.5)	15 (13.9)	73 (67.6)	2 (1.9)
60-64	140	6 (4.3)	5 (3.6)	19 (13.6)	108 (77.1)	2 (1.4)
65-69	107	2 (1.9)	.	10 (9.3)	93 (86.9)	2 (1.9)
70-74	20	.	.	2 (10.0)	18 (90.0)
75-79	5	.	.	.	5 (100.0)
80-84	1	.	.	.	1 (100.0)
85-
全年齢	630	63 (10.0)	37 (5.9)	84 (13.3)	438 (69.5)	8 (1.3)

8. 筋萎縮性側索硬化症

	1988年度				1992年度			
	受給者数 (100%)	9年継続	4年継続	継続なし	中止後再開	受給者数 (100%)	5年継続	継続なし
		889297	8892×	88×	88×	9297	92×	
男 0-4	1	.	.	1 (100.0)	.	1	1 (100.0)	.
5-9	6	1 (16.7)	3 (50.0)	2 (33.3)	.	3	2 (66.7)	1 (33.3)
10-14	2	2 (100.0)	.	.	.	3	1 (33.3)	2 (66.7)
15-19	5	3 (60.0)	.	2 (40.0)	.	8	5 (62.5)	3 (37.5)
20-24	14	6 (42.9)	4 (28.6)	4 (28.6)	.	7	5 (71.4)	2 (28.6)
25-29	12	7 (58.3)	.	5 (41.7)	.	13	7 (53.8)	6 (46.2)
30-34	31	7 (22.6)	7 (22.6)	17 (54.8)	.	19	12 (63.2)	7 (36.8)
35-39	51	18 (35.3)	10 (19.6)	23 (45.1)	.	45	20 (44.4)	25 (55.6)
40-44	91	31 (34.1)	15 (16.5)	44 (48.4)	1 (1.1)	96	49 (51.0)	47 (49.0)
45-49	133	43 (32.3)	20 (15.0)	67 (50.4)	3 (2.3)	164	82 (50.0)	82 (50.0)
50-54	214	60 (28.0)	35 (16.4)	113 (52.8)	6 (2.8)	218	104 (47.7)	114 (52.3)
55-59	275	39 (14.2)	60 (21.8)	171 (62.2)	5 (1.8)	348	133 (38.2)	215 (61.8)
60-64	320	49 (15.3)	57 (17.8)	211 (65.9)	3 (0.9)	389	98 (25.2)	291 (74.8)
65-69	241	21 (8.7)	43 (17.8)	175 (72.6)	2 (0.8)	381	107 (28.1)	274 (71.9)
70-74	103	8 (7.8)	6 (5.8)	89 (86.4)	.	182	34 (18.7)	148 (81.3)
75-79	35	1 (2.9)	1 (2.9)	33 (94.3)	.	69	11 (15.9)	58 (84.1)
80-84	5	.	1 (20.0)	4 (80.0)	.	13	1 (7.7)	12 (92.3)
85-	3	.	3 (100.0)
全年齢	1,539	296 (19.2)	262 (17.0)	961 (62.4)	20 (1.3)	1,962	672 (34.3)	1,290 (65.7)
女 0-4	1	1 (100.0)	.	.	.	1	.	1 (100.0)
5-9	1	1 (100.0)	.	.	.	2	2 (100.0)	.
10-14	2	.	.	2 (100.0)	.	1	1 (100.0)	.
15-19	4	1 (25.0)	2 (50.0)	1 (25.0)
20-24	6	3 (50.0)	.	3 (50.0)	.	4	2 (50.0)	2 (50.0)
25-29	3	1 (33.3)	.	1 (33.3)	1 (33.3)	4	3 (75.0)	1 (25.0)
30-34	13	6 (46.2)	4 (30.8)	3 (23.1)	.	14	8 (57.1)	6 (42.9)
35-39	39	15 (38.5)	9 (23.1)	15 (38.5)	.	31	18 (58.1)	13 (41.9)
40-44	48	17 (35.4)	7 (14.6)	24 (50.0)	.	53	26 (49.1)	27 (50.9)
45-49	97	24 (24.7)	20 (20.6)	53 (54.6)	.	85	40 (47.1)	45 (52.9)
50-54	108	25 (23.1)	24 (22.2)	58 (53.7)	1 (0.9)	130	51 (39.2)	79 (60.8)
55-59	174	29 (16.7)	42 (24.1)	102 (58.6)	1 (0.6)	177	63 (35.6)	114 (64.4)
60-64	204	19 (9.3)	26 (12.7)	158 (77.5)	1 (0.5)	218	54 (24.8)	164 (75.2)
65-69	162	11 (6.8)	18 (11.1)	131 (80.9)	2 (1.2)	224	48 (21.4)	176 (78.6)
70-74	66	3 (4.5)	2 (3.0)	60 (90.9)	1 (1.5)	118	21 (17.8)	97 (82.2)
75-79	32	.	2 (6.3)	30 (93.8)	.	50	5 (10.0)	45 (90.0)
80-84	6	.	.	6 (100.0)	.	18	.	18 (100.0)
85-	2	.	.	2 (100.0)	.	1	.	1 (100.0)
全年齢	968	156 (16.1)	156 (16.1)	649 (67.0)	7 (0.7)	1,131	342 (30.2)	789 (69.8)


8. 筋萎縮性側索硬化症

図1-5-1-8 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




8. 筋萎縮性側索硬化症

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



□ 5年継続 □ 継続なし

9. 強皮症・皮膚筋炎及び多発性筋炎

表1-5-1-9 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開 8488×97 84×9297 84×92× 84×97
		84889297		848892×		8488××		84××		
男 0-4	7	3 (42.9)	.	.	1 (14.3)	2 (28.6)	1 (14.3)			
5-9	50	20 (40.0)	3 (6.0)	9 (18.0)	18 (36.0)	.	.			
10-14	64	18 (28.1)	10 (15.6)	18 (28.1)	17 (26.6)	1 (1.6)				
15-19	48	11 (22.9)	4 (8.3)	10 (20.8)	20 (41.7)	3 (6.3)				
20-24	23	7 (30.4)	1 (4.3)	4 (17.4)	10 (43.5)	1 (4.3)				
25-29	34	10 (29.4)	7 (20.6)	6 (17.6)	8 (23.5)	3 (8.8)				
30-34	50	20 (40.0)	14 (28.0)	6 (12.0)	9 (18.0)	1 (2.0)				
35-39	79	39 (49.4)	9 (11.4)	11 (13.9)	19 (24.1)	1 (1.3)				
40-44	102	42 (41.2)	10 (9.8)	14 (13.7)	32 (31.4)	4 (3.9)				
45-49	120	64 (53.3)	11 (9.2)	8 (6.7)	34 (28.3)	3 (2.5)				
50-54	150	59 (39.3)	29 (19.3)	20 (13.3)	38 (25.3)	4 (2.7)				
55-59	146	44 (30.1)	20 (13.7)	30 (20.5)	45 (30.8)	7 (4.8)				
60-64	118	20 (16.9)	21 (17.8)	36 (30.5)	39 (33.1)	2 (1.7)				
65-69	102	12 (11.8)	10 (9.8)	23 (22.5)	53 (52.0)	4 (3.9)				
70-74	21	.	3 (14.3)	1 (4.8)	16 (76.2)	1 (4.8)				
75-79	4	.	.	1 (25.0)	3 (75.0)	.				
80-84				
85-				
全年齢	1,118	369 (33.0)	152 (13.6)	198 (17.7)	363 (32.5)	36 (3.2)				
女 0-4	24	7 (29.2)	2 (8.3)	5 (20.8)	7 (29.2)	3 (12.5)				
5-9	44	18 (40.9)	5 (11.4)	8 (18.2)	13 (29.5)	.				
10-14	106	35 (33.0)	20 (18.9)	17 (16.0)	29 (27.4)	5 (4.7)				
15-19	107	37 (34.6)	10 (9.3)	18 (16.8)	36 (33.6)	6 (5.6)				
20-24	101	54 (53.5)	10 (9.9)	11 (10.9)	21 (20.8)	5 (5.0)				
25-29	186	87 (46.8)	23 (12.4)	17 (9.1)	43 (23.1)	16 (8.6)				
30-34	388	214 (55.2)	41 (10.6)	43 (11.1)	78 (20.1)	12 (3.1)				
35-39	601	338 (56.2)	67 (11.1)	62 (10.3)	97 (16.1)	37 (6.2)				
40-44	819	481 (58.7)	105 (12.8)	77 (9.4)	114 (13.9)	42 (5.1)				
45-49	980	556 (56.7)	110 (11.2)	102 (10.4)	157 (16.0)	55 (5.6)				
50-54	986	529 (53.7)	135 (13.7)	110 (11.2)	177 (18.0)	35 (3.5)				
55-59	940	401 (42.7)	169 (18.0)	136 (14.5)	191 (20.3)	43 (4.6)				
60-64	686	220 (32.1)	128 (18.7)	121 (17.6)	195 (28.4)	22 (3.2)				
65-69	347	48 (13.8)	45 (13.0)	69 (19.9)	166 (47.8)	19 (5.5)				
70-74	52	7 (13.5)	5 (9.6)	12 (23.1)	27 (51.9)	1 (1.9)				
75-79	5	.	1 (20.0)	.	4 (80.0)	.				
80-84	3	.	.	.	3 (100.0)	.				
85-				
全年齢	6,375	3,032 (47.6)	876 (13.7)	808 (12.7)	1,358 (21.3)	301 (4.7)				

9. 強皮症・皮膚筋炎及び多発性筋炎

	1988年度 受給者数 (100%)		9年継続		4年継続		継続なし		中止後再開		1992年度 受給者数 (100%)		5年継続		継続なし	
			889297		8892×		88×		88×97			9297		92×		
男 0-4	16	5	(31.3)	3	(18.8)	7	(43.8)	1	(6.3)	9	6	(66.7)	3	(33.3)		
5-9	53	26	(49.1)	13	(24.5)	13	(24.5)	1	(1.9)	49	33	(67.3)	16	(32.7)		
10-14	84	34	(40.5)	28	(33.3)	21	(25.0)	1	(1.2)	73	50	(68.5)	23	(31.5)		
15-19	74	26	(35.1)	16	(21.6)	32	(43.2)	.	.	100	53	(53.0)	47	(47.0)		
20-24	53	22	(41.5)	7	(13.2)	23	(43.4)	1	(1.9)	84	52	(61.9)	32	(38.1)		
25-29	48	28	(58.3)	6	(12.5)	13	(27.1)	1	(2.1)	67	50	(74.6)	17	(25.4)		
30-34	74	39	(52.7)	14	(18.9)	20	(27.0)	1	(1.4)	97	70	(72.2)	27	(27.8)		
35-39	138	71	(51.4)	30	(21.7)	30	(21.7)	7	(5.1)	155	100	(64.5)	55	(35.5)		
40-44	181	120	(66.3)	21	(11.6)	37	(20.4)	3	(1.7)	253	171	(67.6)	82	(32.4)		
45-49	228	127	(55.7)	39	(17.1)	59	(25.9)	3	(1.3)	292	229	(78.4)	63	(21.6)		
50-54	252	148	(58.7)	36	(14.3)	64	(25.4)	4	(1.6)	361	248	(68.7)	113	(31.3)		
55-59	332	156	(47.0)	79	(23.8)	95	(28.6)	2	(0.6)	446	305	(68.4)	141	(31.6)		
60-64	297	109	(36.7)	66	(22.2)	115	(38.7)	7	(2.4)	492	315	(64.0)	177	(36.0)		
65-69	181	51	(28.2)	44	(24.3)	84	(46.4)	2	(1.1)	359	195	(54.3)	164	(45.7)		
70-74	89	14	(15.7)	25	(28.1)	47	(52.8)	3	(3.4)	163	73	(44.8)	90	(55.2)		
75-79	18	1	(5.6)	4	(22.2)	13	(72.2)	.	.	67	23	(34.3)	44	(65.7)		
80-84	4	.	.	1	(25.0)	3	(75.0)	.	.	13	2	(15.4)	11	(84.6)		
85-	2	.	.	1	(50.0)	1	(50.0)	.	.	3	.	.	3	(100.0)		
全年齢	2,124	977	(46.0)	433	(20.4)	677	(31.9)	37	(1.7)	3,083	1,975	(64.1)	1,108	(35.9)		
女 0-4	15	8	(53.3)	3	(20.0)	3	(20.0)	1	(6.7)	14	8	(57.1)	6	(42.9)		
5-9	62	42	(67.7)	8	(12.9)	11	(17.7)	1	(1.6)	54	39	(72.2)	15	(27.8)		
10-14	109	49	(45.0)	28	(25.7)	31	(28.4)	1	(0.9)	126	82	(65.1)	44	(34.9)		
15-19	146	69	(47.3)	32	(21.9)	39	(26.7)	6	(4.1)	171	108	(63.2)	63	(36.8)		
20-24	176	95	(54.0)	35	(19.9)	43	(24.4)	3	(1.7)	230	149	(64.8)	81	(35.2)		
25-29	263	167	(63.5)	36	(13.7)	52	(19.8)	8	(3.0)	299	216	(72.2)	83	(27.8)		
30-34	365	241	(66.0)	61	(16.7)	59	(16.2)	4	(1.1)	445	330	(74.2)	115	(25.8)		
35-39	776	523	(67.4)	106	(13.7)	135	(17.4)	12	(1.5)	718	552	(76.9)	166	(23.1)		
40-44	1,032	741	(71.8)	144	(14.0)	133	(12.9)	14	(1.4)	1,390	1,120	(80.6)	270	(19.4)		
45-49	1,368	959	(70.1)	198	(14.5)	188	(13.7)	23	(1.7)	1,739	1,382	(79.5)	357	(20.5)		
50-54	1,621	1,111	(68.5)	257	(15.9)	234	(14.4)	19	(1.2)	2,213	1,788	(80.8)	425	(19.2)		
55-59	1,477	954	(64.6)	250	(16.9)	261	(17.7)	12	(0.8)	2,378	1,866	(78.5)	512	(21.5)		
60-64	1,253	641	(51.2)	272	(21.7)	296	(23.6)	44	(3.5)	2,026	1,521	(75.1)	505	(24.9)		
65-69	755	277	(36.7)	187	(24.8)	269	(35.6)	22	(2.9)	1,460	914	(62.6)	546	(37.4)		
70-74	207	53	(25.6)	57	(27.5)	93	(44.9)	4	(1.9)	585	309	(52.8)	276	(47.2)		
75-79	48	9	(18.8)	15	(31.3)	23	(47.9)	1	(2.1)	177	75	(42.4)	102	(57.6)		
80-84	11	1	(9.1)	2	(18.2)	8	(72.7)	.	.	45	13	(28.9)	32	(71.1)		
85-	2	2	(100.0)	.	.	4	1	(25.0)	3	(75.0)		
全年齢	9,686	5,940	(61.3)	1,691	(17.5)	1,880	(19.4)	175	(1.8)	14,074	10,473	(74.4)	3,601	(25.6)		


9. 強皮症・皮膚筋炎及び多発性筋炎

図1-5-1-9 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




9. 強皮症・皮膚筋炎及び多発性筋炎

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



□ 5年継続 □ 継続なし

10. 特発性血小板減少性紫斑病

表1-5-1-10 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
		84889297	848892×	8488××	84××	8488×97 84×9297 84×92× 84××97					
男 0-4	544	49 (9.0)	36 (6.6)	100 (18.4)	343 (63.1)	16 (2.9)					
5-9	682	82 (12.0)	84 (12.3)	113 (16.6)	376 (55.1)	27 (4.0)					
10-14	447	56 (12.5)	47 (10.5)	113 (25.3)	225 (50.3)	6 (1.3)					
15-19	174	29 (16.7)	12 (6.9)	39 (22.4)	84 (48.3)	10 (5.7)					
20-24	57	14 (24.6)	2 (3.5)	12 (21.1)	28 (49.1)	1 (1.8)					
25-29	40	13 (32.5)	6 (15.0)	9 (22.5)	9 (22.5)	3 (7.5)					
30-34	55	21 (38.2)	6 (10.9)	7 (12.7)	16 (29.1)	5 (9.1)					
35-39	56	25 (44.6)	6 (10.7)	5 (8.9)	15 (26.8)	5 (8.9)					
40-44	49	21 (42.9)	5 (10.2)	9 (18.4)	11 (22.4)	3 (6.1)					
45-49	61	27 (44.3)	6 (9.8)	10 (16.4)	15 (24.6)	3 (4.9)					
50-54	64	24 (37.5)	5 (7.8)	11 (17.2)	22 (34.4)	2 (3.1)					
55-59	65	21 (32.3)	6 (9.2)	13 (20.0)	21 (32.3)	4 (6.2)					
60-64	72	14 (19.4)	19 (26.4)	14 (19.4)	22 (30.6)	3 (4.2)					
65-69	67	10 (14.9)	10 (14.9)	11 (16.4)	34 (50.7)	2 (3.0)					
70-74	18	1 (5.6)	2 (11.1)	5 (27.8)	10 (55.6)	.					
75-79	3	.	1 (33.3)	2 (66.7)	.	.					
80-84					
85-					
全年齢	2,454	407 (16.6)	253 (10.3)	473 (19.3)	1,231 (50.2)	90 (3.7)					
女 0-4	428	43 (10.0)	33 (7.7)	59 (13.8)	279 (65.2)	14 (3.3)					
5-9	645	81 (12.6)	80 (12.4)	110 (17.1)	338 (52.4)	36 (5.6)					
10-14	429	83 (19.3)	61 (14.2)	81 (18.9)	178 (41.5)	26 (6.1)					
15-19	280	90 (32.1)	18 (6.4)	36 (12.9)	119 (42.5)	17 (6.1)					
20-24	238	79 (33.2)	25 (10.5)	44 (18.5)	72 (30.3)	18 (7.6)					
25-29	342	138 (40.4)	30 (8.8)	53 (15.5)	105 (30.7)	16 (4.7)					
30-34	527	241 (45.7)	51 (9.7)	60 (11.4)	136 (25.8)	39 (7.4)					
35-39	514	263 (51.2)	59 (11.5)	59 (11.5)	98 (19.1)	35 (6.8)					
40-44	400	224 (56.0)	46 (11.5)	42 (10.5)	72 (18.0)	16 (4.0)					
45-49	382	217 (56.8)	42 (11.0)	40 (10.5)	66 (17.3)	17 (4.5)					
50-54	384	206 (53.6)	47 (12.2)	47 (12.2)	66 (17.2)	18 (4.7)					
55-59	375	176 (46.9)	57 (15.2)	45 (12.0)	74 (19.7)	23 (6.1)					
60-64	302	93 (30.8)	54 (17.9)	70 (23.2)	78 (25.8)	7 (2.3)					
65-69	169	33 (19.5)	27 (16.0)	40 (23.7)	62 (36.7)	7 (4.1)					
70-74	32	2 (6.3)	4 (12.5)	8 (25.0)	16 (50.0)	2 (6.3)					
75-79	5	1 (20.0)	.	1 (20.0)	3 (60.0)	.					
80-84	3	.	.	.	3 (100.0)	.					
85-					
全年齢	5,455	1,970 (36.1)	634 (11.6)	795 (14.6)	1,765 (32.4)	291 (5.3)					

10. 特発性血小板減少性紫斑病

	1988年度 受給者数 (100%)	9年継続		4年継続		継続なし		中止後再開		1992年度 受給者数 (100%)	5年継続		継続なし	
		889297		8892×		88×		88×			9297		92×	
男 0-4	807	98	(12.1)	101	(12.5)	601	(74.5)	7	(0.9)	790	183	(23.2)	607	(76.8)
5-9	892	202	(22.6)	183	(20.5)	497	(55.7)	10	(1.1)	912	349	(38.3)	563	(61.7)
10-14	530	127	(24.0)	138	(26.0)	263	(49.6)	2	(0.4)	637	276	(43.3)	361	(56.7)
15-19	309	77	(24.9)	56	(18.1)	173	(56.0)	3	(1.0)	384	177	(46.1)	207	(53.9)
20-24	132	53	(40.2)	17	(12.9)	60	(45.5)	2	(1.5)	216	106	(49.1)	110	(50.9)
25-29	114	44	(38.6)	18	(15.8)	51	(44.7)	1	(0.9)	177	101	(57.1)	76	(42.9)
30-34	127	57	(44.9)	21	(16.5)	48	(37.8)	1	(0.8)	165	102	(61.8)	63	(38.2)
35-39	171	89	(52.0)	30	(17.5)	50	(29.2)	2	(1.2)	229	153	(66.8)	76	(33.2)
40-44	169	88	(52.1)	24	(14.2)	55	(32.5)	2	(1.2)	325	218	(67.1)	107	(32.9)
45-49	151	83	(55.0)	28	(18.5)	39	(25.8)	1	(0.7)	296	207	(69.9)	89	(30.1)
50-54	210	114	(54.3)	38	(18.1)	52	(24.8)	6	(2.9)	308	209	(67.9)	99	(32.1)
55-59	199	90	(45.2)	45	(22.6)	61	(30.7)	3	(1.5)	414	280	(67.6)	134	(32.4)
60-64	205	92	(44.9)	45	(22.0)	63	(30.7)	5	(2.4)	375	232	(61.9)	143	(38.1)
65-69	137	40	(29.2)	41	(29.9)	54	(39.4)	2	(1.5)	341	190	(55.7)	151	(44.3)
70-74	81	20	(24.7)	24	(29.6)	37	(45.7)	.	.	204	92	(45.1)	112	(54.9)
75-79	45	5	(11.1)	10	(22.2)	28	(62.2)	2	(4.4)	109	48	(44.0)	61	(56.0)
80-84	13	.	.	4	(30.8)	9	(69.2)	.	.	47	12	(25.5)	35	(74.5)
85-	7	1	(14.3)	2	(28.6)	4	(57.1)	.	.	15	2	(13.3)	13	(86.7)
全年齢	4,299	1,280	(29.8)	825	(19.2)	2,145	(49.9)	49	(1.1)	5,944	2,937	(49.4)	3,007	(50.6)
女 0-4	553	93	(16.8)	103	(18.6)	354	(64.0)	3	(0.5)	604	165	(27.3)	439	(72.7)
5-9	802	212	(26.4)	169	(21.1)	418	(52.1)	3	(0.4)	869	339	(39.0)	530	(61.0)
10-14	622	176	(28.3)	178	(28.6)	259	(41.6)	9	(1.4)	686	361	(52.6)	325	(47.4)
15-19	440	170	(38.6)	105	(23.9)	151	(34.3)	14	(3.2)	661	338	(51.1)	323	(48.9)
20-24	410	198	(48.3)	66	(16.1)	127	(31.0)	19	(4.6)	663	393	(59.3)	270	(40.7)
25-29	594	314	(52.9)	89	(15.0)	183	(30.8)	8	(1.3)	780	519	(66.5)	261	(33.5)
30-34	691	369	(53.4)	119	(17.2)	184	(26.6)	19	(2.7)	965	670	(69.4)	295	(30.6)
35-39	843	485	(57.5)	150	(17.8)	188	(22.3)	20	(2.4)	947	683	(72.1)	264	(27.9)
40-44	806	516	(64.0)	133	(16.5)	139	(17.2)	18	(2.2)	1,264	956	(75.6)	308	(24.4)
45-49	713	474	(66.5)	119	(16.7)	112	(15.7)	8	(1.1)	1,172	905	(77.2)	267	(22.8)
50-54	745	480	(64.4)	121	(16.2)	129	(17.3)	15	(2.0)	1,162	908	(78.1)	254	(21.9)
55-59	728	476	(65.4)	117	(16.1)	126	(17.3)	9	(1.2)	1,199	937	(78.1)	262	(21.9)
60-64	638	352	(55.2)	125	(19.6)	143	(22.4)	18	(2.8)	1,083	837	(77.3)	246	(22.7)
65-69	429	148	(34.5)	105	(24.5)	168	(39.2)	8	(1.9)	913	606	(66.4)	307	(33.6)
70-74	175	51	(29.1)	40	(22.9)	83	(47.4)	1	(0.6)	429	234	(54.5)	195	(45.5)
75-79	72	9	(12.5)	14	(19.4)	45	(62.5)	4	(5.6)	153	80	(52.3)	73	(47.7)
80-84	18	3	(16.7)	7	(38.9)	8	(44.4)	.	.	68	25	(36.8)	43	(63.2)
85-	7	1	(14.3)	1	(14.3)	5	(71.4)	.	.	20	4	(20.0)	16	(80.0)
全年齢	9,286	4,527	(48.8)	1,761	(19.0)	2,822	(30.4)	176	(1.9)	13,638	8,960	(65.7)	4,678	(34.3)


10. 特発性血小板減少性紫斑病

図1-5-1-10 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




10. 特発性血小板減少性紫斑病

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



▨ 5年継続 □ 継続なし

1 1. 結節性動脈周囲炎

表1-5-1-11 84、88、92年度受給者の各調査年度までの継続状況、性・年齢階級別


性・年齢階級	1984年度 受給者数 (100%)	13年継続		8年継続		4年継続		継続なし		中止後再開	
		84889297		848892×		8488××		84××		8488×97 84×9297 84×92× 84××97	
男 0-4	105	3 (2.9)	.	.	.	7 (6.7)	95 (90.5)
5-9	34	1 (2.9)	2 (5.9)	2 (5.9)	2 (5.9)	28 (82.4)	1 (2.9)
10-14	15	2 (13.3)	3 (20.0)	.	.	8 (53.3)	2 (13.3)
15-19	10	1 (10.0)	1 (10.0)	2 (20.0)	2 (20.0)	6 (60.0)
20-24	3	3 (100.0)
25-29	8	2 (25.0)	1 (12.5)	3 (37.5)	1 (12.5)	1 (12.5)	1 (12.5)
30-34	15	6 (40.0)	1 (6.7)	1 (6.7)	6 (40.0)	1 (6.7)	1 (6.7)
35-39	28	16 (57.1)	2 (7.1)	2 (7.1)	7 (25.0)	1 (3.6)	1 (3.6)
40-44	20	6 (30.0)	5 (25.0)	1 (5.0)	6 (30.0)	2 (10.0)	2 (10.0)
45-49	20	8 (40.0)	4 (20.0)	2 (10.0)	5 (25.0)	1 (5.0)	1 (5.0)
50-54	29	14 (48.3)	2 (6.9)	5 (17.2)	7 (24.1)	1 (3.4)	1 (3.4)
55-59	41	9 (22.0)	7 (17.1)	10 (24.4)	14 (34.1)	1 (2.4)	1 (2.4)
60-64	34	2 (5.9)	6 (17.6)	7 (20.6)	16 (47.1)	3 (8.8)	3 (8.8)
65-69	25	1 (4.0)	4 (16.0)	2 (8.0)	17 (68.0)	1 (4.0)	1 (4.0)
70-74	2	.	1 (50.0)	.	1 (50.0)
75-79
80-84
85-
全年齢	389	71 (18.3)	39 (10.0)	44 (11.3)	220 (56.6)	15 (3.9)	15 (3.9)
女 0-4	61	2 (3.3)	1 (1.6)	1 (1.6)	57 (93.4)
5-9	39	1 (2.6)	1 (2.6)	2 (5.1)	35 (89.7)
10-14	4	1 (25.0)	.	.	3 (75.0)
15-19	9	4 (44.4)	1 (11.1)	.	4 (44.4)
20-24	10	6 (60.0)	1 (10.0)	1 (10.0)	.	2 (20.0)	2 (20.0)
25-29	13	4 (30.8)	1 (7.7)	3 (23.1)	4 (30.8)	1 (7.7)	1 (7.7)
30-34	18	9 (50.0)	1 (5.6)	2 (11.1)	6 (33.3)
35-39	28	16 (57.1)	2 (7.1)	2 (7.1)	6 (21.4)	2 (7.1)	2 (7.1)
40-44	43	23 (53.5)	3 (7.0)	7 (16.3)	8 (18.6)	2 (4.7)	2 (4.7)
45-49	49	30 (61.2)	6 (12.2)	2 (4.1)	11 (22.4)
50-54	58	30 (51.7)	10 (17.2)	4 (6.9)	12 (20.7)	2 (3.4)	2 (3.4)
55-59	57	22 (38.6)	10 (17.5)	7 (12.3)	18 (31.6)
60-64	52	16 (30.8)	11 (21.2)	7 (13.5)	16 (30.8)	2 (3.8)	2 (3.8)
65-69	37	6 (16.2)	5 (13.5)	6 (16.2)	19 (51.4)	1 (2.7)	1 (2.7)
70-74	3	.	.	.	2 (66.7)	1 (33.3)	1 (33.3)
75-79	1	.	.	1 (100.0)
80-84
85-
全年齢	482	170 (35.3)	53 (11.0)	45 (9.3)	204 (41.7)	13 (2.7)	13 (2.7)

1 1. 結節性動脈周囲炎

	1988年度				1992年度			
	受給者数 (100%)	9年継続 889297	4年継続 8892×	継続なし 88×	中止後再開 88×97	受給者数 (100%)	5年継続 9297	継続なし 92×
男 0-4	2	1 (50.0)	1 (50.0)
5-9	16	5 (31.3)	2 (12.5)	9 (56.3)	.	5	3 (60.0)	2 (40.0)
10-14	6	1 (16.7)	3 (50.0)	2 (33.3)	.	13	8 (61.5)	5 (38.5)
15-19	15	4 (26.7)	6 (40.0)	4 (26.7)	1 (6.7)	15	8 (53.3)	7 (46.7)
20-24	16	7 (43.8)	2 (12.5)	7 (43.8)	.	16	10 (62.5)	6 (37.5)
25-29	15	8 (53.3)	2 (13.3)	3 (20.0)	2 (13.3)	15	10 (66.7)	5 (33.3)
30-34	17	10 (58.8)	2 (11.8)	4 (23.5)	1 (5.9)	26	17 (65.4)	9 (34.6)
35-39	24	16 (66.7)	3 (12.5)	4 (16.7)	1 (4.2)	26	20 (76.9)	6 (23.1)
40-44	50	33 (66.0)	7 (14.0)	10 (20.0)	.	59	43 (72.9)	16 (27.1)
45-49	35	19 (54.3)	10 (28.6)	6 (17.1)	.	56	45 (80.4)	11 (19.6)
50-54	53	27 (50.9)	9 (17.0)	17 (32.1)	.	72	49 (68.1)	23 (31.9)
55-59	71	40 (56.3)	15 (21.1)	16 (22.5)	.	74	53 (71.6)	21 (28.4)
60-64	78	23 (29.5)	20 (25.6)	33 (42.3)	2 (2.6)	122	82 (67.2)	40 (32.8)
65-69	46	10 (21.7)	15 (32.6)	21 (45.7)	.	106	53 (50.0)	53 (50.0)
70-74	16	4 (25.0)	6 (37.5)	6 (37.5)	.	40	16 (40.0)	24 (60.0)
75-79	1	.	.	1 (100.0)	.	17	7 (41.2)	10 (58.8)
80-84	4	.	4 (100.0)
85-	1	.	.	1 (100.0)
全年齢	462	208 (45.0)	103 (22.3)	144 (31.2)	7 (1.5)	666	424 (63.7)	242 (36.3)
女 0-4	5	2 (40.0)	.	3 (60.0)	.	1	1 (100.0)	.
5-9	9	3 (33.3)	2 (22.2)	4 (44.4)	.	4	3 (75.0)	1 (25.0)
10-14	6	2 (33.3)	1 (16.7)	3 (50.0)	.	11	8 (72.7)	3 (27.3)
15-19	9	2 (22.2)	1 (11.1)	6 (66.7)	.	10	7 (70.0)	3 (30.0)
20-24	18	10 (55.6)	2 (11.1)	5 (27.8)	1 (5.6)	19	14 (73.7)	5 (26.3)
25-29	26	14 (53.8)	6 (23.1)	6 (23.1)	.	33	28 (84.8)	5 (15.2)
30-34	24	16 (66.7)	2 (8.3)	6 (25.0)	.	45	36 (80.0)	9 (20.0)
35-39	35	20 (57.1)	6 (17.1)	9 (25.7)	.	41	28 (68.3)	13 (31.7)
40-44	51	34 (66.7)	5 (9.8)	11 (21.6)	1 (2.0)	59	45 (76.3)	14 (23.7)
45-49	63	43 (68.3)	11 (17.5)	8 (12.7)	1 (1.6)	81	67 (82.7)	14 (17.3)
50-54	87	59 (67.8)	14 (16.1)	13 (14.9)	1 (1.1)	94	75 (79.8)	19 (20.2)
55-59	86	55 (64.0)	21 (24.4)	10 (11.6)	.	116	90 (77.6)	26 (22.4)
60-64	90	40 (44.4)	24 (26.7)	22 (24.4)	4 (4.4)	154	97 (63.0)	57 (37.0)
65-69	76	31 (40.8)	20 (26.3)	22 (28.9)	3 (3.9)	110	58 (52.7)	52 (47.3)
70-74	29	9 (31.0)	5 (17.2)	14 (48.3)	1 (3.4)	82	49 (59.8)	33 (40.2)
75-79	2	.	.	2 (100.0)	.	24	15 (62.5)	9 (37.5)
80-84	2	.	.	2 (100.0)	.	1	.	1 (100.0)
85-
全年齢	618	340 (55.0)	120 (19.4)	146 (23.6)	12 (1.9)	885	621 (70.2)	264 (29.8)


1 1 . 結節性動脈周囲炎

図1-5-1-1 1 84、88、92年度受給者の各調査年度までの継続状況，性・年齢階級別




11. 結節性動脈周囲炎

1988年度受給者 継続状況



■ 9年継続 □ 4年継続 □ 継続なし ▨ 中止後再開

1992年度受給者 継続状況



■ 5年継続 □ 継続なし