



4. 全身性エリテマトーデス

1992年度受給者 受給状況



■ 84年度受給 □ 88年度受給 □ 過去の受給なし

1988年度受給者 受給状況



■ 84年度受給 □ 過去の受給なし

5. スモン

表2-5-1-5 97、92、88年度受給者の過去の各調査年度における受給状況、性・年齢階級別


性・年齢階級	1997年度 受給者数 (100%)	84年度受給		88年度受給		92年度受給		過去の受給なし	
		84? ? 97		×88? 97		××9297		×××97	
男 0-4
5-9
10-14
15-19
20-24
25-29
30-34	1	.	.	1 (100.0)
35-39	2	1 (50.0)	1 (50.0)	.
40-44	5	2 (40.0)	.	.	.	1 (20.0)	.	2 (40.0)	.
45-49	8	2 (25.0)	.	3 (37.5)	.	1 (12.5)	.	2 (25.0)	.
50-54	19	10 (52.6)	.	2 (10.5)	.	2 (10.5)	.	5 (26.3)	.
55-59	34	14 (41.2)	.	10 (29.4)	.	5 (14.7)	.	5 (14.7)	.
60-64	75	37 (49.3)	.	10 (13.3)	.	6 (8.0)	.	22 (29.3)	.
65-69	104	54 (51.9)	.	19 (18.3)	.	10 (9.6)	.	21 (20.2)	.
70-74	99	62 (62.6)	.	21 (21.2)	.	2 (2.0)	.	14 (14.1)	.
75-79	55	34 (61.8)	.	5 (9.1)	.	4 (7.3)	.	12 (21.8)	.
80-84	37	24 (64.9)	.	5 (13.5)	.	1 (2.7)	.	7 (18.9)	.
85-	21	7 (33.3)	.	5 (23.8)	.	3 (14.3)	.	6 (28.6)	.
全年齢	460	247 (53.7)	.	81 (17.6)	.	35 (7.6)	.	97 (21.1)	.
女 0-4
5-9
10-14
15-19
20-24
25-29
30-34	4	3 (75.0)	1 (25.0)	.
35-39	1	1 (100.0)
40-44	4	1 (25.0)	.	2 (50.0)	.	.	.	1 (25.0)	.
45-49	20	13 (65.0)	.	3 (15.0)	.	2 (10.0)	.	2 (10.0)	.
50-54	57	30 (52.6)	.	13 (22.8)	.	5 (8.8)	.	9 (15.8)	.
55-59	117	71 (60.7)	.	16 (13.7)	.	10 (8.5)	.	20 (17.1)	.
60-64	185	134 (72.4)	.	23 (12.4)	.	10 (5.4)	.	18 (9.7)	.
65-69	284	198 (69.7)	.	33 (11.6)	.	13 (4.6)	.	40 (14.1)	.
70-74	289	214 (74.0)	.	33 (11.4)	.	14 (4.8)	.	28 (9.7)	.
75-79	278	225 (80.9)	.	22 (7.9)	.	10 (3.6)	.	21 (7.6)	.
80-84	178	111 (62.4)	.	15 (8.4)	.	4 (2.2)	.	48 (27.0)	.
85-	151	52 (34.4)	.	29 (19.2)	.	15 (9.9)	.	55 (36.4)	.
全年齢	1,568	1,053 (67.2)	.	189 (12.1)	.	83 (5.3)	.	243 (15.5)	.

5. スモン

性・年齢	1992年度 受給者数 (100%)	84年度受給 84?92	88年度受給 ×8892	過去の受給なし ××92	1988年度 受給者数 (100%)	84年度受給 8488	過去の受給なし ×88
	男						
0-4
5-9
10-14
15-19
20-24	1	.	1 (100.0)
25-29	2	.	1 (50.0)	1 (50.0)	1	1 (100.0)	.
30-34	1	1 (100.0)	.	.	2	2 (100.0)	.
35-39	5	3 (60.0)	.	2 (40.0)	7	4 (57.1)	3 (42.9)
40-44	9	5 (55.6)	3 (33.3)	1 (11.1)	12	9 (75.0)	3 (25.0)
45-49	13	6 (46.2)	4 (30.8)	3 (23.1)	35	21 (60.0)	14 (40.0)
50-54	32	18 (56.3)	8 (25.0)	6 (18.8)	51	40 (78.4)	11 (21.6)
55-59	60	39 (65.0)	12 (20.0)	9 (15.0)	85	62 (72.9)	23 (27.1)
60-64	98	66 (67.3)	20 (20.4)	12 (12.2)	122	95 (77.9)	27 (22.1)
65-69	97	68 (70.1)	18 (18.6)	11 (11.3)	74	62 (83.8)	12 (16.2)
70-74	58	44 (75.9)	7 (12.1)	7 (12.1)	71	55 (77.5)	16 (22.5)
75-79	52	40 (76.9)	9 (17.3)	3 (5.8)	25	14 (56.0)	11 (44.0)
80-84	19	9 (47.4)	7 (36.8)	3 (15.8)	14	6 (42.9)	8 (57.1)
85-	8	5 (62.5)	2 (25.0)	1 (12.5)	4	2 (50.0)	2 (50.0)
全年齢	454	304 (67.0)	91 (20.0)	59 (13.0)	504	373 (74.0)	131 (26.0)
女							
0-4
5-9
10-14
15-19
20-24	4	3 (75.0)	1 (25.0)
25-29	3	3 (100.0)	.	.	2	1 (50.0)	1 (50.0)
30-34	2	2 (100.0)	.	.	2	1 (50.0)	1 (50.0)
35-39	2	.	2 (100.0)	.	20	14 (70.0)	6 (30.0)
40-44	27	19 (70.4)	6 (22.2)	2 (7.4)	42	26 (61.9)	16 (38.1)
45-49	53	31 (58.5)	15 (28.3)	7 (13.2)	84	67 (79.8)	17 (20.2)
50-54	99	73 (73.7)	13 (13.1)	13 (13.1)	173	147 (85.0)	26 (15.0)
55-59	179	143 (79.9)	22 (12.3)	14 (7.8)	262	216 (82.4)	46 (17.6)
60-64	295	232 (78.6)	40 (13.6)	23 (7.8)	362	303 (83.7)	59 (16.3)
65-69	320	249 (77.8)	36 (11.3)	35 (10.9)	334	290 (86.8)	44 (13.2)
70-74	253	218 (86.2)	21 (8.3)	14 (5.5)	232	188 (81.0)	44 (19.0)
75-79	173	133 (76.9)	24 (13.9)	16 (9.2)	131	84 (64.1)	47 (35.9)
80-84	105	54 (51.4)	30 (28.6)	21 (20.0)	54	28 (51.9)	26 (48.1)
85-	44	19 (43.2)	14 (31.8)	11 (25.0)	20	7 (35.0)	13 (65.0)
全年齢	1,555	1,176 (75.6)	223 (14.3)	156 (10.0)	1,722	1,375 (79.8)	347 (20.2)


5. スモン

図2-5-1-5 97、92、88年度受給者の過去の各調査年度における受給状況，性・年齢階級別




5. スモン

1992年度受給者 受給状況



■ 84年度受給 □ 88年度受給 □ 過去の受給なし

1988年度受給者 受給状況



■ 84年度受給 □ 過去の受給なし

6. 再生不良性貧血

表2-5-1-6 97、92、88年度受給者の過去の各調査年度における受給状況、性・年齢階級別


性・年齢階級	1997年度 受給者数 (100%)	84年度受給		88年度受給		92年度受給		過去の受給なし	
		84? ? 97		×88? 97		××9297		×××97	
男 0-4	42	42	(100.0)
5-9	100	.	.	1	(1.0)	21	(21.0)	78	(78.0)
10-14	181	3	(1.7)	22	(12.2)	50	(27.6)	106	(58.6)
15-19	270	34	(12.6)	40	(14.8)	75	(27.8)	121	(44.8)
20-24	326	58	(17.8)	77	(23.6)	75	(23.0)	116	(35.6)
25-29	299	55	(18.4)	51	(17.1)	57	(19.1)	136	(45.5)
30-34	232	45	(19.4)	42	(18.1)	60	(25.9)	85	(36.6)
35-39	208	36	(17.3)	40	(19.2)	62	(29.8)	70	(33.7)
40-44	197	33	(16.8)	39	(19.8)	40	(20.3)	85	(43.1)
45-49	213	38	(17.8)	38	(17.8)	48	(22.5)	89	(41.8)
50-54	226	38	(16.8)	35	(15.5)	52	(23.0)	101	(44.7)
55-59	276	25	(9.1)	45	(16.3)	57	(20.7)	149	(54.0)
60-64	315	40	(12.7)	42	(13.3)	64	(20.3)	169	(53.7)
65-69	385	39	(10.1)	42	(10.9)	74	(19.2)	230	(59.7)
70-74	321	30	(9.3)	33	(10.3)	60	(18.7)	198	(61.7)
75-79	183	26	(14.2)	22	(12.0)	36	(19.7)	99	(54.1)
80-84	91	14	(15.4)	5	(5.5)	15	(16.5)	57	(62.6)
85-	40	2	(5.0)	3	(7.5)	5	(12.5)	30	(75.0)
全年齢	3,905	516	(13.2)	577	(14.8)	851	(21.8)	1,961	(50.2)
女 0-4	26	26	(100.0)
5-9	109	21	(19.3)	88	(80.7)
10-14	178	2	(1.1)	12	(6.7)	55	(30.9)	109	(61.2)
15-19	229	21	(9.2)	37	(16.2)	66	(28.8)	105	(45.9)
20-24	307	57	(18.6)	59	(19.2)	65	(21.2)	126	(41.0)
25-29	328	68	(20.7)	56	(17.1)	58	(17.7)	146	(44.5)
30-34	302	70	(23.2)	61	(20.2)	62	(20.5)	109	(36.1)
35-39	281	59	(21.0)	53	(18.9)	71	(25.3)	98	(34.9)
40-44	299	84	(28.1)	56	(18.7)	56	(18.7)	103	(34.4)
45-49	396	118	(29.8)	59	(14.9)	81	(20.5)	138	(34.8)
50-54	371	99	(26.7)	38	(10.2)	73	(19.7)	161	(43.4)
55-59	471	93	(19.7)	76	(16.1)	76	(16.1)	226	(48.0)
60-64	525	97	(18.5)	60	(11.4)	107	(20.4)	261	(49.7)
65-69	607	126	(20.8)	81	(13.3)	119	(19.6)	281	(46.3)
70-74	592	129	(21.8)	84	(14.2)	107	(18.1)	272	(45.9)
75-79	330	88	(26.7)	41	(12.4)	52	(15.8)	149	(45.2)
80-84	186	39	(21.0)	22	(11.8)	20	(10.8)	105	(56.5)
85-	66	3	(4.5)	7	(10.6)	12	(18.2)	44	(66.7)
全年齢	5,603	1,153	(20.6)	802	(14.3)	1,101	(19.7)	2,547	(45.5)

6. 再生不良性貧血

性・年齢	1992年度 受給者数 (100%)	84年度受給			88年度受給			過去の受給なし			1988年度 受給者数 (100%)	84年度受給			過去の受給なし		
		84	?	92	×	8892	×	×	92	8488		×	88				
男 0-4	51				1	(2.0)	50	(98.0)		58	1	(1.7)	57	(98.3)			
5-9	123	6	(4.9)	40	(32.5)	77	(62.6)		137	48	(35.0)	89	(65.0)				
10-14	211	42	(19.9)	61	(28.9)	108	(51.2)		261	104	(39.8)	157	(60.2)				
15-19	335	80	(23.9)	126	(37.6)	129	(38.5)		285	153	(53.7)	132	(46.3)				
20-24	277	87	(31.4)	72	(26.0)	118	(42.6)		190	101	(53.2)	89	(46.8)				
25-29	216	67	(31.0)	54	(25.0)	95	(44.0)		170	77	(45.3)	93	(54.7)				
30-34	204	55	(27.0)	57	(27.9)	92	(45.1)		143	61	(42.7)	82	(57.3)				
35-39	164	47	(28.7)	60	(36.6)	57	(34.8)		167	77	(46.1)	90	(53.9)				
40-44	213	60	(28.2)	62	(29.1)	91	(42.7)		144	62	(43.1)	82	(56.9)				
45-49	179	43	(24.0)	50	(27.9)	86	(48.0)		145	60	(41.4)	85	(58.6)				
50-54	193	39	(20.2)	62	(32.1)	92	(47.7)		169	61	(36.1)	108	(63.9)				
55-59	245	56	(22.9)	61	(24.9)	128	(52.2)		184	71	(38.6)	113	(61.4)				
60-64	249	44	(17.7)	69	(27.7)	136	(54.6)		221	63	(28.5)	158	(71.5)				
65-69	275	44	(16.0)	78	(28.4)	153	(55.6)		202	75	(37.1)	127	(62.9)				
70-74	161	32	(19.9)	34	(21.1)	95	(59.0)		95	41	(43.2)	54	(56.8)				
75-79	70	21	(30.0)	11	(15.7)	38	(54.3)		47	7	(14.9)	40	(85.1)				
80-84	36	4	(11.1)	10	(27.8)	22	(61.1)		13			13	(100.0)				
85-	9			1	(11.1)	8	(88.9)		4			4	(100.0)				
全年齢	3,211	727	(22.6)	909	(28.3)	1,575	(49.1)		2,635	1,062	(40.3)	1,573	(59.7)				
女 0-4	37			1	(2.7)	36	(97.3)		47	2	(4.3)	45	(95.7)				
5-9	123	5	(4.1)	27	(22.0)	91	(74.0)		124	37	(29.8)	87	(70.2)				
10-14	184	27	(14.7)	55	(29.9)	102	(55.4)		216	101	(46.8)	115	(53.2)				
15-19	266	78	(29.3)	92	(34.6)	96	(36.1)		250	129	(51.6)	121	(48.4)				
20-24	242	88	(36.4)	62	(25.6)	92	(38.0)		195	86	(44.1)	109	(55.9)				
25-29	256	76	(29.7)	83	(32.4)	97	(37.9)		189	96	(50.8)	93	(49.2)				
30-34	240	81	(33.8)	59	(24.6)	100	(41.7)		240	129	(53.8)	111	(46.3)				
35-39	258	102	(39.5)	72	(27.9)	84	(32.6)		252	150	(59.5)	102	(40.5)				
40-44	323	130	(40.2)	76	(23.5)	117	(36.2)		225	126	(56.0)	99	(44.0)				
45-49	276	111	(40.2)	59	(21.4)	106	(38.4)		274	148	(54.0)	126	(46.0)				
50-54	334	121	(36.2)	91	(27.2)	122	(36.5)		278	150	(54.0)	128	(46.0)				
55-59	395	134	(33.9)	93	(23.5)	168	(42.5)		374	208	(55.6)	166	(44.4)				
60-64	461	158	(34.3)	104	(22.6)	199	(43.2)		413	215	(52.1)	198	(47.9)				
65-69	515	182	(35.3)	129	(25.0)	204	(39.6)		389	221	(56.8)	168	(43.2)				
70-74	318	134	(42.1)	76	(23.9)	108	(34.0)		187	104	(55.6)	83	(44.4)				
75-79	136	59	(43.4)	27	(19.9)	50	(36.8)		56	12	(21.4)	44	(78.6)				
80-84	47	6	(12.8)	12	(25.5)	29	(61.7)		20			20	(100.0)				
85-	20			7	(35.0)	13	(65.0)		5			5	(100.0)				
全年齢	4,431	1,492	(33.7)	1,125	(25.4)	1,814	(40.9)		3,734	1,914	(51.3)	1,820	(48.7)				


6. 再生不良性貧血

図2-5-1-6 97、92、88年度受給者の過去の各調査年度における受給状況，性・年齢階級別




6. 再生不良性貧血

1992年度受給者 受給状況



■ 84年度受給 □ 88年度受給 □ 過去の受給なし

1988年度受給者 受給状況



■ 84年度受給 □ 過去の受給なし

7. サルコイドーシス

表 2-5-1-7 97、92、88年度受給者の過去の各調査年度+B42受給状況，性・年齢階級別


性・年齢階級	1997年度 受給者数 (100%)	84年度受給		88年度受給		92年度受給		過去の受給なし	
		84? ?97		×88?97		××9297		×××97	
男 0-4
5-9	3	3	(100.0)
10-14	9	.	.	1	(11.1)	.	.	8	(88.9)
15-19	25	1	(4.0)	.	.	6	(24.0)	18	(72.0)
20-24	284	.	.	4	(1.4)	5	(1.8)	275	(96.8)
25-29	816	4	(0.5)	12	(1.5)	79	(9.7)	721	(88.4)
30-34	775	11	(1.4)	65	(8.4)	213	(27.5)	486	(62.7)
35-39	531	32	(6.0)	95	(17.9)	124	(23.4)	280	(52.7)
40-44	363	40	(11.0)	74	(20.4)	75	(20.7)	174	(47.9)
45-49	376	42	(11.2)	60	(16.0)	101	(26.9)	173	(46.0)
50-54	347	40	(11.5)	39	(11.2)	70	(20.2)	198	(57.1)
55-59	381	28	(7.3)	51	(13.4)	89	(23.4)	213	(55.9)
60-64	441	36	(8.2)	60	(13.6)	109	(24.7)	236	(53.5)
65-69	393	25	(6.4)	59	(15.0)	103	(26.2)	206	(52.4)
70-74	225	18	(8.0)	40	(17.8)	50	(22.2)	117	(52.0)
75-79	99	18	(18.2)	19	(19.2)	20	(20.2)	42	(42.4)
80-84	15	3	(20.0)	3	(20.0)	4	(26.7)	5	(33.3)
85-	4	1	(25.0)	3	(75.0)
全年齢	5,087	298	(5.9)	582	(11.4)	1,049	(20.6)	3,158	(62.1)
女 0-4
5-9	4	2	(50.0)	2	(50.0)
10-14	13	.	.	1	(7.7)	.	.	12	(92.3)
15-19	36	1	(2.8)	3	(8.3)	8	(22.2)	24	(66.7)
20-24	136	4	(2.9)	5	(3.7)	19	(14.0)	108	(79.4)
25-29	488	11	(2.3)	7	(1.4)	60	(12.3)	410	(84.0)
30-34	633	19	(3.0)	53	(8.4)	157	(24.8)	404	(63.8)
35-39	634	55	(8.7)	107	(16.9)	138	(21.8)	334	(52.7)
40-44	574	87	(15.2)	107	(18.6)	128	(22.3)	252	(43.9)
45-49	726	125	(17.2)	95	(13.1)	133	(18.3)	373	(51.4)
50-54	1,003	108	(10.8)	96	(9.6)	161	(16.1)	638	(63.6)
55-59	1,426	105	(7.4)	125	(8.8)	344	(24.1)	852	(59.7)
60-64	1,834	180	(9.8)	253	(13.8)	476	(26.0)	925	(50.4)
65-69	1,785	269	(15.1)	283	(15.9)	409	(22.9)	824	(46.2)
70-74	1,059	236	(22.3)	193	(18.2)	238	(22.5)	392	(37.0)
75-79	438	135	(30.8)	74	(16.9)	82	(18.7)	147	(33.6)
80-84	134	37	(27.6)	21	(15.7)	26	(19.4)	50	(37.3)
85-	29	4	(13.8)	4	(13.8)	8	(27.6)	13	(44.8)
全年齢	10,952	1,376	(12.6)	1,427	(13.0)	2,389	(21.8)	5,760	(52.6)

7. サルコイドーシス

性・年齢	1992年度 受給者数 (100%)	1988年度受給			1988年度受給者数 (100%)		
		84年度受給 84?92	88年度受給 ×8892	過去の受給なし ××92	84年度受給 8488	過去の受給なし ×88	
男 0-4	1	.	.	1 (100.0)	1	.	1 (100.0)
5-9	3	.	1 (33.3)	2 (66.7)	2	1 (50.0)	1 (50.0)
10-14	13	1 (7.7)	.	12 (92.3)	13	2 (15.4)	11 (84.6)
15-19	36	4 (11.1)	10 (27.8)	22 (61.1)	41	11 (26.8)	30 (73.2)
20-24	238	8 (3.4)	18 (7.6)	212 (89.1)	224	27 (12.1)	197 (87.9)
25-29	532	14 (2.6)	113 (21.2)	405 (76.1)	335	54 (16.1)	281 (83.9)
30-34	400	46 (11.5)	144 (36.0)	210 (52.5)	231	71 (30.7)	160 (69.3)
35-39	277	59 (21.3)	94 (33.9)	124 (44.8)	181	66 (36.5)	115 (63.5)
40-44	258	50 (19.4)	69 (26.7)	139 (53.9)	115	47 (40.9)	68 (59.1)
45-49	200	40 (20.0)	53 (26.5)	107 (53.5)	138	48 (34.8)	90 (65.2)
50-54	224	35 (15.6)	64 (28.6)	125 (55.8)	146	49 (33.6)	97 (66.4)
55-59	279	49 (17.6)	82 (29.4)	148 (53.0)	151	47 (31.1)	104 (68.9)
60-64	253	32 (12.6)	70 (27.7)	151 (59.7)	118	35 (29.7)	83 (70.3)
65-69	159	24 (15.1)	52 (32.7)	83 (52.2)	88	37 (42.0)	51 (58.0)
70-74	79	24 (30.4)	25 (31.6)	30 (38.0)	27	11 (40.7)	16 (59.3)
75-79	20	6 (30.0)	8 (40.0)	6 (30.0)	4	.	4 (100.0)
80-84	2	.	1 (50.0)	1 (50.0)	.	.	.
85-
全年齢	2,974	392 (13.2)	804 (27.0)	1,778 (59.8)	1,815	506 (27.9)	1,309 (72.1)
女 0-4	2	.	.	2 (100.0)	1	.	1 (100.0)
5-9	2	.	1 (50.0)	1 (50.0)	5	.	5 (100.0)
10-14	19	1 (5.3)	5 (26.3)	13 (68.4)	22	6 (27.3)	16 (72.7)
15-19	46	4 (8.7)	9 (19.6)	33 (71.7)	49	17 (34.7)	32 (65.3)
20-24	156	16 (10.3)	22 (14.1)	118 (75.6)	143	29 (20.3)	114 (79.7)
25-29	370	23 (6.2)	76 (20.5)	271 (73.2)	300	83 (27.7)	217 (72.3)
30-34	433	69 (15.9)	137 (31.6)	227 (52.4)	329	121 (36.8)	208 (63.2)
35-39	435	108 (24.8)	135 (31.0)	192 (44.1)	323	162 (50.2)	161 (49.8)
40-44	467	147 (31.5)	118 (25.3)	202 (43.3)	306	145 (47.4)	161 (52.6)
45-49	459	117 (25.5)	125 (27.2)	217 (47.3)	328	137 (41.8)	191 (58.2)
50-54	717	123 (17.2)	163 (22.7)	431 (60.1)	578	228 (39.4)	350 (60.6)
55-59	1,094	220 (20.1)	302 (27.6)	572 (52.3)	713	346 (48.5)	367 (51.5)
60-64	1,131	328 (29.0)	325 (28.7)	478 (42.3)	686	347 (50.6)	339 (49.4)
65-69	889	302 (34.0)	246 (27.7)	341 (38.4)	442	267 (60.4)	175 (39.6)
70-74	405	171 (42.2)	104 (25.7)	130 (32.1)	158	90 (57.0)	68 (43.0)
75-79	121	54 (44.6)	27 (22.3)	40 (33.1)	35	16 (45.7)	19 (54.3)
80-84	23	2 (8.7)	7 (30.4)	14 (60.9)	7	.	7 (100.0)
85-	3	.	1 (33.3)	2 (66.7)	2	.	2 (100.0)
全年齢	6,772	1,685 (24.9)	1,803 (26.6)	3,284 (48.5)	4,427	1,994 (45.0)	2,433 (55.0)


7. サルコイドーシス

図2-5-1-7 97、92、88年度受給者の過去の各調査年度における受給状況、性・年齢階級別




7. サルコイドーシス

1992年度受給者 受給状況



■ 84年度受給 □ 88年度受給 ◻ 過去を受給なし

1988年度受給者 受給状況



■ 84年度受給 □ 88年度受給 ◻ 過去を受給なし

8. 筋萎縮性側索硬化症

表2-5-1-8 97、92、88年度受給者の過去の各調査年度における受給状況、性・年齢階級別


性・年齢階級	1997年度 受給者数 (100%)	84年度受給		88年度受給		92年度受給		過去の受給なし	
		84? ?97		×88?97		××9297		×××97	
男 0-4
5-9	2	1 (50.0)		1 (50.0)	
10-14	4	.	.	1 (25.0)		1 (25.0)		2 (50.0)	
15-19	3	1 (33.3)		2 (66.7)	
20-24	8	1 (12.5)		1 (12.5)		3 (37.5)		3 (37.5)	
25-29	13	2 (15.4)		1 (7.7)		1 (7.7)		9 (69.2)	
30-34	31	1 (3.2)		5 (16.1)		2 (6.5)		23 (74.2)	
35-39	43	3 (7.0)		5 (11.6)		2 (4.7)		33 (76.7)	
40-44	75	1 (1.3)		5 (6.7)		13 (17.3)		56 (74.7)	
45-49	216	4 (1.9)		16 (7.4)		27 (12.5)		169 (78.2)	
50-54	282	10 (3.5)		24 (8.5)		47 (16.7)		201 (71.3)	
55-59	451	17 (3.8)		31 (6.9)		65 (14.4)		338 (74.9)	
60-64	548	25 (4.6)		40 (7.3)		68 (12.4)		415 (75.7)	
65-69	559	13 (2.3)		30 (5.4)		65 (11.6)		451 (80.7)	
70-74	424	25 (5.9)		30 (7.1)		57 (13.4)		312 (73.6)	
75-79	216	13 (6.0)		9 (4.2)		16 (7.4)		178 (82.4)	
80-84	64	5 (7.8)		4 (6.3)		5 (7.8)		50 (78.1)	
85-	19	1 (5.3)		.		.		18 (94.7)	
全年齢	2,958	121 (4.1)		202 (6.8)		374 (12.6)		2,261 (76.4)	
女 0-4	1	1 (100.0)	
5-9	1	1 (100.0)	
10-14	2	.	.	1 (50.0)		1 (50.0)		.	
15-19	1	.	.	1 (100.0)		.		.	
20-24	6	6 (100.0)	
25-29	13	2 (15.4)		.		.		11 (84.6)	
30-34	9	1 (11.1)		1 (11.1)		1 (11.1)		6 (66.7)	
35-39	30	.		2 (6.7)		6 (20.0)		22 (73.3)	
40-44	66	3 (4.5)		3 (4.5)		12 (18.2)		48 (72.7)	
45-49	126	7 (5.6)		11 (8.7)		9 (7.1)		99 (78.6)	
50-54	161	10 (6.2)		9 (5.6)		19 (11.8)		123 (76.4)	
55-59	218	6 (2.8)		22 (10.1)		22 (10.1)		168 (77.1)	
60-64	288	9 (3.1)		16 (5.6)		32 (11.1)		231 (80.2)	
65-69	332	14 (4.2)		14 (4.2)		32 (9.6)		272 (81.9)	
70-74	301	8 (2.7)		12 (4.0)		29 (9.6)		252 (83.7)	
75-79	202	5 (2.5)		7 (3.5)		13 (6.4)		177 (87.6)	
80-84	62	3 (4.8)		2 (3.2)		2 (3.2)		55 (88.7)	
85-	7	.		.		.		7 (100.0)	
全年齢	1,826	68 (3.7)		101 (5.5)		178 (9.7)		1,479 (81.0)	

8. 筋萎縮性側索硬化症

性・年齢	1992年度 受給者数 (100%)	84年度受給		88年度受給		過去の受給なし		1988年度 受給者数 (100%)	84年度受給		過去の受給なし	
		84	92	×88	92	×92	84		88	84	88	
男 0-4	1	1	(100.0)	1	.	.	1	(100.0)
5-9	3	1	(33.3)	1	(33.3)	1	(33.3)	6	2	(33.3)	4	(66.7)
10-14	3	.	.	2	(66.7)	1	(33.3)	2	1	(50.0)	1	(50.0)
15-19	8	1	(12.5)	1	(12.5)	6	(75.0)	5	2	(40.0)	3	(60.0)
20-24	7	2	(28.6)	3	(42.9)	2	(28.6)	14	4	(28.6)	10	(71.4)
25-29	13	3	(23.1)	5	(38.5)	5	(38.5)	12	2	(16.7)	10	(83.3)
30-34	19	3	(15.8)	8	(42.1)	8	(42.1)	31	5	(16.1)	26	(83.9)
35-39	45	3	(6.7)	10	(22.2)	32	(71.1)	51	14	(27.5)	37	(72.5)
40-44	96	5	(5.2)	25	(26.0)	66	(68.8)	91	17	(18.7)	74	(81.3)
45-49	164	12	(7.3)	36	(22.0)	116	(70.7)	133	26	(19.5)	107	(80.5)
50-54	218	21	(9.6)	48	(22.0)	149	(68.3)	214	46	(21.5)	168	(78.5)
55-59	348	35	(10.1)	64	(18.4)	249	(71.6)	275	56	(20.4)	219	(79.6)
60-64	389	39	(10.0)	61	(15.7)	289	(74.3)	320	66	(20.6)	254	(79.4)
65-69	381	35	(9.2)	68	(17.8)	278	(73.0)	241	56	(23.2)	185	(76.8)
70-74	182	27	(14.8)	32	(17.6)	123	(67.6)	103	18	(17.5)	85	(82.5)
75-79	69	6	(8.7)	7	(10.1)	56	(81.2)	35	1	(2.9)	34	(97.1)
80-84	13	1	(7.7)	2	(15.4)	10	(76.9)	5	.	.	5	(100.0)
85-	3	3	(100.0)
全年齢	1,962	194	(9.9)	373	(19.0)	1,395	(71.1)	1,539	316	(20.5)	1,223	(79.5)
女 0-4	1	1	(100.0)	1	.	.	1	(100.0)
5-9	2	.	.	1	(50.0)	1	(50.0)	1	.	.	1	(100.0)
10-14	1	.	.	1	(100.0)	.	.	2	.	.	2	(100.0)
15-19	4	2	(50.0)	2	(50.0)
20-24	4	2	(50.0)	2	(50.0)	.	.	6	3	(50.0)	3	(50.0)
25-29	4	1	(25.0)	1	(25.0)	2	(50.0)	3	.	.	3	(100.0)
30-34	14	.	.	2	(14.3)	12	(85.7)	13	6	(46.2)	7	(53.8)
35-39	31	6	(19.4)	6	(19.4)	19	(61.3)	39	10	(25.6)	29	(74.4)
40-44	53	10	(18.9)	14	(26.4)	29	(54.7)	48	16	(33.3)	32	(66.7)
45-49	85	13	(15.3)	19	(22.4)	53	(62.4)	97	16	(16.5)	81	(83.5)
50-54	130	10	(7.7)	35	(26.9)	85	(65.4)	108	30	(27.8)	78	(72.2)
55-59	177	20	(11.3)	35	(19.8)	122	(68.9)	174	35	(20.1)	139	(79.9)
60-64	218	25	(11.5)	42	(19.3)	151	(69.3)	204	29	(14.2)	175	(85.8)
65-69	224	9	(4.0)	33	(14.7)	182	(81.3)	162	27	(16.7)	135	(83.3)
70-74	118	10	(8.5)	16	(13.6)	92	(78.0)	66	8	(12.1)	58	(87.9)
75-79	50	3	(6.0)	3	(6.0)	44	(88.0)	32	1	(3.1)	31	(96.9)
80-84	18	.	.	2	(11.1)	16	(88.9)	6	.	.	6	(100.0)
85-	1	1	(100.0)	2	.	.	2	(100.0)
全年齢	1,131	109	(9.6)	212	(18.7)	810	(71.6)	968	183	(18.9)	785	(81.1)


8. 筋萎縮性側索硬化症

図2-5-1-8 97、92、88年度受給者の過去の各調査年度における受給状況，性・年齢階級別




8. 筋萎縮性側索硬化症

1992年度受給者 受給状況



■ 84年度受給 □ 88年度受給 □ 過去の受給なし

1988年度受給者 受給状況



■ 84年度受給 □ 過去の受給なし

9. 強皮症・皮膚筋炎及び多発性筋炎

表2-5-1-9 97、92、88年度受給者の過去の各調査年度における受給状況、性・年齢階級別

性・年齢階級	1997年度 受給者数 (100%)	84年度受給		88年度受給		92年度受給		過去の受給なし	
		84??97		×88?97		××9297		×××97	
男 0-4	8	8	(100.0)
5-9	34	5	(14.7)	29	(85.3)
10-14	67	.	.	6	(9.0)	26	(38.8)	35	(52.2)
15-19	100	12	(12.0)	21	(21.0)	19	(19.0)	48	(48.0)
20-24	96	19	(19.8)	12	(12.5)	22	(22.9)	43	(44.8)
25-29	119	14	(11.8)	12	(10.1)	29	(24.4)	64	(53.8)
30-34	116	14	(12.1)	11	(9.5)	23	(19.8)	68	(58.6)
35-39	167	12	(7.2)	21	(12.6)	39	(23.4)	95	(56.9)
40-44	249	16	(6.4)	29	(11.6)	54	(21.7)	150	(60.2)
45-49	412	22	(5.3)	57	(13.8)	99	(24.0)	234	(56.8)
50-54	493	44	(8.9)	81	(16.4)	104	(21.1)	264	(53.5)
55-59	604	48	(7.9)	82	(13.6)	116	(19.2)	358	(59.3)
60-64	739	66	(8.9)	88	(11.9)	161	(21.8)	424	(57.4)
65-69	706	61	(8.6)	100	(14.2)	165	(23.4)	380	(53.8)
70-74	444	40	(9.0)	71	(16.0)	96	(21.6)	237	(53.4)
75-79	164	23	(14.0)	30	(18.3)	29	(17.7)	82	(50.0)
80-84	66	9	(13.6)	6	(9.1)	14	(21.2)	37	(56.1)
85-	8	.	.	1	(12.5)	1	(12.5)	6	(75.0)
全年齢	4,592	400	(8.7)	628	(13.7)	1,002	(21.8)	2,562	(55.8)
女 0-4	7	7	(100.0)
5-9	51	7	(13.7)	44	(86.3)
10-14	97	1	(1.0)	12	(12.4)	24	(24.7)	60	(61.9)
15-19	161	12	(7.5)	30	(18.6)	43	(26.7)	76	(47.2)
20-24	230	19	(8.3)	34	(14.8)	48	(20.9)	129	(56.1)
25-29	376	39	(10.4)	39	(10.4)	79	(21.0)	219	(58.2)
30-34	459	46	(10.0)	59	(12.9)	114	(24.8)	240	(52.3)
35-39	679	70	(10.3)	111	(16.3)	145	(21.4)	353	(52.0)
40-44	1,105	132	(11.9)	139	(12.6)	262	(23.7)	572	(51.8)
45-49	2,250	298	(13.2)	290	(12.9)	508	(22.6)	1,154	(51.3)
50-54	2,692	381	(14.2)	410	(15.2)	590	(21.9)	1,311	(48.7)
55-59	3,275	537	(16.4)	506	(15.5)	753	(23.0)	1,479	(45.2)
60-64	3,301	618	(18.7)	532	(16.1)	759	(23.0)	1,392	(42.2)
65-69	2,697	537	(19.9)	452	(16.8)	585	(21.7)	1,123	(41.6)
70-74	1,670	391	(23.4)	275	(16.5)	343	(20.5)	661	(39.6)
75-79	659	168	(25.5)	100	(15.2)	91	(13.8)	300	(45.5)
80-84	215	37	(17.2)	19	(8.8)	32	(14.9)	127	(59.1)
85-	42	1	(2.4)	7	(16.7)	8	(19.0)	26	(61.9)
全年齢	19,966	3,287	(16.5)	3,015	(15.1)	4,391	(22.0)	9,273	(46.4)

9. 強皮症・皮膚筋炎及び多発性筋炎

性・年齢	1992年度 受給者数 (100%)	84年度受給		88年度受給		過去の受給なし		1988年度 受給者数 (100%)	84年度受給		過去の受給なし	
		84	792	×	8892	×	×92		8488	×	88	
男 0-4	9	9	(100.0)	16	.	.	16	(100.0)
5-9	49	.	.	13	(26.5)	36	(73.5)	53	13	(24.5)	40	(75.5)
10-14	73	13	(17.8)	32	(43.8)	28	(38.4)	84	35	(41.7)	49	(58.3)
15-19	100	25	(25.0)	32	(32.0)	43	(43.0)	74	38	(51.4)	36	(48.6)
20-24	84	19	(22.6)	21	(25.0)	44	(52.4)	53	24	(45.3)	29	(54.7)
25-29	67	16	(23.9)	16	(23.9)	35	(52.2)	48	17	(35.4)	31	(64.6)
30-34	97	14	(14.4)	21	(21.6)	62	(63.9)	74	23	(31.1)	51	(68.9)
35-39	155	20	(12.9)	41	(26.5)	94	(60.6)	138	49	(35.5)	89	(64.5)
40-44	253	36	(14.2)	66	(26.1)	151	(59.7)	181	56	(30.9)	125	(69.1)
45-49	292	49	(16.8)	97	(33.2)	146	(50.0)	228	73	(32.0)	155	(68.0)
50-54	361	63	(17.5)	108	(29.9)	190	(52.6)	252	84	(33.3)	168	(66.7)
55-59	446	75	(16.8)	114	(25.6)	257	(57.6)	332	109	(32.8)	223	(67.2)
60-64	492	88	(17.9)	148	(30.1)	256	(52.0)	297	94	(31.6)	203	(68.4)
65-69	359	64	(17.8)	100	(27.9)	195	(54.3)	181	72	(39.8)	109	(60.2)
70-74	163	36	(22.1)	48	(29.4)	79	(48.5)	89	40	(44.9)	49	(55.1)
75-79	67	17	(25.4)	18	(26.9)	32	(47.8)	18	.	.	18	(100.0)
80-84	13	.	.	6	(46.2)	7	(53.8)	4	1	(25.0)	3	(75.0)
85-	3	.	.	1	(33.3)	2	(66.7)	2	.	.	2	(100.0)
全年齢	3,083	535	(17.4)	882	(28.6)	1,666	(54.0)	2,124	728	(34.3)	1,396	(65.7)
女 0-4	14	14	(100.0)	15	.	.	15	(100.0)
5-9	54	2	(3.7)	16	(29.6)	36	(66.7)	62	19	(30.6)	43	(69.4)
10-14	126	15	(11.9)	37	(29.4)	74	(58.7)	109	39	(35.8)	70	(64.2)
15-19	171	33	(19.3)	54	(31.6)	84	(49.1)	146	72	(49.3)	74	(50.7)
20-24	230	57	(24.8)	48	(20.9)	125	(54.3)	176	86	(37.5)	110	(62.5)
25-29	299	55	(18.4)	83	(27.8)	161	(53.8)	263	87	(33.1)	176	(66.9)
30-34	445	80	(18.0)	137	(30.8)	228	(51.2)	365	155	(42.5)	210	(57.5)
35-39	718	162	(22.6)	192	(26.7)	364	(50.7)	776	379	(48.8)	397	(51.2)
40-44	1,390	367	(26.4)	332	(23.9)	691	(49.7)	1,032	473	(45.8)	559	(54.2)
45-49	1,739	455	(26.2)	491	(28.2)	793	(45.6)	1,368	679	(49.6)	689	(50.4)
50-54	2,213	630	(28.5)	599	(27.1)	984	(44.5)	1,621	822	(50.7)	799	(49.3)
55-59	2,378	748	(31.5)	643	(27.0)	987	(41.5)	1,477	755	(51.1)	722	(48.9)
60-64	2,026	666	(32.9)	546	(26.9)	814	(40.2)	1,253	696	(55.5)	557	(44.5)
65-69	1,460	512	(35.1)	336	(23.0)	612	(41.9)	755	426	(56.4)	329	(43.6)
70-74	585	247	(42.2)	156	(26.7)	182	(31.1)	207	113	(54.6)	94	(45.4)
75-79	177	53	(29.9)	41	(23.2)	83	(46.9)	48	11	(22.9)	37	(77.1)
80-84	45	4	(8.9)	15	(33.3)	26	(57.8)	11	.	.	11	(100.0)
85-	4	.	.	2	(50.0)	2	(50.0)	2	.	.	2	(100.0)
全年齢	14,074	4,086	(29.0)	3,728	(26.5)	6,260	(44.5)	9,686	4,792	(49.5)	4,894	(50.5)

9. 強皮症・皮膚筋炎及び多発性筋炎

図2-5-1-9 97、92、88年度受給者の過去の各調査年度における受給状況、性・年齢階級別

