

Abiru N., Sun F., Kawasaki E., Yamasaki H., Oshima K., Nagayama Y., Mizuguchi H., Hayakawa T., Miao D., Liu E., Eisenbarth G.S., Eguchi K.	<i>In vivo</i> expression of B:9-23 peptide/I-A ^{g7} complex may abrogate the inhibition of diabetes induced by RGD-fiber-mutant adenovirus in NOD mice.	Annals N. Y. Acad. Sci.	in press
Nakanishi M., Eguchi A., Akuta T., Nagoshi E., Fujita S., Okabe J., Senda T., Hasegawa M.	Basic Peptides as Functional Components of Non-viral Gene Transfer Vehicles.	Current Protein & Peptide Science	in press
K. SATOH, A. IWATA, M. MURATA, M. HIKATA, T. HAYAKAWA, T. YAMAGUCHI	Virus Concentration Using Polyethyleneimine- conjugated Magnetic Beads for Improvement of Sensitivity in Nucleic Acid Amplification Tests.	J. Virol. Method.	in press
Kita A., Uotani S., Kuwahara H., Takahashi R., Oshima K., Yamasaki H., Mizuguchi H., Hayakawa T., Nagayama Y., Yamaguchi Y., Eguchi K.	Vanadate enhances leptin-induced activation of JAK/STAT pathway in CHO cells.	Biochem. Biophys. Res. Commun.	302, 805-809 (2003)
Uchida E., Mizuguchi H., Ishi-Watabe A., Hayakawa T.	Comparison of the efficiency and safety of non-viral vector-mediated gene transfer into a wide range of human cells.	Biol. Pharm. Bull.	25, 891-897 (2002)
Akuta T., Eguchi A., Okuyama H., Senda T., Inokuchi H., Suzuki Y., Nagoshi E., Mizuguchi H., Hayakawa T., Takeda K., Hasegawa M., Nakanishi M.	Enhancement of phage-mediated gene transfer by nuclear localization signal.	Biochem. Biophys. Res. Commun.	297, 779-786 (2002)
Mizuguchi H., Hayakawa T.	Enhanced anti-tumor effect and reduced vector dissemination with fiber-modified adenovirus vectors expressing herpes simplex virus thymidine kinase.	Cancer Gene Ther.	9, 236-242 (2002)
Mizuguchi H., Hayakawa T.	The tet-off system is more effective than the tet-on system for regulating transgene expression in a single adenovirus vector.	J. Gene Med.	4, 240-247 (2002)
Kawamata Y., Nakao K., Nagayama Y., Mizuguchi H., Hayakawa T., Sato T., Ishii N.	Receptor-independent augmentation of adenovirus-mediated gene transfer with chitosan in vitro.	Biomaterials.	23, 4573-4579 (2002)
Mizuguchi H., Hayakawa T.	Adenovirus vectors containing chimeric type 5 and type 35 fiber proteins exhibit altered and expanded tropism and increase the size limit of foreign genes.	Gene	285, 69-77 (2002)
Okada Y., Okada N., Nakagawa S., Mizuguchi H., Takahashi K., Mizuno N., Fujita T., Yamamoto A., Hayakawa T., Mayumi T.	Tumor necrosis factor α -gene therapy for an established murine melanoma using RGD (Arg-Gly-Asp) fiber-mutant adenovirus vectors.	Jpn. J. Cancer Res.	93, 436-444 (2002)

Nagayama Y, Kita-Furuyama M., Ando T., Nakao K., Mizuguchi H., Hayakawa T., Eguchi K., Niwa M.	A Novel murine model of graves' hyperthyroidism with intramuscular injection of adenovirus expressing the thyrotropin receptor.	<i>J. Immunol.</i>	168, 2789-2794 (2002)
Omori M., Mizuguchi H., Ohsawa K., Kohsaka S., Hayakawa T., Abe K., Shibasaki F.	Modification of a fiber protein in an adenovirus vector improves in vitro gene transfer efficiency to microglial cell line.	<i>Neurosci. Lett.</i>	324, 145-148 (2002)
Xu Z.L., Mizuguchi H., Ishii-Watabe I., Uchida E., Mayumi T., Hayakawa T.	Strength evaluation of transcriptional regulatory elements for transgene expression by adenovirus vector.	<i>J. Control. Rel.</i>	81, 155-163 (2002)
Mizuguchi H., Koizumi N., Hosono T., Ishii-Watabe A., Uchida E., Utoguchi N., Watanabe Y., Hayakawa T.	CAR- or αv integrin-binding ablated adenovirus vectors, but not fiber-modified vectors containing RGD peptide, do not change the systemic gene transfer properties in mice.	<i>Gene Ther.</i>	9, 769-776 (2002)
Takahashi M., Seki N., Ozaki T., Kato M., Kuno T., Nakagawa T., Watanabe K., Miyazaki K., Ohira M., Hayashi S., Hosoda M., Tokita H., Mizuguchi H., Hayakawa T., Todo S., Nakagawara A.	Identification of the p33ING1-regulated genes which include cyclin B1 and proto-oncogene DEK by using cDNA microarray in a mouse mammary epithelial cell line NmuMG.	<i>Cancer Res.</i>	62, 2203-2209 (2002)
Nakagawa T., Takahashi M., Ozaki T., Watanabe K., Todo S., Mizuguchi H., Hayakawa T., Nakagawara A.	Autoinhibitory regulation of p73 by Dnp73 to modulate cell survival and death through p73-specific target element within the Dnp73 promoter.	<i>Mol. Cell. Biol.</i>	22, 2575-2585 (2002)
Okada Y., Okada N., Nakagawa S., Mizuguchi H., Kanehira M., Nishino N., Takahashi K., Mizuno N., Hayakawa T., Mayumi T.	Fiber-mutant technique can augment gene transduction efficacy and anti-tumor effects against established murine melanoma by cytokine-gene therapy using adenovirus vectors.	<i>Cancer Lett.</i>	177, 57-63 (2002)
水口裕之・早川堯夫	遺伝子機能解析のための遺伝子導入ベクター-ウイルスベクターを中心として-	蛋白質核酸酵素	印刷中
水口裕之・早川堯夫	アデノウイルスベクター：最近の進歩	分子細胞治療	印刷中
水口裕之・早川堯夫	in vitro ライゲーションを利用したアデノウイルスベクターの作製・増殖法	実験医学	20, 1799-1804 (2002)
早川堯夫・石井(渡部)明子	生物薬品の開発の現状とトランスレーショナルリサーチへの条件	医学のあゆみ	200, 539-543 (2002)
早川堯夫・石井(渡部)明子	先端的バイオロジクス開発の現状と新たなバイオ創薬に向けての課題	医薬品研究	33, 693-729 (2002)
早川堯夫・山口照英・石井(渡部)明子・押澤 正	核酸増幅法によるウイルスゲノム等検出に関するフィージビリティスタディ	医薬品研究	33, 275-284 (2002)

20020423

以降は雑誌/図書に掲載された論文となりますので、
P.78-P.80の「研究成果の刊行に関する一覧表」をご参照ください。