

| | | | | | |
|---|--|-------------------------------|--------|-----------|------|
| 大河内 仁志 | 皮膚に存在する多能性幹細胞 | Medical Practice | 19 | 1220-1221 | 2002 |
| 大河内 仁志 | 皮膚の幹細胞について | 日本臨床皮膚科医学雑誌 | 72 | 92-96 | 2002 |
| Yano, S., Komine M, Fujimoto, M, Okochi, H., Tamaki, K | Interleukin 15 induces the Signals of Epidermal proliferation through E- RK and PI3-kinase in human epidermal kera- tinocyte cell line, HaCaT | Biochem. Biophys. Res. Commun | 301 | 841-847 | 2003 |
| Mitsui H, Komine M, Shirai A, Kanda N, Asahina A, Okochi H, Hitomi S, Kimura S, Tamaki K. | Chronic active EB virus infection complicated with IgG3 subclass deficiency: an adult case treated with intravenous immunoglobulin and IFN-alpha | Acta Derm Venereol | 83(1) | 31-5 | 2003 |
| Mitsui H, Komine M, Watanabe T, Kikuchi K, Okochi H, Tamaki K | Does Hermansky-Pudlak syndrome predispose to systemic lupus erythematosus? | Br J Dermatol | 146(5) | 908-11 | 2002 |
| Yano S, Nakamura K, Okochi H, Tamaki K: | Analysis of the expression of cutaneous lymphocyte-associated antigen on the peripheral blood and cutaneous lymphocytes of alopecia areata patients. | Acta Derm Venereol | 82 | 82-85 | 2002 |

| | | | | | |
|--|--|---------------------------|-----------------|-------------|------|
| Iwamoto R, Yamazaki S, Asakura M, Takashima S, Hasuwa H, Miyado K, Adachi S, Kitakaze M, Hashimoto K. Raab G, Klagsbrun M Nanba D ,Higasiyama S,Hori M,Mekada E | HB-EGF and ErbB signaling is essential for heart function | Proc Natl Acad Sci | in press | | |
| Shirakata Y, Tamai K,Naka- oka H,Tokuma- ru S, Sayama k Murakami S, Hashimoto K | Severe Palmo-plantar Hyperkeratosis in Koebner Epider- molysis Bullosa Simplex. | Dermatol | in press | | |
| Wada T,Shira- kata Y,Taka- hashi H,Mura- kami S, Iizuka H, Suzuki H, Hashimoto K | A Japanese Case of Segmental Darier's Disease Caused by Mosaicism for the ATP2A2 Mutation | Br J Derma- tol | in press | | |
| Sayama K,Ya- masaki K,Ha- nakawa Y,Shi- rakata Y,To- kumaru S,Iju- in T, Takenawa T, Hashimoto K. | Phosphatidyl inositol 3 kinase is a key regulator of early phase differentiation in keratinocytes. | J Biol Chem. | 277 | 40390-40396 | 2002 |

| | | | | | |
|--|---|------------------------------|----------|-----------|---------|
| Hanakawa Y, Amagai M, Shirakata Y, Yahata Y, Tokumaru S, Yamasaki K, Tohyama M, Sayama K, Hashimoto K | Differentiatl effects of desmoglein 1 and desmoglein 3 on desmosome formation | J Invest Dermatol | 119 | 1231-1236 | 2002 |
| Tsuda T, Thoyama M, Yamasaki K, Shirakata Y, Yahata Y, Tokumaru S, Sayama K, Hashimoto K | Lack of evidence for TARC/CCL17 production by normal human keratinocytes in vitro. | J Dermatol Sci | in press | | |
| Kobayashi M, Amagai M, Kuroda-Kinoshita K, Hashimoto T, Shirakata Y, Hashimoto K , Nishikawa T | BP180 ELISA using bacterial recombinant NC16a protein as a diagnostic and monitoring tool for bullous pemphigoid. | J Dermatol Sci | 30 | 224-232 | 2002 |
| Hattori N, Komine M, Yano S, Kaneko T, Hanakawa Y, Hashimoto K , Tamaki K | Interferon-gamma, a strong suppressor of cell proliferation, induces upregulation of keratin K6, one of the inflammatory- and proliferation-associated keratins. | J Invest Dermatol | 119 | 403-410 | 2002 |
| 白方裕司、橋本公三 | 下腿潰瘍の治療：培養皮膚による治療 | 皮膚病診療 | 24 | 1006-1011 | 2002.09 |
| 橋本公三 | 上皮再生の現状と新たな展望 | Medical Science Digest | 28 | 8-10 | 2002.12 |

| | | | | | |
|---|---|-----------------------------------|----------|-------|------|
| Nagashima T, Ueda Y, Hanazono Y, Kume A, Shibata H, Ageyama N, Terao K, Ozawa K, Hasegawa M | In vivo expansion of gene-modified hematopoietic cells by a novel selective amplifier gene utilizing the erythropoietin receptor as a molecular switch | Biochem Biophys Res Comm | in press | | |
| Hanazono Y, Asano T, Ueda Y, Ozawa K | Genetic manipulation of primate embryonic and hematopoietic stem cells with simian lentivirus vectors | Trends in Cardiovascular Medicine | in press | | |
| Kume A, Koremoto M, Xu R, Okada T, Mizukami H, Hanazono Y, Hasegawa M, Ozawa K | <i>In vivo</i> expansion of transduced murine hematopoietic cells with a selective amplifier gene. | J Gene Med | in press | | |
| Kanazawa T, Mizukami H, Okada T, Hanazono Y, Kume A, Nishino H, Takeuchi K, Kitamura K, Ichimura K, Ozawa K | Suicide gene therapy using AAV-HSVtk/ganciclovir in combination with irradiation results in regression of human head and neck cancer xenografts in nude mice. | Gene Ther | 10 | 51-58 | 2003 |

| | | | | | |
|--|---|---------------|----|---------|------|
| Lu YY, Wang LJ, Muramatsu S, Ikeguchi K, Fujimoto K, Okada T, Mizukami H, Matsushita T, Hanazono Y, Kume A, Nagatsu T, Ozawa K, Nakano I | Intramuscular injection of AAV-GDNF results in sustained expression of transgenic GDNF, and its delivery to spinal motoneurons by retrograde transport. | Neurosci Res | 45 | 33-40 | 2003 |
| Ageyama N, Hanazono Y, Shibata H, Ohto K, Ono F, Nagashima T, Ueda Y, Donahue RE, Hasegawa M, Ozawa K, Yoshikawa Y, Terao K | Safe and efficient methods for cynomolgus monkey autologous hematopoietic stem cell transplantation for biomedical researches. | Comp Med | 52 | 445-451 | 2002 |
| Kume A, Hanazono Y, Mizukami H, Okada T, Ozawa K | Selective expansion of transduced cells for hematopoietic stem cell gene therapy. | Int J Hematol | 76 | 299-304 | 2002 |
| Okada T, Nomoto T, Shimazaki K, Lijun W, Lu Y, Matsushita T, Mizukami H, Urabe M, Hanazono Y, Kume A, Muramatsu S, Nakano I, Ozawa K | Adeno-associated virus vectors for gene transfer to the brain. | Methods | 28 | 237-247 | 2002 |

| | | | | | |
|--|---|------------------------|------|-----------|------|
| Wang LJ, Lu YY, Muramatsu S, Ikeguchi K, Fujimoto K, Okada T, Mizukami H, Matsushita T, Hanazono Y, Kume A, Nagatsu T, Ozawa K, Nakano I | Neuroprotective effects of glial cell line-derived neurotrophic factor mediated by an adeno-associated virus vector in a transgenic animal model of amyotrophic lateral sclerosis | J Neurosci | 22 | 6920-6928 | 2002 |
| Hanazono Y, Terao K, Shibata H, Nagashima T, Ageyama N, Asano T, Ueda Y, Kato I, Kume A, Hasegawa M., Ozawa K | Introduction of the GFP gene into hematopoietic stem cells results in prolonged discrepancy of in vivo transduction levels between bone marrow progenitors and peripheral blood cells in nonhuman primates. | J Gene Med | 2002 | 470-477 | 2002 |
| Asano T, Hanazono Y, Ueda Y, Muramatsu S, Kume A, Suemori H, Suzuki Y, Kondo Y, Harii K, Hasegawa M, Nakatsuji N, Ozawa K | Highly efficient gene transfer into primate embryonic stem cells with a simian lentivirus vector. | Mol. Ther | 6 | 162-168 | 2002 |
| Kume A, Koremoto M, Mizukami H, Okada T, Hanazono Y, Sugamura K, Ozawa K | Selective growth advantage of wild-type lymphocytes in X-linked SCID recipients | Bone Marrow Transplant | 30 | 113-118 | 2002 |

| | | | | | |
|--|--|---------------------|-----|---------------|------|
| Hanazono Y, Nagashima T, Takatoku M, Shibata H, Ageyama N, Asano T, Ueda Y, Dunbar CE, Kume A, Terao K, Hasegawa M, Ozawa K | <i>In vivo</i> selective expansion of gene-modified hematopoietic cells in a nonhuman primate model. | Gene Ther | 9 | 1055-106 4 | 2002 |
| Okada T, Shimazaki K, Nomoto T, Matsushita T, Mizukami H, Urabe M, Hanazono Y, Kume A, Tobita K, Ozawa K, Kawai N | AAV mediated gene transfer for gene therapy of ischemia-induced neuronal death. | Methods Enzymol. | 346 | 378-393 | 2002 |
| Wang L, Muramatsu S, Lu Y, Ikeguchi K, Fujimoto K, Okada K, Mizukami H, Hanazono Y, Kume A, Urano F, Ichinose H, Nagatsu T, Nakano I, Ozawa K | Delayed delivery of AAV-GDNF prevents nigral neurodegeneration and promotes functional recovery in a rat model of Parkinson's disease. | Gene Ther | 9 | 381-389 | 2002 |

20020473

以降は雑誌/図書に掲載された論文となりますので、
P.38-P.45の「研究成果の刊行に関する一覧表」をご参照ください。