

- Tamaki K, Kaneko F. TGF- β 1-mediated regulation of thymus and activation regulated chemokine synthesis and secretion by HaCaT cells co-stimulated with TNF- α and IFN- γ . *J Dermatol Sci.* 2002; 30(2):154-60.
- 3) Nagaoka Y, Nakamura K, Yasaka N, Watanabe T, Asahina A, Tamaki K. Identification and Characterization of the Low-Affinity Receptor for Immunoglobulin E (Fc ϵ RII/CD23) on Murine Langerhans Cells. *J Invest Dermatol.* 2002. 119(1), 130-6.
- 4) Tsunemi Y, Saeki H, Nakamura K, Sekiya T, Hirai K, Fujita H, Asano N, Tanida Y, Kakinuma T, Wakugawa M, Torii H, Tamaki K. Eotaxin gene single nucleotide polymorphisms in the promoter and exon regions are not associated with susceptibility to atopic dermatitis, but two of them in the promoter region are associated with IgE levels in patients with atopic dermatitis. *J Dermatol Sci.* 29, 222-228, 2002.
- 5) Tsunemi Y, Saeki H, Nakamura K, Kakinuma T, Fujita H, Asano N, Tanida Y, Wakugawa M, Torii H, Asahina A, Tamaki K. Interleukin-13 (IL-13) gene polymorphisms in Japanese patients with atopic dermatitis. *J Dermatol Sci.* in press
- 6) Kakinuma T, Nakamura K, Wakugawa M, Mitsui H, Tada Y, Saeki H, Torii H, Komine M, Asahina A, Tamaki K. Serum macrophage - derived chemokine (MDC) levels are closely related with the disease activity of atopic dermatitis. *Clin Exp Immunol.* 2002; 127(2):270-3.
- 7) Kakinuma T, Nakamura K, Wakugawa M, Yano S, Saeki H, Torii H, Komine M, Asahina A, Tamaki K. IL-4, but not IL-13, modulate TARC (thymus and activation regulated chemokine) / CCL17 and IP-10 (Interferon -induced protein of 10kDa)/ CXCL10 released by TNF-alpha and IFN-gamma in HaCaT cell line. *Cytokine.* 2002; 20(1):1-6.
- 8) Tsunemi Y, Saeki H, Nakamura K, Sekiya T, Hirai K, Fujita H, Asano N, Kishimoto M, Tanida Y, Kakinuma T, Mitsui H, Tada Y, Wakugawa M, Torii H, Komine M, Asahina A, Tamaki K. Interleukin-12 p40 gene (IL12B) 3'-untranslated region polymorphism is associated with susceptibility to atopic dermatitis and psoriasis vulgaris. *J Dermatol Sci.* 2002; 30(2):161-6.
- 9) Tsunemi Y, Saeki H, Nakamura K, Sekiya T, Hirai K, Kakinuma T, Fujita H, Asano N, Tanida Y, Wakugawa M, Torii H, Tamaki K. Interleukin-13 gene polymorphism G4257A is associated with atopic dermatitis in Japanese patients. *J Dermatol Sci.* 2002 ;30(2):100-7.
- 10) Kobayashi N, Taguchi-Nakamura H, Goto M, Nakamura T, Nakamura K, Sugiura W, Iwamoto A, Kitamura Y. Polymorphisms and haplotypes of the CD209L gene and their association with the clinical course of HIV- positive Japanese patients. *Jpn J Infect Dis* 55: 131-, 2002.

H.知的財産権の出願、登録状況

- 1 特許申請 なし
- 2 実用新案登録 なし