

研究成果の刊行に関する一覧表

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Furukawa, F., <u>Nishikawa, A.</u> , Chihara, T., Shimpo, K., Beppu, H., Kuzuya, H., Lee, I-S., Hirose M.	Chemopreventive effects of <i>Aloe arborescens</i> on <i>N</i> - nitrosobis(2-oxopropyl)amine-i nduced pancreatic carcinogenesis in hamsters.	Cancer Lett.	178	117- 122	2002
Miyauchi, M., Nakamura, H., Furukawa, F., Son, H-Y., <u>Nishikawa</u> <u>A.</u> , Hirose, M.	Promoting effects of combined treatment of antioxidants with sodium nitrite on forestomach carcinogenesis in rats after initiation with <i>N</i> -methyl- <i>N'</i> -nitro- <i>N</i> - nitrosoguanidine.	Cancer Lett.	178	19-24	2002
Hirose, M., <u>Nishikawa, A.</u> , Shibutani, M., Imai, T., Shirai, T.	Chemoprevention of heterocyclic amine-induced mammary carcinogenesis in rats.	Environ. Mol. Mutagen.	39	271- 278	2002
Kasahara, K., <u>Nishikawa, A.</u> , Furukawa, F., Ikezaki, S., Tanakamaru, Z., Lee, I-S., Imazawa, T., Hirose, M.	A chronic toxicity study of josamycin in F344 rats.	Food Chem. Toxicol.	40	1017- 1022	2002
Okazaki, K., Yamagishi, M., Son, H-Y., Imazawa, T., Furukawa, F., Nakamura, H., <u>Nishikawa, A.</u> , Masegi, T., Hirose, M.	Simultaneous treatment with benzyl isothiocyanate, a bladder promoter, inhibits rat urinary bladder carcinogenesis by <i>N</i> butyl- <i>N</i> (4- hydroxybutyl)nitrosamine.	Nutr. Cancer	42	211- 216	2002

研究成果の刊行に関する一覧表

Shibutani, M., Takahashi, N., Kobayashi, T., Uneyama, C., Masutomi, N., <u>Nishikawa, A.</u> , Hirose, M.	Molecular profiling of genes up-regulated during promotion by phenobarbital-treatment in a medium-term rat liver bioassay.	Carcinogenesis	23	1047-1055	2002
Yamagishi, M., Natsume, M., Osakabe, N., Nakamura, H., Furukawa, F., Imazawa, T., <u>Nishikawa, A.</u> , Hirose, M.	Effects of cacao liquor proanthocyanidins on PhIP-induced mutagenesis <i>in vitro</i> , and <i>in vivo</i> mammary and pancreatic tumorigenesis in female Sprague-Dawley rats.	Cancer Lett.	185	123-130	2002
Umemura, T., Kodama, Y., Hioki, K., Nomura, T., <u>Nishikawa, A.</u> , Hirose, M., Kurokawa, Y.	The mouse rasH2/BHT model as an <i>in vivo</i> rapid assay for lung carcinogens.	Jpn. J. Cancer Res.	93	861-866	2002
Mori, Y., Koide, A., Kobayashi, Y., Furukawa, F., Hirose, M., <u>Nishikawa, A.</u>	Effects of cigarette smoke and a heterocyclic amine, MeIQx on cytochrome P-450, mutagenic activation of various carcinogens and glucuronidation in rat liver.	Mutagenesis	18	87-93	2003
Son, H-Y., <u>Nishikawa, A.</u> , Okazaki, K., Kanki, K., Yamagishi, M., Imazawa, T., Umemura, T., Hirose, M.	Prolonged effects of $\beta$ -estradiol 3-benzoate on thyroid tumorigenesis in gonadectomized rats pretreated with <i>N</i> -bis(2-hydroxypropyl) nitrosamine.	Cancer Lett.	190	21-29	2003

## 研究成果の刊行に関する一覧表

Yamagishi, M., Natsume, M., Osakabe, N., Okazaki, K., Furukawa, F., Imazawa, T., <u>Nishikawa, A.</u> , Hirose, M.	Chemoprevention of lung carcinogenesis by cacao liquor proanthocyanidins in a male rat multi-organ carcinogenesis model.	Cancer Lett.	191	49-57	2003
Yanaida, Y., Kohno, H., Yoshida, K., Hirose, Y., Yamada, Y., Mori, H., <u>Tanaka, T.</u>	Dietary silymarin suppresses 4-nitroquinoline 1-oxide-induced tongue carcinogenesis in male F344 rats.	Carcinogenesis	23	787-794	2002
Kohno, H., Maeda, M., Tanino, M., Tsukio, Y., Ueda, N., Wada, K., Sugie, S., Mori, H., <u>Tanaka, T.</u>	A bitter diterpenoid furanolactone columbin from <i>Calumbae Radix</i> inhibits azoxymethane-induced rat colon carcinogenesis.	Cancer Lett.	183	131-139	2002
<u>Tanaka, T.</u> , Kohno, H., Yamada, Y., Hirose, Y., Sugie, S., Mori, H.	Modifying effects of dietary capsaicin and rotenone on 4-nitroquinoline 1-oxide-induced rat tongue carcinogenesis.	Carcinogenesis	23	1361-1367	2002
Yamada, Y., Hata, K., Hirose, Y., Hara, A., Sugie, S., Kuno, T., Yoshimi, N., <u>Tanaka, T.</u> , Mori, H.	Microadenomatous lesions involving loss of <i>Apc</i> heterozygosity in the colon of adult <i>Apc<sup>Min/+</sup></i> mice.	Cancer Res.	62	6367-6370	2002
Iwai, S., Karim, R., Kitano, M., Sukata, T., Wei, M., Morimura, K., <u>Wanibuchi, H.</u> , Fukushima, S.	Role of oxidative DNA damage caused by carbon tetrachloride-induced liver injury - enhancement of MeIQx-induced glutathione <i>S</i> -transferase placental form-positive foci in rats.	Cancer Lett.	179	15-24	2002

## 研究成果の刊行に関する一覧表

Kakuni, M., Morimura, K., <u>Wanibuchi, H.</u> Ogawa, M., Min, W., Hayashi, S., Fukushima, S.	Food restriction inhibits the growth of intestinal polyps in multiple intestinal neoplasia mouse.	Jpn. J. Cancer Res.	93	236-241	2002
Nishikawa, T., <u>Wanibuchi, H.</u> Ogawa, M., Kinoshita, A., Morimura, K., Hiroi, T., Funae, Y., Kishida, H., Nakae, D., Fukushima, S.	Promoting effects of monomethylarsonic acid, dimethylarsinic acid and trimethylarsine oxide on induction of rat liver preneoplastic glutathione <i>S</i> -transferase placental form positive foci: a possible reactive species mechanism.	Int. J. Cancer	100	136-139	2002
Wei, M., <u>Wanibuchi, H.</u> , Morimura, K., Iwai, S., Yoshida, K., Endo, G., Nakae, D., Fukushima, S.	Carcinogenicity of dimethylarsinic acid in male F344 rats and genetic alterations in induced urinary bladder tumors.	Carcinogenesis	23	1387-1397	2002
Kaneko, M., Morimura, K., Nishikawa, T., <u>Wanibuchi, H.</u> Takada, N., Osugi, H., Kinoshita, H., Fukushima, S.	Different genetic alterations in rat forestomach tumors induced by genotoxic and non-genotoxic carcinogens.	Carcinogenesis	23	1729-1735	2002

研究成果の刊行に関する一覧表

<p>Fukushima, S., <u>Wanibuchi, H.</u> Morimura, K., Wei, M., Nakae, D., Konishi, Y., Tsuda, H., Uehara, N., Imaida, K., Shirai, T., Tatematsu, M., Tsukamoto, T., Hirose, M., Furukawa, F., Wakabayashi, K., Totsuka, Y.</p>	<p>Lack of a dose-response relationship for carcinogenicity in the rat liver with low doses of 2-amino-3,8-dimethyl- imidazo[4,5-<i>f</i>]quinoxaline or <i>N</i>-nitrosodiethylamine.</p>	<p>Jpn. J. Cancer Res.</p>	<p>93</p>	<p>1076- 1082</p>	<p>2002</p>
<p>Salim, E. I., <u>Wanibuchi, H.</u> Morimura, K., Wei, M., Mitsuhashi, M., Yoshida, K., Endo, G., Fukushima, S.</p>	<p>Carcinogenicity of dimethylarsinic acid in <i>p53</i> heterozygous knockout and wild-type C57BL/6J mice.</p>	<p>Carcinogenesis</p>	<p>24</p>	<p>335- 342</p>	<p>2003</p>
<p>Fukushima, S., <u>Wanibuchi, H.</u> Morimura, K., Wei, M., Nakae, D., Konishi, Y., Tsuda, H., Takasuka, N., Imaida, K., Shirai, T., Tatematsu, M., Tsukamoto, T., Hirose, M., Furukawa, F.</p>	<p>Lack of initiation activity in rat liver of low doses of 2-amino-3,8-dimethyl- imidazo[4,5-<i>f</i>]quinoxaline.</p>	<p>Cancer Lett.</p>	<p>191</p>	<p>35-40</p>	<p>2003</p>
<p>Seike, N., <u>Wanibuchi, H.</u> Morimura, K., Wei, M., Nishikawa, T., Hirata, K., Yoshikawa, J. and Fukushima, S.</p>	<p>Enhancement of lung carcinogenesis by nonylphenol and genistein in a F344 rat multiorgan carcinogenesis model.</p>	<p>Cancer Lett.</p>	<p>192</p>	<p>25-36</p>	<p>2003</p>

## 研究成果の刊行に関する一覧表

Karim, R., <u>Wanibuchi, H.</u> , Wei, M., Morimura, K., Salim, E.I. and Fukushima, S.	Enhancing risk of ethanol on MeIQx-induced rat hepatocarcinogenesis is accompanied with increased levels of cellular proliferation and oxidative stress.	Cancer Lett.	192	37-47	2003
Kuno T, Yamada Y, <u>Hirose Y.</u> , Katayama M, Sakata K, Hara A, Saji S, Mori H.	Induction of apoptosis by sulindac in azoxymethane-induced possible colonic premalignant lesions in rats.	Jpn. J. Cancer Res.	93	242- 246	2002
Yamada Y, Oyama T, <u>Hirose Y.</u> , Hara A, Sugie S, Yoshida K, Yoshimi N, Mori H.	beta-Catenin mutation is selected during malignant transformation in colon carcinogenesis.	Carcinogenesis	24	91-97	2003
<u>Hirose Y.</u> , Kuno T, Yamada Y, Sakata K, Katayama M, Yoshida K, Qiao Z, Hata K, Yoshimi N, Mori H.	Azoxymethane-induced beta-catenin-accumulated crypts in colonic mucosa of rodents as an intermediate biomarker for colon carcinogenesis.	Carcinogenesis	24	107- 111	2003
T. Shirai, M. Asamoto, S. Takahashi, <u>K.</u> <u>Imaida.</u>	Diet and prostate cancer.	Toxicology	181- 182	89-94	2002
T. Shirai, K. Kato, M. Futakuchi, S. Takahashi, S. Suzuki, <u>K. Imaida.</u> M. Asamoto.	Organ differences in the enhancing potential of 2-amino-1-methyl-6- phenylimidazo[4,5-b]pyridine on carcinogenicity in the prostate, colon and pancreas.	Mutat Res.	506- 507	129- 136	2002

研究成果の刊行に関する一覧表

M. Futakuchi, M. Hirose, <u>K. Imaida</u> , S. Takahashi, K. Ogawa, M. Asamoto, T. Miki, T. Shirai.	Chemoprevention of 2-amio-1-methyl-6-phenylimidazo[4,5- <i>b</i> ]pyridine-induced colon carcinogenesis by 1- <i>O</i> -hexyl-2,3,5-trimethyl hydroquinone after initiation with 1,2-dimethylhydrazine in F344 rats.	Carcinogenesis	23	283-287	2002
S. Takahashi, S. Inaguma, Y.M. Cho, <u>K. Imaida</u> , J. Wang, O. Fujiwara, T. Shirai.	Lack of mutation induction with exposure to 1.5 GHz electromagnetic near fields used for cellular phones in brains of Big Blue mice.	Cancer Res.	62	1956-1960	2002
N. Niho, M. Shibutani, K. Toyoda, H. Sato, A. Hirose, <u>K. Imaida</u> , M. Takahashi, Y. Hayashi, M. Hirose.	Dose- and time-response studies of sodium <i>o</i> -phenylphenate urinary bladder carcinogenicity in rats.	Food Chem. Toxicol.	40	715-722	2002
Puatanachokchai R, Kishida H, Denda A, Murata N, Konishi Y, Vinitketkumnuen U and <u>Nakae D</u>	Inhibitory effects of lemon grass ( <i>Cymbopogon citratus</i> , Stapf) extract on the early phase of hepatocarcinogenesis initiated with diethyl-nitrosamine in male Fischer 344 rats.	Cancer Lett.	183	9-15	2002
Kawai Y, Kato Y, <u>Nakae D</u> , Kusuoka O, Konishi Y, Uchida K and Osawa T	Immunohistochemical detection of a substituted 1, <i>N</i> <sup>2</sup> -ethenodeoxyguanosine adduct by $\omega$ -6 poly-unsaturated fatty acid hydroperoxides in the livers of rats fed a choline-deficient, L-amino acid-defined diet.	Carcinogenesis	23	485-489	2002

## 研究成果の刊行に関する一覧表

Yoshida M, Katsuda S, Tanimoto T, Asai S, <u>Nakae D</u> , Kurokawa Y, Taya K and Maekawa A	Induction of different types of uterine adenocarcinomas in Donryu rats due to neonatal exposure to high-dose <i>p-tert</i> -octylphenol for different periods.	Carcinogenesis	23	1745-1750	2002
<u>Nakae D</u> , Kishida H, Enami T, Konishi Y, Hensley KL, Floyd RA and Kotake Y	Effects of <i>N-tert</i> butyl nitron and its derivatives on the early phase of hepatocarcinogenesis in rats fed a choline-deficient, <i>L</i> -amino acid-defined diet.	Cancer Sci	94	26-31	2003