

**"Daphnia sp., Acute Immobilisation Test  
and Reproduction Test"**

2. DIN 38 412, Part 1: Allgemeine Hinweise zur Planung, Durchführung und Auswertung biologischer Testverfahren, (June 1982), and DIN 38 412, Part 2: Bestimmung der Wirkung von Wasserinhaltsstoffen auf Kleinkrebse (Daphnien-Kurzzeitest), (L 11), Normenausschuss Wasserwesen (NAW) beim Deutschen Institut für Normung e.V. (DIN), Arbeitsgruppe "Biotests": Testverfahren mit Wasserorganismen (Gruppe L), Berlin (March 1982).
  3. U.S. EPA: Registration of Pesticides in the United States - Proposed Guidelines, *Federal Register*, 43, N° 132 (July 10, 1978).
  4. AFNOR T 90 301: Qualité d'eau - Détermination de l'inhibition de la mobilité de *Daphnia magna* Straus (Cladocera, Crustacea) (French standard - 1974).
  5. NEN 6501 (Dutch standard on acute test).
  6. NEN 6502 (Dutch standard on reproduction test).
- Other
7. U.S. EPA: *Environmental Effects Test Guidelines*, ES-1, EPA 560/6-82-002, Office of Toxic Substances, U.S. Environmental Protection Agency, Washington, D.C (1982).
  8. D.M.M. Adema, in *Degradability, Ecotoxicity and Bioaccumulation: The Determination of the Possible Effects of Chemicals and Wastes on the Aquatic Environment*, Chapter 5, Government Publishing Office, The Hague (1980).
  9. D.M.M. Adema, *Hydrobiologia* 519 (2), 125-134 (1978).
  10. K.E. Biesinger and G.M. Christensen, *J. Fish. Res. Board of Canada* 29 (12), 1691-1700 (1972).
  11. Commission of the European Communities, Study D.8369, Inter-laboratory Test Programme concerning the Study of the Ecotoxicity of a Chemical Substance with respect to *Daphnia* (1979).
  12. J.H. Canton and D.M.M. Adema, *Hydrobiologia* 519 (2), 135-140 (1978).
  13. J. Finney, *Statistical Methods in Biological Assay*, Griffin, Weycombe, U.K. (1978).

**"Daphnia sp., Acute Immobilisation Test  
and Reproduction Test"**

14. J.T. Litchfield and F. Wilcoxon, *J. Pharmacol. Exper. Ther.* 96, 99-113 (1949).
15. C.E. Stephan, in *Aquatic Toxicology and Hazard Evaluation: Methods for Calculating an LC 50*, (edited by F.L. Mayer and J.L. Hamelink), ASTM STP 634, pp. 65-84, American Society for Testing and Materials (1977).
16. R.W. Wimmer and M.P. Farrel, *J. Fish Res. Bd. of Canada* 33, 1685-1691 (1976).

## OECD GUIDELINE FOR TESTING OF CHEMICALS

Adopted by the Council on 17<sup>th</sup> July 1992

### Fish, Acute Toxicity Test

#### INTRODUCTION

1. This new version of the guideline, originally adopted in 1981 and first updated in 1984, is based on a proposal from the United Kingdom to reduce the numbers of fish in tests of acute aquatic toxicity. The proposal was discussed at a meeting of OECD experts convened at Medmenham (United Kingdom) in November 1988.
2. The main differences in comparison with the earlier versions are the reduction in group-size allowing the use of seven fish per group, the extension of the concentration range by allowing a spacing factor of 2.2 instead of 2 and the introduction of a limit test at 100 mg/l of test substance.

#### PRINCIPLE OF THE TEST

3. The fish are exposed to the test substance preferably for a period of 96 hours. Mortalities are recorded at 24, 48, 72 and 96 hours and the concentrations which kill 50 per cent of the fish (LC50) are determined where possible.

#### INFORMATION ON THE TEST SUBSTANCE

4. It is necessary to know the water solubility of the substance under the conditions of the test. A reliable analytical method for the quantification of the substance in the test solutions must also be available.
5. Useful information includes the structural formula, purity of the substance, stability in water and light,  $pK_a$ ,  $P_{ow}$ , vapour pressure and results of a test for ready biodegradability (see Guideline 301). Solubility and vapour pressure can be used to calculate Henry's constant which will indicate if losses of the test substance may occur.

#### VALIDITY OF THE TEST

6. For a test to be valid the following conditions should be fulfilled:
  - the mortality in the control(s) should not exceed 10 per cent (or one fish if less than ten are used) at the end of the test;

- constant conditions should be maintained as far as possible throughout the test and, if necessary, *semi-static or flow-through procedures should be used (see Annex 1 for definitions)*;
- the dissolved oxygen concentration must have been at least 60 per cent of the air saturation value *throughout the test*;
- there must be evidence that the concentration of the substance being tested has been satisfactorily maintained, and preferably it should be at least 80 per cent of the *nominal concentration throughout the test*. If the deviation from the nominal concentration is greater than 20 per cent, results should be based on the measured concentration.

### **DESCRIPTION OF THE METHOD**

#### **Apparatus**

7. Normal laboratory equipment and especially the following is necessary :
- (a) oxygen meter;
  - (b) equipment for determination of hardness of water;
  - (c) adequate apparatus for temperature control;
  - (d) tanks made of chemically inert material and of a suitable capacity in relation to the recommended loading.

#### **Selection of species**

8. One or more species may be used, the choice being at the discretion of the testing laboratory. It is suggested that the species used be selected on the basis of such important practical criteria as, for example, their ready availability throughout the year, ease of maintenance, convenience for testing and any relevant economic, biological or ecological factors. The fish should be in good health and free from any apparent malformation.

9. Examples of fish recommended for testing are given in the Table. The fish mentioned in the Table are easy to rear and/or widely available throughout the year. They can be bred and cultivated either in fish farms or in the laboratory, under disease- and parasite-controlled conditions, so that the test fish will be healthy and of known parentage. These fish are available in many parts of the world. If other species fulfilling the above criteria are used, the test method should be adapted in such a way as to provide suitable test conditions.

#### **Holding of fish**

10. All fish must be obtained and held in the laboratory for at least 12 days before they are used for testing. They must be held in water of the quality to be used in the test for at least seven days immediately before testing and under the following conditions:

- Light: 12 to 16 hours photoperiod daily;
- Temperature: appropriate to the species (see Table);
- Oxygen  
concentration: at least 80 per cent of air saturation value;

Feeding: three times per week or daily until 24 hours before the test is started.

11. Following a 48-hour settling-in period, mortalities are recorded and the following criteria applied:

- mortalities of greater than 10 per cent of population in seven days: rejection of entire batch;
- mortalities of between 5 and 10 per cent of population: acclimatisation continued for seven additional days;
- mortalities of less than 5 per cent of population: acceptance of batch.

#### Water

12. Good quality natural water or reconstituted water (see Annex 2) is preferred, although drinking water (dechlorinated if necessary) may also be used. Waters with total hardness of between 10 and 250 mg CaCO<sub>3</sub> per liter, and with a pH 6.0 to 8.5 are preferable. The reagents used for the preparation of reconstituted water should be of analytical grade and the deionised or distilled water should be of conductivity equal to or less than 10  $\mu\text{Scm}^{-1}$ .

#### Test solutions

13. Test solutions of the chosen concentrations are prepared by dilution of a stock solution. Stock solutions of substances of low water solubility may be prepared by ultrasonic dispersion or other suitable physical means. If necessary, vehicles such as organic solvents, emulsifiers or dispersants of low toxicity to fish may be used. When such vehicles are used an additional control should be exposed to the same concentration of the vehicle as that used in the most concentrated solution of the test substance. The concentration of organic solvents, emulsifiers or dispersants should not exceed 100 mg/l.

14. The test should be carried out without adjustment of pH. If there is evidence of marked change in the pH of the tank water after addition of the test substance, it is advisable that the test be repeated, adjusting the pH of the stock solution to that of the tank water before addition of the test substance. This pH adjustment should be made in such a way that the stock solution concentration is not changed to any significant extent and that no chemical reaction or precipitation of the test substance is caused. HCl and NaOH are preferred.

### PROCEDURE

#### Conditions of exposure

15. Duration: preferably 96 hours.
- Loading: maximum loading of 1.0 g fish/litre for static and semi-static tests is recommended; for flow-through systems higher loading can be accepted.
- Light: 12 to 16 hours photoperiod daily.
- Temperature: appropriate to the species (see Table) and constant within a range of 2°C.

Oxygen concentration: not less than 60 per cent of the air saturation value. Aeration can be used provided that it does not lead to a significant loss of test substance.

Feeding: none.

Disturbance: disturbances that may change the behaviour of the fish should be avoided.

#### **Number of fish**

16. At least 7 fish must be used at each test concentration and in the controls.

#### **Test concentrations**

17. At least five concentrations in a geometric series with a factor preferably not exceeding 2.2. A range-finding test properly conducted before the definitive test enables the choice of the appropriate concentration range.

#### **Controls**

18. One blank and, if relevant, one control containing the solubilising agent are run in addition to the test series.

#### **Observations**

19. The fish are inspected at least after 24, 48, 72 and 96 hours. Fish are considered dead if there is no visible movement (e.g. gill movements) and if touching of the caudal peduncle produces no reaction. Dead fish are removed when observed and mortalities are recorded. Observations at three and six hours after the start of the test are desirable. Records are kept of visible abnormalities (e.g. loss of equilibrium, swimming behaviour, respiratory function, pigmentation, etc.). Measurement of pH, dissolved oxygen and temperature should be carried out at least daily.

#### **LIMIT TEST**

20. Using the procedures described in this Guideline, a limit test may be performed at 100 mg(active ingredient)/l in order to demonstrate that the LC50 is greater than this concentration. The limit test should be performed using a minimum of 7 fish, with the same number in the control(s). (Binomial theory dictates that when 10 fish are used with zero mortality, there is a 99.9 % confidence that the LC50 is greater than 100 mg/l. With 7, 8 or 9 fish, the absence of mortality provides at least 99% confidence that the LC50 is greater than the concentration used in the limit test.) If any mortalities occur, a full study should be conducted. If sublethal effects are observed, these should be recorded.

#### **DATA AND REPORTING**

##### **Treatment of results**

21. The cumulative percentage mortality for each exposure period is plotted against concentration on logarithmic probability paper. Normal statistical procedures are then employed to calculate the LC50 for the appropriate exposure period. Confidence limits ( $p = 0.95$ ) for the calculated LC50 values are determined using standard procedures (1)(2)(3)(4)(5).

22. Where the data obtained are inadequate for the use of standard methods of calculating the LC50, the highest concentration causing no mortality and the lowest concentration producing 100 per cent mortality should be used as an approximation for the LC50 (this being considered the geometric mean of these two concentrations).

#### Test report

23. The test report must include the following information:

##### Test substance:

- physical nature and, where relevant, physicochemical properties;
- identification data.

##### Test fish:

- scientific name, strain, size, supplier, any pretreatment, etc.

##### Test conditions:

- test procedure used (e.g. static, semi-static, flow-through; aeration; fish loading; etc.);
- water quality characteristics (pH, hardness, temperature);
- dissolved oxygen concentration, pH values and temperature of the test solutions at 24 hour intervals (in semi-static systems the pH should be measured prior to and after water renewal);
- methods of preparation of stock and test solutions;
- concentrations used;
- information on concentrations of the test substance in the test solutions;
- number of fish in each test solution.

##### Results:

- maximum concentration causing no mortality within the period of the test;
- minimum concentration causing 100 per cent mortality within the period of the test;
- cumulative mortality at each concentration at the recommended observation times;
- LC50 values, with 95 per cent confidence limits, at each of the recommended observation times, if possible;
- graph of the concentration-mortality curve at the end of the test;
- statistical procedures used for determining the LC50 values;
- mortality in the controls;
- incidents in the course of the test which might have influenced the results;
- abnormal responses of the fish.

##### Discussion of the results.

TABLE: FISH SPECIES RECOMMENDED FOR TESTING

Recommended species	Recommended test temperature range (°C)	Recommended total length of test fish (cm) <sup>1</sup>
<b><u>Brachydanio rerio</u></b> (Teleostei, Cyprinidae) (Hamilton-Buchanan) Zebra-fish	21 - 25	2.0 ± 1.0
<b><u>Pimephales promelas</u></b> (Teleostei, Cyprinidae) (Rafinesque) Fathead Minnow	21 - 25	2.0 ± 1.0
<b><u>Cyprinus carpio</u></b> (Teleostei, Cyprinidae) (Linnaeus) Common carp	20 - 24	3.0 ± 1.0
<b><u>Oryzias latipes</u></b> (Teleostei, Cyprinodontidae) (Temminck and Schlegel) Ricefish	21 - 25	2.0 ± 1.0
<b><u>Poecilia reticulata</u></b> (Teleostei, Poeciliidae) (Peters) Guppy	21 - 25	2.0 ± 1.0
<b><u>Lepomis macrochirus</u></b> (Teleostei, Centrarchidae) (Rafinesque) Bluegill	21 - 25	2.0 ± 1.0
<b><u>Oncorhynchus mykiss</u></b> (Teleostei, Salmonidae) (Walbaum) Rainbow trout	13 - 17	5.0 ± 1.0

<sup>1</sup> If fish of sizes other than those recommended are used, this should be reported together with the rationale.



LITERATURE

- (1) Litchfield J.T. and Wilcoxon F. (1949). A simplified method of evaluating dose-effect experiments. *J. Pharmacol and Exper. Ther.*, 96, 99-113.
- (2) Sprague J.B. (1969). Measurement of pollutant toxicity to fish. I Bioassay methods for acute toxicity. *Water Res.* 3, 793-821.
- (3) Sprague J.B. (1970). Measurement of pollutant toxicity to fish. II Utilising and applying bioassay results. *Water Res.* 4, 3-32.
- (4) Stephan C.E. (1977). Methods for calculating an LC50. In *Aquatic Toxicology and Hazard Evaluation* (edited by Mayer F.I. and Hamelink J.L.). ASTM STP 634, pp 65-84, American Society for Testing and Materials.
- (5) Finney D.J. (1978). *Statistical Methods in Biological Assay*. Griffin, Weycombe, U.K.

ANNEX 1DEFINITIONS

Static test is a test with aquatic organisms in which no flow of test solution occurs. Solutions remain unchanged throughout the duration of the test.

Semi-static test is a test without flow of solution, but with occasional batchwise renewal of the test solution after prolonged periods (e.g. 24 hours).

Flow-through test is a test in which solutions are automatically and continually renewed in the test chambers, the displaced solutions running to waste.

LC50 in this Test Guideline is the median lethal concentration, i.e. that concentration of the test substance in water which kills 50 per cent of a test batch of fish within a particular period of exposure (which must be stated).

ANNEX 2EXAMPLE OF A SUITABLE RECONSTITUTED WATER (ISO 6341-1982)

- (a) Calcium chloride solution  
Dissolve 11.76 g  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$  in deionised water; make up to 1 litre with deionised water
- (b) Magnesium sulphate solution  
Dissolve 4.93 g  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  in deionised water; make up to 1 litre with deionised water
- (c) Sodium bicarbonate solution  
Dissolve 2.59 g  $\text{NaHCO}_3$  in deionised water; make up to 1 litre with deionised water
- (d) Potassium chloride solution  
Dissolve 0.23 g  $\text{KCl}$  in deionised water; make up to 1 litre with deionised water

All chemicals must be of analytical grade.

The conductivity of the distilled or deionised water should not exceed  $10 \mu\text{Scm}^{-1}$ .

25 ml each of solutions (a) to (d) are mixed and the total volume made up to 1 litre with deionised water. The sum of the calcium and magnesium ions in this solutions is 2.5 mmol/l. The proportion Ca:Mg ions is 4:1 and Na:K ions 10:1. The acid capacity  $K_{\text{S4.3}}$  of this solution is 0.8 mmol/l.

Aerate the dilution water until oxygen saturation is achieved, then store it for about two days without further aeration before use.



204

Adopted:  
4 April 1984

## OECD GUIDELINE FOR TESTING OF CHEMICALS

**"Fish, Prolonged Toxicity Test:  
14-day Study"****1. INTRODUCTORY INFORMATION**• Prerequisites

- Water solubility
- Vapour pressure

• Guidance information

- Structural formula
- Purity of the test substance
- Methods of analysis for the quantification of the substance in water
- Chemical stability in water and light
- n-Octanol/water partition coefficient
- Results of a ready biodegradability test (see Test Guideline 301)

• Qualifying statements

- Constant conditions should be maintained as far as possible throughout the test. A flow-through procedure should normally be used; if adequate, a semi-static procedure may be adopted.
- For chemicals with limited solubility under the test conditions it may not be possible to determine values called for in this Test Guideline.
- This Test Guideline is only suitable for freshwater fish species.

• Standard documents

There are no relevant international standards.

---

*Users of this Test Guideline should consult the Preface,  
in particular paragraphs 3, 4, 7 and 8.*

---

---

**"Fish, Prolonged Toxicity Test:  
14-day Study"**

## 2. METHOD

### A. INTRODUCTION, PURPOSE, SCOPE, RELEVANCE, APPLICATION AND LIMITS OF TEST

This Test Guideline presents guidance for measurement of lethal and other observed effects in fish exposed to test substances. It may be used in place of Test Guideline 203 (Fish, Acute Toxicity Test) if a longer observation period is considered useful and appropriate and the reporting of additional information deemed necessary.

#### • Definitions

Semi-static test is a test without flow, but with occasional batchwise renewal of the test solution after prolonged periods (e.g. every 24 hours).

Flow-through test in this Test Guideline is a test in which water is renewed continuously in the test chambers, the test substance being transported with the water used to renew the test medium.

Threshold level of lethal effect is the lowest concentration of the test substance in the test solution at which the substance has a lethal effect.

Threshold level of observed effects is the lowest concentration of the test substance in the test solution at which the substance is observed to have an effect other than a lethal one on a significant number of test fish.

NOEC (no observed effect concentration) is the highest tested concentration of a test substance at which no statistically significant lethal or other effect is observed.

#### • Reference substances

No reference substances are recommended for this test. However, if a reference substance has been tested, the results should be given.

---

**"Fish, Prolonged Toxicity Test:  
14-day Study"**

- Principle of the test method

Threshold levels of lethal and other observed effects and NOEC are determined at intervals during the test period, which is at least fourteen days. If necessary, the test period should be extended beyond 14 days by one or two weeks.

- Conditions for the validity of the test

- The mortality in the controls should not exceed 10 per cent at the end of the test.
- The dissolved oxygen concentration should be at least 60 per cent of the air saturation value throughout the test.
- In semi-static procedures, aeration can be used, provided it does not lead to a significant loss of test substance.
- There should be evidence that the concentration of the substance being tested has been satisfactorily maintained (it should be at least 80 per cent of the nominal concentration) over the test period. The results should be based on measured concentrations if the deviation from the nominal concentration is greater than 20 per cent.

## B. DESCRIPTION OF THE TEST PROCEDURE

- Preparations

*Equipment*

Normal laboratory equipment and especially the following is necessary:

- Equipment for determination of temperature, pH, oxygen concentration and hardness of water
- Adequate apparatus for temperature control
- Test tanks made of chemically inert material and of a suitable capacity

*Solutions of the test substance*

Stock solutions of the appropriate concentrations are prepared by dissolving the appropriate amount of the test substance in the required volume of dilution water.

---

## "Fish, Prolonged Toxicity Test: 14-day Study"

Stock solutions of test substances of low water solubility may be prepared by mechanical dispersion or, if necessary, by use of vehicles, such as organic solvents, emulsifiers or dispersants of low toxicity to fish. The concentration of organic solvents, emulsifiers or dispersants should preferably not exceed 100 mg/l in the test solution.

Test solutions of chosen concentrations are prepared by dilution of the stock solution.

The test should be carried out without adjustment of pH. If there is evidence of marked change in the pH of the tank water after addition of the test substance, it is advised that the test be repeated, adjusting the pH of the stock solution to that of the tank water before addition of the test substance. This pH adjustment should be made in such a way that the stock solution concentration is not changed to any significant extent and that no chemical reaction or physical precipitation of the test substance is caused. HCl or NaOH are preferred.

### • Experimental animals

#### *Selection of species*

One or more species may be used, the choice being at the discretion of the testing laboratory. However, it is recommended that the species used for this test be selected from those recommended for the Acute Toxicity Test (Test Guideline 203). The species used should be selected on the basis of such important practical criteria as, for example, their ready availability throughout the year, their ease of maintenance, their convenience for testing and any relevant economic, biological or ecological factors. The fish should be in good health and free from any apparent malformation.

The fish mentioned in Test Guideline 203 are easy to rear and/or widely available throughout the year. They can be bred and cultivated either in fish farms or in the laboratory, under disease and parasite-controlled conditions, so that the test animal will be healthy and of known parentage. These fish are available in many parts of the world.

If other species fulfilling the above criteria are used, the test method should be adapted in such a way as to provide suitable test conditions.

---

**"Fish, Prolonged Toxicity Test:  
14-day Study"**

***Holding***

- Acclimatisation:** At least 12 to 15 days. All fish must be exposed to water of the quality to be used in the test for at least seven days before they are used. Any disturbances that may change the behaviour of the fish should be avoided.
- Water:** Drinking water supply (dechlorinated if necessary), good quality natural water or reconstituted water (see Test Guideline 203). Waters with a total hardness of between 50 and 250 mg of CaCO<sub>3</sub> per litre, and with a pH 6.0 to 8.5 are preferable. The reagents used for the preparation of the dilution water should be of analytical grade and the deionised or distilled water should be of conductivity equal to or less than 10 µS<sub>cm</sub><sup>-1</sup>.
- Light:** 12 to 16 hours photoperiod daily.
- Temperature:** Appropriate to the species (see Test Guideline 203).
- Oxygen concentration:** At least 80 per cent of air saturation value.
- Prophylactic treatment:** Prophylactic treatments should be avoided but reported when used.
- Feeding:** Once daily
- Mortality:** Following a 48-hours settling-in period, mortalities are recorded and the following criteria applied:
- greater than 10 per cent of population in seven days: rejection of entire batch
  - between 5 and 10 per cent of population: acclimatisation continued for seven additional days
  - less than 5 per cent of population acceptance of batch



---

**"Fish, Prolonged Toxicity Test:  
14-day Study"**

• Performance of the test

If a vehicle is used in the preparation of the stock solution of the test substance, it is necessary to run, in addition to the control group, a control group of fish exposed to the highest concentration of the vehicle used in the test.

*In the flow-through test, the concentration of the substance in the test solution may be determined at the beginning of the test; in the semi-static test at the beginning, immediately prior to the first renewal of the test solution and at the termination of the test. Appropriate procedures other than analysis for giving evidence that adequate concentrations of the test substance have been maintained can also be used.*

*Conditions of exposure*

- Duration: Normally 14 days, but can be extended by one or two weeks.
- Tanks: Of suitable capacity in relation to the recommended loading.
- Loading: For semi-static tests maximum loading of 1.0 g fish/litre is recommended; for flow-through systems higher loading can be acceptable.
- Number of animals: At least 10 for each concentration and control.
- Test concentrations: The test concentrations chosen must permit the determination both of the threshold levels for the lethal and other observable effects and of the NOEC value. Concentrations of the substance in excess of 100 mg/l need not be tested if a threshold level has not been reached up to this concentration.
- Water: Drinking water supply (dechlorinated if necessary), good quality natural water or reconstituted water (see Test Guideline 203). Waters with a total hardness of between 50 and 250 mg of CaCO<sub>3</sub> per litre, and with a pH 6.0 to 8.5 are preferable.

---

**"Fish, Prolonged Toxicity Test:  
14-day Study"**

The reagents used for the preparation of the dilution water should be of analytical grade and the deionised or distilled water should be of conductivity equal to or less than  $10 \mu\text{Scm}^{-1}$ .

- Light: 12 to 16 hours photoperiod daily.
- Temperature: Appropriate to the species (see Test Guideline 203) constant within  $\pm 2^\circ\text{C}$ .
- Oxygen concentration: Not less than 60 per cent of the maximum air saturation value throughout the test.
- Feeding: Either several times daily (the quantity of feed administered should not exceed the amount ingested immediately by the fish) or daily (the quantity of food being kept constant - e.g. 2 per cent dry weight related to the initial fish weight).
- Cleaning: Inside surfaces of the test tank in the flow-through test must be cleaned if necessary and the remaining excrement removed, at least twice weekly; in the semi-static test the test tank is replaced by a clean one each time the water is changed.

***Observations***

Observed effects are defined as follows:

Lethal effects: a fish is presumed to be dead if no respiratory movement and no reaction to a slight mechanical stimulus can be detected.

Effects other than lethal effects: these include all effects observed on the appearance, size and behaviour of the fish that make them clearly distinguishable from the control animals, e.g. different swimming behaviour, different reaction to external stimuli, changes in appearance of the fish, reduction or cessation of food intake, changes in length or body weight.

The fish are inspected at least once a day for mortality. Dead fish are removed when observed and mortalities are recorded.

---

**"Fish, Prolonged Toxicity Test:  
14-day Study"**

It is desirable that daily records be kept of all observed effects, but a minimum of three observation sessions per week must be conducted.

Measurements of pH, dissolved oxygen and temperature must be carried out at least twice a week.

Representative samples of the test population should be weighed and measured before the test starts. All survivors should be weighed and measured at the termination of the test. Fish should not be weighed or measured during the test as unnecessary handling may lead to damage and/or mortality.

### **3. DATA AND REPORTING**

#### **• Interpretation of results**

If it is observed that the stability or homogeneity of the test solutions cannot be maintained, care should be taken in the interpretation of the results and note made that these may not be reproducible.

#### **• Test report**

The test report should include the following information:

Test substance: *chemical identification data.*

Test organisms: *scientific name, strain, size, supplier, any pretreatment, etc.*

Test conditions:

- test procedure used (e.g. semi-static or flow-through, aeration, fish loading, etc.)
- water quality characteristics (treatment, including dechlorination, dissolved oxygen concentration, pH, hardness, temperature, any other information available)
- *dissolved oxygen concentration, pH values, temperature and total hardness of the test solutions at each of the recommended observation times*

---

**"Fish, Prolonged Toxicity Test:  
14-day Study"**

- methods of preparation of stock and test solutions
- concentrations used
- information on the maintenance of the concentration of the test substance in the test solutions
- number of fish at each test concentration

Values from the fish acute toxicity test

Results:

- observed effects at each concentration for each observation time in tabular form
- concentrations that produce lethal or other effects can be presented graphically with respect to time
- Threshold level of lethal effect
- Threshold level of observed effects
- NOEC
- cumulative mortality at each concentration and for each recommended observation time if possible
- mortality in the controls
- behavioural observation of the fish
- incidents in the course of the test which might have influenced the results
- any deviation from the Test Guideline

#### **4. LITERATURE**

1. D.M.M. Adema, in *Degradability, Ecotoxicity and Bioaccumulation*, Chapter 5, Government Publishing Office, The Hague (1980).
2. R. Bathe, *Arch. Toxicol. Suppl.* 2, 417-423 (1979).