

Fujimoto S, Matsushima A, Yoshitani K, Oya M, Shimono D, Takeda T, Kurose T, Yamada Y, Seino Y..	ype-1 diabetes mellitus with insufficient serum immunoreactive insulin elevation after subcutaneous NPH-insulin injection.	Diabetes Res Clin Pract	60 (1)	69-73	2003
Taniguchi A, Nishimura F, Murayama Y, Nagasaki S, Fukushima M, Sakai M, Yoshii S, Kuroe A, Suzuki H, Iwamoto Y, Soga Y, Okumura T, Ogura M, Yamada Y, Seino Y, Nakai Y	Porphyromonas gingivalis infection is associated with carotid atherosclerosis in non-obese Japanese type 2 diabetic patients.	Metabolism	52 (2)	142-145	2003
Pamir N, Lynn FC, Buchan AM, Ehses J, Hinke SA, Pospisilik JA, Miyawaki K, Yamada Y, Seino Y, McIntosh CH, Pederson RA.	Glucose-dependent insulino-tropic polypeptide receptor null mice (GIPR-/-) exhibit compensatory changes in the enteroinsular axis.	Am J Physiol	284 (5)	E931-939	2003
Y. Horikawa, N. Oda, L. Yu, K. Fujiwara, M. Makino, Y. Seino, M. Itoh and J. Takeda.	Genetic variations in CAPN10 are not a major factor in the occurrence of type 2 diabetes in Japanese.	J. Clin. Endocrinol. Metab	88	244-247	2003
M. Hayashi, H. Yamada, S. Uehara, R. Morimoto, A. Muroyama, S. Yatsushiro, J. Takeda, A. Yamamoto and Y. Moriyama.	Secretory granule- mediated co- secretion of L- glutamate and glucagon triggers glutamatergic signal transmission in islets of Langerhans	J. Biol. Chem	278	1966-1974	2003

Q. Zhu, K. Yamagata, A. Miura, N. Shihara, Y. Horikawa, J. Takeda, J. Miyagawa and Y. Matsuzawa.	T130I mutation in HNF-4 γ gene is a loss-of-function mutation in hepatocytes and is associated with late-onset Type II diabetes mellitus in Japanese subjects.	Diabetologica	46	567-573	2003
T. Tanaka, K. Ikari, K. Furushima, A. Okada, H. Tanaka, K. Furukawa, K. Yoshida, T. Ikeda, S. Ikegawa, S. Hunt, J. Takeda, S. Toh, S. Harata, T. Nakajima and I. Inoue	Genomewide linkage and linkage disequilibrium analyses pinpoint the ossification of the posterior longitudinal ligament of the spine to COL6A1 locus on chromosome 21.	Am. J. Hum. Genet.	73	812-822	2003
J. Lin, H. Wang, T. Narita, R. Kikuno, O. Ohara, N. Shihara, T. Nishigori, Y. Horikawa and J. Takeda.	Expression profile of mRNAs from human pancreatic islet tumors.	J. Mol. Endocrinol.	31	519-528	2003
Y. Shimamoto, J. Ishida, K. Yamagata, T. Saito, H. Kato, T. Matsuoka, K. Hirota, H. Daitoku, M. Nangaku, K. Yamagata, H. Fuji, J. Takeda and A. Fukamizu.	Inhibitory effect of small heterodimer partner hepatocyte nuclear factor-4 mediates bile acid-induced repression of human angiotensinogen gene	J. Biol. Chem.	279	7770-7779	2004
M. Enya, Y. Kanoh, T. Mune, M. Ishizawa, H. Sarui, M. Yamamoto, N. Takeda, K. Yasuda, M. Yasujiama, S. Tsutaya and J. Takeda.	Depressive state and paresthesia dramatically improved by intravenous MgSO ₄ in Gitelman's syndrome.	Intern. Med.	in press		

J-H. Kim, H-J. Kim, H-A. Seong, K-C. Park, A. Sanyal, J. Takeda, H. Ha, M. Shong, M-J. Tsai and H-S. Choi.	Orphan nuclear receptor SHP, a novel corepressor for a basic Helix-Loop-Helix (bHLH) transcription factor BETA2/NeuroD.	Mol. Endocrinol	in press		
N. Shihara, Y. Horikawa, T. Onishi, M. Ono, K. Kashimada and J. Takeda., ,	Identification of a de novo case of hepatocyte nuclear factor-1? deficiency with highly varied phenotypes.	Diabetologia	in press		

20030409

以降は雑誌/図書等に掲載された論文となりますので、
「研究成果の刊行に関する一覧表」をご参照ください。