

研究成果の一覧表

雑誌

発表者名	論文タイトル名	発表誌名	巻号	ページ	出版年
Salim, E I, Wanibuchi, H, Morimura, K, Wei, M, Mitsuhashi, M, Yoshida, K, Endo, G and Fukushima, S	Carcinogenicity of dimethylarsinic acid in p53 heterozygous knockout and wild-type C57BL/6J mice	Carcinogenesis	24	335-342	2003
Fukushima, S, Wanibuchi, H, Morimura, K, Wei, M, Nakae, D, Konishi, Y, Tsuda, H, Takasuka, N, Imaida, K, Shirai, T, Tatemetsu, M, Tsukamoto, T, Hirose, M and Furukawa, F	Lack of initiation activity in rat liver of low doses of 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline	Cancer Lett	191	35-40	2003
Mitsuhashi, M, Wanibuchi, H, Wei, M, Doi, K, Morimura, K, Masuca, C, Wada, S, Nakatani, T, Kakizoe, T and Fukushima, S	Lack of inhibition of BBN-induced bladder carcinogenesis in C57BL/6 mice by intravesical instillation of KRN 7000	J Toxicol Pathol	16	19-23	2003
Seike, N, Wanibuchi, H, Morimura, K, Wei, M, Nishikawa, T, Hirata, K, Yoshikawa, J and Fukushima, S	Enhancement of lung carcinogenesis by nonylphenol and genistein in a F344 rat multiorgan carcinogenesis model	Cancer Lett	192	25-36	2003
Karim, R, Wanibuchi, H, Wei, M, Morimura, K, Salim, E I and Fukushima, S	Enhancing risk of ethanol on MeIQx-induced rat hepatocarcinogenesis is accompanied with increased levels of cellular proliferation and oxidative stress	Cancer Lett	192	37-47	2003
Mitsuhashi, M, Wanibuchi, H, Wei, M, Doi, K, Morimura, K, Masuda, C, Wada, S, Nakatani, T, Kakizoe, T and Fukushima, S	No inhibition of urinary bladder carcinogenesis in rats with intravesical instillation of α -galactosylceramide	Asian Pacific J Cancer Prev	4	45-50	2003
Romanenko, A, Morimura, K, Wanibuchi, H, Wei, M, Zaparin, W, Vinnichenko, W, Kinoshita, A, Vozianov, A and Fukushima, S	Urinary bladder lesions induced by persistent chronic low-dose ionizing radiation	Cancer Sci	94	328-333	2003
Ogawa, M, Wanibuchi, H, Nishikawa, M, Yano, Y, Otani, S, Totsuka, Y, Wakabayashi, K, Nakae, D and Fukushima, S	Post-initiation inhibition of MeIQx hepatocarcinogenesis in rats by cysteine	Osaka City Med J	49	21-30	2003

Kinoshita, A , Wanibuchi, H , Morimura, K , Wei, M , Shen, J , Imaoka, S , Funae, Y and Fukushima, S	Phenobarbital at low dose exerts hormesis in rat hepatocarcinogenesis by reducing oxidative DNA damage, altering cell proliferation, apoptosis and gene expression	Carcinogenesis	24	1389-1399	2003
Fukushima, S , Morimura, K and Wanibuchi, H	Theresholds for carcinogenicity of environmental carcinogens	J Toxicol Sci	28	241	2003
Shen, J , Wanibuchi, H , Salim, E I , Wei, M , Kinoshita, A , Yoshida, K., Endo, G and Fukushima, S	Liver tumorigenicity of trimethylarsine oxide in male Fischer 344 rats-association with oxidative DNA damage and enhanced cell proliferation	Carcinogenesis	24	1827-1835	2003
Shen, J , Wanibuchi, H , Salim, E I , Wei, M , Doi, K , Yoshida, K , Endo, G , Morimura, K and Fukushima, S	Induction of glutathione S - transferase placental form positive foci in liver and epithelial hyperplasia in urinary bladder, but no tumor development in male Fischer 344 rats treated with monomethylarsonic acid for 104 weeks	Toxicology and Applied Pharmacology	193	335-345	2003
Mitsuhashi, M , Wanibuchi, H , Morimura, K , Doi, K , Wei, M , Wada, S , Nakatani, T and Fukushima, S	Significance of overexpression of metallothionein in mouse urinary bladder focal lesions induced by treatment with N-butyl-N-(4-hydroxybutyl)-nitrosamine	Cancer Sci	94	1052-1058	2003
Wei, M , Morimura, K , Wanibuchi, H , Shen, J , Doi, K , Mitsuhashi, M , Moku, M , Salim, E I and Fukushima, S	Chemopreventive effect of JTE-522, a selective cyclooxygenase-2 inhibitor, on 1,2-dimethylhydrazine-induced rat colon carcinogenesis	Cancer Lett	202	11-16	2003
K Imaida, S Tamano, A Hagihara, S Fukushima, T Shirai, and N Ito	Application of rat medium-term bioassays for detecting carcinogenic and modifying potentials of endocrine active substances	Pure Appl Chem	75	2491-2495	2003
H Takeuchi, K Saoo, M Yokohira, M Ikeda, H Maeta, M Miyazaki, H Yamazaki, T Kamataki, and K Imaida	Pretreatment with 8-methoxypsoralen, a potent human CYP2A6 inhibitor, strongly inhibits lung tumorigenesis induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in female A/J mice	Cancer Res	63	7581-7583	2003

H Tsuda, S Fukushima, H Wanibuchi, K Morimura, D Nakae, K Imaida, M Tate_ma_tsu, M Hirose, K Wakabayashi, M A Moore	Value of GST-P positive preneoplastic hepatic foci in dose-response studies of hepa-to-carci-nogenesis Evidence for practical thresholds with both genotoxic and non-geno-toxic carci-nogens A review of recent work	Toxicol Pathol	31	80-86	2003
S Fukushima, H Wanibuchi, K Morimura, M Wei, D Nakae, Y Konishi, H Tsuda, N Takasuka, K Imaida, T Shirai, M Tatematsu, T Tsukamoto, M Hirose, F Furukawa	Lack of initiation activity in rat liver of low doses of 2-amino-3,8-dimethyl-imidazo-[4,5-f]-quinoline	Cancer Lett	191	35-40	2003
J Q Li, A Kubo, F Wu, H Usuki, J Fujita, S Bandoh, T Masaki, K Saoo, H Takeuchi, S Kobayashi, K Imaida, H Maeta, T Ishida, S Kuriyama	Cyclin B1, unlike cyclin G1, increases significantly during colorectal carci-no-genesis and during later metastasis to lymph nodes	Int J Oncol	22	1101-1110	2003
I Narama, K Imaida, H Iwata, D Nakae, A Nishikawa, T Harada	A review of nomen--cla-ture and diagnostic criteria for pro-liferative lesions in the liver of rats by a work-ing group of the Japanese Society of Toxi-cologic Patho-logy	J Toxicol Pathol	16	1-17	2003
S Fukushima, H Wanibuchi, K Morimura, S Iwai, D Nakae, H Kishida, H Tsuda, N Uehara, K Imaida, T Shirai, M Tatematsu, T Tsukamoto, M Hirose, F Furukawa	Existence of a threshold for induction of aberrant crypt foci in the rat colon with low doses of 2-amino- 1-methyl-6 phenolimidazo[4,5-b] pyridine	Toxicol Sci	in press	in press	2004
Nakae D, Kishida H, Enami T, Konishi Y, Hensley KL, Floyd RA and Kotake Y	Effects of <i>N-tert</i> -butyl nitron and its derivatives on the early phase of hepatocarcinogenesis in rats fed a choline-deficient, l-amino acid-defined diet	Cancer Sci	94	26-31	2003
Tsuda H, Fukushima S, Wanibuchi H, Morimura K, Nakae D, Imaida K, Tatematsu M, Hirose M, Wakabayashi K and Moore MA	values of GST-P positive preneoplastic hepatic foci in dose-response studies of hepatocarcinogenesis Evidence for practical thresholds with both genotoxic and nongenotoxic carcinogens A review of recent work	Toxicol Pathol	31	80-86	2003

Fukushima S, Wanibuchi H, Morimura K, Wei M, Nakae D, Konishi Y, Tsuda H, Takasuka N, Imaida K, Shirai T, Tatematsu M, Tsukamoto T, Hirose M and Furukawa F	Lack of initiation activity in rat liver of low doses of 2-amino-3,8-domethylimidazo[2,5-f]quinoxaline	Cancer Lett	191	35-40	2003
Narama I, Imaida K, Iwata H, Nakae D, Nihikawa A and Harada T	A review of nomenclature and diagnostic criteria for proliferative lesions in the liver of rats by a working group of the Japanese Society of Toxicologic Pathology	J Toxicol Pathol	16	1-18	2003
Yoshida M, Shimomoto T, Katashima S, Shirai T, Nakae D, Watanabe G, Taya K and Maekawa A	Effects of maternal exposure to nonylphenol on growth and development of the female reproductive system and uterine carcinogenesis in rats	J Toxicol Pathol	16	259-266	2003
Shimomoto T, Yoshida M, Takahashi M, Uematsu F, Maekawa A and Nakae D	A case report of a choroids plexus carcinoma spontaneously occurring in the right lateral ventricle of a 14-week-old, female Donryu rat	Toxicol Pathol	32	264-268	2004
Yoshida M, Katsuda S, Nakae D and Maekawa A	Lack of toxicity or carcinogenicity of S-170, a sucrose fatty acid ester, in F344 rats	Food Chem Toxicol	42	667-676	2004