

34. Faqi AS, Dalsenter PR, Merker HJ, Chahoud I. Reproductive toxicity and tissue concentrations of low doses of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in male offspring rats exposed throughout pregnancy and lactation. *Toxicol Appl Pharmacol* 1998;150(2):383–392.
35. Haavisto T, Nurmela K, Pohjanvirta R, Huusko H, El-Gehani F, Paranko J. Prenatal testosterone and luteinizing hormone levels in male rats exposed during pregnancy to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin and diethylstilbestrol. *Mol Cell Endocrinol* 2001;178(1/2):169–179.
36. Ohsako S, Miyabara Y, Sakaue M, Ishimura R, Kakeyama M, Izumi H, Yonemoto J, Tohyama C. Developmental stage-specific effects of perinatal 2,3,7,8-tetrachlorodibenzo-*p*-dioxin exposure on reproductive organs of male rat offspring. *Toxicol Sci* 2002;66(2):283–292.
37. Karolczak M, Beyer C. Developmental sex differences in estrogen receptor-beta mRNA expression in the mouse hypothalamus/preoptic region. *Neuroendocrinology* 1998;68(4):229–234.
38. Poletti A, Negri-Cesi P, Rabuffetti M, Colciago A, Celotti F, Martini L. Transient expression of the 5α-reductase type 2 isozyme in the rat brain in late fetal and early postnatal life. *Endocrinology* 1998;139(4):2171–2178.
39. Drent HJ, Bouwman CA, Seinen W, Van den Berg M. Effects of some persistent halogenated environmental contaminants on aromatase (CYP19) activity in the human choriocarcinoma cell line JEG-3. *Toxicol Appl Pharmacol* 1998;148(1):50–55.