


HindIII-digested viral DNA and hCpG exhibited significant NF- $\kappa$ B activation, suggesting that undigested viral DNA is incapable of penetrating cells by transfection. No activation of NF- $\kappa$ B was observed in 293T cells cotransfected with a human TLR2 or TLR4 expression plasmid when stimulated with digested AcNPV DNA (data not shown).

Recent work demonstrated that the endogenous expression of TLR3, TLR7, TLR8, and TLR9 was mainly detected in the cytoplasmic vesicles of macrophages (58). To examine the localization of transiently expressed TLR9, we transfected 293T cells with a TLR9 expression plasmid and examined TLR9 expression by immunofluorescence microscopy and cell sorting. The expression of TLR9 in the cytoplasm was three times higher than that at the cell surface (Fig. 6B and C). These results indicate that the introduction of AcNPV DNA into the



cytoplasm is specifically detected by human TLR9 and results in the activation of NF- $\kappa$ B.

AcNPV requires endosomal maturation to induce immune system activation in macrophages. To further explore the role of endocytosis in the signal transduction pathway triggered by AcNPV DNA, we examined the effect of endosomal maturation or acidification inhibitors. As shown in Fig. 7A, chloroquine was able to inhibit immune system activation of RAW264.7 cells treated with AcNPV and mCpG oligonucleotides in a dose-dependent manner, but no inhibition of LPS or PGN activation was observed. Other inhibitors of endoso-

mal maturation or acidification inhibitors. As shown in Fig. 7A, chloroquine was able to inhibit immune system activation of RAW264.7 cells treated with AcNPV and mCpG oligonucleotides in a dose-dependent manner, but no inhibition of LPS or PGN activation was observed. Other inhibitors of endoso-




FIG. 7. AcNPV requires endosomal maturation to induce immune system activation in macrophages. (A) RAW264.7 cells ( $10^6$  cells/well) were stimulated with AcNPV ( $5 \mu\text{g/ml}$ ), mCpG ( $200 \text{ ng/ml}$ ), LPS ( $10 \text{ ng/ml}$ ), or PGN ( $2.5 \mu\text{g/ml}$ ) at the indicated concentrations of chloroquine. After 24 h of incubation, the production of TNF- $\alpha$  in culture supernatants was determined by a sandwich ELISA. Chloroquine was added to the cells 2 h before stimulation. Data are shown as means  $\pm$  SD. (B) RAW264.7 cells ( $10^6$  cells/well) were treated with AcNPV ( $5 \mu\text{g/ml}$ ) or LPS ( $10 \text{ ng/ml}$ ) and with the indicated concentrations of endosomal maturation inhibitors. After 24 h of incubation, the production of TNF- $\alpha$  in culture supernatants was determined by a sandwich ELISA. The inhibitors were added to the cells 2 h before stimulation. Data are shown as means  $\pm$  SD.

mal maturation, such as ammonium chloride, bafilomycin A1, and MDC, inhibited AcNPV-induced, but not LPS-induced, immune system activation (Fig. 7B). Together with our other data, these results indicate that endosomal acidification and/or maturation is a key step in AcNPV-induced immune system activation via TLR9, a process that requires the release of the viral genome into TLR9-expressing cytoplasmic vesicles following the internalization of viral DNA by endocytosis through gp64-mediated membrane fusion.

**AcNPV penetrates macrophages via the phagocytic pathway.** To further confirm that baculovirus was internalized into macrophages, we inoculated RAW264.7 cells with a recombinant baculovirus carrying a luciferase gene under the control of a mammalian promoter, AcCAGLuc (49). As shown in Fig. 8A, the expression of luciferase was observed in 293T cells, but not RAW264.7 cells, that were infected with AcCAGLuc. The viral capsid protein was clearly detected by immunoblotting for both 293T and RAW264.7 cells infected with AcNPV, but the protein level was greatly diminished in RAW264.7 cells by 6 h postinoculation, probably as a result of degradation (Fig. 8B).

These results suggest that baculovirus can penetrate into different cells via gp64-mediated endocytosis but that it translocates into different subcellular compartments in different cells. In 293T cells, the nucleocapsid was apparently able to reach the nucleus, where the reporter gene was efficiently transcribed following uncoating. However, in the immunocompetent RAW264.7 cells, the nucleocapsid appeared to have been trapped by the phagocytic pathway, and degraded viral DNA was then translocated into TLR9-expressing intracellular compartments (58).

## DISCUSSION

We have previously demonstrated that intranasal inoculation with AcNPV induces a strong innate immune response that protects mice from a lethal challenge with influenza virus (1). The lungs of mice inoculated with AcNPV exhibited a marked infiltration of macrophages, which presumably inhibit the growth of influenza virus in the lung tissues. The baculovirus envelope glycoprotein gp64 contains mannose, fucose,




FIG. 8. AcNPV penetrates macrophages through the phagocytic pathway. (A) 293T and RAW264.7 cells ( $10^6$  cells/well) were inoculated with a recombinant baculovirus possessing the luciferase gene under the control of the CAG promoter, AcCAGluc (49) (10 and 20  $\mu$ g/ml). Cells were harvested 24 h after infection, and relative luciferase activities were determined. (B) 293T and RAW264.7 cells ( $10^6$  cells/well) were inoculated with AcCAGluc (40  $\mu$ g/ml), washed extensively after 1 h of adsorption, and harvested after 4 or 6 h of incubation. The presence of the p39 capsid protein in cells inoculated with AcNPV was determined by immunoblotting with an anti-p39 monoclonal antibody.

and *N*-acetyl-glucosamine modifications but no detectable galactose or terminal sialic acid residues (29). The mannose receptor (MR) recognizes a range of carbohydrates present on the surfaces and cell walls of microorganisms. MR is primarily expressed on macrophages and DCs and is involved in MR-mediated endocytosis and phagocytosis. In addition, MR plays a key role in host defense and the induction of innate immunity (8). Therefore, it is tempting to speculate that gp64 interacts with MR through its mannose modifications in macrophages and DCs of mice inoculated with AcNPV. However, our data contradict such a model; instead, we show that it is AcNPV DNA, not the gp64 glycoprotein, that induces immune system activation in a MyD88/TLR9-dependent manner.

Recently, it was shown that plasmacytoid DCs (pDCs) naturally produce IFN- $\alpha$  in response to viruses (30). HSV-1 and -2, whose genomes contain abundant CpG motifs, are able to induce the production of IFN- $\alpha$  in pDCs. The HSV-induced production of IFN- $\alpha$  in pDCs derived from MyD88- and TLR9-deficient mice was completely eliminated (33, 40). The recognition of the HSV genome by TLR9 was shown to be mediated by an endocytic pathway that can be inhibited by chloroquine or bafilomycin A1. In this study, we demonstrated that AcNPV induces proinflammatory cytokines through a

MyD88/TLR9-dependent signaling pathway, whereas signaling molecules other than MyD88 may participate in IFN- $\alpha$  production in response to AcNPV. Recently, MyD88-independent TLR signaling events involving TIR domain-containing adaptor inducing IFN- $\beta$  (TRIF) were described (59). Therefore, it is possible that the TRIF pathway is one means by which AcNPV induces MyD88-independent IFN production. However, future studies are needed to clarify the precise mechanisms of this induction.

While UV irradiation of AcNPV abolishes its ability to stimulate an immune response, the addition of liposomes is able to restore this activity. UV-inactivated HSV is capable of inducing the production of IFN- $\alpha$  in pDCs (40), indicating that viral replication is not required for the HSV-induced immune response. In contrast, UV irradiation of AcNPV abolishes immune stimulation in macrophages, while internalization of the inactivated virus by liposomes restores the activity. These results, in conjunction with our data for AcNPV $\Delta$ 64, indicate that the AcNPV-induced production of cytokines in immunocompetent cells requires a fusion process mediated by gp64 that leads to internalization of the viral genome into the cells.

Recently, several viral envelope glycoproteins were shown to induce immune system activation through TLRs (10, 22, 34, 47). However, gp64 does not directly participate in a TLR-mediated immune response. TLR family members are expressed differentially at very low levels on the surfaces of different immune cells and appear to respond to different stimuli (43). A recent study indicated that LPS and CpG-rich DNA activate TLRs in distinct cellular compartments (3). Internalization and endosomal maturation are required for CpG-rich DNA to activate TLR9, but not for LPS to activate TLR4 on the plasma membrane. We showed here that the inhibition of endosomal maturation by a treatment with chloroquine abolishes the immune system activation of AcNPV in a dose-dependent manner. These results imply that immune system activation by AcNPV through TLR9 requires membrane fusion via gp64 as well as the liberation of the viral genome into cytoplasmic vesicles expressing TLR9.

Interestingly, Lund et al. demonstrated that the TLR7-mediated immune recognition of single-stranded RNAs from vesicular stomatitis virus and influenza virus requires endosomal acidification (41). The recognition of HSV-1 and HSV-2 viral DNAs through a TLR9/MyD88-dependent pathway in pDCs also requires endosomal acidification (40). These data indicate that TLR7 and TLR9 expressed in the endosomal or lysosomal compartments of immunocompetent cells recognize the viral genome entering the cell through receptor-mediated endocytosis or phagocytosis, leading to the secretion of inflammatory cytokines and IFNs. However, the precise mechanisms by which viral genomes translocate to TLR-expressing compartments are still unknown.

Since the first report on the immunostimulatory potential of bacterial DNA, which found that the main immunogenic fraction of mycobacterial lysates consists of genomic DNA (55, 56), substantial progress has been made towards understanding the immunostimulatory potency of CpG-rich DNA motifs, which are more common in bacteria than in vertebrates. For instance, TLR9 was shown to be responsible in vivo for immune system stimulation by oligodeoxynucleotides containing unmethylated CpG motifs (24). Like bacteria, AcNPV contains a significant

number of potentially bioactive CpG motifs. Interestingly, the frequency of CpG motifs in HSV DNA, which has been shown to be involved in the induction of angiogenesis in stromal keratitis (61), was similar to that in *E. coli* DNA. In contrast, the frequency of CpG motifs in the genome of an insect poxvirus was much lower than that for AcNPV (Table 1).

In conclusion, we have demonstrated that AcNPV has the ability to induce innate immune system activation through a MyD88/TLR9-dependent pathway. The molecular mechanisms of viral uptake, intracellular processing, and the induction of potent antiviral activity in immune cells require further investigation. However, the strong immune response induced by AcNPV makes it a promising candidate for a novel, adjuvant-containing vaccine vehicle against infectious diseases. In particular, our findings raise the possibility that AcNPV may be harnessed therapeutically to induce a host immune response against various infectious diseases caused by pathogens invading the respiratory tract.

#### ACKNOWLEDGMENTS

We are grateful to T. H. Chuang for providing us with the TLR9 expression plasmid, D. T. Golenbock for the pELAM Luc plasmid, and G. F. Rohrmann for the p39 monoclonal antibody. We also thank I. Yanase and H. Murase for secretarial work. T.A. and H.H. are Research Fellows of the Japan Society for the Promotion of Science.

This work was supported by grants-in-aid from the Ministry of Health, Labor and Welfare in Japan to Y.M. and from the 21st Century Center of Excellence Program of Japan to Y.M. and S.A.

#### REFERENCES

1. Abe, T., H. Takahashi, H. Hamazaki, N. Miyano-Kurosaki, Y. Matsuura, and H. Takaku. 2003. Baculovirus induces an innate immune response and confers protection from lethal influenza virus infection in mice. *J. Immunol.* 171:1133-1139.
2. Adachi, O., T. Kawai, K. Takeda, M. Matsumoto, H. Tsutsui, M. Sakagami, K. Nakanishi, and S. Akira. 1998. Targeted disruption of the MyD88 gene results in loss of IL-1- and IL-18-mediated function. *Immunity* 9:143-150.
3. Ahmad-Nejad, P., H. Hacker, M. Rutz, S. Bauer, R. M. Vabulas, and H. Wagner. 2002. Bacterial CpG-DNA and lipopolysaccharides activate Toll-like receptor at distinct cellular compartments. *Eur. J. Immunol.* 32:1958-1968.
4. Airene, K. J., M. O. Hiltunen, M. P. Turunen, A. M. Turunen, O. H. Laitinen, M. S. Kutomaa, and S. Y. Herttuala. 2000. Baculovirus-mediated periaxonal gene transfer to rabbit carotid artery. *Gene Ther.* 7:1499-1504.
5. Akira, S., K. Takeda, and T. Kaisho. 2001. Toll-like receptors: critical proteins linking innate and acquired immunity. *Nat. Immunol.* 2:675-680.
6. Alexopoulou, L., A. C. Holt, R. Medzhitov, and R. A. Flavell. 2001. Recognition of double-stranded RNA and activation of NF- $\kappa$ B by Toll-like receptor 3. *Nature* 413:732-738.
7. Aliprantis, A. O., R. B. Yang, M. R. Mark, S. Suggett, B. Devaux, J. D. Radolf, G. R. Klimpel, P. Godowski, and A. Zychlinsky. 1999. Cell activation and apoptosis by bacterial lipoproteins through Toll-like receptor-2. *Science* 285:736-739.
8. Apostolopoulos, V., and I. F. McKenzie. 2001. Role of the mannose receptor in the immune response. *Curr. Mol. Med.* 1:469-474.
9. Barsoum, J., R. Brown, M. McKee, and F. M. Boyce. 1997. Efficient transduction of mammalian cells by a recombinant baculovirus having the vesicular stomatitis virus G glycoprotein. *Hum. Gene Ther.* 8:2011-2018.
10. Bieback, K., E. Lien, I. M. Klagge, E. Avota, J. Schneider-Schaulies, W. P. Duprex, H. Wagner, C. J. Kirschning, V. Ter Meulen, and S. Schneider-Schaulies. 2002. Hemagglutinin protein of wild-type measles virus activates Toll-like receptor 2 signaling. *J. Virol.* 76:8729-8736.
11. Bowie, A., E. Kiss-Toth, J. A. Symons, G. L. Smith, S. K. Dower, and L. A. O. Neill. 2000. A46R and A52R from vaccinia virus are antagonists of host IL-1 and Toll-like receptor signaling. *Proc. Natl. Acad. Sci. USA* 97:10162-10167.
12. Boyce, F. M., and N. L. R. Bucher. 1996. Baculovirus mediated gene transfer into mammalian cells. *Proc. Natl. Acad. Sci. USA* 93:2348-2352.
13. Brightbill, H. D., D. H. Libraty, S. R. Krutzik, R. B. Yang, J. T. Bellse, J. R. Bleharski, M. Maitland, M. V. Norgard, S. E. Plevy, S. T. Smale, P. J. Brennan, B. R. Bloom, P. J. Godowski, and R. L. Modlin. 1999. Host defense mechanisms triggered by microbial lipoproteins through Toll-like receptors. *Science* 285:732-736.
14. Chuang, T. H., J. Lee, L. Kilne, J. C. Mathison, and R. J. Ulevitch. 2002. Toll-like receptor 9 mediates CpG-DNA signaling. *J. Leukoc. Biol.* 71:538-544.
15. Compton, T., E. A. Kurt-Jones, K. W. Boehme, J. Belko, E. Latz, D. T. Golenbock, and R. W. Finberg. 2003. Human cytomegalovirus activates inflammatory cytokine responses via CD14 and Toll-like receptor 2. *J. Virol.* 77:4588-4596.
16. Condreay, J. P., S. M. Witherspoon, W. C. Clay, and T. A. Kost. 1999. Transient and stable gene expression in mammalian cells transduced with a recombinant baculovirus vector. *Proc. Natl. Acad. Sci. USA* 96:127-132.
17. Diebold, S. S., T. Kaisho, H. Hemmi, S. Akira, E. Reis, and C. Sousa. 2004. Innate antiviral responses by means of TLR7-mediated recognition of single-stranded RNA. *Science* 303:1529-1531.
18. Gronowski, A. M., D. M. Hilbert, K. C. F. Sheehan, G. Garotta, and R. D. Schreiber. 1999. Baculovirus stimulates antiviral effect in mammalian cells. *J. Virol.* 73:9944-9951.
19. Hacker, H., H. Mischak, T. Miethke, S. Liptay, R. Schmid, T. Sparwasser, K. Heeg, G. B. Lipford, and H. Wagner. 1998. CpG-DNA-specific activation of antigen-presenting cells requires stress kinase activity and is preceded by non-specific endocytosis and endosomal maturation. *EMBO J.* 17:6230-6240.
20. Hacker, H., R. M. Vabulas, O. Takeuchi, K. Hoshino, S. Akira, and H. Wagner. 2000. Immune cell activation by bacterial CpG-DNA through myeloid differential marker 88 and tumor necrosis factor receptor-associated factor (TRAF) 6. *J. Exp. Med.* 192:595-600.
21. Hayashi, F., K. D. Smith, A. Ozinsky, T. R. Hawn, E. C. Yi, D. R. Goodlett, J. K. Eng, S. Akira, D. M. Underhill, and A. Aderem. 2001. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature* 410:1099-1103.
22. Haynes, L. M., D. D. Moore, E. A. Kurt-Jones, R. W. Finberg, L. J. Anderson, and R. A. Tripp. 2001. Involvement of Toll-like receptor 4 in innate immunity to respiratory syncytial virus. *J. Virol.* 75:10730-10737.
23. Heil, F., H. Hemmi, H. Hochrein, F. Ampenberger, C. Kirschning, S. Akira, G. Lipford, H. Wagner, and S. Bauer. 2004. Species-specific recognition of single-stranded RNA via Toll-like receptor 7 and 8. *Science* 303:1526-1529.
24. Hemmi, H., O. Takeuchi, T. Kawai, T. Kaisho, S. Sato, H. Sanjo, M. Matsumoto, K. Hoshino, H. Wagner, K. Takeda, and S. Akira. 2000. A Toll-like receptor recognizes bacterial DNA. *Nature* 408:740-745.
25. Hemmi, H., T. Kaisho, O. Takeuchi, S. Sato, H. Sanjo, K. Hoshino, T. Horiuchi, H. Tomizawa, K. Takeda, and S. Akira. 2002. Small anti-viral compounds activate immune cells via the TLR7 MyD88-dependent signaling pathway. *Nat. Immunol.* 3:196-200.
26. Hofmann, C., V. Sandig, G. Jennings, M. Rudolph, P. Schlag, and M. Strauss. 1995. Efficient gene transfer into human hepatocytes by baculovirus vectors. *Proc. Natl. Acad. Sci. USA* 92:10099-10103.
27. Hoshino, K., O. Takeuchi, T. Kawai, H. Sanjo, T. Ogawa, Y. Takeda, K. Takeda, and S. Akira. 1999. Cutting edge: Toll-like receptor 4 (TLR4)-deficient mice are hyporesponsive to lipopolysaccharide: evidence for TLR4 as the LPS gene product. *J. Immunol.* 162:3749-3752.
28. Huser, A., M. Rudolph, and C. Hofmann. 2001. Incorporation of decay-accelerating factor into the baculovirus envelope generates complement resistant gene transfer vectors. *Nat. Biotechnol.* 19:451-455.
29. Jarvis, D. L., and E. E. Flinn. 1995. Biochemical analysis of the N-glycosylation pathway in baculovirus-infected lepidopteran insect cells. *Virology* 212:500-511.
30. Kadowaki, N., S. Antonenko, J. Y. Lau, and Y. J. Liu. 2000. Natural interferon alpha/beta-producing cells link innate and adaptive immunity. *J. Exp. Med.* 192:219-226.
31. Kawai, T., O. Takeuchi, T. Fujita, J. Inoue, P. F. Muhlradt, S. Sato, K. Hoshino, and S. Akira. 2001. Lipopolysaccharide stimulates the MyD88-independent pathway and results in activation of IFN-regulatory factor 3 and the expression of a subset of lipopolysaccharide-inducible genes. *J. Immunol.* 167:5887-5894.
32. Krieg, A. M., and H. Wagner. 2000. Causing a commotion in the blood: immunotherapy progresses from bacteria to bacterial DNA. *Immunol. Today* 21:521-526.
33. Krug, A., G. D. Luker, W. Barchet, D. A. Leib, S. Akira, and M. Colonna. 2004. Herpes simplex virus type 1 (HSV-1) activates murine natural interferon-producing cells (IPC) through Toll-like receptor 9. *Blood* 103:1433-1437.
34. Kurt-Jones, E. A., L. Popova, L. Kwinn, L. M. Haynes, L. P. Jones, R. A. Tripp, E. E. Walsh, M. W. Freeman, D. T. Golenbock, L. J. Anderson, and R. W. Finberg. 2000. Pattern recognition receptors TLR4 and CD14 mediate response to respiratory syncytial virus. *Nat. Immunol.* 5:398-401.
35. Kurt-Jones, E. A., M. Chan, S. Zhou, J. Wang, G. Reed, R. Bronson, M. M. Arnold, D. M. Knipe, and R. W. Finberg. 2004. Herpes simplex virus 1 interaction with Toll-like receptor 2 contributes to lethal encephalitis. *Proc. Natl. Acad. Sci. USA* 101:1315-1320.
36. Lee, J., T. H. Chuang, V. Redecke, L. She, P. M. Pitha, D. A. Carson, E. Raz, and H. B. Cottam. 2003. Molecular basis for the immunostimulatory activity of guanine nucleoside analogs: activation of Toll-like receptor 7. *Proc. Natl. Acad. Sci. USA* 100:6646-6651.

37. Lehtolainen, P., K. Tynnela, J. Kannasto, K. J. Alrenne, and S. Y. Herttuala. 2002. Baculoviruses exhibit restricted cell type specificity in rat brain: a comparison of baculovirus- and adenovirus-mediated intracerebral gene transfer *in vivo*. *Gene Ther.* **9**:1693-1699.
38. Lemaitre, B., E. Nicolas, L. Michaut, J. M. Reichhart, and J. A. Hoffmann. 1996. The dorsoventral regulatory gene cassette *spatzle/Toll/cactus* controls the potent antifungal response in *Drosophila* adults. *Cell* **86**:973-983.
39. Luckow, V. A., and M. D. Summers. 1988. Signals important for high-level expression of foreign genes in *Autographa californica* nuclear polyhedrosis virus expression vectors. *Bio/Technology* **6**:47-55.
40. Lund, J., A. Sato, S. Akira, R. Medzhitov, and A. Iwasaki. 2003. Toll-like receptor 9-mediated recognition of herpes simplex virus-2 by plasmacytoid dendritic cells. *J. Exp. Med.* **198**:513-520.
41. Lund, J., L. Alexopoulou, A. Sato, M. Karow, N. C. Adams, N. W. Gale, A. Iwasaki, and R. A. Flavell. 2004. Recognition of single-stranded RNA viruses by Toll-like receptor 7. *Proc. Natl. Acad. Sci. USA* **101**:5598-5603.
42. Matsuura, Y., R. D. Possee, H. A. Overton, and D. H. L. Bishop. 1987. Baculovirus expression vector. The requirement for high level expression of proteins, including glycoproteins. *J. Gen. Virol.* **68**:1233-1250.
43. Muzio, M., D. Bosisio, N. Polentarutti, G. Damico, A. Stoppacciaro, R. Mancinelli, C. van Veer, G. Penton-Rol, L. P. Ruco, P. Allavena, and A. Mantovani. 2000. Differential expression and regulation of Toll-like receptors (TLR) in human leukocytes: selective expression of TLR3 in dendritic cells. *J. Immunol.* **164**:5998-6004.
44. Ozinsky, A., D. M. Underhill, J. D. Fontenot, A. M. Hajjar, K. D. Smith, C. B. Wilson, L. Schroeder, and A. Aderem. 2000. The repertoire for pattern recognition of pathogens by the innate immune system is defined by cooperation between Toll-like receptors. *Proc. Natl. Acad. Sci. USA* **97**:13766-13771.
45. Pieroni, L., D. Malone, and N. L. Monica. 2001. *In vivo* gene transfer in mouse skeletal muscle mediated by baculovirus vectors. *Hum. Gene Ther.* **12**:871-881.
46. Poltorak, A., X. He, I. Smirnova, M. Y. Liu, C. Van Huffel, X. Du, D. Birdwell, E. Alejos, M. Silva, C. Gafanos, M. Freudenberg, P. Ricciardi-Castagnoli, B. Layton, and B. Beutler. 1998. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in *Tlr4* gene. *Science* **282**:2085-2088.
47. Rassa, J. C., J. L. Meyers, Y. Zhang, R. Kudravalli, and S. R. Ross. 2002. Murine retroviruses activate B cells via interaction with Toll-like receptor 4. *Proc. Natl. Acad. Sci. USA* **99**:2281-2286.
48. Sandig, V., C. Hofmann, S. Steinert, G. Jennings, P. Schlag, and M. Strauss. 1996. Gene transfer into hepatocytes and human liver tissue by baculovirus vectors. *Hum. Gene Ther.* **7**:1937-1945.
49. Shoji, I., H. Aizaki, H. Tani, K. Ishii, T. Chiba, I. Saito, T. Miyamura, and Y. Matsuura. 1997. Efficient gene transfer into various mammalian cells, including non-hepatic cells, by baculovirus vectors. *J. Gen. Virol.* **78**:2657-2664.
50. Takeuchi, O., K. Hoshino, T. Kawai, H. Sanjo, H. Takeda, T. Ogawa, K. Takeda, and S. Akira. 1999. Differential roles of TLR2 and TLR4 in recognition of gram-negative and gram-positive bacterial cell wall components. *Immunity* **11**:443-451.
51. Takeuchi, O., and S. Akira. 2001. Toll-like receptors: their physiological role and signal transduction system. *Int. Immunopharmacol.* **1**:625-635.
52. Takeuchi, O., A. Kaufmann, K. Grote, T. Kawai, K. Hoshino, M. Morr, P. F. Muhlradt, and S. Akira. 2000. Cutting edge: preferentially the R-stereoisomer of the mycoplasma lipopeptide macrophage-activating lipopeptide-2 activates immune cells through a Toll-like receptor 2- and MyD88-dependent signaling pathway. *J. Immunol.* **164**:554-557.
53. Tani, H., M. Nishijima, H. Ushijima, T. Miyamura, and Y. Matsuura. 2001. Characterization of cell-surface determinants important for baculovirus infection. *Virology* **279**:343-353.
54. Tani, H., C. K. Lim, C. C. Yap, M. Onishi, M. Nozaki, Y. Nishimune, N. Okahashi, Y. Kitagawa, R. Watanabe, R. Mochizuki, K. Moriishi, and Y. Matsuura. 2003. *In vitro* and *in vivo* gene delivery by recombinant baculoviruses. *J. Virol.* **77**:9799-9808.
55. Tokunaga, T., H. Yamamoto, S. Shimada, H. Abe, T. Fukuda, Y. Fujisawa, Y. Furutani, O. Yano, T. Kataoka, T. Sudo, N. Makiguchi, and T. Suganuma. 1984. Antitumor activity of deoxyribonucleic acid fraction from *Mycobacterium bovis* BCG. I. Isolation, physicochemical characterization and antitumor activity. *JNCI* **72**:955-962.
56. Tokunaga, T., T. Yamamoto, and S. Yamamoto. 1999. How BCG led to the discovery of immunostimulatory DNA. *Jpn. J. Infect. Dis.* **52**:1-11.
57. Underhill, D. M., A. Ozinsky, A. M. Hajjar, A. Stevens, C. B. Wilson, M. Bassett, and A. Aderem. 1999. The Toll-like receptor 2 is recruited to macrophage phagosomes and discriminates between pathogens. *Nature* **401**:811-815.
58. Wagner, H. 2004. The immunobiology of the TLR9 subfamily. *Trends Immunol.* **25**:381-386.
59. Yamamoto, M., S. Sato, H. Hemmi, K. Hoshino, T. Kalsho, H. Sanjo, O. Takeuchi, M. Sugiyama, M. Okabe, K. Takeda, and S. Akira. 2003. Role of adaptor TRIF in the MyD88-independent Toll-like receptor signaling pathway. *Science* **301**:640-643.
60. Zhang, D., G. Zhang, M. S. Hayden, M. B. Greenblatt, C. Bussey, R. A. Flavell, and S. Ghosh. 2004. A Toll-like receptor that prevents infection by uropathogenic bacteria. *Science* **303**:1522-1526.
61. Zheng, M., D. M. Klinman, M. Gierynska, and B. T. Rouse. 2002. DNA containing CpG motifs induces angiogenesis. *Proc. Natl. Acad. Sci. USA* **99**:8944-8949.

## Ligand-Directed Gene Targeting to Mammalian Cells by Pseudotype Baculoviruses†

Yoshinori Kitagawa, Hideki Tani, Chang Kwang Limn, Tomoko M. Matsunaga, Kohji Moriishi, and Yoshiharu Matsuura\*

Research Center for Emerging Infectious Diseases, Research Institute for Microbial Diseases, Osaka University, Osaka, Japan

Received 27 August 2004/Accepted 25 October 2004

The baculovirus *Autographa californica* multiple nucleopolyhedrovirus (AcMNPV) can infect a variety of mammalian cells, as well as insect cells, facilitating its use as a viral vector for gene delivery into mammalian cells. Glycoprotein gp64, a major component of the budded AcMNPV envelope, is involved in viral entry into cells by receptor-mediated endocytosis and subsequent membrane fusion. We examined the potential production of pseudotype baculovirus particles transiently carrying ligands of interest in place of gp64 as a method of ligand-directed gene delivery into target cells. During amplification of a gp64-null pseudotype baculovirus carrying a green fluorescent protein gene in gp64-expressing insect cells, however, we observed the high-frequency appearance of a replication-competent virus incorporating the gp64 gene into the viral genome. To avoid generation of replication-competent revertants, we prepared pseudotype baculoviruses by transfection with recombinant bacmids without further amplification in the gp64-expressing cells. We constructed gp64-null recombinant bacmids carrying cDNAs encoding either vesicular stomatitis virus G protein (VSVG) or measles virus receptors (CD46 or SLAM). The VSVG pseudotype baculovirus efficiently transduced a reporter gene into a variety of mammalian cell lines, while CD46 and SLAM pseudotype baculoviruses allowed ligand-receptor-directed reporter gene transduction into target cells expressing measles virus envelope glycoproteins. Gene transduction mediated by the pseudotype baculoviruses could be inhibited by pretreatment with specific antibodies. These results indicate the possible application of pseudotype baculoviruses in ligand-directed gene delivery into target cells.

The baculovirus *Autographa californica* multiple nucleopolyhedrovirus (AcMNPV) is an insect virus possessing a 134-kb double-stranded circular DNA genome (3). Due to the strong polyhedrin and p10 promoters, baculovirus is commonly used as a tool for the large-scale production of recombinant protein in insect cells (32, 38). Baculovirus is also capable of entering into a variety of mammalian cells to facilitate the expression of foreign genes under the control of the mammalian promoters without replication of the viral genome (8, 21, 61). Therefore, baculovirus is a useful viral vector, not only for the abundant expression of foreign genes in insect cells, but also for efficient gene delivery to mammalian cells (29). AcMNPV has a number of unique beneficial properties as a viral vector, including a large capacity for foreign gene incorporation, easy manipulation, and replication competence in insect cells combined with incompetence in mammalian cells. Therefore, the possibility of generating replication-competent revertants expressing baculoviral gene products, which can often lead to harmful immune responses against mammalian cells, is significantly lower than for other viral vectors presently in use. Furthermore, studies of host responses to baculovirus infection in vivo revealed that AcMNPV can stimulate interferon production in mammalian cell lines, conferring protection from lethal encephalomyocarditis virus infections in mice (18). Intranasal

inoculation with AcMNPV also induces a strong innate immune response, protecting mice from lethal challenges of influenza A or B virus (1). The precise mechanism of protective immune response induction by AcMNPV, however, remains unclear.

Recently, several groups have reported enhanced gene transfer in a variety of cell lines infected with recombinant baculoviruses expressing either foreign viral envelope proteins, such as vesicular stomatitis virus envelope G protein (VSVG), or excess amounts of the endogenous envelope glycoprotein, gp64, on the virion surface (4, 65, 66). Although modification of the virion surface enhances the efficiency of gene transduction into a variety of cell lines, the utility of recombinant baculoviruses in cell-type-specific gene transduction is still unsatisfactory. Ojala et al. demonstrated that, while baculoviruses bearing either a single chain antibody fragment specific for carcinoembryonic antigen or a synthetic immunoglobulin G (IgG) binding domain derived from protein A could specifically bind target cells, cell type-specific gene transduction was unsuccessful (44, 45). Although gp64-null pseudotype baculoviruses expressing a foreign viral envelope protein, such as VSVG or fusion envelope glycoproteins from other baculoviruses, exhibited high infectivity to insect cells, their capacity for gene transduction into mammalian cells has yet to be explored (33, 34). The inefficiency of present gene transfer vectors in gaining entry into cells needing treatment can be problematic, as many therapeutic genes may be deleterious if delivered to bystander cells. Therefore, the development of a ligand-directed gene delivery vector capable of distinguishing between

\* Corresponding author. Mailing address: Research Center for Emerging Infectious Diseases, Research Institute for Microbial Diseases, Osaka University, Osaka 565-0871, Japan. Phone: 81-6-6879-8340. Fax: 81-6-6879-8269. E-mail: matsuura@biken.osaka-u.ac.jp.

† This study is dedicated to the memory of Ikuko Yanase

target and nontarget tissue is essential for both the safety and efficacy of gene therapy.

In this study, we examined the stability of a generated gp64-null pseudotype baculovirus possessing the green fluorescent protein (GFP) gene during passages in insect cells stably expressing the gp64 protein. Replication-competent revertant viruses emerged with high frequency during passage in the cell line, incorporating the gp64 gene into the revertants' viral genomes. To overcome the emergence of revertant viruses during passage, we generated recombinant bacmids lacking the gp64 gene and carrying a ligand of interest and a reporter gene under the control of the polyhedrin and the CAG promoters, respectively. Pseudotype baculoviruses generated from these bacmids exhibited specific ligand-directed gene delivery into target cells. These pseudotype baculovirus vectors may be useful in future clinical gene targeting.

#### MATERIALS AND METHODS

**Cells.** *Spodoptera frugiperda* (Sf9) cells were grown in TC-100 medium (Sigma, St. Louis, Mo.) supplemented with 0.26% tryptose phosphate broth (Difco, Detroit, Mich.) and 10% (vol/vol) fetal bovine serum (FBS) (Sigma) (66). To establish a cell line constitutively expressing gp64, Sf9 cells were transfected with pAFgp64 (see below) and pIB/V5-His (Invitrogen, Carlsbad, Calif.) using UniFactor reagent (B-Bridge, Sunnyvale, Calif.). Thirty-six hours after transfection, Sf9 cells were selected in TC-100 medium containing blasticidin (50 µg/ml; Invitrogen). Resistant cells were stained with anti-gp64 antibodies (AcV1) (kindly provided by P. Faulkner) (22); positive cells were sorted using a FACS-Calibur (Becton Dickinson, Franklin Lakes, N.J.) to establish a cell line, Sf9gp64, stably expressing gp64 at the cell surface. The human embryonic kidney cell line 293T and the hamster kidney cell line BHK, purchased from the American Type Culture Collection, were maintained in Dulbecco's modified Eagle's medium (Sigma) containing 2 mM L-glutamine, penicillin (50 IU/ml), streptomycin (50 µg/ml), and 10% FBS (66).

**Construction of plasmids.** We constructed two expression plasmids, pAF-MCS1 and pAF-MCS2, harboring the A3 actin promoter, a multiple cloning site, and the polyadenylation signal derived from the *Bombix mori* fibroin H-chain gene, for the subcloning of ligand molecules. First, the promoter and polyadenylation signal were excised from pA3Fb-Luc, kindly provided by H. Bando (Hokkaido University, Sapporo, Japan), and inserted into pUC18. To generate pAFgp64, the gp64 gene was excised from pFBgp64 (see below) by digestion with SalI and HindIII. This fragment was then inserted into the SalI-HindIII site of pAF-MCS1. Recombinant baculoviruses were constructed using the transfer vector pFASTBAC1 (Invitrogen). To measure the expression of foreign genes in mammalian cells, the firefly luciferase gene under the control of the CAG promoter (43) was subcloned into pFASTBAC1. To construct the transfer vector pFBCALuc, the CAG-luciferase cassette was excised from pCAGLuc (61) by digestion with SalI, extension with Klenow enzyme, and redigestion with BamHI and inserted into the SnaBI-BamHI site of pFASTBAC1.

pUCgp64locus was generated by cloning the EcoRI-SmaI fragment from AcMNPV genomic DNA (corresponding to 107,325 to 112,041 nt) (3) into the EcoRI-SmaI site of pUC18. To generate pUCgp64, a fragment encoding the gp64 gene was excised from pUCgp64locus by digestion with SpeI and BglII and then cloned into the XbaI-BamHI site of pUC18. The gp64 gene was excised from pUCgp64 by digestion with SalI and KpnI and inserted into the SalI-KpnI site of pFASTBAC1. The resulting plasmid was designated pFBgp64. To generate pFBgp64CALuc, the cassette including the polyhedrin promoter and the gp64 gene was excised from pFBgp64 by digestion with SnaBI and KpnI and cloned into pFBCALuc, which was digested with SalI, extended with Klenow enzyme, and redigested with KpnI. The VSVG gene fragment was excised from pCAG-VSVG (64) by digestion with EcoRI and cloned into the EcoRI site of pFASTBAC1 to create pFBVSVG. pFBGFP was constructed by excision of the GFP gene from pAcVSVG-CAGFP (65) by digestion with EcoRI and subsequent insertion into the EcoRI site of pFASTBAC1. To generate pFBVSVGCALuc and pFBGFPALuc, the DNA fragment encoding the polyhedrin promoter and either the VSVG or GFP gene was excised from pFBVSVG or pFBGFP, respectively, by digestion with SnaBI and XhoI and cloned into pFBCALuc, which was digested with SalI, extended with Klenow enzyme, and redigested with XhoI.

cDNAs encoding human CD46 and signaling lymphocyte activation molecule (SLAM; also known as CDw150) were amplified from the genomic DNAs of CHO/CD46 (kindly provided by T. Seya) (25) and CHO.SLAM (kindly provided by Y. Yanagi) (67) cells, respectively, by PCR. The CD46-Fw (1st) (5'-TTT CCTCCGGAGAAATAACAGC-3') and CD46-Rv (1st) (5'-CTAAGCCAC AGTTGCACTCATG-3') primers were used to amplify CD46 cDNA, and the SLAM-Fw (1st) (5'-TGACACGAAGCTTGCTTCTG-3') and SLAM-Rv (1st) (5'-GTCCGACCTTGTGGTCTCTGGTG-3') primers were used to amplify SLAM cDNA. These PCR products were used as templates for a second PCR with the primers CD46-Fw-HindIII (5'-CCCCAAGCTTCGCGCCGCG CATGGG-3') and CD46-Rv-SalI (5'-TTTTGTGCACTCAGCCTCTCTGCTC TGCTG-3') to amplify CD46 cDNA and SLAM-Fw-HindIII (5'-CCCCAAGC TTCCTCATTGGCTGATGGATC-3') and SLAM-Rv-SalI (5'-AAAAAGTCGA CTCAGCTCTCTGGAAGTGCA-3') to amplify SLAM cDNA. The amplified CD46 and SLAM cDNAs were digested with HindIII and SalI and then cloned into the HindIII-SalI sites of pAF-MCS2 to create pAFCD46 and pAFSLAM, respectively. The CD46 and SLAM cDNAs were excised from pAFCD46 and pAFSLAM, respectively, by digestion with HindIII, extension with Klenow enzyme, and redigestion with XbaI and cloned into pFASTBAC1. To generate pFBCD46CALuc and pFBSLAMCALuc, the DNA fragments encoding the polyhedrin promoter and either the CD46 gene or the SLAM gene were excised from pFBCD46 or pFBSLAM, respectively, by digestion with SnaBI and PvuII and cloned into pFBCALuc. A mutant SLAM gene, SLAMcyto7, possessing a truncated cytoplasmic domain of 7 amino acids, was generated by PCR with the primers SLAM-Fw-SmaI (5'-CCCCCGGGCCCTCATTGGCTGATGGATC-3') and SLAM-7aa-stop-Rv-SalI (5'-GGGGGGTCCACTCAGTTCGTTTT ACCTCTTCTCTCAAC-3'). This PCR product was digested with SmaI and SalI and then cloned into the SmaI-SalI sites of pAF-MCS1 to create pAFSLAMcyto7. To construct pFBSLAMcyto7CALuc, the SLAMcyto7 gene was excised from pAFSLAMcyto7 and substituted for the full-length SLAM gene of pFBSLAMCALuc. All plasmids containing PCR-derived sequences were confirmed by sequence analyses. For infection with pseudotype baculoviruses bearing CD46 or SLAM, we transfected target cells with expression plasmids encoding either the hemagglutinin and fusion proteins of the Edmonston strain (EdH and EdF) or those of the Ichinose strain (IcH and IcF) measles viruses. The pCA-EdH, pCA-EdF, pCA-IcH, and pCA-IcF plasmids were kindly provided by K. Takeuchi (63).

**Construction of pseudotype baculoviruses.** The gp64 gene of the AcMNPV-bacmid (bMON14272; Invitrogen) was replaced with the chloramphenicol acetyltransferase (CAT) gene as described previously with slight modifications (5, 33). Briefly, the CAT gene was amplified by PCR with the Chl-Fw-SpeI (5'-GGAC TAGTCCGAATAAATACCTGTGACGG-3') and Chl-Rv-BglII (5'-GAAG ATCTCGCAATTATTACCTCCACGG-3') primers using the pBT plasmid (Stratagene, La Jolla, Calif.) as a template. Following digestion with SpeI and BglII, the amplified CAT gene replaced the gp64 gene of pUCgp64locus to create p64locus/cat. To construct a gp64-null AcMNPV-bacmid, bMONΔ64/cat, the p64locus/cat plasmid was linearized by digestion with NdeI and cotransfected with bMON14272 into Sf9 cells. Forty-eight hours posttransfection, the cells were washed with cold phosphate-buffered saline and lysed in proteinase K buffer (50 mM Tris-HCl [pH 7.4], 100 mM NaCl, 1 mM EDTA, and 0.5% sodium dodecyl sulfate [SDS]). DNA was purified from cell lysates by phenol-chloroform extraction and then transformed into *Escherichia coli* DH10B competent cells (Invitrogen) by electroporation using a Gene Pulser (Bio-Rad, Hercules, Calif.). Resistant colonies were selected in kanamycin and chloramphenicol. Disruption of the gp64 gene was confirmed by PCR in a bMON14272-transformed colony that was resistant to kanamycin and chloramphenicol (Fig. 1A). To generate DH10BacΔ64/cat, we cotransfected bMONΔ64/cat and the helper plasmid pMON7124 into DH10B cells. To construct recombinant bacmids, DH10BacΔ64/cat was transformed with transfer vectors and selected according to the manufacturer's instructions. To separate recombinant bacmids from the pMON7124 helper plasmid, miniprep bacmid DNA was transformed into DH10B cells by electroporation. To generate pseudotype baculoviruses, bacmids lacking the gp64 gene and possessing both an exogenous ligand gene and the luciferase gene under the polyhedrin and CAG promoters, respectively, were transfected into Sf9 cells. Fifteen micrograms of the bacmid DNA was used to transfect  $5 \times 10^6$  Sf9 cells in a 10-cm-diameter dish by using 30 µl of UniFactor reagent (B-Bridge). Four days after transfection, 500 ml of culture supernatants (50 dishes) was harvested. The resulting pseudotype baculoviruses, AcΔ64/gp64/CALuc, AcΔ64/VSVG/CALuc, AcΔ64/CD46/CALuc, AcΔ64/SLAM/CALuc, and AcΔ64/SLAMcyto7/CALuc, were concentrated ~2,000 times by ultracentrifugation as described previously (66). The number of virus particles was determined from the signal intensity by Western blotting for the capsid protein vp39. Although both AcΔ64/gp64/CALuc and AcΔ64/VSVG/CALuc infected and repli-




FIG. 1. (A) Schematic representations of the gp64 loci of the AcMNPV (bMON14272) and gp64-null AcMNPV (bMON $\Delta$ 64/cat) bacmids. The gp64 gene (BglII/SpeI fragment corresponding to 108,033 to 109,761 nt) (3) of bMON14272 was replaced with the CAT gene by homologous recombination. The arrows indicate the locations of the PCR primers within the gp64 loci. (B) Construction of the recombinant bacmid bMON $\Delta$ 64/ligand/CAluc. The gp64 gene in bMON14272 was replaced with the CAT gene. The desired ligand and luciferase genes were inserted under the control of the polyhedrin and CAG promoters, respectively, within the polyhedrin locus. (C) The bacmids bMON $\Delta$ 64/ligand/CAluc and bMON14272 and plasmids containing the gp64 locus, pUC/gp64 locus, and p64 locus/cat (the gp64 locus with the CAT gene replacement) were amplified by PCR using primers specific for gp64, the gp64 locus, and vp39, a nucleocapsid protein of AcMNPV used as an internal control. Primers for gp64 and vp39 amplified fragments of 1,558 and 900 bp, respectively. The gp64 locus primers generated 2,741- and 2,143-bp fragments corresponding to the wild-type gp64 locus and the mutant locus with the CAT gene replacement shown in panel A, respectively. (D) Sf9p64 and Sf9 cells were transfected with bMON $\Delta$ 64/GFP/CAluc. GFP expression was examined by fluorescence microscopy 4 days posttransfection.



cated in Sf9 cells, it was not possible to determine the infectivity in the cases of the pseudotype baculoviruses possessing ligands incapable of entering into insect cells. To standardize the viral titer, we determined the amount of viral capsid protein vp39 by semiquantitative Western blot analysis. The infectious titer determined by plaque assay in Sf9 cells correlated well with the intensity of the vp39 signal obtained by Western blotting for both AcΔ64/gp64/CALuc and AcΔ64/VSVG/CALuc (data not shown). Pseudotype baculovirus titers are expressed as relative infectious units (RIU) in this study. To confirm the absence of gp64 in the bacmids, we synthesized oligonucleotide primers specific for the gp64 gene, the gp64 locus, and the vp39 gene as follows: for the gp64 gene, gp64-Fw (Bgl) (5'-AAAGATCTACCATGGTAAGCGCTATTGTTT-3') and gp64-Rv (Sal) (5'-TTGTGCGACTTAATATTGTCTATTACGGTTT-3'); for the gp64locus, gp64locus-Fw (5'-GCACGGATTGGGGAGAGGACGGATTTT-3') and gp64locus-Rv (5'-AGCTCGTTATTCAGGTGCCCGCTAC-3'); and for vp39, vp39-Fw (5'-ATATGCCGCTAGTGGCCCGTGGGTATGG-3') and vp39-Rv (5'-GACGGCTATTCCTCACTGCTGCTG-3'). PCR amplification was performed using *Taq* DNA polymerase (Invitrogen) according to the manufacturer's protocol.

**Stability of pseudotype baculoviruses during passage in Sf9 cells.** Culture supernatants from Sf9 cells transfected with recombinant bacmids were harvested 4 days after transfection. After serial passage in Sf9 cells for 4 days, each Sf9 cell supernatant was inoculated into Sf9 cells. The culture supernatants were further inoculated into Sf9 cells to examine the generation of replication-competent revertants during the replication in Sf9 cells. The presence of replication-competent virus in the culture supernatants was assessed by the appearance of cytopathic effect and GFP expression in Sf9 cells. GFP expression in insect cells was observed by fluorescence microscopy (UFX-II; Nikon, Tokyo, Japan). The generation of replication-competent viruses incorporating gp64 was examined by PCR using the viral DNA as a template. The supernatants of Sf9 cells were concentrated by centrifugation at  $18,000 \times g$  for 45 min at 4°C. Viral DNA, purified from replication-competent revertants by phenol-chloroform extraction, was examined by Southern blot analysis. DNA was digested with *Bgl*II or *Pst*I, separated by electrophoresis on a 0.6% agarose gel, and transferred to a Hybond N+ nylon membrane (Amersham Biosciences, Piscataway, N.J.). PCR primers [gp64-Fw (Bgl) and gp64-Rv (Sal) for the gp64 gene or vp39-Fw and vp39-Rv for the vp39 gene] were used to amplify the target fragments for use as hybridization probes. PCR products were purified and labeled using the ECL direct nucleic acid labeling and detection system (Amersham Biosciences) according to the manufacturer's instructions. Fragments containing the gp64 or vp39 gene were visualized using image analyzer LAS-3000 (Fujifilm, Tokyo, Japan).

**Incorporation of ligands into pseudotype particles.** To examine the expression of ligand proteins in insect cells or the incorporation of the ligands into pseudotype particles, cell lysates or purified baculoviruses were separated by SDS-polyacrylamide gel electrophoresis and electroblotted onto Hybond-P polyvinylidene difluoride membranes (Amersham Bioscience). After being blocked in phosphate-buffered saline containing 5% skim milk and 0.05% Tween 20 (Sigma), the membranes were incubated at room temperature for 1 h with a rabbit polyclonal anti-CD46 antibody (H-294; 1:200) (Santa Cruz, Santa Cruz, Calif.) or one of the following mouse monoclonal antibodies: anti-gp64 (AcV5; 1:1,000) (kindly provided by P. Faulkner) (22), anti-VSVG (P5D4; 1:2,000) (Sigma), anti-SLAM (123317; 1:200) (R&D systems, Minneapolis, Minn.), or anti-vp39 (236; 1:2,000) (kindly provided by G. F. Rohrmann) (51). The membranes were then incubated in horseradish peroxidase-conjugated anti-mouse IgG or anti-rabbit IgG antibodies at room temperature for 1 h. Immunoreactive bands were visualized using enhanced-chemiluminescence Super Signal West Femto substrate (Pierce, Rockford, Ill.) (47).

**Reporter gene expression by pseudotype baculoviruses.** AcΔ64/gp64/CALuc and AcΔ64/VSVG/CALuc baculoviruses were inoculated into  $3.0 \times 10^4$  293T and BHK cells. Twenty-four hours after infection, the cells were lysed in Bright-Glo luciferase substrate (Promega, Madison, Wis.) according to the manufacturer's instructions. Relative light units were measured using a luminometer (AB-2200; ATTO Co. Ltd., Tokyo, Japan). To demonstrate ligand-directed gene targeting by AcΔ64/CD46/CALuc, AcΔ64/SLAM/CALuc, and AcΔ64/SLAMcyto7/CALuc baculoviruses,  $3.0 \times 10^4$  BHK cells were cotransfected with either pCA-EdF and pCA-EdH or pCA-IcF and pCA-IcH and then infected with  $5.0 \times 10^6$  RIU of pseudotype baculoviruses at 24 h posttransfection. Luciferase expression was determined after a 24-h incubation.

**Inhibition of gene transduction by specific antibodies against ligands.** To examine ligand-directed gene transduction by pseudotype baculoviruses, we examined the neutralization of gene transduction by antibodies specific for the ligands presented by the pseudotypes. The appropriate dilutions of anti-gp64 (AcV1), anti-VSVG (I1) (kindly provided by M. A. Whitt) (30), anti-CD46

(M75) (Seikagaku Co. Ltd., Tokyo, Japan), or anti-SLAM (IPO-3) (Biodesign International, Saco, Maine) antibodies were preincubated with each virus ( $10^6$  RIU) at 37°C for 60 min and then inoculated into the appropriate target cells. After incubation at 37°C for 24 h, we determined the neutralization by the included antibodies from the reduction of luciferase expression.

**Entry of pseudotype baculovirus into target cells.** BHK cells expressing hemagglutinin and fusion proteins derived from the Edmonston strain of measles virus were preincubated with either ammonium chloride (2, 10, or 50 mM) (Wako Pure Chemical Industries, Osaka, Japan) or chloroquine (20, 100, or 500 μM) (Sigma) for 1 h. The cells were then inoculated with  $1.0 \times 10^6$  RIU of AcΔ64/CD46/CALuc, AcΔ64/gp64/CALuc, or AcΔ64/VSVG/CALuc in the presence of the above-mentioned reagents. The effects of ammonium chloride and chloroquine on gene transduction by pseudotype baculoviruses were determined by the changes in luciferase expression.

**Electron microscopy.** Viral particles purified by ultracentrifugation as described above were put onto carbon-coated copper 400-mesh electron microscopy grids for 15 min. After being washed in water, the grids were negatively stained with 1% (wt/vol) uranyl acetate and examined using a Hitachi (Tokyo, Japan) H-7100 electron microscope at 75 kV. For immunoelectron microscopy, virus particles put onto grids were incubated with murine monoclonal antibodies specific for VSVG (I1) or CD46 (E4.3) (Santa Cruz) and then treated with a gold particle-conjugated anti-mouse IgG antibody (British Biocell International, Ltd., Cardiff, United Kingdom). Samples were stained and observed as described above.

## RESULTS

**Construction of recombinant AcMNPV lacking the gp64 gene.** The gp64 gene of the AcMNPV bacmid, bMON14272, was replaced with the CAT gene by homologous recombination in Sf9 cells using a modification of the methods reported by Bideshi and Federici (5) and Lung et al. (33) (Fig. 1A). We cotransfected bMON14272 and a linearized p64locus/cat plasmid bearing the CAT gene in place of the gp64 gene into Sf9 cells. DNA, extracted from the cells 48 h after transfection, was then transformed into competent DH10B cells. The disruption of the gp64 gene in colonies selected with kanamycin and chloramphenicol was confirmed by PCR (data not shown). We also constructed a recombinant bacmid, bMONΔ64/GFP/CALuc, which contained the insertion of the GFP gene under the control of the polyhedrin promoter and the luciferase gene under the control of the CAG promoter into the polyhedrin locus of the gp64-null bacmid (Fig. 1B). Disruption of gp64 in bMONΔ64/GFP/CALuc was confirmed by PCR using a series of specific primers (Fig. 1C). PCR with primers specific for the vp39 gene, used as an internal control for the AcMNPV bacmid, amplified a 900-bp product from both the bMONΔ64/GFP/CALuc and parent bMON14272 bacmids. The gp64 gene (1,558 bp) was amplified from bMON14272 and pUC/p64locus, but not from bMONΔ64/GFP/CALuc and p64locus/cat. The 2,741- and 2,143-bp fragments corresponding to the wild-type and mutant gp64 genes, respectively, were amplified using gp64 locus-specific primers. The wild-type gene was amplified from bMON14272 and pUC/p64locus, while the mutant gene was amplified from bMONΔ64/GFP/CALuc and p64locus/cat (Fig. 1A and C). These data indicate that the gp64 gene was replaced with the cat gene in bMONΔ64/GFP/CALuc. Previous studies demonstrated that gp64-null AcMNPV could propagate in Sf9<sup>OP64-6</sup> or Sf9<sup>Op1D</sup> cell lines constitutively expressing the gp64 protein of *Orgyia pseudotsugata* NPV (OpNPV) but not in untransfected Sf9 cells (40, 49). We then established a cell line, Sf9gp64, constitutively expressing the gp64 gene derived from AcMNPV. The pAFgp64 plasmid, carrying the gp64 gene of AcMNPV without any flanking sequence, was used to

avoid homologous recombination between the viral genome and the plasmid. To examine the replication competency of gp64-null AcMNPV (Ac $\Delta$ 64/GFP/CAIuc), Sf9 cells were transfected with bMON $\Delta$ 64/GFP/CAIuc. We assessed the propagation of infectious virus by measuring GFP expression by fluorescence microscopy. Forty-eight hours posttransfection, ~10% of the Sf9 cells were GFP positive (data not shown). While Sf9 cells exhibited the spread of infection 96 h posttransfection, Sf9 cells did not (Fig. 1D). These data indicate that Ac $\Delta$ 64/GFP/CAIuc can replicate only in Sf9 cells, not in Sf9 cells.


**Appearance of revertants incorporating the gp64 gene during replication in Sf9 cells.** To determine the stability of Ac $\Delta$ 64/GFP/CAIuc during replication in Sf9 cells, we serially passaged Ac $\Delta$ 64/GFP/CAIuc in Sf9 cells. Culture supernatants of Sf9 cells collected 4 days after transfection with bMON $\Delta$ 64/GFP/CAIuc (passage 1) were inoculated into Sf9 cells. The supernatants were further passaged in Sf9 cells for 4 days. To examine the appearance of replication-competent viruses, the culture supernatants from each passage were inoculated into Sf9 cells. At 4 days postinfection, we examined GFP expression in Sf9 cells by fluorescence microscopy (Fig. 2A). The expression of GFP was observed in Sf9 cells inoculated with Sf9 culture supernatants, irrespective of the passage history. As gp64-negative Ac $\Delta$ 64/GFP/CAIuc baculovirus only transiently carries gp64, progeny viruses produced in Sf9 cells should not be infectious. The supernatants of Sf9 cells inoculated with supernatants recovered after >3 passages (passages 3, 4, and 5) with Sf9 cells exhibited infectivity to Sf9 cells, suggesting the generation of replication-competent revertants incorporating the gp64 gene into the viral genome. To confirm the incorporation of gp64 into the viral genome, virus particles were purified from the supernatants of each Sf9 passage. The presence of the gp64 gene within the viral genome was determined by PCR. We detected the gp64 gene in viruses obtained from the culture supernatants of passages 3, 4, and 5 but not in those from the first and second passages (Fig. 2B). Furthermore, PCR amplification of viral DNA with the gp64 locus-specific primers revealed that a 2,143-bp fragment, corresponding to the mutant form, was detected in the genome of Ac $\Delta$ 64/GFP/CAIuc, while a 2,741-bp fragment, corresponding to the wild-type form, was amplified from Ac14272, irrespective of the number of passages. These results confirmed that the emergence of replication-competent virus during the passage in Sf9 cells is not due to the contamination of the parental virus, Ac14272. The recombinant virus incorporated the gp64 gene into the Ac $\Delta$ 64/GFP/CAIuc genome during propagation in Sf9 cells.

Plasmid DNA can be integrated into multiple sites within the viral genome by nonhomologous recombination upon cotransfection of plasmid DNA with the baculovirus genome in Sf9 cells (71). To determine if gp64 genes integrated into the baculovirus genome by nonhomologous recombination during propagation in Sf9 cells, we analyzed the DNAs of three independent revertant viruses by PCR and Southern blot analyses. Viral DNA was extracted from these revertant viruses and analyzed by PCR as described above (Fig. 3A). We detected the gp64 gene in all revertant viruses and bMON14272 but not in the parental bacmid, bMON $\Delta$ 64/GFP/CAIuc. The gp64 locus primers amplified the mutant 2,143-bp fragment

from all revertant viruses and the parental bMON $\Delta$ 64/GFP/CAIuc bacmid, not the 2,741-bp wild-type fragment that could be amplified from bMON14272. These results confirmed that the three independent revertant viruses, instead of deriving from contaminating wild-type virus, had incorporated the gp64 gene into their genomes exogenously. DNA from the revertants was digested with BglII or PstI, which do not digest sequences within the gp64 or vp39 genes, and hybridized to gp64- or vp39-specific probes (Fig. 3B). If the gp64 gene integrated into the viral genome by nonhomologous recombination, the digested fragments containing the gp64 gene would be of different sizes. Following digestion with BglII, the DNA fragments containing the gp64 gene in the revertants differed in size from each other (Fig. 3B, lanes 3 to 5). When digested with PstI, the sizes of the fragments containing the gp64 gene were similar in revertant clones 2 and 3 (Fig. 3B, lanes 9 to 10), indicating that the gp64 gene may have integrated into nearby sites in the viral genomes of clones 2 and 3. The fragment containing the gp64 gene in revertant clone 1 following digestion with either BglII or PstI was similar to that seen in bMON14272 (Fig. 3B lanes 3 and 8). These results, however, were not due to contamination with bMON14272, as the PCR analysis demonstrated that the gp64 locus of revertant clone 1 was of the mutant type (Fig. 3A). These data suggested that the gp64 gene integrated into the virus genomes of the revertants by nonhomologous recombination. As an internal control, the vp39 gene was detected in fragments of the predicted sizes (31,975 bp when digested with BglII and 29,009 bp when digested with PstI) in all viruses. To determine the sites of integration of the gp64 gene in the genomes of the revertants, we tried to sequence from within the gp64 gene out into the baculovirus genome by using an internal gp64 primer. In revertant 2, the sequences including the actin promoter and the gp64 gene were detected upstream of the polyhedrin promoter, where no homologous sequence was observed. In revertants 1 and 3, however, sequence analyses by the internal primer obtained only sequences of pAFgp64 and could not reach the integration site, due to a large insertion of the plasmid sequence (data not shown).

**Characterization of pseudotype baculovirus carrying VSVG.** Previous studies demonstrated that the gp64 protein plays a critical role in infection of various mammalian cells, as well as insect cells (66). To determine if the pseudotype baculoviruses bearing foreign viral envelope proteins in place of gp64 can infect and express foreign genes within mammalian cells, we constructed a gp64-null pseudotype virus, Ac $\Delta$ 64/VSVG/CAIuc, by the transfection of bMON $\Delta$ 64/VSVG/CAIuc, which encodes the VSVG gene under the control of the polyhedrin promoter and the luciferase gene under the control of the CAG promoter, into Sf9 cells (Fig. 1B). As a control, we also generated Ac $\Delta$ 64/gp64/CAIuc, in which the gp64 gene under the control of the polyhedrin promoter replaced the VSVG gene in the above-mentioned virus. Sf9 cells were transfected with appropriate bacmids and incubated for 4 days. The pseudotype baculoviruses in the culture supernatants were concentrated and purified by ultracentrifugation ( $10^8$  to  $10^9$  RIU/ml). To examine the expression and incorporation of the glycoproteins into virions, we transfected these bacmid constructs into Sf9 cells. The cell lysates and the purified virus particles were examined by Western blot analysis (Fig. 4A). VSVG and gp64

A



B




FIG. 2. Appearance of replication-competent viruses incorporating the gp64 gene during passage in Sf9 cells. (A) Sf9 cells were transfected with bMON $\Delta$ 64/GFP/CAIuc. Culture supernatants were harvested 4 days after transfection and then serially passaged in Sf9 cells at 4-day intervals. Each culture supernatant from Sf9 cells was passaged two more times in Sf9 cells to detect the appearance of replication-competent viruses. GFP expression in Sf9 cells was examined by fluorescence microscopy 4 days after infection. (B) PCR analysis of purified virus particles from the supernatant of each Sf9 cell passage. The gp64 gene was detectable in particles obtained from the third or later passages. The numbers above the lanes represent the passage numbers. The bMON $\Delta$ 64/GFP/CAIuc and bMON14272 bacmids and Ac14272, generated from bMON14272 and passaged in Sf9 cells, were used as controls. M is the culture supernatant of uninfected Sf9 cells concentrated under the same conditions as the virus particles. The primers amplified fragments as detailed in the legend to Fig. 1.

were expressed in the cells transfected with the appropriate bacmids. The proteins were also detected in the purified Ac $\Delta$ 64/VSVG/CAIuc and Ac $\Delta$ 64/gp64/CAIuc viruses, respectively, but not in Ac $\Delta$ 64/GFP/CAIuc.

To assess the efficacy of mammalian cell gene transduction by the pseudotype baculoviruses, 293T and BHK cells were inoculated with various amounts of pseudotype viruses (Fig.

4B). Similar levels of reporter gene expression were observed in a dose-dependent manner in both cell lines following infection with Ac $\Delta$ 64/gp64/CAIuc and Ac $\Delta$ 64/VSVG/CAIuc. Ac $\Delta$ 64/GFP/CAIuc, however, was unable to infect either cell line. To confirm the role of gp64- or VSVG-mediated gene transduction into mammalian cells by the pseudotype baculoviruses, we attempted to neutralize 293T cell infection using





FIG. 3. Incorporation of the gp64 gene into gp64-null baculovirus genomes by nonhomologous recombination. (A) PCR analysis of three independent revertant viruses. In each revertant virus, the gp64 gene and gp64 locus primer pairs produced 1,558- and 2,143-bp fragments, respectively, indicating the presence of the mutant gp64 locus. (B) Southern blot analysis of revertant viruses. Viral DNA was digested with BglII or PstI, separated, and hybridized to gp64- or vp39-specific probes. Fragments containing the gp64 gene were detectable in all of the revertant viruses, but the fragment sizes differed. The vp39 gene, used as an internal control, was detectable in all revertant DNAs and bacmids. The numbers above the lanes represent the revertant clones. The bMONΔ64/GFP/CAIuc and bMON14272 bacmids were used as controls.

specific monoclonal antibodies against gp64 and VSVG. Luciferase expression in 293T cells infected with either AcΔ64/gp64/CAIuc or AcΔ64/VSVG/CAIuc was specifically inhibited by antibodies against gp64 or VSVG, respectively (Fig. 4C). These results indicate that reporter gene expression in mammalian cells inoculated with pseudotype baculoviruses relies on


the interactions mediated by the ligand proteins on the viral particles.

**Ligand-directed gene targeting by pseudotype baculovirus.** To demonstrate the ligand-directed gene transduction of target cells by pseudotype baculoviruses, we constructed pseudotype viruses bearing CD46 or SLAM in place of the gp64

A



B



C




FIG. 4. Characterization of pseudotype baculoviruses bearing VSVG. (A) VSVG and gp64 expression in Sf9 cells transfected with the bMON $\Delta$ 64/gp64/CAIuc, bMON $\Delta$ 64/VSVG/CAIuc, bMON $\Delta$ 64/GFP/CAIuc, or bMON14272 bacmid were examined by Western blot analysis using monoclonal antibodies specific for VSVG (P5D4) and gp64 (AcV5) (left). The incorporation of gp64 and VSVG into pseudotype particles, Ac $\Delta$ 64/gp64/CAIuc, Ac $\Delta$ 64/VSVG/CAIuc, Ac $\Delta$ 64/GFP/CAIuc, or AcMNPV, was examined by Western blot analysis using monoclonal antibodies specific for gp64, VSVG, and vp39 (236) (right). (B) Gene transduction into mammalian cells by pseudotype baculoviruses. 293T or BHK cells ( $3 \times 10^5$ ) were inoculated with various amounts of Ac $\Delta$ 64/gp64/CAIuc, Ac $\Delta$ 64/VSVG/CAIuc, or Ac $\Delta$ 64/GFP/CAIuc. The pseudotype titers are expressed as RIU. Luciferase expression was determined 24 h after infection. The results shown are the means of three independent assays, while the error bars represent the standard deviations. RLU, relative light units. (C) Neutralization of gene transduction into mammalian cells by pseudotype baculoviruses by antibodies specific for the particle ligands. Ac $\Delta$ 64/gp64/CAIuc or Ac $\Delta$ 64/VSVG/CAIuc ( $10^6$  RIU) was preincubated with the indicated dilutions of monoclonal antibodies specific for gp64 (AcV1) or VSVG (I1), respectively, for 60 min at 37°C. Residual activity, determined as luciferase expression in 293T cells 24 h postinfection, is expressed as the relative percentages of expression. The results shown are the means of three independent assays, with the error bars representing the standard deviations.

protein. The receptor usage of measles virus has been well characterized; while laboratory strains of measles virus, such as the Edmonston strain, can use either CD46 or SLAM as receptors, wild-type strains, such as the Ichinose strain, can only use SLAM for entry (15, 24, 42, 48, 67). Expression of these receptor molecules in Sf9 cells transfected with the bMON $\Delta$ 64/CD46/CALuc or bMON $\Delta$ 64/SLAM/CALuc bacmid (Fig. 1B) and subsequent incorporation of the receptors into progeny particles (Ac $\Delta$ 64/CD46/CALuc and Ac $\Delta$ 64/SLAM/CALuc, respectively) were confirmed by Western blotting (Fig. 5A). CD46 was detected in cells transfected with bMON $\Delta$ 64/CD46/CALuc and in the purified particles of Ac $\Delta$ 64/CD46/CALuc, whereas SLAM was detected in cells transfected with the bacmid but not in the particles of Ac $\Delta$ 64/SLAM/CALuc.

The gp64, CD46, and SLAM proteins are all type I membrane proteins. SLAM has a 77-amino-acid cytoplasmic domain (23), while gp64 and CD46 have only 7- and 33-amino-acid tails, respectively (49, 60). Therefore, we speculated that SLAM may be only inefficiently incorporated into baculovirus particles, due to its large cytoplasmic domain. To examine the effect of the cytoplasmic domain length on incorporation into baculovirus particles, we constructed a mutant SLAM molecule, SLAMcyto7, with a deletion in the C-terminal cytoplasmic domain that preserves only the seven membrane-proximal amino acids. Western blot analysis revealed that SLAMcyto7 was efficiently expressed in Sf9 cells transfected with bMON $\Delta$ 64/SLAMcyto7/CALuc and subsequently incorporated into Ac $\Delta$ 64/SLAMcyto7/CALuc particles at levels similar to those seen for CD46 inclusion into Ac $\Delta$ 64/CD46/CALuc (Fig. 5A).

To determine the efficiency of ligand-directed gene delivery, BHK cells were cotransfected with expression plasmids encoding the measles virus H and F glycoproteins of the Edmonston (EdH and EdF) or Ichinose (IcH and IcF) strain. These cells were inoculated with pseudotype baculoviruses (Fig. 5B). Ac $\Delta$ 64/CD46/CALuc exhibited gene delivery specifically to cells expressing EdH and EdF, but not IcH and IcF. Although the efficiency of gene transduction was 10 times lower than that seen with Ac $\Delta$ 64/CD46/CALuc, Ac $\Delta$ 64/SLAMcyto7/CALuc could also deliver a reporter gene to cells expressing the Edmonston and Ichinose strain glycoproteins but not to control cells. While Ac $\Delta$ 64/gp64/CALuc and Ac $\Delta$ 64/VSVG/CALuc could effectively deliver a reporter gene to all of the cells examined, Ac $\Delta$ 64/SLAM/CALuc was ineffective against all of the cell lines tested, likely due to the lack of SLAM incorporation into the virions.

To confirm ligand-directed gene delivery by Ac $\Delta$ 64/CD46/CALuc and Ac $\Delta$ 64/SLAMcyto7/CALuc to cells expressing appropriate measles virus glycoproteins, we tested the neutralization of gene transduction by specific monoclonal antibodies against CD46 and SLAM (Fig. 5C). Gene transduction of target cells by either Ac $\Delta$ 64/CD46/CALuc or Ac $\Delta$ 64/SLAMcyto7/CALuc, but not by Ac $\Delta$ 64/gp64/CALuc, could be inhibited in a dose-dependent manner by anti-CD46 and anti-SLAM monoclonal antibodies, respectively. These results indicate that pseudotype baculoviruses can deliver foreign genes to target cells in a ligand-directed manner.

**Entry pathway of the pseudotype baculoviruses.** Virus entry occurs either by the direct fusion of viral envelope proteins with the host plasma membrane at neutral pH, as seen for measles virus, or following receptor-mediated endocytosis, as seen for AcMNPV and VSV, in which envelope glycoproteins

undergo conformational changes into a fusion-competent state, leading to fusion between viral and host membranes at low pH within endosomes (6, 36, 69). Ammonium chloride and chloroquine, which inhibit endosomal acidification, have been used as entry inhibitors for viruses that penetrate cells through receptor-mediated endocytosis (7). To examine the entry pathways used by the pseudotype baculoviruses, we examined the infectivity of Ac $\Delta$ 64/gp64/CALuc, Ac $\Delta$ 64/VSVG/CALuc, and Ac $\Delta$ 64/CD46/CALuc to BHK cells expressing EdH and EdF in the presence or absence of ammonium chloride or chloroquine (Fig. 6). Although these compounds inhibited gene transduction of BHK cells inoculated with Ac $\Delta$ 64/gp64/CALuc or Ac $\Delta$ 64/VSVG/CALuc in a dose-dependent manner, gene transduction by Ac $\Delta$ 64/CD46/CALuc was not inhibited. In contrast, ammonium chloride treatment enhanced gene expression following Ac $\Delta$ 64/CD46/CALuc infection. These results indicate that the pseudotype baculoviruses utilize entry pathways conferred by the nature of the ligand protein replacing gp64.

**Morphology of pseudotype baculovirus.** To address any alterations in pseudotype baculovirus morphology, we examined the AcMNPV, Ac $\Delta$ 64/VSVG/CALuc, and Ac $\Delta$ 64/CD46/CALuc virus particles by transmission electron microscopy (Fig. 7A to C). All of the pseudotype baculoviruses exhibited rod shapes and similar sizes, indistinguishable from the wild-type baculovirus. To examine the incorporation of exogenous ligands into the virion, purified Ac $\Delta$ 64/VSVG/CALuc and Ac $\Delta$ 64/CD46/CALuc particles were examined by immunoelectron microscopy using specific monoclonal antibodies against VSVG and CD46 (Fig. 7E and F). In both pseudotype viruses, gold particles were detected on the virion surface from the stalk to the head domains, indicating that VSVG and CD46 were incorporated into the Ac $\Delta$ 64/VSVG/CALuc and Ac $\Delta$ 64/CD46/CALuc virus particles, respectively.

## DISCUSSION

Baculovirus is a useful tool for gene delivery to mammalian cells due to the large capacity of the virus to incorporate foreign genes, the wide host range, and the lack of replication in mammalian cells, providing minimal toxicity (29, 53, 55, 61, 68). The gp64 envelope glycoprotein, involved in attachment to both insect and mammalian cells, is required for low-pH-triggered membrane fusion following endocytosis during virus entry (6, 8, 12, 14, 20, 21, 31, 40, 41, 66, 69, 70). We have previously demonstrated that the interaction of gp64 with cell surface phospholipids may be important in baculovirus infection of mammalian cells (66). As the recombinant baculoviruses used for gene delivery to mammalian cells in this system retained the gp64 envelope glycoprotein, it was difficult to deliver foreign genes to specific target cells.

To generate a ligand-directed gene-targeting baculovirus vector, we established a bacmid system to produce recombinant baculoviruses in which the gp64 gene was replaced with other ligand genes of interest. The gp64 protein is required for efficient budding from Sf9 cells; the budding of a mutant virus lacking gp64 was reduced to 2% of that seen for wild-type baculovirus (49). Although Ac $\Delta$ 64/gp64/CALuc, a pseudotype virus in which gp64 was reintroduced into the gp64 deletion mutant, incorporated two to three times as much gp64 protein as seen in wild-type baculovirus (Fig. 4A), the infectious titers




FIG. 5. Ligand-directed gene targeting by pseudotype baculoviruses. (A) The expression of gp64, CD46, SLAM, and SLAMcyto7 in Sf9 cells transfected with the bMON $\Delta$ 64/gp64/CALuc, bMON $\Delta$ 64/CD46/CALuc, bMON $\Delta$ 64/SLAM/CALuc, bMON $\Delta$ 64/SLAMcyto7/CALuc, or bMON14272 bacmid was examined by Western blot analysis using monoclonal antibodies specific for gp64 (AcV5), CD46 (H-294), and SLAM (123317), as indicated (left). The incorporation of gp64, CD46, SLAM, and SLAMcyto7 into pseudotype particles, Ac $\Delta$ 64/gp64/CALuc, Ac $\Delta$ 64/CD46/CALuc, Ac $\Delta$ 64/SLAM/CALuc, Ac $\Delta$ 64/SLAMcyto7/CALuc, or AcMNPV, was examined by Western blot analysis using monoclonal antibodies specific for gp64, CD46, SLAM, and vp39 (236), as indicated (right). (B) Ligand-directed gene targeting by pseudotype baculoviruses. BHK cells ( $3 \times 10^6$ ) were co-transfected with expression plasmids encoding measles virus H and F glycoproteins of either the Edmonston (EdH and EdF) or Ichinose (IcH and IcF) strain or with an empty vector and then inoculated with  $5 \times 10^6$  RIU of Ac $\Delta$ 64/gp64/CALuc, Ac $\Delta$ 64/CD46/CALuc, Ac $\Delta$ 64/SLAM/CALuc, Ac $\Delta$ 64/SLAMcyto7/CALuc, or Ac $\Delta$ 64/VSVG/CALuc 24 h after transfection. Luciferase expression was determined 24 h after infection. The results shown are the means of three independent assays, and the error bars represent the standard deviations. RLU, relative light units. (C) Neutralization of ligand-directed gene targeting by antibodies specific for viral ligands. Ac $\Delta$ 64/gp64/CALuc, Ac $\Delta$ 64/CD46/CALuc, or Ac $\Delta$ 64/SLAMcyto7/CALuc ( $10^6$  RIU) was preincubated with various concentrations of monoclonal antibodies specific for CD46 (M75) or SLAM (IPO-3) for 60 min at 37°C. Residual activity was determined by measurement of luciferase expression in BHK cells expressing the H and F glycoproteins of the Edmonston or Ichinose strain 24 h postinfection. The values are expressed as the relative percentages of expression. The results shown are the means of three independent assays, with the error bars representing the standard deviations.




FIG. 6. Effects of lysosomotropic reagents on gene transduction of pseudotype baculoviruses. BHK cells ( $3 \times 10^4$ ) transfected with expression plasmids encoding the measles virus H and F glycoproteins of the Edmonston strain (EdH and EdF) were pretreated with various concentrations of ammonium chloride or chloroquine for 60 min. The cells were then inoculated with  $10^6$  RIU of AcΔ64/gp64/CALuc, AcΔ64/VSVG/CALuc, or AcΔ64/CD46/CALuc in the presence of the lysosomotropic reagents. Luciferase expression was determined 24 h postinfection. The results shown are the averages of three independent assays, with the error bars representing the standard deviations.

of the virus, determined by plaque formation in Sf9 cells, were similar. These results suggest that, while the polyhedrin promoter is sufficient to overexpress and incorporate a ligand of interest into the virion, this is not necessarily the best choice to maintain ligand function. The discrepancy between gp64 incorporation and the infectious titer may be attributed to a limited capacity to incorporate functional gp64 into particles and the timing of ligand gene activation. As the polyhedrin promoter is activated in the late stage of infection, baculoviruses budding

in the early stage of infection may be unable to incorporate the ligand expressed by the polyhedrin promoter. Use of the immediate-early promoter for ligand expression may improve the efficiency of incorporation into virus particles. In support of this possibility, the infectious titer of a recombinant AcMNPV in which the gp64 gene was replaced with the F gene from *Lymantria dispar* NPV under the control of the polyhedrin promoter was ~60-fold lower than that of a virus with the F gene under the control of the gp64 promoter (33).




FIG. 7. Electron micrographs of pseudotype baculoviruses. Purified virus particles of wild-type AcMNPV (A), AcΔ64/VSVG/CALuc (B), and AcΔ64/CD46/CALuc (C) were examined by electron microscopy. A typical rod shape was visible in all of the pseudotype baculoviruses. The VSVG or CD46 proteins were observed on the surfaces of AcΔ64/VSVG/CALuc (E) and AcΔ64/CD46/CALuc (F) by immunoelectron microscopy using specific monoclonal antibodies against VSVG and CD46, respectively. AcMNPV treated with the monoclonal antibody against VSVG was used as a control (D). The bars on the panels represent 100 nm.



The replication competency in Sf9 cells of a gp64-null recombinant baculovirus could be rescued by incorporation of the VSVG gene (33, 34). We confirmed that the recombinant baculovirus deleted the gp64 gene and instead incorporated the VSVG gene under the control of the polyhedrin promoter, was replication competent in insect cells, and exhibited a high level of reporter gene transduction into 293T and BHK cells. The compatibility of VSVG in this system may result from similarities of the structural and functional characteristics of VSVG to those of gp64; both proteins are type I membrane glycoproteins, exist in trimeric complexes, and are capable of inducing membrane fusion at low pHs (6, 50, 54). Although the recombinant baculovirus in which gp64 is replaced with VSVG is able to replicate in insect cells and transduce foreign genes into a wide variety of mammalian cells, VSVG was suggested to recognize phosphatidylserine (58) or other ubiquitously expressed molecules other than phosphatidylserine (13) as a receptor(s), making it difficult to confer cell type specificity to gene delivery using VSVG pseudotype baculoviruses.

To establish ligand-directed gene delivery by pseudotype baculoviruses, it is necessary to propagate replication-deficient pseudotype baculoviruses possessing a ligand of interest in Sf9 cells, a replication-competent cell line stably expressing gp64. After three rounds of passage of the pseudotype virus in Sf9 cells, however, replication-competent revertant viruses that had incorporated the gp64 gene were generated. Sf9 cells were established by transfection of Sf9 with a plasmid encoding the gp64 gene lacking any flanking sequences. Southern blot analysis revealed that the gp64 gene was integrated into the genomes of the baculovirus revertants by nonhomologous recombination. Homologous recombination between the AcMNPV genome and either a transfer vector (32, 38), additional AcMNPVs (19), or *B. mori* NPV can occur in insect cells with high frequency (27, 28). In contrast, our data suggest that the revertant viruses were not generated by homologous recombination but by nonhomologous incorporation of the gp64 gene from the Sf9gp64 chromosome into the viral genome. Nonhomologous recombination between plasmid DNA and the baculovirus genome was previously reported upon cotransfection into insect cells (71). A gp64-null virus could propagate in Sf9<sup>Op1D</sup> cell lines constitutively expressing OpNPV gp64 without incorporating the gp64 gene into the viral genome (33, 34). One possible explanation for our result may be the difference between the gp64 genes of OpNPV and AcMNPV. Although the Sf9gp64 cell line was established by sorting cells expressing a high level of gp64 on the cell surface, the expression level of gp64 might be lower than that of the Sf9<sup>Op1D</sup> cell line, and the lower level of expression of gp64 might result in the selective amplification of revertants. To circumvent the high-frequency incorporation of foreign DNA into the baculoviral genome, we attempted the lipofection of recombinant bacmids into Sf9 cells instead of amplification of pseudotype baculoviruses in Sf9 cells. Although it is not possible to obtain a high pseudotype virus titer by this method, we can generate pure virus stocks without any contaminating replication-competent baculoviruses incorporating the gp64 gene. To prepare a high-titer stock of a replication-deficient pseudotype baculovirus carrying a foreign ligand, however, it is essential to propagate the pseudotype virus in Sf9 cells expressing gp64 without the transfection of plasmid DNA. Expression of gp64 by RNA trans-

fection, RNA viral vectors, or the RNA replicon system may be able to avoid the incorporation of gp64 DNA into the baculovirus genome.

We constructed a pseudotype baculovirus, AcΔ64/CD46/CALuc, bearing human CD46 in place of gp64 on viral particles. CD46 is a multifunctional protein involved in the infection of various microorganisms and the regulation of complement activation (10). CD46, also known as membrane cofactor protein, protects autologous cells from complement attack by serving as a cofactor for factor I-mediated inactivation of C3b and C4b, blocking the complement cascade at C3 activation (2). CD46 also serves as a receptor for human herpes virus 6 (56), group B adenovirus (17, 59, 62), bovine viral diarrhea virus (39), two bacterial strains (*Streptococcus pyogenes* and pathogenic *Neisseria*) (26, 46), and the Edmonston strain of measles virus (15, 42). In this study, we demonstrated that AcΔ64/CD46/CALuc exhibited specific reporter gene transfer to and expression in BHK cells expressing the measles virus H and F glycoproteins of the Edmonston strain but not those expressing the Ichinose strain glycoproteins that require SLAM as a receptor (67). Therefore, a CD46 pseudotype baculovirus bearing a suicide gene may be able to eliminate cells expressing pathogen ligands that utilize CD46 as a receptor. Furthermore, CD46 is frequently overexpressed on cancer cells, possibly serving as a mechanism to overcome lysis by complement (16). In support of the potential utility of this vector, the Edmonston strain of measles virus has a potent and selective oncolytic activity (52). CD46 pseudotype baculovirus may also be applicable for the clearance of tumor cells surviving oncolytic measles virus treatment.

Infections with AcΔ64/gp64/CALuc and AcΔ64/VSVG/CALuc, carrying gp64 and VSVG on the particles, respectively, could be decreased by treatment with ATPase inhibitors, chloroquine, or ammonium chloride, while AcΔ64/CD46/CALuc was resistant to treatment. This finding suggests the possibility of constructing a baculovirus vector capable of both targeting and modulating the viral entry pathway, as seen for VSV (7). This is different from influenza virus vectors, where acidic exposure within endosomes is critical for the dissociation of the matrix protein from the ribonucleocapsid (9, 37). In contrast to AcΔ64/CD46/CALuc, AcΔ64/SLAM/CALuc could not incorporate the full-length SLAM molecule into particles, preventing specific gene transduction. The inability to mediate gene delivery was not due to the absence of an interaction with specific targets, as Sf9 cells expressing SLAM induced membrane fusion with BHK cells expressing the H and F glycoproteins of either the Edmonston or Ichinose strain of measles virus (data not shown). The ligands incorporated efficiently into baculovirus particles, including gp64, VSVG, and CD46, all have relatively short cytoplasmic domains, measuring 7, 29, and 33 amino acids in length, respectively (49, 54, 60). In contrast, SLAM possesses a cytoplasmic tail of 77 amino acids (23). We therefore hypothesized that the length of the ligand cytoplasmic domain may be critical for efficient incorporation into baculovirus particles. AcΔ64/SLAMcyto7/CALuc, possessing a mutant SLAM molecule with the C-terminal 70 amino acids of the cytoplasmic domain deleted, efficiently incorporated the mutant SLAM into particles and exhibited specific gene delivery to BHK cells expressing the H and F glycoproteins of both measles virus strains. Although the mechanism by which SLAM

incorporation into baculovirus particles is enhanced by C-terminal truncation of the cytoplasmic domain is not known, one possibility might be a change in SLAM localization. The gp64 protein localizes to the cell surface but is excluded from lipid raft microdomains (72). As VSVG and CD46 also do not associate with lipid rafts (35, 57), proteins localized to lipid raft microdomains may be excluded during virus assembly and budding. Although the cellular localization of SLAM is not known, further studies will be necessary to clarify the relationship between cell surface localization and incorporation into baculovirus particles and to test the involvement of lipid raft microdomains in this process.

Mangor et al. demonstrated that gp64-null baculoviruses pseudotyped with VSVG were not morphologically distinguishable from budded wild-type AcMNPV particles (34). We confirmed that Ac $\Delta$ 64/VSVG/CAIuc, as well as Ac $\Delta$ 64/CD46/CAIuc and Ac $\Delta$ 64/GFP/CAIuc, exhibited a morphology similar to that of AcMNPV. These results indicate that expression of gp64 is not required for the morphogenesis of a rod-shaped structure for budded AcMNPV particles. Immunogold labeling of Ac $\Delta$ 64/VSVG/CAIuc and Ac $\Delta$ 64/CD46/CAIuc revealed that the VSVG and CD46 proteins were incorporated into and distributed throughout the whole viral surface. These results are consistent with previous observations that VSVG fusion proteins were distributed throughout the stalk and head domains of baculovirus particles, in contrast to gp64, which was primarily localized in the head domain (11, 44).

In this study, we have demonstrated the capability for ligand-directed gene delivery by pseudotype baculoviruses in vitro. For future in vivo applications of baculovirus vectors for gene targeting to specific organs or virus-infected cells as a method of treatment of inherited or infectious diseases, it is imperative to exhaustively study the transcription of baculoviral genes in mammalian cells for certification of safety. In addition, further studies are needed to establish replication-competent cell lines capable of supporting the propagation of pseudotype viruses without the possibility of replication-competent virus breakthrough by incorporation of gp64 and to optimize the conditions necessary for the efficient incorporation of ligands into recombinant baculovirus particles.

#### ACKNOWLEDGMENTS

We thank T. Seya and Y. Yanagi for kindly providing CHO cell lines; P. Faulkner, G. F. Rohrmann, and M. A. Whitt for their generous donation of antibodies; and H. Bando and K. Takeuchi for providing plasmids. We also thank H. Murase and I. Yanase for secretarial work and S. Ogawa for technical assistance.

This work was supported by grants-in-aid from the Ministry of Health, Labor, and Welfare in Japan and the 21st Century Center of Excellence Program of Japan.

#### REFERENCES

- Abe, T., H. Takahashi, H. Hamazaki, N. Miyano-Kurosaki, Y. Matsuura, and H. Takaku. 2003. Baculovirus induces an innate immune response and confers protection from lethal influenza virus infection in mice. *J. Immunol.* 171:1133-1139.
- Adams, E. M., M. C. Brown, M. Nunge, M. Krych, and J. P. Atkinson. 1991. Contribution of the repeating domains of membrane cofactor protein (CD46) of the complement system to ligand binding and cofactor activity. *J. Immunol.* 147:3005-3011.
- Ayres, M. D., S. C. Howard, J. Kuzio, M. Lopez-Ferber, and R. D. Possee. 1994. The complete DNA sequence of *Autographa californica* nuclear polyhedrosis virus. *Virology* 202:586-605.
- Barsoum, J., R. Brown, M. McKee, and F. M. Boyce. 1997. Efficient transduction of mammalian cells by a recombinant baculovirus having the vesicular stomatitis virus G glycoprotein. *Hum. Gene Ther.* 8:2011-2018.
- Bideshi, D. K., and B. A. Federici. 2000. The *Trichoplusia ni* granulovirus helicase is unable to support replication of *Autographa californica* multicapsid nucleopolyhedrovirus in cells and larvae of *T. ni*. *J. Gen. Virol.* 81:1593-1599.
- Blissard, G. W., and J. R. Wenz. 1992. Baculovirus gp64 envelope glycoprotein is sufficient to mediate pH-dependent membrane fusion. *J. Virol.* 66:6829-6835.
- Boritz, E., J. Gerlach, J. E. Johnson, and J. K. Rose. 1999. Replication-competent rhabdoviruses with human immunodeficiency virus type 1 coats and green fluorescent protein: entry by a pH-independent pathway. *J. Virol.* 73:6937-6945.
- Boyce, F. M., and N. L. Bucher. 1996. Baculovirus-mediated gene transfer into mammalian cells. *Proc. Natl. Acad. Sci. USA* 93:2348-2352.
- Bui, M., G. Whittaker, and A. Helenius. 1996. Effect of M1 protein and low pH on nuclear transport of influenza virus ribonucleoproteins. *J. Virol.* 70:8391-8401.
- Cattaneo, R. 2004. Four viruses, two bacteria, and one receptor: membrane cofactor protein (CD46) as pathogens' magnet. *J. Virol.* 78:4385-4388.
- Chapple, S. D., and I. M. Jones. 2002. Non-polar distribution of green fluorescent protein on the surface of *Autographa californica* nucleopolyhedrovirus using a heterologous membrane anchor. *J. Biotechnol.* 95:269-275.
- Chernomordik, L., E. Leikina, M. S. Cho, and J. Zimmerberg. 1995. Control of baculovirus gp64-induced syncytium formation by membrane lipid composition. *J. Virol.* 69:3049-3058.
- Coit, D. A., and A. D. Miller. 2004. Phosphatidylserine is not the cell surface receptor for vesicular stomatitis virus. *J. Virol.* 78:10920-10926.
- Condreay, J. P., S. M. Witherspoon, W. C. Clay, and T. A. Kost. 1999. Transient and stable gene expression in mammalian cells transduced with a recombinant baculovirus vector. *Proc. Natl. Acad. Sci. USA* 96:127-132.
- Dörig, R. E., A. Marcell, A. Chopra, and C. D. Richardson. 1993. The human CD46 molecule is a receptor for measles virus (Edmonston strain). *Cell* 75:295-305.
- Fishelson, Z., N. Donin, S. Zell, S. Schultz, and M. Kirschfink. 2003. Obstacles to cancer immunotherapy: expression of membrane complement regulatory proteins (mCRPs) in tumors. *Mol. Immunol.* 40:109-123.
- Gaggar, A., D. M. Shayakhmetov, and A. Lieber. 2003. CD46 is a cellular receptor for group B adenoviruses. *Nat. Med.* 9:1408-1412.
- Gronowski, A. M., D. M. Hilbert, K. C. Sheehan, G. Garotta, and R. D. Schreiber. 1999. Baculovirus stimulates antiviral effects in mammalian cells. *J. Virol.* 73:9944-9951.
- Hajós, J. P., J. Pijnenburg, M. Usmany, D. Zuidema, P. Závodszy, and J. M. Vlak. 2000. High frequency recombination between homologous baculoviruses in cell culture. *Arch. Virol.* 145:159-164.
- Hefferon, K., A. Oomens, S. Monsma, C. Finnerty, and G. W. Blissard. 1999. Host cell receptor binding by baculovirus GP64 and kinetics of virion entry. *Virology* 258:455-468.
- Hofmann, C., V. Sandig, G. Jennings, M. Rudolph, P. Schlag, and M. Strauss. 1995. Efficient gene transfer into human hepatocytes by baculovirus vectors. *Proc. Natl. Acad. Sci. USA* 92:10099-10103.
- Hohmann, A. W., and P. Faulkner. 1983. Monoclonal antibodies to baculovirus structural proteins: determination of specificities by Western blot analysis. *Virology* 125:432-444.
- Howie, D., M. Simarro, J. Sayos, M. Guirado, J. Sancho, and C. Terhorst. 2002. Molecular dissection of the signaling and costimulatory functions of CD150 (SLAM): CD150/SAP binding and CD150-mediated costimulation. *Blood* 99:957-965.
- Hsu, E. C., C. Iorio, F. Sarangl, A. A. Khine, and C. D. Richardson. 2001. CDw150 (SLAM) is a receptor for a lymphotropic strain of measles virus and may account for the immunosuppressive properties of this virus. *Virology* 279:9-21.
- Iwata, K., T. Seya, Y. Yanagi, J. M. Pesando, P. M. Johnson, M. Okabe, S. Ueda, H. Ariga, and S. Nagasawa. 1995. Diversity of sites for measles virus binding and for inactivation of complement C3b and C4b on membrane cofactor protein CD46. *J. Biol. Chem.* 270:15148-15152.
- Källström, H., M. K. Liszewski, J. P. Atkinson, and A. B. Jonsson. 1997. Membrane cofactor protein (MCP or CD46) is a cellular pilus receptor for pathogenic *Neisseria*. *Mol. Microbiol.* 25:639-647.
- Kamita, S. G., S. Maeda, and B. D. Hammock. 2003. High-frequency homologous recombination between baculoviruses involves DNA replication. *J. Virol.* 77:13053-13061.
- Kondo, A., and S. Maeda. 1991. Host range expansion by recombination of the baculoviruses *Bombyx mori* nuclear polyhedrosis virus and *Autographa californica* nuclear polyhedrosis virus. *J. Virol.* 65:3625-3632.
- Kost, T. A., and J. P. Condreay. 2002. Recombinant baculoviruses as mammalian cell gene-delivery vectors. *Trends Biotechnol.* 20:173-180.
- Lefrançois, L., and D. S. Lyles. 1982. The interaction of antibody with the major surface glycoprotein of vesicular stomatitis virus. II. Monoclonal antibodies of nonneutralizing and cross-reactive epitopes of Indiana and New Jersey serotypes. *Virology* 121:168-174.
- Leikina, E., H. O. Onaran, and J. Zimmerberg. 1992. Acidic pH induces

- fusion of cells infected with baculovirus to form syncytia. *FEBS Lett.* **304**: 221–224.
32. Luckow, V. A., and M. D. Summers. 1988. Trends in the development of baculovirus expression vectors. *Biotechnology* **6**:47–55.
  33. Lung, O., M. Westenberg, J. M. Vlak, D. Zuidema, and G. W. Blissard. 2002. Pseudotyping *Autographa californica* multicapsid nucleopolyhedrovirus (AcMNPV): F proteins from group II NPVs are functionally analogous to AcMNPV GP64. *J. Virol.* **76**:5729–5736.
  34. Mangor, J. T., S. A. Monsma, M. C. Johnson, and G. W. Blissard. 2001. A GP64-null baculovirus pseudotyped with vesicular stomatitis virus G protein. *J. Virol.* **75**:2544–2556.
  35. Manié, S. N., S. Debreyne, S. Vincent, and D. Gerlier. 2000. Measles virus structural components are enriched into lipid raft microdomains: a potential cellular location for virus assembly. *J. Virol.* **74**:305–311.
  36. Marsh, M. 1984. The entry of enveloped viruses into cells by endocytosis. *Biochem. J.* **218**:1–10.
  37. Martin, K., and A. Helenius. 1991. Nuclear transport of influenza virus ribonucleoproteins: the viral matrix protein (M1) promotes export and inhibits import. *Cell* **67**:117–130.
  38. Matsuura, Y., R. D. Possee, H. A. Overton, and D. H. Bishop. 1987. Baculovirus expression vectors: the requirements for high level expression of proteins, including glycoproteins. *J. Gen. Virol.* **68**:1233–1250.
  39. Maurer, K., T. Krey, V. Moening, H. J. Thiel, and T. Rümmer. 2004. CD46 is a cellular receptor for bovine viral diarrhea virus. *J. Virol.* **78**:1792–1799.
  40. Monsma, S. A., A. G. Oomens, and G. W. Blissard. 1996. The GP64 envelope fusion protein is an essential baculovirus protein required for cell-to-cell transmission of infection. *J. Virol.* **70**:4607–4616.
  41. Monsma, S. A., and G. W. Blissard. 1995. Identification of a membrane fusion domain and an oligomerization domain in the baculovirus GP64 envelope fusion protein. *J. Virol.* **69**:2583–2595.
  42. Naniche, D., G. Varior-Krishnan, F. Cervoni, T. F. Wild, B. Rossi, C. Rabbourdin-Combe, and D. Gerlier. 1993. Human membrane cofactor protein (CD46) acts as a cellular receptor for measles virus. *J. Virol.* **67**:6025–6032.
  43. Niwa, H., K. Yamamura, and J. Miyazaki. 1991. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* **108**:193–199.
  44. Ojala, K., J. Koski, W. Ernst, R. Grabherr, I. Jones, and C. Oker-Blom. 2004. Improved display of synthetic IgG-binding domains on the baculovirus surface. *Technol. Cancer Res. Treat.* **3**:77–84.
  45. Ojala, K., D. G. Mottershead, A. Suokko, and C. Oker-Blom. 2001. Specific binding of baculoviruses displaying gp64 fusion proteins to mammalian cells. *Biochem. Biophys. Res. Commun.* **284**:777–784.
  46. Okada, N., M. K. Liszewski, J. P. Atkinson, and M. Caparon. 1995. Membrane cofactor protein (CD46) is a keratinocyte receptor for the M protein of the group A streptococcus. *Proc. Natl. Acad. Sci. USA* **92**:2489–2493.
  47. Okamoto, K., K. Moriishi, T. Miyamura, and Y. Matsuura. 2004. Intramembrane proteolysis and endoplasmic reticulum retention of hepatitis C virus core protein. *J. Virol.* **78**:6370–6380.
  48. Ono, N., H. Tatsuo, Y. Hidaka, T. Aoki, H. Minagawa, and Y. Yanagi. 2001. Measles viruses on throat swabs from measles patients use signaling lymphocytic activation molecule (CDw150) but not CD46 as a cellular receptor. *J. Virol.* **75**:4399–4401.
  49. Oomens, A. G., and G. W. Blissard. 1999. Requirement for GP64 to drive efficient budding of *Autographa californica* multicapsid nucleopolyhedrovirus. *Virology* **254**:297–314.
  50. Oomens, A. G., S. A. Monsma, and G. W. Blissard. 1995. The baculovirus GP64 envelope fusion protein: synthesis, oligomerization, and processing. *Virology* **209**:592–603.
  51. Pearson, M. N., R. L. Russell, G. F. Rohrmann, and G. S. Beaudreau. 1988. p39, a major baculovirus structural protein: immunocytochemical characterization and genetic location. *Virology* **167**:407–413.
  52. Peng, K. W., C. J. TenEyck, E. Galanis, K. R. Kalli, L. C. Hartmann, and S. J. Russell. 2002. Intraperitoneal therapy of ovarian cancer using an engineered measles virus. *Cancer Res.* **62**:4656–4662.
  53. Pieroni, L., and N. La Monica. 2001. Towards the use of baculovirus as a gene therapy vector. *Curr. Opin. Mol. Ther.* **3**:464–467.
  54. Rose, J. K., and M. A. Whitt. 2001. Rhabdoviruses, p. 1221–1244. In D. M. Knipe and P. M. Howley (ed.), *Fields virology*, 4th ed. Lippincott Williams and Wilkins, Philadelphia, Pa.
  55. Sandig, V., and M. Strauss. 1996. Liver-directed gene transfer and application to therapy. *J. Mol. Med.* **74**:205–212.
  56. Santoro, F., P. E. Kennedy, G. Locatelli, M. S. Malnati, E. A. Berger, and P. Lusso. 1999. CD46 is a cellular receptor for human herpesvirus 6. *Cell* **99**: 817–827.
  57. Scheifele, P., A. Rietveld, T. Wilk, and K. Simons. 1999. Influenza viruses select ordered lipid domains during budding from the plasma membrane. *J. Biol. Chem.* **274**:2038–2044.
  58. Schlegel, R., T. S. Tralka, M. C. Willingham, and I. Pastan. 1983. Inhibition of VSV binding and infectivity by phosphatidylserine: is phosphatidylserine a VSV-binding site? *Cell* **32**:639–646.
  59. Segerman, A., J. P. Atkinson, M. Marttila, V. Dennerquist, G. Wadell, and N. Arnberg. 2003. Adenovirus type 11 uses CD46 as a cellular receptor. *J. Virol.* **77**:9183–9191.
  60. Seya, T., A. Hirano, M. Matsumoto, M. Nomura, and S. Ueda. 1999. Human membrane cofactor protein (MCP, CD46): multiple isoforms and functions. *Int. J. Biochem. Cell Biol.* **31**:1255–1260.
  61. Shoji, I., H. Alzaki, H. Tani, K. Ishii, T. Chiba, I. Saito, T. Miyamura, and Y. Matsuura. 1997. Efficient gene transfer into various mammalian cells, including non-hepatic cells, by baculovirus vectors. *J. Gen. Virol.* **78**:2657–2664.
  62. Sirena, D., B. Lillienfeld, M. Eisenhut, S. Kälin, K. Boucke, R. R. Beerli, L. Vogt, C. Ruedi, M. F. Bachmann, U. F. Greber, and S. Hemmi. 2004. The human membrane cofactor CD46 is a receptor for species B adenovirus serotype 3. *J. Virol.* **78**:4454–4462.
  63. Takeuchi, K., M. Takeda, N. Miyajima, F. Kobune, K. Tanabayashi, and M. Tashiro. 2002. Recombinant wild-type and Edmonston strain measles viruses bearing heterologous H proteins: role of H protein in cell fusion and host cell specificity. *J. Virol.* **76**:4891–4900.
  64. Takikawa, S., K. Ishii, H. Alzaki, T. Suzuki, H. Asakura, Y. Matsuura, and T. Miyamura. 2000. Cell fusion activity of hepatitis C virus envelope proteins. *J. Virol.* **74**:5066–5074.
  65. Tani, H., C. K. Limn, C. C. Yap, M. Onishi, M. Nozaki, Y. Nishimune, N. Okahashi, Y. Kitagawa, R. Watanabe, R. Mochizuki, K. Moriishi, and Y. Matsuura. 2003. In vitro and in vivo gene delivery by recombinant baculoviruses. *J. Virol.* **77**:9799–9808.
  66. Tani, H., M. Nishijima, H. Ushijima, T. Miyamura, and Y. Matsuura. 2001. Characterization of cell-surface determinants important for baculovirus infection. *Virology* **279**:343–353.
  67. Tatsuo, H., N. Ono, K. Tanaka, and Y. Yanagi. 2000. SLAM (CDw150) is a cellular receptor for measles virus. *Nature* **406**:893–897.
  68. Tjia, S. T., G. M. zu Altneschilde, and W. Doerfler. 1983. *Autographa californica* nuclear polyhedrosis virus (AcNPV) DNA does not persist in mass cultures of mammalian cells. *Virology* **125**:107–117.
  69. Volkman, L. E., and P. A. Goldsmith. 1985. Mechanism of neutralization of budded *Autographa californica* nuclear polyhedrosis virus by a monoclonal antibody: inhibition of entry by adsorptive endocytosis. *Virology* **143**:185–195.
  70. Volkman, L. E., P. A. Goldsmith, R. T. Hess, and P. Faulkner. 1984. Neutralization of budded *Autographa californica* NPV by a monoclonal antibody: identification of the target antigen. *Virology* **133**:354–362.
  71. Wu, Y., G. Liu, and E. B. Carstens. 1999. Replication, integration, and packaging of plasmid DNA following cotransfection with baculovirus viral DNA. *J. Virol.* **73**:5473–5480.
  72. Zhang, S. X., Y. Han, and G. W. Blissard. 2003. Palmitoylation of the *Autographa californica* multicapsid nucleopolyhedrovirus envelope glycoprotein GP64: mapping, functional studies, and lipid rafts. *J. Virol.* **77**:6265–6273.



## Challenges in creating a vaccine to prevent hepatitis E

Bryan J. Maloney<sup>a</sup>, Naokazu Takeda<sup>b</sup>, Yuriko Suzuki<sup>b</sup>, Yasushi Ami<sup>b</sup>, Tian Cheng Li<sup>b</sup>,  
Tatsuo Miyamura<sup>b</sup>, Charles J. Arntzen<sup>c</sup>, Hugh S. Mason<sup>c,\*</sup>

<sup>a</sup> Boyce Thompson Institute for Plant Research, Ithaca, NY, USA

<sup>b</sup> National Institute of Infectious Diseases, Tokyo, Japan

<sup>c</sup> Biodesign Institute at Arizona State University, Life Sciences E Tower,  
Tempe, AZ 85287-4501, USA

Available online 19 November 2004

### Summary

Recombinant hepatitis E virus capsid protein (HEV CP) assembles orally immunogenic virus-like particles (VLP) when expressed in an insect cell system. We used plant expression cassettes, pHEV101 and pHEV110, for transformation of potato to express HEV CP, and 10 independent transgenic lines of HEV101 and 6 lines of HEV110 were obtained. ELISA for HEV CP was performed on tuber extracts. Accumulation of HEV CP in tubers varied from about 5 to 30 µg/g fresh tuber depending on the transgenic plant line. We further compared the expression levels with the yield of tubers for each line. Tuber yield varied less than expression levels, and ranged from about 600 to 1000 g per pot. Although Western blot showed that apparently intact HEV CP accumulated, we observed very limited assembly of virus-like particles in potato tubers. Oral immunization of mice with transgenic potatoes failed to elicit detectable anti-CP antibody response in serum, suggesting that VLP assembly is a key factor in orally delivered HEV CP vaccines.

© 2004 Elsevier Ltd. All rights reserved.

**Keywords:** Hepatitis E; HEV; VLP; Transgenic potato

### 1. Introduction

Hepatitis E virus (HEV) is a causative agent of hepatitis E that occurs in many developing countries [1], and this virus is currently classified into a tentative genus, “Hepatitis E-like viruses.” HEV is transmitted mainly by the fecal-oral route, and large epidemics due to this virus are often associated with contaminated water [2,3]. Hepatitis E has been formerly known as an enterically transmitted non-A, non-B hepatitis [4]. The fact that HEV can survive in the intestinal tract suggests that the virus is relatively stable to acid and mild alkaline conditions.

HEV contains a single-stranded positive-sense approximately 7.5 kb RNA molecule that is 3′ polyadenylated and includes three open reading frames (ORFs). ORF1, mapped in the 5′ half of the genome, is thought to encode viral non-structural proteins. ORF2, located at the 3′-terminus of the

genome, encodes a 72 kDa protein for the putative viral capsid. ORF3, with unknown function, is mapped between ORF1 and ORF2 [5]. In the absence of an appropriate cell culture for HEV propagation, research has focused on the expression of the ORF2 protein in heterologous systems. Recently, virus-like particles (VLP) of recombinant hepatitis E virus (rHEV) were produced by using a baculovirus system carrying an N-terminally truncated ORF2 gene of the Burma strain [6]. Thus, rHEV VLP were formed in Tn5 cells and could be collected from the culture supernatant.

In order to evaluate the potential of rHEV VLP as an oral immunogen, we analyzed the immune responses in mice and monkeys after oral administration [7,8]. The animals were orally inoculated with purified rHEV VLP without adjuvant. ELISA indicated that oral immunization with rHEV VLP induced immune responses in both mice and monkeys. In addition, the monkeys were completely protected from infection when challenge was carried out with native HEV, suggesting that rHEV VLPs are a potential mucosal vaccine for HEV infection.

\* Corresponding author. Tel.: +1 480 727 8228; fax: +1 480 727 6194.

E-mail address: [hugh.mason@asu.edu](mailto:hugh.mason@asu.edu) (H.S. Mason).