

- mamoto M. Transcription factor Nrf2 regulates inflammation by mediating the effect of 15-deoxy- $\Delta^{12,14}$ -prostaglandin J_2 . *Mol Cell Biol* 24: 36–45, 2004.
42. Kang KW, Choi SH, and Kim SG. Peroxynitrite activates NF-E2-related factor 2/antioxidant response element through the pathway of phosphatidylinositol 3-kinase: the role of nitric oxide synthase in rat glutathione S-transferase A2 induction. *Nitric Oxide* 7: 244–253, 2002.
 43. Kang M-I, Kobayashi A, Wakabayashi N, Kim S-G, and Yamamoto M. Scaffolding of Keap1 to the actin cytoskeleton controls the function of Nrf2 as key regulator of cytoprotective phase 2 genes. *Proc Natl Acad Sci USA* 101: 2046–2051, 2004.
 44. Kataoka K, Igarashi K, Itoh K, Fujiwara KT, Noda M, Yamamoto M, and Nishizawa M. Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NF-E2 transcription factor. *Mol Cell Biol* 15: 2180–2190, 1995.
 45. Katoh Y, Itoh K, Yoshida E, Miyagishi M, Fukamizu A, and Yamamoto M. Two domains of Nrf2 cooperatively bind CBP, a CREB binding protein, and synergistically activate transcription. *Genes Cells* 6: 857–868, 2001.
 46. Kim YC, Masutani H, Yamaguchi Y, Itoh K, Yamamoto M, and Yodoi J. Hemin-induced activation of the thioredoxin gene by Nrf2. A differential regulation of the antioxidant responsive element by a switch of its binding factors. *J Biol Chem* 276: 18399–18406, 2001.
 47. Kobayashi A, Ito E, Toki T, Kogame K, Takahashi S, Igarashi K, Hayashi N, and Yamamoto M. Molecular cloning and functional characterization of a new Cap'n'-collar family transcription factor Nrf3. *J Biol Chem* 274: 6443–6452, 1999.
 48. Kobayashi M, Itoh K, Suzuki T, Osanai H, Nishikawa K, Katoh Y, Takagi Y, and Yamamoto M. Identification of the interactive interface and phylogenetic conservation of the Nrf2-Keap1 system. *Genes Cells* 7: 807–820, 2002.
 49. Kotlo KU, Yehiely F, Efimova E, Harasty H, Hesabi B, Shchors K, Einat P, Rozen A, Berent E, and Deiss LP. Nrf2 is an inhibitor of the Fas pathway as identified by Achilles' Heel Method, a new function-based approach to gene identification in human cells. *Oncogene* 22: 797–806, 2003.
 50. Kuroha T, Takahashi S, Komeno T, Itoh K, Nagasawa T, and Yamamoto M. Ablation of Nrf2 function does not increase the erythroid or megakaryocytic cell lineage dysfunction caused by p45 NF-E2 gene disruption. *J Biochem (Tokyo)* 123: 376–379, 1998.
 51. Kusunoki H, Motohashi H, Katsuoka F, Morohashi A, Yamamoto M, and Tanaka T. Solution structure of the DNA-binding domain of MafG. *Nat Struct Biol* 9: 252–256, 2002.
 52. Kwak MK, Egner PA, Dolan PM, Ramos-Gomez M, Groopman JD, Itoh K, Yamamoto M, and Kensler TW. Role of phase 2 enzyme induction in chemoprotection by dithiolethiones. *Mutat Res* 480–481: 305–315, 2001.
 53. Kwak MK, Itoh K, Yamamoto M, Sutter TR, and Kensler TW. Role of transcription factor Nrf2 in the induction of hepatic phase 2 and antioxidant enzymes in vivo by the cancer chemoprotective agent, 3H-1,2-dimethiole-3-thione. *Mol Med* 7: 135–145, 2001.
 54. Kwak MK, Itoh K, Yamamoto M, and Kensler TW. Enhanced expression of the transcription factor Nrf2 by cancer chemopreventive agents: role of antioxidant response element-like sequences in the nrf2 promoter. *Mol Cell Biol* 22: 2883–2892, 2002.
 55. Kwak MK, Wakabayashi N, Greenlaw JL, Yamamoto M, and Kensler TW. Antioxidants enhance mammalian proteasome expression through the Keap1-Nrf2 signaling pathway. *Mol Cell Biol* 23: 8786–8794, 2003.
 56. Kwak MK, Wakabayashi N, Itoh K, Motohashi H, Yamamoto M, and Kensler TW. Modulation of gene expression by cancer chemopreventive dithiolethiones through the Keap1-Nrf2 pathway. Identification of novel gene clusters for cell survival. *J Biol Chem* 278: 8135–8145, 2003.
 57. Kwong M, Kan YW, and Chan JY. The CNC basic leucine zipper factor, Nrfl, is essential for cell survival in response to oxidative stress-inducing agents. Role for Nrfl in γ -*gcs*₂ and *gss* expression in mouse fibroblasts. *J Biol Chem* 274: 37491–37498, 1999.
 58. Lee JM, Calkins MJ, Chan K, Kan YW, and Johnson JA. Identification of the NF-E2-related factor-2-dependent genes conferring protection against oxidative stress in primary cortical astrocytes using oligonucleotide microarray analysis. *J Biol Chem* 278: 12029–12038, 2003.
 59. Lee JM, Shih AY, Murphy TH, and Johnson JA. NF-E2-related factor-2 mediates neuroprotection against mitochondrial complex I inhibitors and increased concentrations of intracellular calcium in primary cortical neurons. *J Biol Chem* 278: 37948–37956, 2003.
 60. Leung I., Kwong M, Hou S, Lee C, and Chan JY. Deficiency of the Nrfl and Nrf2 transcription factors results in early embryonic lethality and severe oxidative stress. *J Biol Chem* 278: 48021–48029, 2003.
 61. Li J, Lee JM, and Johnson JA. Microarray analysis reveals an antioxidant responsive element-driven gene set involved in conferring protection from an oxidative stress-induced apoptosis in IMR-32 cells. *J Biol Chem* 277: 388–394, 2002.
 62. Martin F, van Deursen JM, Shivdasani RA, Jackson CW, Troutman AG, and Ney PA. Erythroid maturation and globin gene expression in mice with combined deficiency of NF-E2 and nrf-2. *Blood* 91: 3459–3466, 1998.
 63. McMahon M, Itoh K, Yamamoto M, and Hayes JD. Keap1-dependent proteasomal degradation of transcription factor Nrf2 contributes to the negative regulation of antioxidant response element-driven gene expression. *J Biol Chem* 278: 21592–21600, 2003.
 64. Mohler J, Mahaffey JW, Deutsch E, and Vani K. Control of *Drosophila* head segment identity by the bZIP homeotic gene *cnc*. *Development* 121: 237–247, 1995.
 65. Moi P, Chan K, Asunis I, Cao A, and Kan YW. Isolation of NF-E2-related factor 2 (Nrf2), a NF-E2-like basic leucine zipper transcriptional activator that binds to the tandem NF-E2/AP1 repeat of the β -globin locus control region. *Proc Natl Acad Sci USA* 91: 9926–9930, 1994.
 66. Moran JA, Dahl EL, and Mulcahy RT. Differential induction of *mafF*, *mafG* and *mafK* expression by electrophile-response-element activators. *Biochem J* 361: 371–377, 2002.

67. Morito N, Yoh K, Itoh K, Hirayama A, Koyama A, Yamamoto M, and Takahashi S. Nrf2 regulates the sensitivity of death receptor signals by affecting intracellular glutathione levels. *Oncogene* 22: 9275–9281, 2003.
68. Motohashi H, O'Connor T, Katsuoka F, Engel JD, and Yamamoto M. Integration and diversity of the regulatory network composed of Maf and CNC families of transcription factors. *Gene* 294: 1–12, 2002.
69. Motohashi H, Katsuoka F, Engel JD, and Yamamoto M. Small Maf proteins serve as transcriptional cofactors for keratinocyte differentiation in the Keap1-Nrf2 regulatory pathway. *Proc Natl Acad Sci U S A* 101: 6379–6384, 2004.
70. Muto A, Tashiro S, Tsuchiya H, Kume A, Kanno M, Ito E, Yamamoto M, and Igarashi K. Activation of Maf/AP-1 repressor Bach2 by oxidative stress promotes apoptosis and its interaction with promyelocytic leukemia nuclear bodies. *J Biol Chem* 277: 20724–20733, 2002.
71. Nakaso K, Yano H, Fukuhara Y, Takeshima T, Wada-Isoe K, and Nakashima K. PI3K is a key molecule in the Nrf2-mediated regulation of antioxidant proteins by hemin in human neuroblastoma cells. *FEBS Lett* 546: 181–184, 2003.
72. Nguyen T, Sherratt PJ, Huang HC, Yang CS, and Pickett CB. Increased protein stability as a mechanism that enhances Nrf2-mediated transcriptional activation of the antioxidant response element. Degradation of Nrf2 by the 26 S proteasome. *J Biol Chem* 278: 4536–4541, 2003.
73. Numazawa S, Ishikawa M, Yoshida A, Tanaka S, and Yoshida T. Atypical protein kinase C mediates activation of NF-E2-related factor 2 in response to oxidative stress. *Am J Physiol Cell Physiol* 285: C334–C342, 2003.
74. Pietsch EC, Chan JY, Torti FM, and Torti SV. Nrf2 mediates the induction of ferritin H in response to xenobiotics and cancer chemopreventive dithiolethiones. *J Biol Chem* 278: 2361–2369, 2003.
75. Pintard L, Willis JH, Willems A, Johnson JLF, Srayko M, Kurz T, Glaser S, Mains PE, Tyers M, Bowerman B, and Peter M. The BTB protein MEL-26 is a substrate-specific adaptor of the CUL-3 ubiquitin-ligase. *Nature* 425: 311–316, 2003.
76. Ramos-Gomez M, Kwak MK, Dolan PM, Itoh K, Yamamoto M, Talalay P, and Kensler TW. Sensitivity to carcinogenesis is increased and chemoprotective efficacy of enzyme inducers is lost in *nrf2* transcription factor-deficient mice. *Proc Natl Acad Sci U S A* 98: 3410–3415, 2001.
77. Ramos-Gomez M, Dolan PM, Itoh K, Yamamoto M, and Kensler TW. Interactive effects of *nrf2* genotype and oltipraz on benzo[a]pyrene-DNA adducts and tumor yield in mice. *Carcinogenesis* 24: 461–467, 2003.
78. Sasaki H, Sato H, Kuriyama-Matsumura K, Sato K, Maebara K, Wang H, Tamba M, Itoh K, Yamamoto M, and Bannai S. Electrophile response element-mediated induction of the cystine/glutamate exchange transporter gene expression. *J Biol Chem* 277: 44765–44771, 2002.
79. Sekhar KR, Spitz DR, Harris S, Nguyen TT, Meredith MJ, Holt JT, Gius D, Marnett LJ, Summar ML, Freeman ML, and Guis D. Redox-sensitive interaction between KIAA0132 and Nrf2 mediates indomethacin-induced expression of γ -glutamylcysteine synthetase. *Free Radic Biol Med* 32: 650–662, 2002.
80. Sekhar KR, Yan XX, and Freeman ML. Nrf2 degradation by the ubiquitin proteasome pathway is inhibited by KIAA0132, the human homolog to INrf2. *Oncogene* 21: 6829–6834, 2002.
81. Stewart D, Killeen E, Naquin R, Alam S, and Alam J. Degradation of transcription factor Nrf2 via the ubiquitin-proteasome pathway and stabilization by cadmium. *J Biol Chem* 278: 2396–2402, 2003.
82. Sun J, Hoshino H, Takaku K, Nakajima O, Muto A, Suzuki H, Tashiro S, Takahashi S, Shibahara S, Alam J, Taketo MM, Yamamoto M, and Igarashi K. Hemoprotein Bach1 regulates enhancer availability of heme oxygenase-1 gene. *EMBO J* 21: 5216–5224, 2002.
83. Suzuki T, Blank V, Sesay JS, and Crawford DR. Maf genes are involved in multiple stress response in human. *Biochem Biophys Res Commun* 280: 4–8, 2001.
84. Takagi Y, Kobayashi M, Li L, Suzuki T, Nishikawa K, and Yamamoto M. MafT, a new member of the small Maf protein family in zebrafish. *Biochem Biophys Res Commun* 320: 62–69, 2004.
85. Thimmulappa RK, Mai KH, Srisuna S, Kensler TW, Yamamoto M, and Biswal S. Identification of Nrf2-regulated genes induced by the chemopreventive agent sulforaphane by oligonucleotide microarray. *Cancer Res* 62: 5196–5203, 2002.
86. Toki T, Itoh J, Kitazawa J, Arai K, Hatakeyama K, Akasaka J, Igarashi K, Nomura N, Yokoyama M, Yamamoto M, and Ito E. Human small Maf proteins form heterodimers with CNC family transcription factors and recognize the NF-E2 motif. *Oncogene* 14: 1901–1910, 1997.
87. Toone WM and Jones N. AP-1 transcription factors in yeast. *Curr Opin Genet Dev* 9: 55–61, 1999.
88. Venugopal R and Jaiswal AK. Nrf2 and Nrf1 in association with Jun proteins regulate antioxidant response element-mediated expression and coordinated induction of genes encoding detoxifying enzymes. *Oncogene* 17: 3145–3156, 1998.
89. Veraksa A, McGinnis N, Li X, Mohler J, and McGinnis W. Cap 'n' collar B cooperates with a small Maf subunit to specify pharyngeal development and suppress deformed homeotic function in the *Drosophila* head. *Development* 127: 4023–4037, 2000.
90. Wakabayashi N, Itoh K, Wakabayashi J, Motohashi H, Noda S, Takahashi S, Imakado S, Kotsuji T, Otsuka F, Roop DR, Harada T, Engel JD, and Yamamoto M. Keap1-null mutation leads to postnatal lethality due to constitutive Nrf2 activation. *Nat Genet* 35: 238–245, 2003.
91. Wakabayashi N, Dinkova-Kostova AT, Holtzclaw WD, Kang MI, Kobayashi A, Yamamoto M, Kensler TW, and Talalay P. Protection against electrophiles and oxidant stress by induction of the phase 2 response: fate of the cysteine residues of the Keap1 sensor modified by inducers. *Proc Natl Acad Sci U S A* 101: 2040–2045, 2004.
92. Wild AC, Moinova HR, and Mulcahy RT. Regulation of γ -glutamylcysteine synthetase subunit gene expression by the transcription factor Nrf2. *J Biol Chem* 274: 33627–33636, 1999.

93. Xu L, Wei Y, Reboul J, Vaglio P, Shin TH, Vidal M, Elledge SJ, and Harper JW. BTB proteins are substrate-specific adaptors in an SCF-like modular ubiquitin ligase containing CUL-3. *Nature* 425: 316-321, 2003.
94. Yoh K, Itoh K, Enomoto A, Hirayama A, Yamaguchi N, Kobayashi M, Morito N, Koyama A, Yamamoto M, and Takahashi S. Nrf2-deficient female mice develop lupus-like autoimmune nephritis. *Kidney Int* 60: 1343-1353, 2001.
95. Yu R, Chen C, Mo YY, Hebbar V, Owuor ED, Tan TH, and Kong ANT. Activation of mitogen-activated protein kinase pathways induces antioxidant response element-mediated gene expression via a Nrf2-dependent mechanism. *J Biol Chem* 275: 39907-39913, 2000.
96. Zhang DD and Hannink M. Distinct cysteine residues in Keap1 are required for Keap1-dependent ubiquitination of Nrf2 and for stabilization of Nrf2 by chemopreventive agents and oxidative stress. *Mol Cell Biol* 23: 8137-8151, 2003.
97. Zhu M and Fahl WE. Functional characterization of transcription regulators that interact with the electrophile response element. *Biochem Biophys Res Commun* 289: 212-219, 2001.
98. Zipper LM and Mulcahy RT. Inhibition of ERK and p38 MAP kinases inhibits binding of Nrf2 and induction of GCS genes. *Biochem Biophys Res Commun* 278: 484-492, 2000.
99. Zipper LM and Mulcahy RT. The Keap1 BTB/POZ dimerization function is required to sequester Nrf2 in cytoplasm. *J Biol Chem* 277: 36544-36552, 2002.
100. Zipper LM and Mulcahy RT. Erk activation is required for Nrf2 nuclear localization during pyrrolidine dithiocarbamate induction of glutamate cysteine ligase modulatory gene expression in HepG2 cells. *Toxicol Sci* 73: 124-134, 2003.
101. Kobayashi A, Kang M-I, Okawa H, Ohtsuji M, Zenke Y, Chiba T, Igarashi K, and Yamamoto M. Oxidative stress sensor Keap1 functions as an adaptor for Cul3-based E3 ligase to regulate proteasomal degradation of Nrf2. *Mol Cell Biol* 24: 7130-7139, 2004.

Address reprint requests to:

Makoto Kobayashi, Ph.D.

Institute of Basic Medical Sciences

University of Tsukuba

Tsukuba 305-8575, Japan

E-mail: kobayash@tara.tsukuba.ac.jp

Received for publication April 30, 2004; accepted October 9, 2004.