

Rosenbauer F, Wagner K, Zhang P, Knobeloch K-P, <u>Iwama A</u> , and Tenen DG.	pDP4, a novel glycoprotein secreted by mature granulocytes is regulated by transcription factor PU.1..	<i>Blood</i>	103	4294- 4301	2004
Kubo-Akashi C, Iseki M, Kwon SM, Takizawa H, Takatsu K, <u>Takaki S</u> .	Roles of a conserved family of adaptor proteins, Lnk, SH2-B, and APS, for mast cell development, growth, and functions: APS-deficiency causes augmented degranulation and reduced actin assembly.	<i>Biochem Biophys Res Commun.</i>	315	356-362	2004
Iseki M, Kubo C, Kwon SM, Yamaguchi A, Kataoka Y, Yoshida N, Takatsu K, <u>Takaki S</u> .	Increased numbers of B-1 cells and enhanced responses against TI-2 antigen in mice lacking APS, an adaptor molecule containing PH and SH2 domains.	<i>Mol Cell Biol.</i>	24	2243- 2250	2004
Moon BG, <u>Takaki S</u> , Miyake K, Takatsu K.	The role of IL-5 for mature B-1 cells in homeostatic proliferation, cell survival, and Ig production.	<i>J Immunol.</i>	172	6020- 6029	2004
Moon BG, <u>Takaki S</u> , Nishizumi H, Yamamoto T, Takatsu K.	Abrogation of autoimmune disease in Lyn-deficient mice by the deletion of IL-5 receptor alpha chain gene.	<i>Cell Immunol.</i>	228	110-118	2004
Tamura T, Ariga H, Kinashi T, Uehara S, Kikuchi T, Nakada M, Tokunaga T, Xu W, Kariyone A, Saito T, Kitamura T, Maxwell G, <u>Takaki S</u> , Takatsu K.	The role of antigenic peptide in CD4+ T helper phenotype development in a T cell receptor transgenic model.	<i>Int Immunol.</i>	16	1691- 1699	2004
Iseki M, Kubo- Akashi C, Kwon SM, Yamaguchi A, Takatsu K, <u>Takaki S</u> .	APS, an adaptor molecule containing PH and SH2 domains, has a negative regulatory role in B cell proliferation.	<i>Biochem Biophys Res Commun.</i>	330	1005- 1013	2005