

## Role of direct cytotoxic effects of NSAIDs in the induction of gastric lesions

Wataru Tomisato, Shinji Tsutsumi, Tatsuya Hoshino, Hyun-Jung Hwang, Mitsunobu Mio<sup>1</sup>, Tomofusa Tsuchiya, Tohru Mizushima\*

Faculty of Pharmaceutical Sciences, Okayama University, Okayama 700-8530, Japan

Received 20 June 2003; accepted 17 September 2003

### Abstract

A major clinical problem encountered with the use of non-steroidal anti-inflammatory drugs (NSAIDs), is gastrointestinal complications. We previously reported that NSAIDs induce both necrosis and apoptosis *in vitro*. We here examined the cyclooxygenase (COX) dependency of this cytotoxic effect of NSAIDs and its involvement in NSAID-induced gastric lesions. Necrosis and apoptosis by NSAIDs was observed with all selective COX-2 inhibitors except rofecoxib and was not inhibited by exogenously added prostaglandin E<sub>2</sub>, suggesting that cytotoxicity of NSAIDs seems to be independent of the inhibition of COX. Intravenously administered indomethacin, which completely inhibited COX activity at gastric mucosa, did not produce gastric lesions. Orally administered selective COX-2 inhibitors, which did not inhibit COX at gastric mucosa, also did not produce gastric lesions. Interestingly, a combination of the oral administration of each of all selective COX-2 inhibitors except rofecoxib with the intravenous administration of indomethacin clearly produced gastric lesions. These results suggest that in addition to COX inhibition by NSAIDs, direct cytotoxicity of NSAIDs may be involved in NSAID-induced gastric lesions.

© 2003 Elsevier Inc. All rights reserved.

**Keywords:** Direct cytotoxicity; Cyclooxygenase; NSAIDs; Gastric lesions; Prostaglandin E<sub>2</sub>; Selective cyclooxygenase-2 inhibitors

### 1. Introduction

NSAIDs are one of the most frequently used classes of medicines in the world and account for nearly 5% of all prescribed medications [1]. However, NSAID administration is associated with gastrointestinal complications, such as gastric ulcers and bleeding, which sometimes become life-threatening diseases [2]. About 15–30% of chronic users of NSAIDs have gastrointestinal ulcers and bleeding [3–6]. In the United States, about 16,500 people per year die as a result of NSAID-associated gastrointestinal complications [7]. Therefore, the molecular mechanism governing NSAID-induced gastrointestinal damage needs to

be elucidated in order to develop new NSAIDs that do not have these side effects.

Inhibition of COX by NSAIDs, which is responsible for their anti-inflammatory activity was previously thought to be fully responsible for their gastrointestinal side effects [8]. This is because COX is an enzyme essential for the synthesis of prostaglandins (PGs), which have a strong cytoprotective effect on the gastrointestinal mucosa [9]. There are at least two subtypes of COX, COX-1 and COX-2, which are responsible for the majority of COX activity at the gastric mucosa and tissues with inflammation, respectively [10,11]. Therefore, it is reasonable to speculate that selective COX-2 inhibitors have anti-inflammatory activity without gastrointestinal side effects [10]. In fact, a greatly reduced incidence of gastroduodenal lesions was reported for selective COX-2 inhibitors (such as celecoxib and rofecoxib) both in animal and clinical data [12–15], however, recently published paper showed no difference in serious gastrointestinal complications between celecoxib and two non-selective NSAIDs [16]. Two lines of evidence have worked against the idea that the gastrointestinal side

\* Corresponding author. Tel.: +81-86-2517958; fax: +81-86-2517958.

E-mail address: [mizushima@pharm.okayama-u.ac.jp](mailto:mizushima@pharm.okayama-u.ac.jp) (T. Mizushima).

<sup>1</sup> Present address: School of Pharmacy, Shujitsu University, Okayama 703-8515, Japan.

**Abbreviations:** NSAID, non-steroidal anti-inflammatory drug; COX, cyclooxygenase; PG, prostaglandin; FBS, fetal bovine serum; MTT, 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyl tetrazolium bromide; PGI<sub>2</sub>, prostacyclin.

effects of NSAIDs are caused only by the inhibition of COX-1. The first is that a selective COX-1 inhibitor (SC-560) induced no gastric injury even at high dosages, and COX-1 knockout mice showed no detectable gastric ulcers [17–20]. The second line of evidence is that the increased incidence of gastrointestinal lesions and the decrease in PG levels induced by NSAIDs are not always linked with each other. For example, higher doses of NSAIDs were required for producing gastric lesions than were required for inhibiting COX at the gastric mucosa [21,22]. The former contradiction can be explained by the recent proposal that inhibition of both COX-1 and COX-2 was necessary for NSAID-induced gastrointestinal lesions [17]. However, the latter contradiction cannot be explained by this idea, suggesting therefore that NSAID-induced gastrointestinal lesions involve additional mechanisms [23]. Understanding the additional mechanisms is necessary in order to establish an alternative method for development of gastro-intestinally safe NSAIDs other than simply increasing their COX-2 selectivity. This new class of NSAIDs may be clinically beneficial because clinical disadvantages (i.e. risk of cardiovascular thrombotic disease) of selective COX-2 inhibitors were recently suggested [24,25].

In addition to various possibilities proposed for this additional mechanism (such as reduced blood flow, hypermotility, and activation of neutrophils) [26–28], a direct cytotoxic effect of NSAIDs (topical irritant property) on gastric mucosal cells was also proposed to be involved in NSAID-induced gastric lesions [23,29]. As for this direct cytotoxic effect, we previously reported that NSAIDs (indomethacin and aspirin) induced both necrosis and apoptosis in primary cultures of guinea pig gastric mucosal cells [30]. In this study, we suggested that the direct cytotoxic effect of NSAIDs is independent of the inhibition of COX and suggest that in addition to COX inhibition by NSAIDs, direct cytotoxic effect of NSAIDs is involved in NSAID-induced gastric lesions *in vivo* by use of a combination of the oral administration of selective COX-2 inhibitors with the intravenous administration of non-selective NSAIDs.

## 2. Materials and methods

### 2.1. Chemicals, media, and animals

FBS and trypan blue were from Gibco Co. Indomethacin, aspirin, and NS-398 were from Wako Co. Ibuprofen, diclofenac, and 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) were from Sigma Co. Celecoxib was from LKT Laboratories Inc. Rofecoxib was synthesized in our laboratory. We confirmed that this rofecoxib has bioequivalence with that made by Merck, by measuring their inhibitory effect on inflammatory PG synthesis. Etodolac was gift kindly provided by Nippon Shinyaku Co. The ELISA kits for PGE<sub>2</sub> and 6-keto-PGF<sub>1 $\alpha$</sub>

quantitation were from Cayman Chemical Co. Male Wistar rats weighing 160–190 g and male guinea pigs weighing 200–300 g were purchased from Shimizu Co. The experiments and procedures described here were carried out in accordance with the Guide for the Care and Use of Laboratory Animals as adopted and promulgated by the National Institute of Health and were approved by the Animal Care Committee of Okayama University.

### 2.2. *In vitro* assay of cytotoxicity and DNA fragmentation

Gastric mucosal cells were isolated from guinea pig fundic glands, as described previously [31,32]. Isolated gastric mucosal cells were cultured for 12 hr in RPMI 1640 containing 0.3% (v/v) FBS, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin in type-I collagen-coated plastic culture plates under the conditions of 5% CO<sub>2</sub>/95% air and 37°. After removing non-adherent cells by washing with RPMI 1640, cells that were attached to the plate at about 50% confluence were used. Guinea pig gastric mucosal cells prepared under these conditions were previously characterized, with the majority (about 90%) of cells being identified as pit cells [31]. NSAIDs were dissolved in DMSO and control experiments (without NSAIDs) were performed in the presence of same concentrations of DMSO. Cells were exposed to NSAIDs by changing the entire bathing medium. Cell viability was determined by the trypan blue exclusion test or the MTT method as previously described [33].

Apoptotic DNA fragmentation was monitored as previously described [34]. Cells were collected using a rubber policeman and suspended in 70  $\mu$ L of lysis buffer, consisting of 50 mM Tris-HCl (pH 7.8), 10 mM EDTA, and 0.5% sodium-*N*-lauroylsarcosinate. Proteinase K was added to a final concentration of 1 mg/mL, and the lysate was incubated at 50° for 2 hr. RNase A was then added to a final concentration of 0.5 mg/mL and incubated at 50° for 30 min. These samples were analyzed by 2% agarose gel electrophoresis in the presence of 0.5  $\mu$ g/mL ethidium bromide.

### 2.3. Gastric damage assay

Rats (24 hr fasted) were orally administered with selective COX-2 inhibitors or non-selective NSAIDs with 1% methylcellulose in a volume of 5 mL/kg. Control rats received an equal volume of the vehicle (1% methylcellulose). In some experiments, 1 hr before the oral administration, indomethacin (dissolved in PBS), aspirin (dissolved in PBS), or vehicle (PBS) was administered intravenously *via* the tail vein. Six hours after the oral administration, the rats were anesthetized and the stomach was removed and scored for hemorrhagic damage by an observer unaware of the treatment that the rats had received. The score involved measuring the area of all

lesions in millimeters and summing the values to give an overall gastric lesion index. Determination of PGE<sub>2</sub> levels at the gastric mucosa was done by ELISA as previously described [35].

2.4. Caspase activity assay

The activity of caspase 3 was determined as described previously [30]. Briefly, cells were collected by centrifugation and suspended in extraction buffer (50 mM PIPES

(pH 7.0), 50 mM KCl, 5 mM EGTA, 2 mM MgCl<sub>2</sub>, and 1 mM DTT). Suspensions were sonicated and centrifuged, after which the supernatants were incubated with fluorogenic peptide substrate (Ac-DEVD-MCA) in reaction buffer (100 mM HEPES-KOH (pH 7.5), 10% sucrose, 0.1% CHAPS, and 1 mg/mL BSA) for 15 min at 37°. The release of AMC was determined using a fluorescence spectrophotometer. One unit of protease activity was defined, as the amount of enzyme required to release 1 pmol AMC/min.




Fig. 1. Necrosis and apoptosis induced by various NSAIDs. Cultured guinea pig gastric mucosal cells were incubated with indicated concentrations of various NSAIDs for 1 hr (A, C) or 16 hr (B, C). Cell viability was determined by the MTT method (A, B). Chromosomal DNA was extracted and analyzed by 2% agarose gel electrophoresis (C). Values are mean ± SEM (N = 3).




Fig. 1. (Continued).

### 2.5. Statistical analysis

All values are expressed as the mean  $\pm$  standard error (SEM). One-way ANOVA followed by Scheffe's multiple comparison was used for evaluation of differences between the groups. A Student's *t* test for unpaired results was performed for the evaluation of differences between two groups. Differences were considered to be significant for values of  $P < 0.05$ .

## 3. Results

### 3.1. *In vitro* necrosis and apoptosis induced by various NSAIDs, and their relationship with COX inhibition

We previously reported that short-term (1 hr) treatment of primary cultures of guinea pig gastric mucosal cells with

high concentrations of NSAIDs (indomethacin, 2.5 mM) and long-term (16 hr) treatment of these cells with low concentrations of NSAIDs (indomethacin, 1 mM) induced necrosis and apoptosis, respectively [30]. In the present study, selective COX-2 inhibitors (etodolac, NS-398, celecoxib, and rofecoxib) and non-selective NSAIDs (or slightly selective NSAIDs for COX-1 or COX-2) (indomethacin, diclofenac, and ibuprofen) were tested for their ability to induce necrosis and apoptosis *in vitro*. Necrosis and apoptosis were assessed on the basis of the presence and absence, respectively, of apoptotic DNA fragmentation and caspase 3 activation. The decrease in cell viability with short-term (1 hr) NSAIDs treatment (Fig. 1A) is not associated with apoptotic DNA fragmentation (Fig. 1C) and caspase 3 activation (Table 1), suggesting that it is mediated by necrosis. To confirm this finding, we carried out double-staining experiments with propidium iodide and Hoechst 33342. Since necrotic cells lose

Table 1  
Activation of caspase 3 by NSAIDs

Incubation (hr)	NSAIDs	Caspase 3-like activity (U/mg protein)
1	Control	10 ± 3
	4 mM indomethacin	11 ± 1
	4 mM diclofenac	13 ± 6
	10 mM ibuprofen	13 ± 2
	10 mM etodolac	10 ± 3
	2 mM NS-398	11 ± 5
	0.5 mM celecoxib	10 ± 5
	6 mM rofecoxib	10 ± 4
16	Control	14 ± 4
	1 mM indomethacin	525 ± 23 <sup>***</sup>
	1 mM diclofenac	564 ± 50 <sup>***</sup>
	3 mM ibuprofen	591 ± 32 <sup>***</sup>
	3 mM etodolac	455 ± 29 <sup>***</sup>
	0.25 mM NS-398	473 ± 33 <sup>***</sup>
	0.125 mM celecoxib	637 ± 61 <sup>***</sup>
	3 mM rofecoxib	29 ± 8

Cultured guinea pig gastric mucosal cells were incubated with indicated concentrations of various NSAIDs for 1 or 16 hr. Activities of caspase 3 were examined by the use of specific fluorogenic peptide substrates (Ac-DEVD-MCA). Values are mean ± SEM (N = 3).

<sup>\*\*\*</sup>P < 0.001.

their membrane integrity, propidium iodide staining causes pink nuclear staining in necrotic cells, whereas living cells and apoptotic cells are not stained with propidium iodide. We previously reported that 1 hr treatment with 2.5 mM indomethacin caused pink nuclear staining [30]. We here performed the same type of experiment for other NSAIDs used in Fig. 1 and found that 1 hr treatment with 4 mM indomethacin, 4 mM diclofenac, 10 mM ibuprofen, 10 mM etodolac, 2 mM NS-398, or 0.5 mM celecoxib caused pink nuclear staining (data not shown), strongly

suggesting that cell death shown in Fig. 1A is mediated through necrosis. In contrast, the decrease in cell viability with long-term (16 hr) NSAIDs treatment (Fig. 1B) is associated with apoptotic DNA fragmentation (Fig. 1C) and caspase 3 activation (Table 1), suggesting that it is mediated by apoptosis. Among all of NSAIDs tested here, only rofecoxib induced neither necrosis nor apoptosis (Fig. 1). Interestingly, apoptosis induced by NS-398 was greatest at a concentration of 0.25 mM and the higher concentrations of NS-398 caused the less induction. We have no explanation for this phenomenon at present. We consider that the concentrations of NSAIDs required for necrosis and apoptosis *in vitro* are possible *in vivo* associating with gastric ulceration in animal models, as discussed in our previous paper [30]. However, it is unclear whether gastric mucosal cells can be exposed to NSAIDs as long as 16 hr in animal models. Furthermore, it is also unclear whether these concentrations are relevant for clinical use of NSAIDs. There seemed to be no direct relationship between the cytotoxicity (concentrations of NSAIDs required for necrosis and apoptosis) and the selectivity for COX-2 of NSAIDs.

The level of PGE<sub>2</sub> in the medium upon treatment of cells with various NSAIDs was measured. Compared to non-selective NSAIDs, higher concentrations of selective COX-2 inhibitors were required for inhibiting PGE<sub>2</sub> synthesis, being consistent with the idea that the majority of COX activity is derived from COX-1 activity in gastric mucosal cells [10,11]. Comparing results (Figs. 1 and 2), it is clear that there is no relationship between them in terms of their cytotoxicity and their ability to inhibit PGE<sub>2</sub> synthesis. For example, celecoxib was the strongest compound in terms of cytotoxicity, but the weakest for inhibiting PGE<sub>2</sub> synthesis. Therefore, it seems that the cytotoxic




Fig. 2. Inhibition of PGE<sub>2</sub> synthesis by exposure to various NSAIDs in cultured cells. Cultured guinea pig gastric mucosal cells were incubated with indicated concentrations of various NSAIDs for 30 min, following which the levels of PGE<sub>2</sub> in the media were determined by ELISA. Values are expressed as relative to control (without NSAIDs) ( $2 \times 10^{-9}$  M).

effects of NSAIDs (necrosis and apoptosis) are independent of their ability to inhibit COX. For further confirmation of this point, we examined the effect of exogenously added PGE<sub>2</sub> on necrosis and apoptosis induced by indomethacin. Exogenously added PGE<sub>2</sub> did not affect the extent of cell death by short-term or long-term treatment with indomethacin (necrosis or apoptosis, respectively) even at higher concentrations of PGE<sub>2</sub> than is present endogenously in medium (10<sup>-9</sup> M) (Fig. 3).

It was recently reported that prostacyclin (PGI<sub>2</sub>) protects cells from apoptosis [36,37]. Therefore, we also measured the level of PGI<sub>2</sub> in the medium. Since PGI<sub>2</sub> is very unstable in medium, we determined the level of 6-keto-PGF<sub>1α</sub> (metabolite of PGI<sub>2</sub>) instead of PGI<sub>2</sub>. In the absence of NSAIDs, the concentration of 6-keto-PGF<sub>1α</sub> in the medium was 0.7 nM. The IC<sub>50</sub> value of indomethacin and celecoxib for inhibiting 6-keto-PGF<sub>1α</sub> synthesis was about 5 × 10<sup>-9</sup> M. We also examined the effect of exogenously added PGI<sub>2</sub> on necrosis and apoptosis induced

by indomethacin. Due to the instability of PGI<sub>2</sub> in medium, we used carbaprostacyclin (stable analogue of PGI<sub>2</sub>) instead of PGI<sub>2</sub>. Exogenously added carbaprostacyclin did not affect the extent of cell death by short-term or long-term treatment with indomethacin (necrosis or apoptosis, respectively) (Fig. 3C and D). These results suggest that inhibition of PGI<sub>2</sub> synthesis by NSAIDs is not involved in NSAID-induced necrosis and apoptosis.

### 3.2. Development of gastric lesions by a combination of the oral administration of selective COX-2 inhibitors with the intravenous administration of non-selective NSAIDs

We considered that not only COX inhibition (inhibition of PG synthesis) but also the COX-independent direct cytotoxic effect of NSAIDs is involved in the development of gastrointestinal lesions *in vivo*. For testing this idea by pharmacological experiments, it is necessary to separate




Fig. 3. Effect of PGE<sub>2</sub> on necrosis and apoptosis induced by NSAIDs. Cultured guinea pig gastric mucosal cells were incubated with 2.5 mM (A, C) or 0.9 mM (B, D) indomethacin for 1 hr (A, C) or 16 hr (B, D) in the presence of indicated concentrations of PGE<sub>2</sub> (A, B) or carbaprostacyclin (C, D). Cell viability was determined by the trypan blue exclusion test. Values are mean ± SEM (N = 3). Similar results were obtained by MTT assay.




Fig. 4. Production of gastric lesions in rats. Rats were intravenously administered with 3 mg/kg indomethacin (A), both 3 mg/kg indomethacin and indicated dose of etodolac (B), 100 mg/kg aspirin (C), or vehicle. After 1 hr, animals were orally administered with NSAIDs as indicated or vehicle (A, C) (no oral administration (B)). After 6 hr, the stomach was removed and scored for hemorrhagic damage. Values are mean  $\pm$  SEM (N = 6). \*\*\* $P$  < 0.001; \*\* $P$  < 0.01; \* $P$  < 0.05 (both intravenously and orally administered groups vs. only orally administered groups). n.d.; not detected.




Fig. 4. (Continued).

these two properties of NSAIDs (i.e. COX inhibition and direct cytotoxicity) in the model of NSAID-induced gastric lesions *in vivo*. We tried to achieve this by employing intravenous administration of a non-selective NSAID (indomethacin) and oral administration of selective COX-2 inhibitors in rats. Intravenous administration of non-selective NSAIDs may cause inhibition of both COX-1 and COX-2 (thus inhibition of PG synthesis) at the gastric mucosa without any direct cytotoxicity to the gastric mucosa, because the concentration of NSAIDs at the gastric mucosa following intravenous administration is much lower compared to when NSAIDs are orally administered. On the other hand, oral administration of selective COX-2 inhibitors (except rofecoxib) may cause direct cytotoxicity to the gastric mucosa without inhibition of COX-1, and thus PG synthesis may be maintained.

Intravenous administration of indomethacin (3 mg/kg) in rats did not produce gastric lesions (Fig. 4A) even though the level of PGE<sub>2</sub> at the gastric mucosa was reduced by more than 90% (Table 2). These data suggest that inhibition of COX is not sufficient to produce gastric lesions. On the other hand, oral administration of selective COX-2 inhibitors (etodolac, NS-398, celecoxib, and rofecoxib) did not by themselves (i.e. without intravenous administration of indomethacin) produce gastric lesions (Fig. 4A). PGE<sub>2</sub> synthesis

at the gastric mucosa was not inhibited by the oral administration of these selective COX-2 inhibitors (except for weak inhibition by 15 mg/kg etodolac) (Table 1). Therefore, the absence of gastric lesions only by oral administration of these selective COX-2 inhibitors can be explained by the fact that inhibition of PG synthesis is required for the development of gastric lesions by NSAIDs.

Table 2  
Inhibition of gastric PGE<sub>2</sub> synthesis by NSAIDs *in vivo*

NSAIDs	Gastric PGE <sub>2</sub> (ng/g tissue)
Control	26.8 ± 1.0
3 mg/kg indomethacin i.v.	1.4 ± 0.4 <sup>***</sup>
100 mg/kg aspirin i.v.	1.4 ± 0.2 <sup>***</sup>
3.8 mg/kg indomethacin p.o.	2.3 ± 0.4 <sup>***</sup>
3.8 mg/kg diclofenac p.o.	12.2 ± 1.1 <sup>***</sup>
7.5 mg/kg diclofenac p.o.	3.5 ± 1.5 <sup>***</sup>
7.5 mg/kg etodolac p.o.	18.2 ± 3.8
15 mg/kg etodolac p.o.	5.1 ± 1.4 <sup>***</sup>
30 mg/kg NS-398 p.o.	20.2 ± 3.2
15 mg/kg celecoxib p.o.	25.3 ± 1.3
30 mg/kg rofecoxib p.o.	19.9 ± 4.1

Rats were intravenously (i.v.) or orally (p.o.) administered with indicated doses of NSAIDs. After 6 hr (p.o.) or 7 hr (i.v.), the level of PGE<sub>2</sub> in gastric mucosa was determined by ELISA. Values are mean ± SEM (N = 4–6).

<sup>\*\*\*</sup>P < 0.001.

Interestingly, a combination of intravenous administration of indomethacin and oral administration of COX-2-selective inhibitors (except rofecoxib) clearly produced gastric lesions (Fig. 4A). On the other hand, a combination of intravenous administration of indomethacin and oral administration of rofecoxib did not significantly produce gastric lesions (Fig. 4A). We repeated experiments in Fig. 4A using piroxicam instead of indomethacin and obtained similar results (data not shown). Since among all of COX-2-selective inhibitors, only rofecoxib did not show direct cytotoxicity *in vitro* (Fig. 1), results in Fig. 4A suggest that direct cytotoxicity of NSAIDs is involved in production of gastric lesions. However, since it was recently proposed that inhibition of both COX-1 and COX-2 is required for the development of gastric lesions by NSAIDs [17,19,20], and indomethacin has a weak selectivity for COX-1 (1:0.3) [38], one can argue that the inhibition of COX-2 by intravenously administered indomethacin was not enough and that orally administered selective COX-2 inhibitors inhibited the remaining COX-2 activity, thereby resulting in the development of gastric lesions. However, this possibility was ruled out by an experiment which showed that intravenous administration of both indomethacin (3 mg/kg) and etodolac (15 mg/kg), which must inhibit both COX-1 and COX-2 [39], did not produce gastric lesions (Fig. 4B). We also showed that a combination of intravenous administration of indomethacin and oral administration of SC-560 (selective COX-1 inhibitor) produced gastric lesions (data not shown). Furthermore, we confirmed that the level of PGE<sub>2</sub> at the gastric mucosa with intravenous administration of indomethacin was not further decreased by oral administration of COX-2-selective inhibitors (data not shown). These combined results support our idea that not only COX inhibition (inhibition of PG synthesis) but also the COX-independent direct cytotoxic effect of NSAIDs is involved in the development of gastric lesions *in vivo*.

On the other hand, oral administration of indomethacin did produce gastric lesions without intravenous administration of indomethacin (Fig. 4A). Based on the hypothesis described above, these data can be explained by the fact that orally administered indomethacin had not only a direct cytotoxic effect but also resulted in COX inhibition (inhibition of PG synthesis) (Table 2), and thus produced gastric lesions without intravenous administration of indomethacin. Production of gastric lesions by oral administration of 3.8 but not 7.5 mg/kg diclofenac was increased by the prior intravenous administration of indomethacin (Fig. 4A), which may be related to the fact that orally administration of 7.5 mg/kg but not 3.8 mg/kg diclofenac inhibited PG synthesis completely (about 90%) (Table 2).

We also intravenously administered aspirin, a non-selective NSAID, instead of indomethacin (Fig. 4C). Administration of aspirin alone in this way did not produce gastric lesions, but lesions were produced when etodolac was administered orally in conjunction with intravenously

administered aspirin (Fig. 4C). Similar results were obtained for NS-398 and celecoxib (data not shown). This result not only supports our hypothesis but also provides us with an important suggestion for the clinical use of selective COX-2 inhibitors (see Section 4).

#### 4. Discussion

In this study, we have shown that the cytotoxic effects (necrosis and apoptosis) of NSAIDs on gastric mucosal cells *in vitro* are independent of COX inhibition by those NSAIDs. Furthermore, *in vivo* analysis using both oral and intravenous administration of NSAIDs suggested that not only COX inhibition but also the COX-independent direct cytotoxic effect of NSAIDs is involved in the development of gastric lesions. Both increase in aggressive factors and decrease in defensive factors cause gastropathy. As for NSAID-induced gastropathy, the decrease in defensive factors by NSAIDs (inhibition of PG synthesis) had been paid much attention. Results in this paper suggested that increase in aggressive factors by NSAIDs (direct cytotoxic effect of NSAIDs) is also involved in NSAID-induced gastropathy. This finding can be used to explain the previously unsolved issue that the decrease in PG levels and gastrointestinal lesions by NSAIDs are not always linked (see Section 1). Our findings can explain the fact that higher doses of NSAIDs were required for producing gastric lesions than those for inhibiting COX at the gastric mucosa [21,22], because higher concentrations were required for inducing the direct cytotoxic effect of NSAIDs than were required for inhibiting PG synthesis (Figs. 1 and 2). On the other hand, the fact that parenterally administered NSAIDs causes gastric and duodenal lesions [40–42] and that immunoneutralization of PGs causes gastric lesions [43] show that the direct cytotoxicity of NSAIDs is not essential for the development of gastric lesions by NSAIDs. Furthermore, at present, it is possible that mechanisms other than direct cytotoxicity and COX inhibition (such as increase in gastric motility by NSAIDs) are involved in results in Fig. 4. For example, the beneficial roles of COX-2 at gastric mucosa, such as stimulation of wound healing and resolution of inflammation, were reported [44]. Therefore, inhibition of these beneficial roles of COX-2 by oral administration of selective COX-2 inhibitors may be partly involved in results in Fig. 4. The mechanism of the direct cytotoxicity of NSAIDs and lack of the direct cytotoxicity in rofecoxib are also unclear at present. Furthermore, we have no direct evidence that necrosis and apoptosis are induced, accompanying with production of gastric lesions by NSAIDs *in vivo*.

A recently raised issue concerning the use of selective COX-2 inhibitors is their potential risk for cardiovascular thrombotic events [24,25], although there are still discussions on this point. PGI<sub>2</sub>, a potent anti-aggregator of

platelets and a vasodilator, is mainly produced by COX-2 in vascular endothelial cells, while thromboxane A<sub>2</sub>, a potent aggregator of platelets and a vasoconstrictor, is mainly produced by COX-1 in platelets [45–47]. Therefore, selective COX-2 inhibitors, but not non-selective NSAIDs, may lead to increased prothrombotic activity. Recent genetic studies using knockout mice for the receptor of PGI<sub>2</sub> or thromboxane A<sub>2</sub> supported this notion [48,49]. Furthermore, both animal and clinical data suggest that, compared to non-selective NSAIDs, selective COX-2 inhibitors increase cardiovascular thrombotic events [24,50,51]. Therefore, the method for decreasing the gastric side effects of NSAIDs other than increasing their selectivity for COX-2 may be useful in order to develop safer NSAIDs for both gastrointestinal and cardiovascular. Considering our hypothesis described above, NSAIDs that do not exhibit direct cytotoxicity on gastric mucosal cells (i.e. NSAIDs that do not induce necrosis and apoptosis in gastric mucosal cells) may be safe for the gastrointestinal tract even if they do not have high selectivity for COX-2.

Low doses of aspirin are widely used for preventing thrombosis. Therefore, it is not unusual that patients who use aspirin chronically for preventing thrombosis are further administered with selective COX-2 inhibitors as anti-inflammatory drugs. Results (Fig. 4C) suggest that the oral administration of selective COX-2 inhibitors into chronic aspirin users causes gastric lesions, even though such administration into non-aspirin users is safe. Similar results were reported recently using simultaneous oral administration of aspirin and selective COX-2 inhibitors [52]. In fact, the Celecoxib Long-term Arthritis Safety Study (CLASS) showed that, compared to non-selective NSAIDs, celecoxib (at dosages greater than those indicated clinically) was clearly associated with a lower incidence of symptomatic lesions and lesion complication for patients not taking aspirin concomitantly. The difference in gastric side effects between them (non-selective NSAIDs and celecoxib) was not so clear for patients taking aspirin concomitantly [13]. Therefore, much attention should be paid to the concomitant use of both aspirin and selective COX-2 inhibitors.

#### Acknowledgments

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Health, Labour, and Welfare of Japan.

#### References

- [1] Smalley WE, Ray WA, Daugherty JR, Griffin MR. Nonsteroidal anti-inflammatory drugs and the incidence of hospitalizations for peptic ulcer disease in elderly persons. *Am J Epidemiol* 1995;141: 539–45.
- [2] Hawkey CJ. Nonsteroidal anti-inflammatory drug gastropathy. *Gastroenterology* 2000;119:521–35.
- [3] Barrier CH, Hirschowitz BI. Controversies in the detection and management of nonsteroidal anti-inflammatory drug-induced side effects of the upper gastrointestinal tract. *Arthritis Rheum* 1989;32: 926–32.
- [4] Gabriel SE, Jaakkimainen L, Bombardier C. Risk for serious gastrointestinal complications related to use of nonsteroidal anti-inflammatory drugs. A meta-analysis. *Ann Intern Med* 1991;115:787–96.
- [5] Fries JF, Miller SR, Spitz PW, Williams CA, Hubert HB, Bloch DA. Toward an epidemiology of gastropathy associated with nonsteroidal anti-inflammatory drug use. *Gastroenterology* 1989;96:647–55.
- [6] Kurata JH, Abbey DE. The effect of chronic aspirin use on duodenal and gastric ulcer hospitalizations. *J Clin Gastroenterol* 1990;12:260–6.
- [7] Singh G. Recent considerations in nonsteroidal anti-inflammatory drug gastropathy. *Am J Med* 1998;105:31S–8S.
- [8] Vane JR, Botting RM. Mechanism of action of anti-inflammatory drugs. *Scand J Rheumatol Suppl* 1996;102:9–21.
- [9] Miller TA. Protective effects of prostaglandins against gastric mucosal damage: current knowledge and proposed mechanisms. *Am J Physiol* 1983;245:G601–23.
- [10] Vane J. Towards a better aspirin. *Nature* 1994;367:215–6.
- [11] Smith CJ, Zhang Y, Koboldt CM, Muhammad J, Zweifel BS, Shaffer A, Talley JJ, Masferrer JL, Seibert K, Isakson PC. Pharmacological analysis of cyclooxygenase-1 in inflammation. *Proc Natl Acad Sci USA* 1998;95:13313–8.
- [12] Chan CC, Boyce S, Brideau C, Charleson S, Cromlish W, Ethier D, Evans J, Ford HA, Forrest MJ, Gauthier JY, Gordon R, Gresser M, Guay J, Kargman S, Kennedy B, Leblanc Y, Leger S, Mancini J, O'Neill GP, Ouellet M, Patrick D, Percival MD, Perrier H, Prasit P, Rodger I, Tagari P, Therien M, Vickers P, Visco D, Wang Z, Webb J, Wong E, Xu LJ, Young RN, Zamboni R, Riendeau D. Rofecoxib [Vioxx, MK-0966; 4-(4'-methylsulfonylphenyl)-3-phenyl-2-(5H)-furanone]: a potent and orally active cyclooxygenase-2 inhibitor. Pharmacological and biochemical profiles. *J Pharmacol Exp Ther* 1999; 290:551–60.
- [13] Silverstein FE, Faich G, Goldstein JL, Simon LS, Pincus T, Whelton A, Makuch R, Eisen G, Agrawal NM, Stenson WF, Burr AM, Zhao WW, Kent JD, Lefkowitz JB, Verburg KM, Geis GS. Gastrointestinal toxicity with celecoxib vs. nonsteroidal anti-inflammatory drugs for osteoarthritis and rheumatoid arthritis: the CLASS study: a randomized controlled trial. Celecoxib Long-term Arthritis Safety Study. *JAMA* 2000;284:1247–55.
- [14] Bombardier C, Laine L, Reicin A, Shapiro D, Burgos VR, Davis B, Day R, Ferraz MB, Hawkey CJ, Hochberg MC, Kvien TK, Schnitzer TJ. Comparison of upper gastrointestinal toxicity of rofecoxib and naproxen in patients with rheumatoid arthritis. VIGOR Study Group. *New Engl J Med* 2000;343:1520–8 [2 pages following 1528].
- [15] FitzGerald GA, Patrono C. The coxibs, selective inhibitors of cyclooxygenase-2. *New Engl J Med* 2001;345:433–42.
- [16] Juni P, Rutjes AW, Dieppe PA. Are selective COX 2 inhibitors superior to traditional non steroidal anti-inflammatory drugs? *BMJ* 2002;324: 1287–8.
- [17] Wallace JL, McKnight W, Reuter BK, Vergnolle N. NSAID-induced gastric damage in rats: requirement for inhibition of both cyclooxygenase 1 and 2. *Gastroenterology* 2000;119:706–14.
- [18] Langenbach R, Morham SG, Tiano HF, Loftin CD, Ghanayem BI, Chulada PC, Mahler JF, Lee CA, Goulding EH, Kluckman KD, Kim HS, Smithies O. Prostaglandin synthase 1 gene disruption in mice reduces arachidonic acid-induced inflammation and indomethacin-induced gastric ulceration. *Cell* 1995;83:483–92.
- [19] Tanaka A, Araki H, Hase S, Komoike Y, Takeuchi K. Up-regulation of COX-2 by inhibition of COX-1 in the rat: a key to NSAID-induced gastric injury. *Aliment Pharmacol Ther* 2002;16(Suppl 2):90–101.
- [20] Gretzer B, Maricic N, Respondek M, Schuligoi R, Peskar BM. Effects of specific inhibition of cyclo-oxygenase-1 and cyclo-oxygenase-2 in

- the rat stomach with normal mucosa and after acid challenge. *Br J Pharmacol* 2001;132:1565–73.
- [21] Ligumsky M, Golanska EM, Hansen DG, Kauffman GJ. Aspirin can inhibit gastric mucosal cyclo-oxygenase without causing lesions in rat. *Gastroenterology* 1983;84:756–61.
- [22] Ligumsky M, Sestieri M, Karmeli F, Zimmerman J, Okon E, Rachmilewitz D. Rectal administration of nonsteroidal antiinflammatory drugs. Effect on rat gastric ulcerogenicity and prostaglandin E2 synthesis. *Gastroenterology* 1990;98:1245–9.
- [23] Lichtenberger LM. Where is the evidence that cyclooxygenase inhibition is the primary cause of nonsteroidal anti-inflammatory drug (NSAID)-induced gastrointestinal injury? Topical injury revisited. *Biochem Pharmacol* 2001;61:631–7.
- [24] Mukherjee D, Nissen SE, Topol EJ. Risk of cardiovascular events associated with selective COX-2 inhibitors. *JAMA* 2001;286:954–9.
- [25] Mukherjee D. Selective cyclooxygenase-2 (COX-2) inhibitors and potential risk of cardiovascular events. *Biochem Pharmacol* 2002;63:817–21.
- [26] Takeuchi K, Ueki S, Okabe S. Importance of gastric motility in the pathogenesis of indomethacin-induced gastric lesions in rats. *Dig Dis Sci* 1986;31:1114–22.
- [27] Asako H, Kubes P, Wallace J, Gaginella T, Wolf RE, Granger DN. Indomethacin-induced leukocyte adhesion in mesenteric venules: role of lipoxygenase products. *Am J Physiol* 1992;262:G903–8.
- [28] Ashley SW, Sonnenschein LA, Cheung LY. Focal gastric mucosal blood flow at the site of aspirin-induced ulceration. *Am J Surg* 1985;149:53–9.
- [29] Lichtenberger LM, Wang ZM, Romero JJ, Ulloa C, Perez JC, Giraud MN, Barreto JC. Non-steroidal anti-inflammatory drugs (NSAIDs) associate with zwitterionic phospholipids: insight into the mechanism and reversal of NSAID-induced gastrointestinal injury. *Nat Med* 1995;1:154–8.
- [30] Tomisato W, Tsutsumi S, Rokutan K, Tsuchiya T, Mizushima T. NSAIDs induce both necrosis and apoptosis in guinea pig gastric mucosal cells in primary culture. *Am J Physiol Gastrointest Liver Physiol* 2001;281:G1092–100.
- [31] Hirakawa T, Rokutan K, Nikawa T, Kishi K. Geranylgeranylacetone induces heat shock proteins in cultured guinea pig gastric mucosal cells and rat gastric mucosa. *Gastroenterology* 1996;111:345–57.
- [32] Tomisato W, Takahashi N, Komoto C, Rokutan K, Tsuchiya T, Mizushima T. Geranylgeranylacetone protects cultured guinea pig gastric mucosal cells from indomethacin. *Dig Dis Sci* 2000;45:1674–9.
- [33] Tomisato W, Tsutsumi S, Tsuchiya T, Mizushima T. Geranylgeranylacetone protects guinea pig gastric mucosal cells from gastric stressor-induced necrosis by induction of heat-shock proteins. *Biol Pharm Bull* 2001;24:887–91.
- [34] Tsutsumi S, Tomisato W, Takano T, Rokutan K, Tsuchiya T, Mizushima T. Gastric irritant-induced apoptosis in guinea pig gastric mucosal cells in primary culture. *Biochim Biophys Acta* 2002;1589:168–80.
- [35] Futaki N, Arai I, Hamasaka Y, Takahashi S, Higuchi S, Otomo S. Selective inhibition of NS-398 on prostanoid production in inflamed tissue in rat carrageenan-air-pouch inflammation. *J Pharm Pharmacol* 1993;45:753–5.
- [36] Chen MF, Chen JC, Chiu DF, Ng CJ, Shyr MH, Chen HM. Prostacyclin analogue (OP-2507) induces delayed ex vivo neutrophil apoptosis and attenuates reperfusion-induced hepatic microcirculatory derangement in rats. *Shock* 2001;16:473–8.
- [37] Cutler NS, Graves-Deal R, LaFleur BJ, Gao Z, Boman BM, Whitehead RH, Terry E, Morrow JD, Coffey RJ. Stromal production of prostacyclin confers an antiapoptotic effect to colonic epithelial cells. *Cancer Res* 2003;63:1748–51.
- [38] Kawai S, Nishida S, Kato M, Furumaya Y, Okamoto R, Koshino T, Mizushima Y. Comparison of cyclooxygenase-1 and -2 inhibitory activities of various nonsteroidal anti-inflammatory drugs using human platelets and synovial cells. *Eur J Pharmacol* 1998;347:87–94.
- [39] Wallace JL, Bak A, McKnight W, Asfaha S, Sharkey KA, MacNaughton WK. Cyclooxygenase 1 contributes to inflammatory responses in rats and mice: implications for gastrointestinal toxicity. *Gastroenterology* 1998;115:101–9.
- [40] Henry D, Dobson A, Turner C. Variability in the risk of major gastrointestinal complications from nonaspirin nonsteroidal anti-inflammatory drugs. *Gastroenterology* 1993;105:1078–88.
- [41] Komoike Y, Takeeda M, Tanaka A, Kato S, Takeuchi K. Prevention by parenteral aspirin of indomethacin-induced gastric lesions in rats: mediation by salicylic acid. *Dig Dis Sci* 2002;47:1538–45.
- [42] Wallace JL, McKnight GW. Characterization of a simple animal model for nonsteroidal anti-inflammatory drug induced antral ulcer. *Can J Physiol Pharmacol* 1993;71:447–52.
- [43] Redfern JS, Feldman M. Role of endogenous prostaglandins in preventing gastrointestinal ulceration: induction of ulcers by antibodies to prostaglandins. *Gastroenterology* 1989;96:596–605.
- [44] Mizuno H, Sakamoto C, Matsuda K, Wada K, Uchida T, Noguchi H, Akamatsu T, Kasuga M. Induction of cyclooxygenase 2 in gastric mucosal lesions and its inhibition by the specific antagonist delays healing in mice. *Gastroenterology* 1997;112:387–97.
- [45] McAdam BF, Catella LF, Mardini IA, Kapoor S, Lawson JA, FitzGerald GA. Systemic biosynthesis of prostacyclin by cyclooxygenase (COX)-2: the human pharmacology of a selective inhibitor of COX-2. *Proc Natl Acad Sci USA* 1999;96:272–7.
- [46] Catella LF, McAdam B, Morrison BW, Kapoor S, Kujubu D, Antes L, Lasseter KC, Quan H, Gertz BJ, FitzGerald GA. Effects of specific inhibition of cyclooxygenase-2 on sodium balance, hemodynamics, and vasoactive eicosanoids. *J Pharmacol Exp Ther* 1999;289:735–41.
- [47] Belton O, Byrne D, Kearney D, Leahy A, Fitzgerald DJ. Cyclooxygenase-1 and -2-dependent prostacyclin formation in patients with atherosclerosis. *Circulation* 2000;102:840–5.
- [48] Cheng Y, Austin SC, Rocca B, Koller BH, Coffman TM, Grosser T, Lawson JA, FitzGerald GA. Role of prostacyclin in the cardiovascular response to thromboxane A2. *Science* 2002;296:539–41.
- [49] Murata T, Ushikubi F, Matsuoka T, Hirata M, Yamasaki A, Sugimoto Y, Ichikawa A, Aze Y, Tanaka T, Yoshida N, Ueno A, Oh IS, Narumiya S. Altered pain perception and inflammatory response in mice lacking prostacyclin receptor. *Nature* 1997;388:678–82.
- [50] Hennen JK, Huang J, Barrett TD, Driscoll EM, Willens DE, Park AM, Crofford LJ, Lucchesi BR. Effects of selective cyclooxygenase-2 inhibition on vascular responses and thrombosis in canine coronary arteries. *Circulation* 2001;104:820–5.
- [51] Dowd NP, Scully M, Adderley SR, Cunningham AJ, Fitzgerald DJ. Inhibition of cyclooxygenase-2 aggravates doxorubicin-mediated cardiac injury in vivo. *J Clin Invest* 2001;108:585–90.
- [52] Fiorucci S, de Lima Jr OM, Mencarelli A, Palazzetti B, Distrutti E, McKnight W, Dickey M, Ma L, Romano M, Morelli A, Wallace JL. Cyclooxygenase-2-derived lipoxin A4 increases gastric resistance to aspirin-induced damage. *Gastroenterology* 2002;123:1598–606.

## Induction of Claudin-4 by Nonsteroidal Anti-inflammatory Drugs and Its Contribution to Their Chemopreventive Effect

Shinji Mima,<sup>1</sup> Shinji Tsutsumi,<sup>1</sup> Hironori Ushijima,<sup>1</sup> Miho Takeda,<sup>1</sup> Ikue Fukuda,<sup>1</sup> Kazumi Yokomizo,<sup>1</sup> Keitarou Suzuki,<sup>1</sup> Kuniaki Sano,<sup>2</sup> Tohru Nakanishi,<sup>4</sup> Wataru Tomisato,<sup>2</sup> Tomofusa Tsuchiya,<sup>2</sup> and Tohru Mizushima<sup>1</sup>

<sup>1</sup>Graduate School of Medical and Pharmaceutical Sciences, Kumamoto University, Kumamoto, Japan and <sup>2</sup>Faculty of Pharmaceutical Sciences and <sup>3</sup>Graduate School of Medicine and Dentistry, Okayama University, <sup>4</sup>Department of Clinical Pharmacy, Shujitsu University School of Pharmacy, Okayama, Japan

### Abstract

Nonsteroidal anti-inflammatory drugs (NSAID) have shown chemopreventive effects in both preclinical and clinical studies; however, the precise molecular mechanism governing this response remains unclear. We used DNA microarray techniques to search for genes whose expression is induced by the NSAID indomethacin in human gastric carcinoma (AGS) cells. Among identified genes, we focused on those related to tight junction function (*claudin-4*, *claudin-1*, and *occludin*), particularly *claudin-4*. Induction of *claudin-4* by indomethacin was confirmed at both mRNA and protein levels. NSAIDs, other than indomethacin (diclofenac and celecoxib), also induced *claudin-4*. All of the tested NSAIDs increased the intracellular  $Ca^{2+}$  concentration. Other drugs that increased the intracellular  $Ca^{2+}$  concentration (thapsigargin and ionomycin) also induced *claudin-4*. Furthermore, an intracellular  $Ca^{2+}$  chelator [1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid] inhibited the indomethacin-dependent induction of *claudin-4*. These results strongly suggest that induction of *claudin-4* by indomethacin is mediated through an increase in the intracellular  $Ca^{2+}$  concentration. Overexpression of *claudin-4* in AGS cells did not affect cell growth or the induction of apoptosis by indomethacin. On the other hand, addition of indomethacin or overexpression of *claudin-4* inhibited cell migration. Colony formation in soft agar was also inhibited. Suppression of *claudin-4* expression by small interfering RNA restored the migration activity of AGS cells in the presence of indomethacin. Based on these results, we consider that the induction of *claudin-4* and other tight junction-related genes by NSAIDs may be involved in the chemopreventive effect of NSAIDs through the suppression of anchorage-independent growth and cell migration. (Cancer Res 2005; 65(5): 1-9)

### Introduction

Nonsteroidal anti-inflammatory drugs (NSAID) are the most widely used therapeutic agents in the treatment of pain, inflammation, and fever (1). Recent epidemiologic studies clearly show that NSAID use is associated with a reduced risk of cancer, and preclinical and clinical studies have shown that some NSAIDs are effective for the treatment and prevention of cancer. This effect

is particularly well documented in relation to colon and rectal cancer. Recent studies have also shown that NSAID use reduces the risk of stomach cancer (2, 3). Several different effects of NSAIDs on cancer cells, such as stimulation of apoptosis, cell growth suppression, inhibition of angiogenesis, and inhibition of metastasis, have been proposed to play important roles in NSAID-mediated chemoprevention (4, 5). However, the precise molecular mechanisms governing these effects of NSAIDs have not been elucidated.

The anti-inflammatory action of NSAIDs is mediated through its inhibition of cyclooxygenase (COX). COX is an enzyme essential for the synthesis of prostaglandins, which have a strong propensity for inducing inflammation. Prostaglandins, such as prostaglandin  $E_2$  ( $PGE_2$ ), inhibit apoptosis and stimulate cell growth, angiogenesis, and metastasis (6-8). Furthermore, overexpression of COX-2 (a subtype of COX) has been reported in various tumor cells and tissues (9, 10). Therefore, the inhibition of COX by NSAIDs was thought previously to be the sole explanation for their chemopreventive effect. However, several lines of evidence suggest that chemoprevention by NSAIDs also involves COX-independent mechanisms. Sulindac sulfone, a derivative of the NSAID sulindac, does not inhibit COX activity and has been shown to display antitumor activity *in vivo* as well as induce apoptosis and inhibit cell growth in tumor cells *in vitro* (11, 12). Moreover, the induction by NSAIDs of apoptosis and the inhibition of cell growth in COX-null fibroblasts and tumor cells in which COX expression was absent have been reported (13, 14). Therefore, it is important that the COX-independent mechanisms for chemoprevention by NSAIDs are elucidated to develop more effective NSAIDs.

Tight junctions are the most apical intercellular structure in epithelial and endothelial cells and create a physiologic barrier separating the apical and basolateral spaces; in other words, they create a paracellular permeability barrier. Tight junctions contain the transmembrane proteins occludin and claudin, which are connected to the cytoskeleton via zonula occludens (ZO-1; ref. 15). Several studies have shown a correlation between a reduction in tight junction function and tumor progression. A loss of tight junction structure is frequently observed in epithelium-derived cancers, whereas some tumor-promoting agents are known to disrupt tight junctions (16, 17). Furthermore, overexpression of tight junction-related proteins (such as *claudin-1*, *claudin-4*, and *occludin*) in cancer cells has been reported to induce apoptosis and suppress the invasive potential of these cells (18, 19).

NSAIDs affect the expression of several genes in a COX-independent manner. For example, NSAIDs induce NAG-1, a transforming growth factor- $\beta$  superfamily member protein, which is involved in the induction of apoptosis by NSAIDs (20). We reported recently that NSAIDs induce CCAAT/enhancer binding protein homologous transcription factor, which is involved

Requests for reprints: Tohru Mizushima, Graduate School of Medical and Pharmaceutical Sciences, Kumamoto University, Kumamoto 862-0973, Japan. Phone: 81-96-371-4323; Fax: 81-96-371-4323; E-mail: mizu@gpo.kumamoto-u.ac.jp.  
©2005 American Association for Cancer Research.

evaluation of differences between the groups. The Student's *t* test for unpaired results was done for the evaluation of differences between two groups, which were considered to be significant for values of *P* < 0.05.

**Results**


**DNA Microarray Analysis for Gene Expression in the Presence of Indomethacin.** We used the DNA microarray technique and AGS cells to identify genes whose expression is altered by indomethacin. AGS cells were treated with 0.3 mmol/L indomethacin for 4 hours before microarray analysis. As shown in Fig. 1A, this treatment did not affect cell viability. We did microarray analysis four times (four hybridizations) and selected genes that were induced by indomethacin based on the criteria that the induction was observed in all four hybridizations and that the mean value (fold change) of four hybridizations was >2.0. As shown in

Table 1, 34 genes were identified. Induction of some of these genes, such as *CCAAT/enhancer binding protein β* and *prostate differentiation factor (NAG-1)*, by NSAIDs in other cancer cell types has been reported previously (20, 27). Among these genes, we focused our attention on genes related to tight junction function (*claudin-1*, *claudin-4*, and *occludin*), particularly on *claudin-4*, because the induction was relatively clear, its expression in gastric mucosal cells has been confirmed previously (28), and a recent report showed that overexpression of claudin-4 suppressed anchorage-independent growth and the invasive potential of pancreatic cancer cells (19). Nineteen genes were identified whose expression was repressed by the indomethacin treatment (data not shown).

Changes in the indomethacin-induced expression of these genes were then verified by RT-PCR. As shown in Fig. 1B, the induction of claudin-1, claudin-4, and occludin was confirmed. Results of the

F1

T1



**Figure 1.** Induction of tight junction-related genes by NSAIDs. AGS (A-D and G) or MKN-45, KATO-III, Caco-2, and HCT-15 (E-G) cells were incubated with indicated concentrations of NSAIDs for 4 hours (A-C) or 24 hours (D-F). Results for cells cultured without NSAIDs (G). Cell viability was determined by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide method. Columns, mean (n = 3); bars, SE (A). Total RNA was extracted and subjected to RT-PCR by use of a specific primer for each gene. GAPDH (*G3PDH*; B) or actin (G) was used as a control. Reaction products were analyzed by agarose (1%) gel electrophoresis (B and G). Total RNA samples were analyzed by Northern blotting experiments using a specific DNA probe for claudin-4. Bands of rRNA (28S) stained with ethidium bromide (C). Whole cell extracts (2.5 μg protein) were analyzed by immunoblotting with an antibody against claudin-4 or actin (D-F).

Table 1. List of genes overexpressed in AGS cells following treatment with indomethacin

Gene name	Accession no.	Function	Fold change
Claudin-1	AF115546	Tight junction	2.00
Claudin-4	AK026651	Tight junction	2.54
Occludin	U49184	Tight junction	2.24
Tissue factor pathway inhibitor 2	NM_006528	Blood coagulation	2.46
Zinedin	AF212940	Calmodulin binding protein	2.01
Arginine-rich protein	AA582041	Carcinogenicity	2.00
Human urokinase-type plasminogen receptor, exon 7	U09937	Cell surface plasminogen activation	2.81
Chromobox homologue 4 ( <i>Drosophila</i> Pc class)	AF013956	Cellular memory system	2.30
Human low-density lipoprotein receptor gene, exon 18	L00352	Cholesterol homeostasis	3.43
Low-density lipoprotein receptor (familial hypercholesterolemia)	NM_000527	Cholesterol homeostasis	2.75
Epithelial protein lost in neoplasm $\beta$	AA594624	Cytoskeleton	2.58
Keratin 8	AJ978932	Cytoskeleton	2.37
Immediate early response 3	AJ022951	Differentiation	2.36
Prostate differentiation factor	AB000584	Differentiation	2.00
Procollagen-proline	J02783	Disulfide isomerase/oxidoreductase	2.50
Glucosidase $\beta$ , acid (includes glucosylceramidase)	AF023268	Glucocerebrosidase	2.01
Tumor necrosis factor- $\alpha$ -induced protein 6	M31165	Hyaluronan binding protein family	2.14
Basigin	X64364	Immunoglobulin superfamily	2.15
Solute carrier family 7	M80244	L-Amino acid transporter	3.00
Cathepsin D	M11233	Lysosomal proteinase	2.40
Pim-1 oncogene	M24779	Protein kinase	2.52
Cytochrome <i>c</i> oxidase subunit VIII	J04823	Respiratory	2.90
3,4-Dihydroxy-L-phenylalanine decarboxylase (aromatic L-amino acid decarboxylase)	M76180	Synthesis of dopamine and serotonin	2.27
CCAAT/enhancer binding protein $\beta$	W93514	Transcription factor	4.76
Predicted using Genefinder, preliminary prediction	CAB60892	Tumor protein p53	2.18
ATPase, H <sup>+</sup> transporting, lysosomal (vacuolar proton pump) 21 kDa	A1567477	Vacuolar proton pump	2.39
Ribosomal protein S21	BE221408	Unknown	2.34
Human genomic DNA, chromosome 22q11.2, BCRL2 region	AP000553	Unknown	2.23
Ubiquinol-cytochrome <i>c</i> reductase (6.4 kDa) subunit	AW163002	Unknown	2.13
IFN-induced transmembrane protein 3 (1-8U)	X57352	Unknown	2.10
Conserved hypothetical protein	AAF96700	Unknown	2.07
KIAA0316 gene product	AB002314	Unknown	2.06
Sequence 100 from patent WO9951727	AX015425	Unknown	2.03
Ribosomal protein S28	AW161288	Unknown	2.03

NOTE: Fold changes in gene expression by indomethacin compared with untreated cells. Mean values from four independent hybridizations. AGS cells were treated with or without 0.3 mmol/L indomethacin for 4 hours and subjected to DNA microarray analysis.

real-time RT-PCR experiments used to determine the extent of the induction yielded fold changes in copy number of 2.3, 3.0, and 1.5 for claudin-1, claudin-4, and occludin mRNA, respectively, in response to treatment of cells for 4 hours with 0.3 mmol/L indomethacin. In addition, the induction by indomethacin of claudin-4 mRNA or claudin-4 protein was confirmed using Northern blot analysis (Fig. 1C) and immunoblot analysis (Fig. 1D), respectively.

We then examined whether the induction of claudin-4 by indomethacin is specific to AGS cells or is a general property also observed in other cell types. We used MKN-45 and KATO-III cells (derived from gastric cancer tissue) and Caco-2 and HCT-15 cells (derived from colon cancer tissue) to test this effect. As shown in Fig. 1E, indomethacin induced claudin-4 in each of the cell lines tested, with the concentration of indomethacin required for the induction being similar for each cell line.

Diclofenac, another NSAID, also induced claudin-4 in a dose-dependent manner (Fig. 1D). Some NSAIDs are specific in their

effect on COX, which exists in two forms, COX-1 and COX-2. Celecoxib, a COX-2-specific NSAID, induced claudin-4 not only in AGS cells (Fig. 1D) but also in the other cell lines tested (Fig. 1F). These results suggest that NSAIDs induce claudin-4 irrespective of whether they are specific for COX-2. It has been reported that both COX-1 and COX-2 mRNA are expressed in AGS, MKN-45, and Caco-2 cells, whereas COX-2 mRNA expression is very low in KATO-III and HCT-15 cells (29-33). COX-1 mRNA expression was confirmed by RT-PCR in each of the cell lines tested, whereas COX-2 mRNA expression was detected only in AGS, MKN-45, and Caco-2 cells (Fig. 1G). Therefore, COX-2-specific NSAIDs (in this case, celecoxib) induce claudin-4 not only in COX-2-expressing cells but also in cells lacking COX-2 expression. Furthermore, whereas indomethacin inhibited both COX-1 and COX-2 at a concentration of <1 mol/L (34), the induction of claudin-4 required higher concentrations (Fig. 1). These findings strongly suggest that NSAIDs induce claudin-4 independently of COX-inhibition.

**Mechanism for Induction of Claudin-4 by Indomethacin.** For further confirmation that NSAIDs induce claudin-4 independently of COX-inhibition, we examined the effect of PGE<sub>2</sub>, a major prostaglandin in the gastric mucosa, on the induction of claudin-4 by indomethacin. PGE<sub>2</sub> (0.1-10 μmol/L) did not affect the level of claudin-4 in the presence and absence of indomethacin (Fig. 2A). We determined previously the level of PGE<sub>2</sub> in the culture medium of AGS cells to be ~10 nmol/L (23). Therefore, inhibition of PGE<sub>2</sub> synthesis by indomethacin does not seem to be involved in the induction of claudin-4 by indomethacin.


Recent studies suggest that indomethacin and other NSAIDs act as agonists of the peroxisome proliferator-activated receptor-γ (35). To test the contribution of this activity to the induction of claudin-4 by indomethacin, we examined the effect of a peroxisome proliferator-activated receptor-γ antagonist (GW9662) on the induction of claudin-4 by indomethacin. As shown in Fig. 2B, GW9662 did not inhibit but rather slightly heightened the induction of claudin-4 by indomethacin. The different concentrations of GW9662 tested did not affect cell viability (data not shown), but based on data from a previous report, these concentrations are considered sufficient to inhibit agonist binding to peroxisome proliferator-activated receptor-γ (36). Therefore, peroxisome proliferator-activated receptor-γ does not seem to be associated with the induction of claudin-4 by indomethacin.

It has been reported that some NSAIDs increase reactive oxygen species production (37). To test the contribution of reactive oxygen species to the induction of claudin-4 by indomethacin, we examined the effects of the antioxidants *N*-acetylcysteine and SOD. As shown in Fig. 2C and D, neither *N*-acetylcysteine nor SOD affected claudin-4 expression in either the presence or the absence of indomethacin. Activation of the extracellular signal-regulated kinase pathway—one

of the mitogen-activated protein kinase pathways—has been reported to stimulate the expression of claudin-4. Although some NSAIDs have been reported to activate the extracellular signal-regulated kinase pathway (19, 38), an inhibitor of extracellular signal-regulated kinase (PD98059) did not affect the expression of claudin-4 in either the presence or the absence of indomethacin (Fig. 2E). *N*-acetylcysteine, SOD, and PD98059 did not affect cell viability at the concentrations used (data not shown). These results suggest that neither reactive oxygen species nor extracellular signal-regulated kinase is responsible for the induction of claudin-4 by indomethacin.

Some NSAIDs have been reported to increase the intracellular Ca<sup>2+</sup> concentration, [Ca<sup>2+</sup>]<sub>i</sub> (39, 40). In this study, we tested whether an increase in [Ca<sup>2+</sup>]<sub>i</sub> by NSAIDs is responsible for the induction of claudin-4. Firstly, we confirmed that a NSAID-induced increase in [Ca<sup>2+</sup>]<sub>i</sub> occurred under the same conditions as those in which the induction of claudin-4 in AGS cells was observed. As shown in Fig. 3A, all NSAIDs tested (indomethacin, diclofenac, and celecoxib) increased [Ca<sup>2+</sup>]<sub>i</sub> at the same NSAID concentrations that caused the induction of claudin-4.

Some drugs that are known to increase [Ca<sup>2+</sup>]<sub>i</sub> were examined for their capacity to induce claudin-4 expression. The actions of thapsigargin, an inhibitor of the sarcoplasmic/endoplasmic reticulum Ca<sup>2+</sup> ATPase, and the Ca<sup>2+</sup> ionophore ionomycin were thus tested on AGS cells. As shown in Fig. 3A-C, in addition to increasing [Ca<sup>2+</sup>]<sub>i</sub>, both thapsigargin and ionomycin induced claudin-4 in a dose-dependent manner. Furthermore, an intracellular Ca<sup>2+</sup> chelator, 1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid, was found to inhibit the induction of claudin-4 not only by ionomycin but also by indomethacin (Fig. 3D). 1,2-Bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid did not affect cell viability at the concentration used in these experiments (data not



**Figure 2.** Mechanism for the induction of claudin-4 by indomethacin. AGS cells were incubated with or without 0.3 mmol/L indomethacin for 24 hours in the presence of indicated concentrations of PGE<sub>2</sub> (A), GW9662 (B), *N*-acetylcysteine (NAC; C), SOD (D), or PD98059 (E). Levels of claudin-4 and actin were estimated by immunoblotting experiments as described in Fig. 1. One unit of SOD was evaluated based on its inhibitory effect on the reduction of cytochrome *c* as described in the manufacturer's instructions.




Figure 3. Involvement of the intracellular  $Ca^{2+}$  concentration in the induction of claudin-4 by NSAIDs. The intracellular  $Ca^{2+}$  concentration,  $[Ca^{2+}]_i$ , was monitored by a fluo-3/AM assay system as described in Materials and Methods. Indomethacin (0.3 mmol/L), celecoxib (0.02 mmol/L), diclofenac (0.1 mmol/L), thapsigargin (5 nmol/L), or ionomycin (0.5  $\mu$ mol/L) were added to fluo-3/AM-loaded cells at indicated concentrations, and the time course of changes to fluo-3 fluorescence was monitored (A). AGS cells were incubated with indicated concentrations of thapsigargin (B) or ionomycin (C) for 24 hours. AGS cells were incubated with 0.3 mmol/L indomethacin (IND) or 2  $\mu$ mol/L ionomycin (ION) in the presence of indicated concentrations of 1,2-bis(2-aminophenoxy)ethane-*N,N,N',N'*-tetraacetic acid (BAPTA-AM) for 24 hours (D). Levels of claudin-4 and actin were estimated by immunoblotting experiments as described in Fig. 1 (B-D).

shown). These results strongly suggest that induction of claudin-4 by indomethacin is mediated via an increase in  $[Ca^{2+}]_i$ .

**Role of Claudin-4 Induction in the *In vitro* Antitumor Action of NSAIDs.** As described in Introduction, various mechanisms have been proposed for the chemopreventive action of NSAIDs; these include the inhibition of cell growth, stimulation of apoptosis, and inhibition of metastasis. Here, we examined the contribution that NSAID induction of claudin-4 makes to the antitumor effect of NSAIDs *in vitro*. We constructed stable transfectants of AGS cells that continuously overexpress claudin-4 and selected four clones (clones 1, 6, 7, and 11) in which the level of expression of claudin-4 varied (clone 7 > clone 11 > clone 1 > clone 6; Fig. 4A).

Figure 4B shows the cell growth curve for each clone. The growth of cells from each clone was indistinguishable from that of

the mock transfectant control, demonstrating that overexpression of claudin-4 did not affect the growth of AGS cells. Therefore, induction of claudin-4 by NSAIDs does not seem to be involved in the inhibition of cell growth by NSAIDs.

We also examined the effect of overexpression of claudin-4 on the induction of apoptosis. In the absence of indomethacin, the cell viability of each clone, as determined by the trypan blue exclusion test, was close to 100%, showing that expression of claudin-4 does not affect cell viability. As shown in Fig. 4C, the dose-response curve for the decrease in cell viability by indomethacin was indistinguishable between each of the claudin-4-overexpressing clones and the mock transfectant control. Further, we confirmed that the cell death (Fig. 4C) was mediated by apoptosis as evidenced by apoptotic DNA fragmentation, activation of caspase-3, and chromatin condensation (data not


shown). The results presented in Fig. 4C show that claudin-4 overexpression does not affect the indomethacin-induced cell apoptosis. Therefore, the induction of claudin-4 by NSAIDs does not seem to be involved in NSAID-mediated apoptosis.

The anchorage-independent growth of tumor cells, which can be measured by colony formation in soft agar, is important for tumor progression. NSAIDs are known to inhibit colony formation of some cancer cells in soft agar (13); recently, it was reported that overexpression of claudin-4 in pancreatic cancer cells inhibited colony formation in soft agar (19). In this study, we examined the effect of claudin-4 overexpression and the presence of indomethacin on the anchorage-independent growth of AGS cells. We first examined the colony-forming ability of each of the claudin-4-overexpressing clones in soft agar. All clones showed less activity for colony formation in soft agar than the mock transfectant control (Fig. 4D), which is consistent with previous results obtained using pancreatic cancer cells (19). We compared the extent of inhibition of colony formation in soft agar with the degree of claudin-4 overproduction in these clones and found a close correlation between the two (Fig. 4A and D).


We also examined the effect of indomethacin on colony formation of AGS cells in soft agar. Because a long incubation period (10 days) was required for this assay, relatively low concentrations of indomethacin were used. As shown in Fig. 4E, indomethacin (100  $\mu\text{M}$ ) significantly decreased the colony-forming ability of AGS cells in soft agar. Real-time RT-PCR experiments confirmed that claudin-4 mRNA expression in AGS cells was induced at the concentration of indomethacin used (Fig. 4F). These results suggest that the induction of claudin-4 is involved in the indomethacin-dependent inhibition of AGS cell colony formation in soft agar.

The migration activity of tumor cells is also very important for tumor progression. We examined the relationship between expression of claudin-4 and migration activity in AGS cells. Wound healing assays were carried out in which the cell-free area was measured at the time a wound was made and then 24 hours later. Because neither claudin-4 overexpression nor addition of NSAIDs affected the growth of AGS cells (Fig. 4B; data not shown), a smaller cell-free area is indicative of a higher activity for cell migration. As shown in Fig. 5A, claudin-4-overexpressing cells (clone 7) showed less cell migration activity

F5



**Figure 4.** Effect of overexpression of claudin-4 on cell growth, apoptosis, and colony formation of AGS cells in soft agar. The extent of expression of claudin-4 in each clone (stable transfectant of claudin-4 expression plasmid) was estimated by immunoblotting experiments as described in Fig. 1 (A). Cells of each clone were cultured for indicated periods, and cell numbers were determined by direct cell counting (B). Cells of each clone were cultured in the presence of indicated concentrations of indomethacin for 24 hours and cell viability was determined by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide method (C). Cells of each clone (D) or nontransfected AGS cells (E) were layered over soft agar in the presence (E) or absence (D) of indicated concentrations of indomethacin. After 10 days, cells were stained with crystal violet and colonies were counted (D and E). Expression of claudin-4 mRNA after treatment of cells with indicated concentrations of indomethacin for 24 hours was monitored by real-time RT-PCR (F). Points, mean ( $n = 3$ ); bars, SE (C). Columns, mean ( $n = 3$ ); bars, SE (D and E). \*\*\*,  $P < 0.001$ ; \*\*,  $P < 0.01$  (D and E).



**Figure 5.** Effect of claudin-4 overexpression or of indomethacin on AGS cell migration. AGS cells of stable transfectant of claudin-4 expression plasmid (clone 7 in Fig. 4) and mock transfectant control AGS cells (A) or AGS cells transfected or nontransfected with siRNA for claudin-4 (B) were wounded and cultured for 24 hours in the presence (B) or absence (A) of indicated concentrations of indomethacin. The cell-free area was measured after 24 hours of incubation and expressed as relative to that before incubation. Columns, mean ( $n = 3$ ); bars, SE. \*\*\*,  $P < 0.001$  or ###,  $P < 0.001$ ; \* $P < 0.05$  (A and B). AGS cells transfected or nontransfected with siRNA for claudin-4 were cultured for 24 hours in the presence or absence of 0.3 mmol/L indomethacin for 24 hours. Levels of claudin-4 and actin were estimated by immunoblotting experiments as described in Fig. 1.

than the mock transfectant control. Furthermore, transfection of siRNA for claudin-4 stimulated the migration activity of AGS cells even in the absence of indomethacin (Fig. 5B). We confirmed that the transfection almost completely inhibited the expression of claudin-4 in AGS cells (Fig. 5C). These results suggest that the migration activity of AGS cells decreases as claudin-4 expression increases.

As shown in Fig. 5B, indomethacin inhibited the activity of AGS cells for cell migration and this inhibitory effect was almost completely suppressed by the transfection of siRNA for claudin-4. We confirmed that transfection of siRNA almost completely inhibited the induction of claudin-4 by indomethacin (Fig. 5C). Taken together, these results support the hypothesis that inhibition of cell migration by indomethacin is mediated through the induction of claudin-4.

## Discussion

We have shown here that some tight junction-related genes, especially *claudin-4*, are induced by NSAIDs. Although NSAIDs and tight junctions are closely associated in relation to cancer progression, this is the first time that a connection between NSAIDs and tight junctions has been shown at the molecular level.

It is known that various factors disrupt or stimulate the function of tight junctions. For example, tumor necrosis factor- $\alpha$ , transforming growth factor- $\alpha$ , and interleukin-1 disrupt tight junctions, whereas transforming growth factor- $\beta$ , interleukin-10, and PGE<sub>2</sub> are known to stimulate the function of tight junctions (41). However, the effect of these factors on the expression of components of tight junctions (such as claudin-4) has not been examined to the same extent. It seems that the alteration of tight junction function is not always correlated with an alteration in the expression of tight junction components. For example, we have found that PGE<sub>2</sub>, which is known to stimulate the function of tight junctions, does not induce claudin-4. Because the expression of claudin-4 affects various aspects of cancer progression (see below), we consider that the effect of cancer-promoting agents or anticancer drugs on claudin-4 expression should be examined more extensively.

As for a mechanism of claudin-4 induction by NSAIDs, we postulate that it is mediated by an increase in  $[Ca^{2+}]_i$ , based on the following observations: (a) NSAIDs increased  $[Ca^{2+}]_i$  and induced claudin-4 simultaneously, (b) thapsigargin and ionomycin increased  $[Ca^{2+}]_i$  and induced claudin-4, and (c) the intracellular  $Ca^{2+}$  chelator [1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid] attenuated the indomethacin-dependent induction of claudin-4. As for the mechanism for the increase in  $[Ca^{2+}]_i$  by NSAIDs, both inhibition of sarcoplasmic/endoplasmic reticulum  $Ca^{2+}$  ATPase (endoplasmic reticulum-located  $Ca^{2+}$  pump that is responsible for accumulation of  $Ca^{2+}$  in the endoplasmic reticulum) and stimulation of the influx of extracellular  $Ca^{2+}$  have been proposed (40). We found recently that all of the NSAIDs tested permeabilize the membranes of both erythrocytes and liposomes (42). This activity of NSAIDs was found to be closely related to their ability to increase  $[Ca^{2+}]_i$ , suggesting that NSAIDs permeabilize membranes and stimulate the influx of extracellular  $Ca^{2+}$  (42).

NSAIDs seem to achieve their chemopreventive effect via several mechanisms, such as stimulation of apoptosis, cell growth suppression, inhibition of angiogenesis, and inhibition of metastasis (4, 5). In this study, we examined the contribution of claudin-4 induction to the antitumor activity of NSAIDs *in vitro*. Experiments using claudin-4-overproducing AGS cells and siRNA for claudin-4 suggested that NSAID-induced claudin-4 is involved in the NSAID-dependent suppression of anchorage-independent tumor growth and tumor cell migration but not in stimulation of apoptosis and cell growth suppression. As for cell migration, this is the first evidence showing not only that NSAIDs inhibit of cancer cell migration but also that claudin-4 is involved in cell migration. It was reported recently that overexpression of claudin-4 suppressed the invasive potential of pancreatic cancer cells (19); therefore, if NSAIDs also induce claudin-4 *in vivo*, then suppression of the invasive potential of tumor cells by NSAID-induced claudin-4 may be one of the mechanisms involved in the inhibition of metastasis by NSAIDs. It is also possible that the induction of claudin-4 by NSAIDs contributes to their antitumor activity through other

mechanisms. Tight junctions act as a barrier for diffusion of molecules that include nutrients and growth factors. It is well known that the constitutive accessibility of nutrients and growth factors is very important for tumor progression. Therefore, if NSAIDs also induce claudin-4 *in vivo*, then the supply of nutrients and growth factors to a tumor may be retarded or inhibited, thereby suppressing tumor progression.

## Acknowledgments

Received 8/3/2004; revised 11/22/2004; accepted 12/20/2004.

Grant support: Ministry of Health, Labor, and Welfare of Japan Grants-in-Aid for Scientific Research, Suzuken Memorial Foundation, and Japan Research Foundation for Clinical Pharmacology.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

## References

- Smalley WE, Ray WA, Daugherty JR, Griffin MR. Nonsteroidal anti-inflammatory drugs and the incidence of hospitalizations for peptic ulcer disease in elderly persons. *Am J Epidemiol* 1995;141:539-45.
- Farrow DC, Vaughan TL, Hansten PD, et al. Use of aspirin and other nonsteroidal anti-inflammatory drugs and risk of esophageal and gastric cancer. *Cancer Epidemiol Biomarkers Prev* 1998;7:97-102.
- Sorensen HT, Friis S, Norgaard B, et al. Risk of cancer in a large cohort of nonaspirin NSAID users: a population-based study. *Br J Cancer* 2003;88:1687-92.
- Gupta RA, Dubois RN. Colorectal cancer prevention and treatment by inhibition of cyclooxygenase-2. *Nat Rev Cancer* 2001;1:11-21.
- Kismet K, Akay MT, Abbasoglu O, Ercan A. Celecoxib: a potent cyclooxygenase-2 inhibitor in cancer prevention. *Cancer Detect Prev* 2004;28:127-42.
- Hoshino T, Tsutsumi S, Tomisato W, Hwang HJ, Tsuchiya T, Mizushima T. Prostaglandin E<sub>2</sub> protects gastric mucosal cells from apoptosis via EP2 and EP4 receptor activation. *J Biol Chem* 2003;278:12752-8.
- Tsuji M, Kawano S, Sawaoka H, Hori M, DuBois RN. Cyclooxygenase regulates angiogenesis induced by colon cancer cells. *Cell* 1998;93:705-16.
- Rolland PH, Martin PM, Jacquemier J, Rolland AM, Toga M. Prostaglandin in human breast cancer: evidence suggesting that an elevated prostaglandin production is a marker of high metastatic potential for neoplastic cells. *J Natl Cancer Inst* 1980;64:1061-70.
- Eberhart CE, Coffey RJ, Radhika A, Giardiello FM, Ferrenbach S, DuBois RN. Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology* 1994;107:1183-8.
- Ristimaki A, Honkanen N, Jankala H, Sipponen P, Harkonen M. Expression of cyclooxygenase-2 in human gastric carcinoma. *Cancer Res* 1997;57:1276-80.
- Piazza GA, Alberts DS, Hixson LJ, et al. Sulindac sulfone inhibits azoxymethane-induced colon carcinogenesis in rats without reducing prostaglandin levels. *Cancer Res* 1997;57:2909-15.
- Reddy BS, Kawamori T, Lubet RA, Steele VE, Kelloff GJ, Rao CV. Chemopreventive efficacy of sulindac sulfone against colon cancer depends on time of administration during carcinogenic process. *Cancer Res* 1999;59:3387-91.
- Zhang X, Morham SG, Langenbach R, Young DA. Malignant transformation and antineoplastic actions of nonsteroidal antiinflammatory drugs (NSAIDs) on cyclooxygenase-null embryo fibroblasts. *J Exp Med* 1999;190:451-9.
- Hanif R, Pittas A, Feng Y, et al. Effects of nonsteroidal anti-inflammatory drugs on proliferation and on induction of apoptosis in colon cancer cells by a prostaglandin-independent pathway. *Biochem Pharmacol* 1996;52:237-45.
- Anderson JM, Van Itallie CM. Tight junctions and the molecular basis for regulation of paracellular permeability. *Am J Physiol* 1995;269:G467-5.
- Soler AP, Miller RD, Laughlin KV, Carp NZ, Klurfeld DM, Mullin JM. Increased tight junctional permeability is associated with the development of colon cancer. *Carcinogenesis* 1999;20:1425-31.
- Li D, Mersny RJ. Oncogenic Raf-1 disrupts epithelial tight junctions via downregulation of occludin. *J Cell Biol* 2000;148:791-800.
- Hoebel T, Mucek R, Swisshelm K, Kubbies M. Reexpression of the TJ protein CLDN1 induces apoptosis in breast tumor spheroids. *Int J Cancer* 2004;108:374-83.
- Michl P, Barth C, Buchholz M, et al. Claudin-4 expression decreases invasiveness and metastatic potential of pancreatic cancer. *Cancer Res* 2003;63:6265-71.
- Baek SJ, Kim KS, Nixon JB, Wilson LC, Eling TE. Cyclooxygenase inhibitors regulate the expression of a TGF- $\beta$  superfamily member that has proapoptotic and antitumorigenic activities. *Mol Pharmacol* 2001;59:901-8.
- Tsutsumi S, Gotoh T, Tomisato W, et al. Endoplasmic reticulum stress response is involved in nonsteroidal anti-inflammatory drug-induced apoptosis. *Cell Death Differ* 2004;11:1009-16.
- Tsutsumi S, Tomisato W, Hoshino T, Tsuchiya T, Mizushima T. Transforming growth factor- $\beta$ 1 is responsible for maturation-dependent spontaneous apoptosis of cultured gastric pit cells. *Exp Biol Med (Maywood)* 2002;227:402-11.
- Tsutsumi S, Tomisato W, Takano T, Rokutan K, Tsuchiya T, Mizushima T. Gastric irritant-induced apoptosis in guinea pig gastric mucosal cells in primary culture. *Biochim Biophys Acta* 2002;1589:168-80.
- Kao JB, Harootyan AT, Tsiang RY. Photochemically generated cytosolic calcium pulses and their detection by fluo-3. *J Biol Chem* 1989;264:8179-84.
- Wittchen ES, Huskins J, Stevenson BR. NZO-3 expression causes global changes to actin cytoskeleton in Madin-Darby canine kidney cells: linking a tight junction protein to Rho GTPases. *Mol Biol Cell* 2003;14:1757-68.
- Ciardiello F, Caputo R, Damiano V, et al. Antitumor effects of ZD6474, a small molecule vascular endothelial growth factor receptor tyrosine kinase inhibitor, with additional activity against epidermal growth factor receptor tyrosine kinase. *Clin Cancer Res* 2003;9:1546-56.
- Bottone FG Jr, Martinez JM, Collins JB, Afshari CA, Eling TE. Gene modulation by the cyclooxygenase inhibitor, sulindac sulfide, in human colorectal carcinoma cells: possible link to apoptosis. *J Biol Chem* 2003;278:25790-801.
- Rahner C, Milic LL, Anderson JM. Heterogeneity in expression and subcellular localization of claudins 2, 3, 4, and 5 in the rat liver, pancreas, and gut. *Gastroenterology* 2001;120:411-22.
- Lim JW, Kim H, Kim KH. Nuclear factor- $\kappa$ B regulates cyclooxygenase-2 expression and cell proliferation in human gastric cancer cells. *Lab Invest* 2001;81:349-60.
- Fan XM, Wong BC, Lin MC, et al. Interleukin-1 $\beta$  induces cyclo-oxygenase-2 expression in gastric cancer cells by the p38 and p44/42 mitogen-activated protein kinase signaling pathways. *J Gastroenterol Hepatol* 2001;16:1098-104.
- Tsuji S, Kawano S, Sawaoka H, et al. Evidences for involvement of cyclooxygenase-2 in proliferation of two gastrointestinal cancer cell lines. *Prostaglandins Leukot Essent Fatty Acids* 1996;55:179-83.
- Tsuji M, Kawano S, DuBois RN. Cyclooxygenase-2 expression in human colon cancer cells increases metastatic potential. *Proc Natl Acad Sci U S A* 1997;94:3336-40.
- Merritt G, Allprandis ET, Prada F, Rigas B, Kashfi K. The retinoid fenretinide inhibits proliferation and downregulates cyclooxygenase-2 gene expression in human colon adenocarcinoma cell lines. *Cancer Lett* 2001;164:15-23.
- Kawai S, Nishida S, Kato M, et al. Comparison of cyclooxygenase-1 and -2 inhibitory activities of various nonsteroidal anti-inflammatory drugs using human platelets and synovial cells. *Eur J Pharmacol* 1998;347:87-94.
- Lehmann JM, Lenhard JM, Oliver BB, Ringold GM, Klewer SA. Peroxisome proliferator-activated receptors  $\alpha$  and  $\gamma$  are activated by indomethacin and other nonsteroidal anti-inflammatory drugs. *J Biol Chem* 1997;272:3406-10.
- Huang JT, Welch JS, Ricote M, et al. Interleukin-4-dependent production of PPAR- $\gamma$  ligands in macrophages by 12/15-lipoxygenase. *Nature* 1999;400:378-82.
- Kusuhara H, Komatsu H, Sumichika H, Suguhara K. Reactive oxygen species are involved in the apoptosis induced by nonsteroidal anti-inflammatory drugs in cultured gastric cells. *Eur J Pharmacol* 1999;383:331-7.
- Elder DJ, Halton DE, Playfe LC, Paraskeva C. The MEK/ERK pathway mediates COX-2-selective NSAID-induced apoptosis and induced COX-2 protein expression in colorectal carcinoma cells. *Int J Cancer* 2002;99:323-7.
- Tepperman BL, Soper BD. Effect of extracellular Ca<sup>2+</sup> on indomethacin-induced injury to rabbit dispersed gastric mucosal cells. *Can J Physiol Pharmacol* 1994;72:63-9.
- Wang JL, Lin KL, Chen JS, et al. Effect of celecoxib on Ca<sup>2+</sup> movement and cell proliferation in human osteoblasts. *Biochem Pharmacol* 2004;67:1123-30.
- Sawada N, Murata M, Kikuchi K, et al. Tight junctions and human diseases. *Med Electron Microsc* 2003;36:147-56.
- Tomisato W, Tanaka K, Katsu T, et al. Membrane permeabilization by non-steroidal anti-inflammatory drugs. *Biochem Biophys Res Commun* 2004;323:1032-9.

## Low Direct Cytotoxicity of Nabumetone on Gastric Mucosal Cells

YASUHIRO ARAI<sup>1</sup>, BS, KEN-ICHIRO TANAKA<sup>1</sup>, BS, HIRONORI USHIJIMA<sup>1</sup>, BS,  
WATARU TOMISATO<sup>2</sup>, PhD, SHINJI TSUTSUMI<sup>1</sup>, PhD, MAYUKO ABURAYA<sup>1</sup>, BS,  
TATSUYA HOSHINO<sup>1</sup>, BS, KAZUMI YOKOMIZO<sup>1</sup>, PhD, KEITAROU SUZUKI<sup>1</sup>, PhD,  
TAKASHI KATSU<sup>2</sup>, PhD, TOMOFUSA TSUCHIYA<sup>2</sup>, PhD AND  
TOHRU MIZUSHIMA<sup>1</sup>, PhD

<sup>1</sup>Graduate School of Medical and Pharmaceutical Sciences, Kumamoto University,  
Kumamoto 862-0973, Japan

<sup>2</sup>Faculty of Pharmaceutical Sciences, Okayama University, Okayama 700-8530, Japan

This work was supported by Grants-in-Aid for Scientific Research from the  
Ministry of Health, Labour, and Welfare of Japan, as well as by the Suzuken Memorial  
Foundation, and the Japan Research Foundation for Clinical Pharmacology.

Running head: Cytotoxicity of nabumetone

Address for reprint requests:

Dr. Tohru Mizushima,