



Herbal medicine Sho-saiko-to (TJ-9) increases expression matrix metalloproteinases (MMPs) with reduced expression of tissue inhibitor of metalloproteinases (TIMPs) in rat stellate cell

Isao Sakaida^{a,*}, Koji Hironaka^a, Teruaki Kimura^b, Shuji Terai^a,
Takahiro Yamasaki^a, Kiwamu Okita^a

^a*Department of Gastroenterology and Hepatology, Yamaguchi University, School of Medicine, Minami-Kogushi 1-1-1, Ube, Yamaguchi 755-8505, Japan*

^b*Department of Bioregulatory Function, Yamaguchi University, School of Medicine, Minami-Kogushi 1-1-1, Ube, Yamaguchi 755-8505, Japan*

Received 14 April 2003; accepted 18 September 2003

Abstract

We have reported that Sho-saiko-to (TJ-9) prevents liver fibrosis *in vivo*. To gain further insights into the effect of TJ-9, the matrix metalloproteinases (MMPs)/tissue inhibitors of metalloproteinases (TIMPs) balance was examined. Hepatic stellate cells (HSCs) were isolated from male Wistar rats and cultured with TJ-9 (0–1000 µg/ml) on uncoated plastic dishes for 4 days. To elucidate the effects on the MMPs/TIMPs balance by TJ-9, quantitative analysis of type IV collagen-degrading activity, gelatin zymography and reverse zymography were carried out. Northern blot analysis was performed to determine the expression of MMP-2, 13 and TIMP-1 mRNAs. TJ-9 treatment resulted in dose-dependent upregulation of MMP-2, 13 mRNA and downregulation of TIMP-1 mRNA up to 500 µg/ml. Gelatin zymography, reverse zymography and quantitative analysis of type IV collagen-degrading activity confirmed that TJ-9 increased MMP-2 activity and prevented TIMP-1, 2 activities in a dose-dependent manner. SB203580 diminished the reduction of mRNA as well as the activity of TIMP-1 by TJ-9 and induction of mRNA as well as the activity of MMP-2. These results show that TJ-9 increased MMP-2, 13 activity with reduced TIMP-1, 2 activities on HSCs possibly via P38 pathway.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Herbal medicine; Fibrosis; Signal transduction; Metalloproteinase; Tissue inhibitor of metalloproteinase

* Corresponding author. Tel.: +81-836-22-2241; fax: +81-836-22-2240.

E-mail address: sakaida@po.cc.yamaguchi-u.ac.jp (I. Sakaida).

Introduction

Hepatic stellate cells have now been clearly identified as the primary cellular source involved in the pathogenesis of liver fibrosis (Friedman, 1991). During the development of liver fibrosis, stellate cells undergo activation, a process characterized by increased cell proliferation, morphological transformation into myofibroblast-like cells and synthesis of excessive extracellular matrix components (Friedman, 1991). These cells are also able to synthesize enzymes, called matrix metalloproteinases (MMPs), which can degrade matrix proteins in the extracellular space. The activity of MMPs is carefully regulated by controlling their conversion from pro-enzymes to the catalytic form and by a family of specific inhibitors, called tissue inhibitors of metalloproteinases (TIMPs).

On the other hand, many studies of signal transduction are now underway. The MAPK family of protein kinases includes the extracellular signal-regulated kinases (ERKs) (Boulton et al., 1991) the c-Jun N-terminal kinase/stress-activated protein kinases (JNK/SAPKs) (Derijard et al., 1994; Kyriakis et al., 1994), and p38 (Han et al., 1994). The MAPK signaling pathway is a multistep phosphorylation cascade that transmits signals from the cell's surface to cytosolic and nuclear targets (Cohen, 1997). It has been suggested that not only do MAPKs serve as the signaling pathways production of MMPs (Esparza et al., 1999; Zeigler et al., 1999), but that they also participate in production of TIMPs (Li and Zafarullah, 1998; Eberhardt et al., 2000). However, the role of ERK, SAPK and p38 pathways in regulating production of MMPs and TIMPs by HSCs is not well known at present.

Many agents have been proposed for the prevention and treatment of fibrosis but no specific inhibitor of stellate cell activation has yet been developed.

TJ-9 has been reported, including by us, to prevent fibrosis via the inhibition of HSCs in different animal models of fibrosis due to choline-deficiency (Sakaida et al., 1998a,b), and pig serum (Shimizu et al., 1999), as well as in isolated stellate cells (Kayano et al., 1998). TJ-9, commonly prescribed in Japan as Sho-saiko-to, is the most popular herbal medicine in Japan and has been widely used in the treatment of chronic liver diseases, especially chronic viral hepatitis. TJ-9 consists of an aqueous extract from the roots of scutellaria, glycyrrhiza, bupleurum, and ginseng; the pinella tuber; the jujube fruit; and the ginger rhizome. A number of studies have indicated its cytoprotective effects in experimental liver injuries (Yamamoto et al., 1985; Araki et al., 1988), cancer-preventive effects (Okita et al., 1993, 1994; Tatsuta et al., 1991; Oka et al., 1995), anti-tumor, and apoptotic effects (Matsuzaki et al., 1996; Yano et al., 1994).

From the clinical point of view, treatment for liver fibrosis has been mainly focused on prevention of stellate cell activation instead of acceleration of degradation.

To gain further insights into the effect of TJ-9 on suppression of hepatic fibrosis, we examined changes of the MMPs/TIMPs balance and MAPK pathways of TJ-9 treated HSCs.

Materials and methods

Animals

Male Wistar rats weighting 400 to 450 g (Nippon SLC Co., Ltd., Shizuoka, Japan) were obtained, quarantined for 1 week, and housed in a room under controlled temperature (25 °C), humidity, and

lighting (12 hours light, 12 hours dark). Access to food and tap water was ad libitum. After a 1-week acclimation period on a basal diet (Oriental MF Diet; Oriental Yeast Company, Japan), the rats were used for the experiments.

Isolation and culture of rat hepatic stellate cells

Rat hepatic stellate cells (HSCs) were isolated with Nycodenz as described previously (Kayano et al., 1998).

Yields were $1.0\text{--}1.5 \times 10^7$ cells/rat. Cell viability was always over 95% as determined by the trypan blue exclusion test. Cell purity was more than 90% as assessed by the presence of yellow droplets and desmin immunoreactivity. Then isolated hepatic stellate cells were cultured at a density of 5.0×10^5 cells/ml in monolayer culture on uncoated 60-mm plastic dishes (Iwaki Glass Co., Ltd., Tokyo, Japan). All cultures were incubated at 37 °C in a humidified atmosphere of 5% CO₂ and 95% air. After incubation for 4 h, non-adherent cells were removed with a pipette and the culture medium was replaced with medium containing various concentrations of TJ-9 or medium alone (control). In some experiments, SB203580 or PD98059 (Calbiochem, La Jolla, CA) was added in addition to TJ-9. SB20380 and PD980597 were dissolved in DMSO at the final concentration of 0.1%. SB20238 is known as a highly specific inhibitor of p38 kinase (IC₅₀ = 34nM in vitro, 600 nM in cells) and even 100 μM does not significantly inhibit JNK and p42 MAP kinase, and PD98059 is known as specific inhibitor of ERK kinase (IC₅₀ = 2 μM), according to the manufacturer's instructions.

The medium was changed every 24 h and cell culture was continued up to 4 days (serum was depleted 24 h before the assessment).

The treated HSCs were scraped after being washed with PBS twice and suspended at 4 °C in 0.5 ml of PBS with 0.1% Triton X. An equal amount of total protein (100 μg) was applied for the quantitative analysis of type IV collagen-degrading activity, zymography and reverse zymography.

Preparation of culture medium with Sho-saiko-to (TJ-9)

Sho-saiko-to (TJ-9, powder) was kindly provided by Tsumura Co., (Tokyo, Japan). Water-soluble ingredients of TJ-9 were obtained as described previously (Kayano et al., 1998). The final concentrations of TJ-9 used were 10, 100, 500 and 1000 μg/ml. The PH values of all culture media with or without TJ-9 were adjusted to within the physiological range.

Hybridization probes

The following probes were used in this study. G3PDH (glyceraldehyde-3-phosphate dehydrogenase) (Sakaida et al., 1996, 1998a,b) was purchased from American Type Culture Collection (Rockville, MD, USA). TIMP-1 cDNAs was used as described (Sakaida et al., 1999). MMP-2 cDNA was a generous gift of Dr. Hiroshi Sato (Molecular Virology and Oncology, Cancer Research Institute, Kanazawa University, Ishikawa-Pref., Japan) (Sato et al., 1994). MMP-13 cDNA was the generous gift of Dr Cheryl O. Quinn (Pediatric Research Institute, Department of Pediatrics, Health Sciences Center, Saint Louis University, MO, USA) (Hironaka et al., 2000).

Northern blot analysis

Northern blot analysis was performed after the isolation of 10 µg of total RNA from isolated hepatic stellate cells as described previously (Hironaka et al., 2000). Each signal strength of mRNA was determined after normalization by relevant G3PDH mRNA levels.

Quantitative analysis of type IV collagen-degrading activity

Type IV collagen-degrading activity in conditioned media was determined as described previously (Hironaka et al., 2000) using a Type IV collagenase activity assay kit (YU-18001, Yagai Co., Kamagata, Japan). The principle of the type IV collagenase activity assay kit is based on fluorescent measurement of collagen fragments upon cleavage by gelatinases. It is known that collagen decomposition fragments differ from whole collagen in their temperature of denaturation and ethanol solubility. Upon gelatinase cleavage of fluorescently-labeled collagen type IV, decomposition fragments are produced. These fragments are selectively denatured and extracted with ethanol. The fluorescence intensity of the extracted product is measured and correlates with type IV collagen-degrading activity.

Zymography

As described by Durko et al. (Durko et al., 1997), SDS-polyacrylamide (10%) gels were copolymerized with gelatin at a final concentration of 1 mg/ml. The concentrated conditioned media were mixed 3:1 (v/v) with sample buffer (0.3 M Tris-HCl, pH 6.8, containing 8% SDS, 0.4% bromophenol blue and 40% glycerol), loaded onto the gels without boiling and separated by electrophoresis. The gels were washed for 1 h in a solution of 2.5% Triton X-100 in 40 mM Tris-HCl, pH 7.6, and for 15 min in 10 mM Tris-HCl, pH 8. For the enzymatic reaction to take place, the gels were incubated for 18 h at 37 °C in a solution of 50 mM Tris-HCl, pH 8, containing 10 mM CaCl₂. The gels were stained for 2 h in a 0.5% Coomassie blue R250 solution, then destained in 20% methanol with 10% acetic acid until clear bands (indicating lysis) were apparent on the blue background. Prestained molecular weight markers were resolved on the same gels, separated from other samples after electrophoresis and fixed in 5% acetic acid. To characterize the protease bands, some of the gels were incubated in a buffer (50 mM Tris-HCl, pH 8, with 10 mM CaCl₂) containing 20 mM EDTA (a metalloproteinase inhibitor).

Reverse zymography

Reverse zymography was performed as previously described (Herron et al., 1986). Gelatinase inhibitory activity was detected by incubating standard gelatin zymograms for 1 h at 37 °C in a preparation of purified collagenase (2 unit/ml) before incubation at 37 °C for 18 h in a solution of 50 mM Tris-HCl, pH 8, containing 10 mM CaCl₂. Gels were stained and destained as described above, after which stained bands manifested the presence of gelatinase inhibitory activity corresponding to TIMPs in the polyacrylamide gels.

Statistical analysis

Results are presented as the mean \pm SD. Statistical analysis was performed using Student's t-test and one-way analysis of variance (ANOVA). A p-value of less than 0.05 was considered to be statistically significant.

Ethical considerations

This experiment was reviewed by the Committee of Animal Experiment Ethics in Yamaguchi University School of Medicine and was carried out under the Guidelines for Animal Experiments in Yamaguchi University School of Medicine (No. 105) and Notification (No. 6) of the Japanese Government.

Results

Effects of TJ-9 on MMP-2, 13 and TIMP-1 mRNA expression of activated HSC

Messenger RNA expression was investigated by Northern blot analysis on day 4 after culture. Under control culture conditions (no treatment), HSCs expressed MMP-2, 13 and TIMP-1 mRNAs (Fig. 1A, 1B, lane 1). 500 μ g/ml of TJ-9 significantly elevated MMP-2 transcripts expressed at 3.1 kb and strikingly inhibited TIMP-1 mRNA expression (Fig. 1A, 1B, line 2). Expression of MMP-13 mRNA with stellate cells was dramatically reduced in a time-dependent manner after isolation. Although MMP-13 mRNA expression was not detected after 4-day culture, addition of 500 μ g/ml of TJ-9 increased MMP-13 mRNA expression to a detectable level, though it was still very weak (Fig. 1A, 1B). Reduction of type I procollagen mRNA expression could be reproduced with 500 μ g/ml of TJ-9 as previously reported (Kayano et al., 1998).

Morphologically, TJ-9 seemed to prevent the transformation to myofibroblast-like cells as previously reported (data not shown) (Kayano et al., 1998).

The doses up to 1000 μ g/ml of TJ-9 showed no cytotoxicity with cultured stellate cells.

Effects of TJ-9 on MMP-2 and TIMPs production by activated HSC

When serum-free conditioned media derived from HSCs were analyzed on polyacrylamide gels copolymerized with 1 mg/ml gelatin, the band of lysis was observed in the zymograms. These gelatinolytic activities were completely inhibited when the gels were incubated in the presence of 20mM EDTA (data not shown), indicating that they were mediated by metalloproteinases. TJ-9 conditioned media exhibited gelatinase activities in a dose-dependent manner on gelatin zymograms, seen mainly as a triplet with bands at molecular masses of 70, 66 and 62kDa (Fig. 2A). The 66 kDa gelatinase predominated, and the 62 kDa band was at the limit of detection. These bands migrated in the Mr 62-70 kDa region corresponding to the predicted molecular weights for MMP-2. The appearance of these bands was dependent on the concentration of TJ-9, and gelatinase activity reached a peak at the concentration of 500 or 1000 μ g/ml. Upregulation of MMP-2 production was also confirmed by quantitative measurement using the type IV collagen-degrading assay. Collagen IV




Fig. 1. A. Messenger RNA expression of MMP-13, MMP-2, TIMP-1 and G3PDH. Freshly isolated hepatic stellate cells were cultured for 4 h. The medium was replaced without TJ-9 (lane 1), with 500 μ g/ml TJ-9 (lane 2), with 500 μ g/ml TJ-9 plus 10 μ M SB203580 (lane 3) and with 500 μ g/ml TJ-9 plus 50 μ M PD 98059 (lane 4). The medium was changed every 24 h. After 4 days, mRNA expression was examined. The figure shows a representative example of 5 independent Northern blots. B. Graphic representation of Fig. 1. The results of the densitometric analysis after normalization against hybridization signals for G3PDH are shown mean \pm SD of 5 independent experiments. * $P < 0.01$ vs lane 1 (control), ** $P < 0.01$ vs lane 2 (TJ-9 500 μ g/ml).




Fig. 2. A. Zymography with various concentrations of TJ-9. Freshly isolated hepatic stellate cells were cultured for 4 h. The medium was replaced without TJ-9 (lane 1, 2), or with 10 µg/ml (lanes 3, 4), 100 µg/ml (lanes 5, 6), 500 µg/ml (lanes 7, 8), and 1000 µg/ml (lanes 9, 10) of TJ-9. After 4 days, zymography was performed. B. Quantitative analysis of type IV collagen-degrading activity was done instead of zymography shown in Fig. 2-A. The results are shown as mean \pm SD of 5 independent experiments. * $P < 0.01$ vs 0 (control).

decomposition by serum-free conditioned media derived from HSCs increased dose-dependently and reached a peak at the concentration of 500 µg/ml of TJ-9, consistent with zymograms (Fig. 2B).

TJ-9 markedly reduced the activity of TIMPs from HSC in a dose-dependent manner as shown by a reverse zymogram (Fig. 3). This inhibitory effect on the activity of TIMP-1, 2 reached a peak at the concentration of 500 µg/ml of TJ-9. Thus, the TJ-9-enhanced gelatinolytic activity in culture media could be mainly attributed to the decreased synthesis of TIMPs in addition to the increased MMP-2 production.

Although the changes of MMP-2 or TIMP-1 mRNA expression were not remarkable, the treated cells showed prominent changes of activity, which is more physiologically relevant.




Fig. 3. Reverse zymography with concentrations of TJ-9. Freshly isolated hepatic stellate cells were cultured for 4 h. The medium was replaced without TJ-9 (lane 1), or with 10 µg/ml (lane 2), 100 µg/ml (lane 3), 500 µg/ml (lane 4), and 1000 µg/ml (lane 5) of TJ-9. After 4 days, reverse zymography was performed. The figure shows a representative example of 5 independent experiments. * $P < 0.01$ vs 0 (control).

Effect of inhibitors of MAPKs on expression of MMPs and TIMPs

The functional role of MAP kinases in mediating the down-regulatory effect of TJ-9 on expression of TIMPs and the up-regulatory effect on MMPs expression were examined.




Fig. 4. Effects of SB203580 and PD98059 on MMP-2 and TIMP-1, 2 expression of TJ-9 treated hepatic stellate cells. Freshly isolated hepatic stellate cells were cultured for 4 h. The medium was replaced without TJ-9 (lanes 1, 11), with 500 µg/ml TJ-9 (lanes 2, 12), with 500 µg/ml TJ-9 plus 0.01 µM SB203580 (lanes 3, 13), with 500 µg/ml TJ-9 plus 0.1 µM SB203580 (lanes 4, 14), with 500 µg/ml TJ-9 plus 1 µM SB203580 (lanes 5, 15) with 500 µg/ml TJ-9 plus 10 µM SB203580 (lanes 6, 16), with 500 µg/ml TJ-9 plus 0.05 µM PD98059 (lanes 7, 17), with 500 µg/ml TJ-9 plus 0.5 µM PD98059 (lanes 8, 18), with 500 µg/ml TJ-9 plus 5 µM PD98059 (lanes 9, 19), or with 500 µg/ml TJ-9 plus 50 µM PD98059 (lanes 10, 20). After 4 days, zymography and reverse zymography were performed.

The effect of SB203580 (Cuenda et al., 1995), a specific inhibitor of p38 MAP kinase activation, was examined.

The expression of MMP (MMP-13, MMP-2) mRNAs was significantly reduced (Fig. 1A, 1B [b, c], lane 3). On the other hand, the effect of TJ-9 on the expression of TIMP-1 mRNA was completely diminished by treatment of SB203580 (Fig. 1A, 1B [d], lane 3).

These results were confirmed by the zymogram and reverse zymogram shown in Fig. 4 (lanes 1-6, 11-16). SB203580 reduced the activity of MMP-2 (Fig. 5) and increased the activity of TIMP-1, 2 in a dose-dependent manner in TJ-9 treated HSCs.

To elucidate the role of the ERK pathway in the effects of TJ-9, HSCs were treated with various concentrations of PD98059, a specific inhibitor of MEK activation, which prevents activation of ERK




Fig. 5. Quantitative analysis of type IV collagen-degrading activity was done for the comparison of the zymography shown in Fig. 5. The results are shown as mean \pm SD of 5 independent experiments. * $P < 0.01$ vs 0 (control), ** $P < 0.05$ vs 1 (SB203580 1 μ M).

(Dudley et al., 1995). PD98059 had no noticeable effect on changes in mRNAs level but there was a mild increase in MMP-2 activity measured with zymogram (Fig. 4) and type IV collagen-degrading assay (Fig. 5).

Although SB20380 and PD980597 were dissolved in DMSO at the final concentration of 0.1%, 0.1% DMSO by itself had no effect on the activity of MMP-2 or TIMP-1, 2 (data not shown).

These results showed that the activity of p38 MAP kinase seemed to be essential to prevent TIMP-1, 2 expressions and enhance MMP-2 expression in TJ-9 treated HSCs.

Discussion

We have previously shown that TJ-9 prevents fibrosis by the inhibition of HSCs with reduced type I procollagen mRNA expression in different animal models of fibrosis due to choline-deficiency (Sakaida et al., 1998a,b), and pig serum (Shimizu et al., 1999). Also we have already reported that doses up to 1000 µg/ml of TJ-9 inhibit type I procollagen mRNA and the cell cycle of cultured stellate cell without any cytotoxicity (Kayano et al., 1998). In this study, we examined the effect of TJ-9 on the balance between MMP-2 and TIMP-1,2 with cultured HSCs.

First we showed that 500 µg/ml of TJ-9 markedly reduced TIMP-1 mRNA expression with enhanced mRNA expression of MMP-2. We choose this dose because 500 µg/ml of TJ-9 was most effective for the prevention of type I procollagen mRNA expression, as reported previously (Kayano et al., 1998). Then activity of MMP-2 and TIMPs was examined by zymogram or reverse zymogram. The reverse zymogram clearly indicated that TJ-9 reduced TIMP-1, 2 activities in a dose-dependent manner up to 500 µg/ml, with enhanced MMP-2 activity (Figs. 2, 3). The effect of TJ-9 on MMP-2 activity was also quantitatively examined by a type IV collagen degrading assay showing the peak value at the concentration of 500 µg/ml.

Secondary, the effects of TJ-9, i.e. the inhibition of TIMP-1 expression and induction of MMP-2 expression, were completely abolished by the selective p38 MAP kinase inhibitor SB203580 in a dose-dependent manner, indicating that p38 MAP kinase activity was essential for reduction of TIMP-1 expression and induction of MMP-2 expression.

In contrast, inhibition of the ERK pathway by PD98059, a selective inhibitor of MEK activation, caused only slight enhancement of MMP-2 activity without affecting mRNAs level in HSCs treated with TJ-9 (500 µg/ml).

Our findings are consistent with the recent works reporting that p38 MAP kinase may be closely related with the expression of MMP-2 in human endothelial cell (Lee et al., 2000) or the MMP-9 (Underwood et al., 2000; Simon et al., 1998) or the MMP-13 (Ravanti et al., 1999a,b) expression in different cells other than HSCs. Eberhardt W et al (Eberhardt et al., 2000) reported that IL-1 β induced MMP-9 and TIMP-1 expressions in rat glomerular mesangial cells with increased activity of MPAKs cascades (ERK, SAPK/JNK and p38 MAP kinase) leading to increased activity of NF- κ B and AP1 and Solis-Herruzo JA et al (Solis-Herruzo et al., 1999) also indicated that interleukin-6 induced MMP-13 and TIMP-1 expressions in rat fibroblast through stimulation of AP1. Both reports found that either MMP-9 or MMP-13 expression paralleled with the expression of TIMP-1.

It has been reported (Benyon et al., 1999) that activated hepatic stellate cells increase MMP-2 and TIMP-2 expression in a time-dependent manner on culture dishes until 7 days and that the increase in MMP-2 expression of stellate cells is responsible for degrading the normal basement

membrane, not type I collagen. Thus a question may be raised as to whether the increase in MMP-2 expression with TJ-9 treatment can be pro-fibrogenic in terms of stellate cell activation. However, TJ-9 also induced MMP-13 mRNA expression, which is responsible for degrading interstitial collagens, e.g. type I collagen. It is also known (Iredale et al., 1996) that enhanced TIMP-1,2 expression relative to interstitial collagenase (MMP-1/MMP-13) expression leads to the accumulation of collagens (fibrogenesis) and the reduction of TIMP-1,2 expression relative to interstitial collagenase (MMP-1/MMP-13) expression, even weak expression, can lead to the resolution of liver fibrosis (fibrolysis).

Kato et al. (Kato et al., 1998) previously reported opposite result. The treatment of melanoma cells with TJ-9 (400 µg/ml), a dose similar to that used here, showed the downregulation of MMP-2 and upregulation of TIMP-1. Studies of different cell types, e.g. malignant cells or benign cells, may reconcile these different results but further studies using other cells are necessary.

Current studies reported that an IL-10 associated silencer element (HTS) (5'-CCACTGGCC-CATCGTATAT-3') (-1284 to -1266 bp) in the 5' promoter region of the TIMP-1 gene functions as a negative regulatory element that controls TIMP-1 expression (Wang et al., 1998) and upstream TIMP-1 element-1 (UTE-1) is essential for transcriptional activity of the human TIMP-1 promoter (Trim et al., 2000).

On the other hand, the promoter for the MMP-2 gene contains an AP-2 site and GC boxes (Benbow and Brinckerhoff, 1997). Chen et al. reported that MAP kinase pathways utilize a GC box to regulate gene transcription in rat HSC (Chen and Davis, 1999). It has also been reported that the MAP kinase signaling cascade stimulates Sp1 binding (Merchant et al., 1999).

But further examinations are necessary to clarify the exact relationships and cross-talk of MAPKs cascades in the expression of MMPs and TIMPs.

In conclusion, the results of this study show that TJ-9 treatment leading to upregulation of MMP-2, 13 production and downregulation of TIMP-1,2 production. The manipulation of the balance of TIMPs and MMPs may provide new insights for the treatment of liver fibrosis.

Acknowledgements

This work was supported in part by Grants-in-Aid 11670507 and 13470121 from the Ministry of Education, Science and Culture, Japan.

References

- Araki, N., Noda, T., Ogawa, K., 1988. Cytochemical studies on the effect of intraperitoneal and oral administration of a traditional Chinese medicine (Sho-saiko-to) on the D-galactosamine-induced hepatic injuries of rats. *Acta Histochemia et Cytochemica* 21, 439–453.
- Benyon, R.C., Hovell, C.J., Da Gaca, M., Jones, E.H., Iredale, J.P., Arthur, M.J., 1999. Progelatinase A is produced and activated by rat hepatic stellate cells and promotes their proliferation. *Hepatology* 30, 977–986.
- Boulton, T.G., Nye, S.H., Robins, D.J., Radziejewska, E., Morgenbesser, S.D., DePinho, R.A., Panayotatos, N., 1991. ERKs: a family of protein-serine/threonine kinases that are activated and tyrosine phosphorylated in response to insulin and NGF. *Cell* 65, 663–675.
- Cohen, P., 1997. The search for physiological substrates of MAP and SAP kinases in mammalian cells. *Trends Cell Biology* 7, 353–361.

- Derijard, B., Hibi, M., Wu, I.H., Barret, T., Su, B., Deng, T., Karin, M., 1994. JNK: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* 76, 1025–1037.
- Eberhardt, W., Huwiler, A., Beck, K.F., Walpen, S., Pfeilschifter, J., 2000. Amplification of IL-1beta-induced matrix metalloproteinase-9 expression by superoxide in rat glomerular mesangial cells is mediated by increased activities of NF-kappaB and activating protein-1 and involves activation of the mitogen-activated protein kinase pathways. *Journal of Immunology* 165, 5788–5797.
- Benbow, U., Brinckerhoff, C.E., 1997. The AP-1 site and MMP gene regulation: what is all the fuss about?. *Matrix Biology* 15, 519–526.
- Chen, A., Davis, B.H., 1999. UV irradiation activates JNK and increases alpha1(I) collagen gene expression in rat hepatic stellate cells. *Journal of Biological Chemistry* 274, 158–164.
- Cuenda, A., Rouse, J., Doza, Y.N., Meier, R., Cohen, P., Gallagher, T.F., Young, P.R., 1995. SB 203580 is a specific inhibitor of a MAP kinase homologue which is stimulated by cellular stresses and interleukin-1. *FEBS Letter* 364, 229–233.
- Dudley, D.T., Pang, L., Decker, S.T., Bridges, A.J., Saltiel, A.R., 1995. A synthetic inhibitor of the mitogen-activated protein kinase cascade. *Proceedings of the National Academy of Science U.S.A.* 92, 7686–7689.
- Durko, M., Navab, R., Shibata, H.R., Brodt, P., 1997. Suppression of basement membrane type IV collagen degradation and cell invasion in human melanoma cells expressing an antisense RNA for MMP-1. *Biochimica et Biophysica Acta* 1356, 271–280.
- Esparza, J., Vilardell, C., Calvo, J., Juan, M., Vives, J., Urbano-Marquez, A., Yague, J., 1999. Fibronectin upregulates gelatinase B (MMP-9) and induces coordinated expression of gelatinase A (MMP-2) and its activator MT1-MMP (MMP-14) by human T lymphocyte cell lines. A process repressed through RAS/MAP kinase signaling pathways. *Blood* 94, 2754–2766.
- Friedman, S.L., 1991. The cellular basis of hepatic fibrosis. Mechanisms and treatment strategies. *New England Journal of Medicine* 328, 1828–1835.
- Han, J., Lee, J.D., Bibbs, L., Ulevitch, R.J., 1994. A MAP kinase targeted by endotoxin and hyperosmolarity in mammalian cells. *Science* 265, 808–811.
- Herron, G.S., Banda, M.J., Clark, E.J., Gavrilovic, J., Werb, Z., 1986. Secretion of metalloproteinases by stimulated capillary endothelial cells. *Journal of Biological Chemistry* 261, 2814–2818.
- Hironaka, K., Sakaida, I., Matsumura, Y., Kaino, S., Miyamoto, K., Okita, K., 2000. Enhanced interstitial collagenase (matrix metalloproteinase-13) production of Kupffer cells by gadolinium chloride prevents pig serum-induced rat liver fibrosis. *Biochemical and Biophysical Research Communications* 267, 290–295.
- Iredale, J.P., Benyon, R.C., Arthur, M.J., Ferris, W.F., Alcolado, R., Winwood, P.J., Clark, N., 1996. Tissue inhibitor of metalloproteinase-1 messenger RNA expression is enhanced relative to interstitial collagenase messenger RNA in experimental liver injury and fibrosis. *Hepatology* 24, 176–184.
- Kayano, K., Sakaida, I., Uchida, K., Okita, K., 1998. Inhibitory effects of the herbal medicine Sho-saiko-to (TJ-9) on cell proliferation and procollagen gene expressions in cultured rat hepatic stellate cells. *Journal of Hepatology* 29, 642–649.
- Kato, M., Liu, W., Yi, H., Asai, N., Hayakawa, A., Kozaki, K., Takahashi, M., Nakashima, I., 1998. The herbal medicine Sho-saiko-to inhibits growth and metastasis of malignant melanoma primarily developed in ret-transgenic mice. *Journal of Investigative Dermatology* 111, 640–644.
- Kyriakis, J.M., Banerjee, P., Nikolakaki, E., Dai, T., Rubie, E.A., Ahmed, M.F., Avruch, J., 1994. The stress-activated protein kinase subfamily of c-Jun kinases. *Nature* 369, 156–160.
- Lee, O.H., Bae, S.K., Bae, M.H., Lee, Y.M., Moon, E.J., Cha, H.J., Kwon, Y.G., 2000. Identification of angiogenic properties of insulin-like growth factor II in in vitro angiogenesis models. *British Journal of Cancer* 82, 385–391.
- Li, W.Q., Zafarullah, M., 1998. Oncostatin M up-regulates tissue inhibitor of metalloproteinases-3 gene expression in articular chondrocytes via de novo transcription, protein synthesis, and tyrosine kinase- and mitogen-activated protein kinase-dependent mechanisms. *Journal of Immunology* 161, 5000–5007.
- Matsuzaki, Y., Kurokawa, F., Terai, S., Matsumura, Y., Kobayashi, N., Okita, K., 1996. Cell death induced by baicalein in human hepatocellular carcinoma cell lines. *Japanese Journal of Cancer Research* 87, 170–177.
- Merchant, J.L., Du, M., Todisco, A., 1999. Sp1 phosphorylation by Erk 2 stimulates DNA binding. *Biochemical and Biophysical Research Communications* 254, 454–461.
- Oka, H., Yamamoto, S., Kanno, T., Kuroki, T., Mizoguchi, Y., Kobayashi, K., 1995. Controlled prospective evaluation of Sho-saiko-to in prevention of hepatocellular carcinoma in patients with cirrhosis of the liver. *Cancer* 76, 743–749.

- Okita, K., Li, Q., Murakami, T., Takahashi, M., 1993. Anti-growth effects with components of Sho-saiko-to (TJ-9) on cultured human hepatoma cells. *European Journal of Cancer Prevention* 2, 169–176.
- Okita, K., Kurokawa, F., Yamasaki, T., Furukawa, T., Li, Q., Murakami, T., 1994. The use of Sho-saiko-to (TJ-9) for chemoprevention of chemical hepatocarcinogenesis in rats and discussion of its possible pharmacologic action. *Transgenica* 1, 39–44.
- Ravanti, L., Hakkinen, L., Larjava, H., Saarialho-Kere, U., Foschi, M., Han, J., Kahari, V.M., 1999a. Transforming growth factor-beta induces collagenase-3 expression by human gingival fibroblasts via p38 mitogen-activated protein kinase. *Journal of Biological Chemistry* 274, 37292–37300.
- Ravanti, L., Heino, J., Lopez-Otin, C., Kahari, V.M., 1999b. Induction of collagenase-3 (MMP-13) expression in human skin fibroblasts by three-dimensional collagen is mediated by p38 mitogen-activated protein kinase. *Journal of Biological Chemistry* 274, 2446–2455.
- Sato, H., Takino, T., Okada, Y., Cao, J., Shinagawa, A., Yamamoto, E., Seiki, M., 1994. A matrix metalloproteinase expressed on the surface of invasive tumour cells. *Nature* 370, 61–65.
- Sakaida, I., Matsumura, Y., Kubota, M., Kayano, K., Takenaka, K., Okita, K., 1996. The prolyl 4-hydroxylase inhibitor HOE 077 prevents activation of Ito cells, reducing procollagen gene expression in rat liver fibrosis induced by choline deficient L-amino acid-defined diet. *Hepatology* 23, 755–763.
- Sakaida, I., Matsumura, Y., Akiyama, S., Hayashi, K., Ishige, A., Okita, K., 1998a. Herbal medicine Sho-saiko-to (TJ-9) prevents liver fibrosis and enzyme-altered lesions in rat liver cirrhosis induced by a choline-deficient L-amino acid-defined diet. *Journal of Hepatology* 28, 298–306.
- Sakaida, I., Hironaka, K., Uchida, K., Kayano, K., Okita, K., 1998b. Fibrosis accelerates the development of enzyme-altered lesions in the rat liver. *Hepatology* 28, 1247–1252.
- Sakaida, I., Uchida, K., Hironaka, K., Okita, K., 1999. Prolyl 4-hydroxylase inhibitor (HOE 077) prevents TIMP-1 gene expression in rat liver fibrosis. *Journal of Gastroenterology* 34, 376–377.
- Shimizu, I., Ma, Y.R., Mizobuchi, Y., Liu, F., Miura, T., Nakai, Y., Yasuda, M., Amagaya, S., Kawada, N., Hori, H., Ito, S., 1999. Effects of Sho-saiko-to, a Japanese herbal medicine, on hepatic fibrosis in rats. *Hepatology* 29, 149–160.
- Simon, C., Goepfert, H., Boyd, D., 1998. Inhibition of the p38 mitogen-activated protein kinase by SB 203580 blocks PMA-induced Mr 92,000 type IV collagenase secretion and in vitro invasion. *Cancer Research* 58, 1135–1139.
- Solis-Herruzo, J.A., Rippe, R.A., Schrum, L.W., de La Torre, P., Garcia, I., Jeffrey, J.J., Munoz-Yague, T., 1999. Interleukin-6 increases rat metalloproteinase-13 gene expression through stimulation of activator protein 1 transcription factor in cultured fibroblasts. *Journal of Biological Chemistry* 274, 30919–30926.
- Tatsuta, M., Ishii, H., Baba, M., Nakaizumi, A., Uehara, H., 1991. Inhibition by xiao-chai-hu-tang (TJ-9) of development of hepatic foci induced by N-nitrosomorpholine in Sprague-Dawley rats. *Japanese Journal of Cancer Research* 82, 987–992.
- Trim, J.E., Samra, S.K., Arthur, M.J., Wright, M.C., McAulay, M., Beri, R., Mann, D.A., 2000. Upstream tissue inhibitor of metalloproteinases-1 (TIMP-1) element-1, a novel and essential regulatory DNA motif in the human TIMP-1 gene promoter, directly interacts with a 30-kDa nuclear protein. *Journal of Biological Chemistry* 275, 6657–6663.
- Underwood, D.C., Osborn, R.R., Bochnowicz, S., Webb, E.F., Rieman, D.J., Lee, J.C., Romanic, A.M., 2000. SB 239063, a p38 MAPK inhibitor, reduces neutrophilia, inflammatory cytokines, MMP-9, and fibrosis in lung. *American Journal of Physiology-Lung Cellular and Molecular Physiology* 279, L895–L902.
- Wang, M., Hu, Y., Shima, I., Stearns, M.E., 1998. Identification of positive and negative regulator elements for the tissue inhibitor of metalloproteinase 1 gene. *Oncology Research* 10, 219–233.
- Yamamoto, K., Araki, N., Ogawa, K., 1985. Ultrastructural and ultracytochemical examination of the effects of preadministration of Xia-Chai-Hu-Tang on hepatic disorders induced by D-galactosamine HCl. *Acta Histochemia et Cytochemica* 18, 403–418.
- Yano, H., Mizoguchi, A., Fukuda, K., Haramaki, M., Ogasawara, S., Momosaki, S., Kojiro, M., 1994. The herbal medicine Sho-saiko-to inhibits proliferation of cancer cell lines by inducing apoptosis and arrest at the G0/G1 phase. *Cancer Research* 54, 448–454.
- Zeigler, M.E., Chi, Y., Schmidt, T., Varani, J., 1999. Role of ERK and JNK pathways in regulating cell motility and matrix metalloproteinase 9 production in growth factor-stimulated human epidermal keratinocytes. *Journal of Cellular Physiology* 180, 271–284.

Proteomic profiling of proteins decreased in hepatocellular carcinoma from patients infected with hepatitis C virus

Yuichiro Yokoyama^{1,2}, Yasuhiro Kuramitsu², Motonari Takashima³, Norio Iizuka⁴, Toshifusa Toda⁵, Shuji Terai¹, Isao Sakaida¹, Masaaki Oka³, Kazuyuki Nakamura² and Kiwamu Okita¹

¹Department of Gastroenterology and Hepatology

²Department of Biochemistry and Biomolecular Recognition

³Department of Surgery II

⁴Department of Bioregulatory Function,
Yamaguchi University School of Medicine, Yamaguchi, Japan

⁵Tokyo Metropolitan Institute of Gerontology, Tokyo, Japan

Hepatocellular carcinoma (HCC) is a major cause of death in Japan. It has been suggested that hepatitis C virus (HCV) plays an important role in hepatocarcinogenesis, because of high incidence among the patients. To understand the mechanism of hepatocarcinogenesis after HCV infection, we performed a comparative study on the protein profiles between tumorous and nontumorous specimens from the patients infected with HCV by means of two-dimensional electrophoresis. Eleven spots were decreased in HCC tissues from over 50% of the patients. Eight proteins out of 11 spots were identified using peptide mass fingerprinting with matrix-assisted laser desorption/ionization-time of flight-mass spectrometry. These proteins were liver type aldolase, tropomyosin β -chain, ketohexokinase, enoyl-CoA hydratase, albumin, smoothelin, ferritin light chain, and arginase 1. The intensity of enoyl-CoA hydratase, tropomyosin β -chain, ketohexokinase, liver type aldolase, and arginase 1 was significantly different ($p < 0.05$). The decrease of 8 proteins was characteristic in HCC. We will discuss the implication of these proteins for the loss of function of hepatocytes and for the possibility of carcinogenesis of HCV-related HCC.

Keywords: Hepatitis C virus / Hepatocellular carcinoma / Liver proteome / Two-dimensional gel electrophoresis

Received	24/9/03
Revised	25/11/03
Accepted	09/12/03

1 Introduction

Hepatocellular carcinoma (HCC) can be caused by chronic infection of hepatitis B or C virus. Particularly, in Japan most patients of HCC are hepatitis C virus (HCV)-positive. Accordingly, HCV may play an important role in hepatocarcinogenesis. That is, HCV carriers proceed to chronic hepatitis, to liver cirrhosis, and to HCC after incubation periods of 15, 25, and 30 years in average, respec-

tively [1, 2]. HCC-specific markers, if present, are reliable for its early detection and for better understanding hepatocarcinogenesis. From this aspect, search for HCC-specific transcripts has been performed by means of cDNA microarray [3–10]. Iizuka *et al.* [5] reported that many genes for detoxification and immune response were up-regulated in HCV-related HCC. However, expression of genes and proteins are not always uniform. Therefore, recently many studies of proteome were reported for understanding of mechanism of diseases by two-dimensional gel electrophoresis (2-DE). HCC was no exception [11–14]. Tannapfel *et al.* [15] showed that 5 proteins (insulin growth factor II, a disintegrin and metalloproteases, signal transducers and activators of transcription 3, suppressors of cytokine signaling 3, and cyclin D1) was significantly upregulated and 4 proteins (collagen 1, SMAD 4, fragile histidine triad, and suppressors of cytokine signaling 1) were downregulated in HCC by protein

Correspondence: Prof. Kazuyuki Nakamura, Department of Biochemistry and Biomolecular recognition, Yamaguchi University School of Medicine, 1-1-1 Minamikogushi, Ube, Yamaguchi, 755-8505, Japan

E-mail: nakamura@yamaguchi-u.ac.jp

Fax: +81-836-22-2212

Abbreviations: HBs-Ag, hepatitis B surface antigen; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; HCV-Ab, anti-hepatitis C virus antibody

microarrays. We tried to analyze proteomics using HCV-positive HCC specimens in comparison with the surroundings. We already reported the expression of heat shock protein 70 family members increased in HCV-related HCC tissues [16].

2 Materials and methods

2.1 Tissue specimens

We examined tumorous and paired nontumorous liver specimens from 20 patients who had undergone partial hepatectomy for HCC at the Yamaguchi University Hospital between 1998 and 2000. All were positive for anti-HCV antibody (HCV-Ab) and negative for hepatitis B surface antigen (HBs-Ag). Four patient's HCCs were well differentiated, 15 were moderately differentiated, and the last one was poorly differentiated. Twelve out of the 20 nontumorous tissues indicated cirrhotic liver.

2.2 Sample preparation

Resected liver tissues which were immediately frozen in liquid nitrogen and stored at -80°C were disrupted in lysis buffer (1% NP-40, 1 mM sodium vanadate, 1 mM PMSF, 50 mM Tris, 10 mM NaF, 10 mM EDTA, 165 mM NaCl, 10 $\mu\text{g}/\text{mL}$ leupeptin, 10 $\mu\text{g}/\text{mL}$ aprotinin) using a Potter type homogenizer with a Teflon tip at 4°C for 1 h. The lysate was separated by centrifugation at $15\,000 \times g$ for 30 min to yield supernatant that was stored at -80°C .

2.3 Two-dimensional gel electrophoresis (2-DE)

The supernatant from liver tissues (300 μg) was applied to Immobiline dry strips (pH 3–10, 7 cm; Amersham Pharmacia Biotechnology, Uppsala, Sweden) in a total volume of 125 μL containing 8 M urea, 2% CHAPS and 0.5% IPG buffer (Amersham Pharmacia Biotechnology) and 2.8 mg/mL dithiothreitol (DTT). After rehydration for 14 h, proteins were separated by isoelectrofocusing (IEF) at 20°C and 50 $\mu\text{A}/\text{strip}$ with the following linear voltage increases: 500 V for 1 h, 1000 V for 1 h, and 8000 V for 2 h. The strips were then equilibrated twice in 50 mM Tris containing 6 M urea, 30% glycerol and 2% sodium dodecyl sulfate (SDS) for 10 min. Dithiothreitol was then added, followed by iodoacetamide. The second dimension proceeded on 12.5% nongradient SDS-polyacrylamide gels (24.5 cm \times 11 cm; SDS-PAGE) at two steps: 600 V, 20 mA for 30 min and 600 V, 50 mA for 70 min in a Multiphor horizontal electrophoresis unit (Amersham Pharmacia Biotechnology). Separated protein spots were fixed and stained on the gel with 30% methanol, 10% acetic acid,

and 0.1% Coomassie Brilliant Blue R-250 (CBB) overnight. The gel was destained with 30% methanol and 10% acetic acid for 30 min, and then with 7% acetic acid until the stain of background was clear and colorless.

2.4 Image analysis

The positions of the protein spots on tumorous and nontumorous tissues were recorded using an Agfa ARCUS 1200™ image scanner (Agfa-Gevaert N.V., Mortsel, Belgium) and analyzed with PDQUEST computer software Ver. 7.1 (Bio-Rad Laboratories, Hercules, CA, USA). Spots stained at different intensities were excised from the gels and identified by peptide mass fingerprinting with matrix-assisted laser desorption/ionization-time of flight-mass spectrometry (MALDI-TOF-MS).

2.5 In-gel digestion

The CBB dye was removed by rinsing twice in 60% methanol containing 50 mM ammonium bicarbonate and 5 mM DTT for each 15 min, and twice in 50% acetonitrile containing 50 mM ammonium bicarbonate and 5 mM DTT for each 7 min. The gel piece was dehydrated in 100% acetonitrile, and then reswollen with an in-gel digestion reagent containing 10 $\mu\text{g}/\text{mL}$ sequencing-grade trypsin (Promega, Madison, WI, USA) in 30% acetonitrile of 50 mM ammonium bicarbonate and 5 mM DTT. The in-gel digestion was performed overnight at 30°C .

2.6 Peptide mass fingerprinting

After in-gel digestion, 1 μL of the reaction mixture was removed and mixed with 1 μL of matrix solution (10 mg/mL α -cyano-4-hydroxycinnamic acid in 50% acetonitrile, 40% methanol, 0.1% trifluoroacetic acid) on a MALDI target plate. MALDI-TOF-MS for peptide mass fingerprinting was performed on AXIMA-CFR mass spectrometer (Shimadzu Biotech, Kyoto, Japan) in reflectron mode. The MS-Fit database search engine in the Protein Prospector web site (<http://prospector.ucsf.edu/>) was used for protein identification.

2.7 Immunoblot analysis

Samples of 7 μg were separated by SDS-PAGE at 15 mA. After SDS-PAGE, the fractionated proteins were transferred electrophoretically to a PVDF membrane (Immobilon; Millipore, Bedford, MA, USA) and blocked overnight at 4°C with TBS containing 5% skim milk. Primary antibodies used were anti-aldolase B polyclonal antibody (1:100) and anti-arginase 1 polyclonal antibody (1:100)

(all from Santa Cruz Biotechnology, Santa Cruz, CA, USA). For each, membranes were incubated for 1 h, washed four times with TBS containing 0.05% Tween 20, incubated for 1 h with horseradish peroxidase-conjugated secondary antibody (1:2000) (ICN Pharmaceuticals, Aurora, OH, USA), and developed with a chemiluminescence reagent (ECL; Amersham Pharmacia Biotechnology).

3 Results

We compared the differential expression of proteins by 2-DE of paired tumorous and nontumorous liver tissues from 20 patients with HCC. Staining with CBB R-250 detected about 480 protein spots (Fig. 1). Eleven of these were decreased in tumorous tissues from over 50% of the patients. Peptide mass fingerprinting with MALDI-TOF-MS identified ketohexokinase, arginase 1, liver type aldolase, enoyl-CoA hydratase, tropomyosin β -chain, albumin, smoothelin, and ferritin light chain (Fig. 2). Information about the 11 spots and the 8 identified proteins are summarized in Table 1. The expression of arginase 1 was decreased in 95% of HCC patients. Liver-type aldolase was decreased in 85%, tropomyosin β -chain in 80%, ketohexokinase, enoyl-CoA hydratase and albumin in 70%, smoothelin in 65%, and ferritin light chain in tumorous tissues from 60% patients. We performed pairwise *t*-tests between the means of spot intensity of tumorous tissues and nontumorous tissues. The intensity of enoyl-CoA hydratase, tropomyosin β -chain, ketohexokinase, liver type aldolase, and arginase 1 significantly differed between tumorous and nontumorous tissues (Fig. 3). The




Figure 1. Two-dimensional gel electrophoresis map of human liver tissue. (a) Tumorous tissue, (b) nontumorous tissue of a HCC patient infected with HCV. Proteins (300 μ g) were separated on 12.5% nongradient polyacrylamide gels and stained with CBB. (c) Enlarged 2-DE map from (b); protein spots are numbered from 1 to 11.

Table 1. Protein spots that decreased in HCC tissues

Spot No. ^{a)}	Accession No.	Protein	Number of patients ^{b)} (%)	Spot intensity (average \pm SD)		Ratio of spot intensity ^{c)} (% average \pm SD)
				Tumorous tissue	Nontumorous tissue	
1	P53814	Smoothelin	13 (65%)	2 466.2 \pm 3 333.6	2 872.8 \pm 1 688.4	108.57 \pm 115.09
2	P02792	Ferritin light chain	12 (60%)	2 601.1 \pm 2 590.2	2 936.4 \pm 2 408.1	286.52 \pm 521.40
3	P02768	Albumin	14 (70%)	25 635.1 \pm 77 312.6	12 585.5 \pm 11 585.3	189.58 \pm 401.46
4	P30084	Enoyl-CoA hydratase	14 (70%)	13 110.9 \pm 15 939.1	27 113.4 \pm 19 645.5	86.66 \pm 112.12
5	P07951	Tropomyosin β -chain	16 (80%)	1 780.7 \pm 1 423.4	4 984.9 \pm 5 349.2	59.20 \pm 50.69
6	P50053	Ketohexokinase	14 (70%)	1 842.2 \pm 1 534.1	3 407.6 \pm 1 951.5	107.36 \pm 181.62
7	P05089	Arginase 1	19 (95%)	2 394.0 \pm 2 331.7	7 460.4 \pm 5 742.2	99.36 \pm 299.39
8	P05062	Liver-type aldolase	17 (85%)	2 321.1 \pm 1 766.5	5 822.9 \pm 6 591.3	104.30 \pm 195.60
9	P05062	Liver-type aldolase	17 (85%)	4 705.5 \pm 5 199.0	20 619.2 \pm 26 706.9	48.73 \pm 67.16
10	P05062	Liver-type aldolase	18 (90%)	6 122.1 \pm 3 997.5	60 282.2 \pm 161 480.1	57.40 \pm 107.92
11	P05062	Liver-type aldolase	17 (85%)	22 109.5 \pm 19 933.8	113 699.8 \pm 164 866.2	61.97 \pm 105.57

a) Spot numbers correspond to that in Fig. 1c.

b) Number of patients in which protein expression decreased in tumorous tissues

c) Percentage of spot intensity of tumorous to nontumorous tissues calculated for each sample pair




Figure 2. Downregulated expression of 11 spots and 8 proteins in HCC. Spots of all 11 proteins that decreased in tumorous tissues include smoothelin, ferritin light chain, tropomyosin β -chain, albumin, ketohexokinase, enoyl-CoA hydratase, liver type aldolase, and arginase 1.




Figure 4. Immunoblot analysis of liver-type aldolase (aldolase B) and arginase 1. Expression of the two proteins was confirmed and the intensity of each band was less in tumorous tissues (T) than in nontumorous tissues (Non-T). Three samples (HCV42, HCV49, and HCV182) from 20 were shown. The molecular weight markers are indicated by arrows.

expression of two proteins among five, liver type aldolase and arginase 1, whose antibodies were available, was confirmed by immunoblot analysis. The intensity of each band was less in tumorous tissues than in nontumorous tissues (Fig. 4).

4 Discussion

There were some previous proteomic studies for HCC, and the etiologies of their HCCs were abundant. Many of them were hepatitis B virus (HBV)-related HCCs [11–14]. However, all of the patients of our study were HCV-Ab-positive and HBs-Ag-negative. Since the hepatocarcinogenic mechanism of HCV may be different from HBV, we need to construct several databases of HBV- and HCV-related HCCs. We found that the decrease of keto-




Figure 3. Comparison of spot intensity between tumorous (T) and nontumorous tissues (Non-T). Significantly decreased proteins in tumorous tissues ($p < 0.05$): (a) enoyl-CoA hydratase, (b) ketohexokinase, (c) arginase 1, (d) tropomyosin β -chain, (e) liver type aldolase. Only No. 9 among 4 aldolase spots is shown; * $p < 0.05$, ** $p < 0.01$.

hexokinase, ferritin light chain, and enzymes involved in the glycolysis and mitochondrial β -oxidation pathways concurs with other human HCC proteomic studies. We discovered also the decrease of smoothelin and tropomyosin β -chain. The proteins which we detected in this study are classified as metabolic enzymes and cytoskeletal proteins. Smoothelin and tropomyosin β -chain belong to the latter. Smoothelin is a cytoskeletal protein existing in visceral and vascular smooth muscle cells [17] as 59 kDa smoothelin-A and 110 kDa smoothelin-B [18, 19]. Rensen *et al.* [20] could not detect smoothelin in normal liver by Western blotting. However, we could detect by 2-DE smoothelin on both gels of tumorous and nontumorous tissues from patients infected with HCV. The use of samples infected with HCV may cause this discrepancy. Hepatic stellate cells (HSCs) have fatty droplets including vitamin A and regulate the sinusoidal microcirculation in normal liver [21, 22]. Once liver tissues are injured, HSCs are activated and change their structure and function. Activated HSCs transform themselves into the myofibroblast and lead to overexpression of α -smooth muscle actin [23, 24]. These data indicated that the source of smoothelin might be not hepatocytes but intermediate cells. Although smoothelin may not be specific to cancer cells, interaction between cancer cells and intermediate cells is important. In accordance with progression of hepatocarcinogenesis, cancer tissues destroy hepatic cords. Therefore, smoothelin may become a marker of HCC progression.

Tropomyosin binds to actin filaments in muscle as well as nonmuscle cells, and regulates muscle contraction in conjunction with troponin. Tropomyosin is a heterodimer of α - and β -chains. In nonmuscle cells such as hepatocytes, tropomyosin plays a role of stabilization of actin filaments [25, 26]. It was reported that tropomyosin suppressed the transformation of phenotypes in nonmuscle cells [27–29]. In previous proteomic studies using 2-DE, the expression of tropomyosin-2 decreased in human ovarian cancer tissues [30] and the expression of tropomyosin β -chain decreased in human prostate cancer tissues [31]. In addition, the expression of tropomyosin decreased in several cancer tissues [30–33].

In metabolic enzymes, the expression of liver-type aldolase considerably decreased in our tumorous tissues, and this decrease has never been reported in previous proteomic studies on human HCCs. Aldolase performs the sixth step of glycolysis. Aldolase has three isozymes. Aldolase A exists in muscle, aldolase B in liver, and aldolase C in brain [34]. Weinhouse *et al.* [35] have shown that liver-type aldolase was replaced by nonliver type aldolase in poorly differentiated rat liver tumors. Kinoshita *et al.* [36] have reported that the mRNA of aldolase B was

downexpressed in 18 of 20 HCC patients, and Kimura *et al.* [37] found that the expression level of aldolase B was low and aldolase A was high in hepatoma cell lines. Salvatore *et al.* [38] have shown that the mRNA expression of aldolase A in HCC tissues was higher than that in surrounding cirrhotic tissues. Aldolase A is exclusively expressed in the fetal liver [39]. As the expression of carcinoembryonic antigen (CEA) or α -fetoprotein (AFP) which are expressed during embryonic or fetal terms increase in cancer tissues, the expression of aldolase A might increase in HCC tissues by replacing aldolase B. We found the spots of albumin and smoothelin at different positions from those predicted by the *pI* value and molecular weight on both gels of tumorous and nontumorous tissues. These results might be caused by post-translational modifications or mutation by HCV infection. These changes could be characteristic of HCV-related HCCs.

In this study, we performed the first proteomic analysis of human HCV-related HCCs. A protein expression database constructed by reference to our study may be useful for the discovery of new tumor markers and for producing new diagnostic techniques.

This study was supported a Grant-in-Aid for Scientific Reserch from the Japan Society for the Promotion of Science (No. 13470121 to Shuji Terai, Isao Sakaida, and Kiwamu Okita; No. 13770262 to Shuji Terai) and a Grant-in-Aid for translational research from the Ministry of Health, Labor and Welfare (H-trans-5 to Shuji Terai, Isao Sakaida, Hiroshi Nishina and Kiwamu Okita).

5 References

- [1] Tanaka, E., Kiyosawa, K., *J. Gastroenterol. Hepatol.* 2000, 15, E97–E104.
- [2] Niederau, C., Lange, S., Heintges, T., Erhardt, A., *et al.*, *Hepatology* 1998, 28, 1687–1695.
- [3] Smith, M. W., Yue, Z. N., Geiss, G. K., Sadovnikova, N. Y., *et al.*, *Cancer Res.* 2003, 63, 859–864.
- [4] Chung, E. J., Sung, Y. K., Farooq, M., Kim, Y., *et al.*, *Mol. Cells* 2002, 14, 382–387.
- [5] Iizuka, N., Oka, M., Yamada-Okabe, H., Mori, N., *et al.*, *Cancer Res.* 2002, 62, 3939–3944.
- [6] Xu, X. R., Huang, J., Xu, Z. G., Qian, B. Z., *et al.*, *Proc. Natl. Acad. Sci. USA* 2001, 98, 15089–15094.
- [7] Takeo, S., Arai, H., Kusano, N., Harada, T., *et al.*, *Cancer Genet. Cytogenet.* 2001, 130, 127–132.
- [8] Shirota, Y., Kaneko, S., Honda, M., Kawai, H. F., Kobayashi, K., *Hepatology* 2001, 33, 832–840.
- [9] Okabe, H., Satoh, S., Kato, T., Kitahara, O., *et al.*, *Cancer Res.* 2001, 61, 2129–2137.
- [10] Lau, W. Y., Lai, P. B., Leung, M. F., Leung, B. C., *et al.*, *Oncol. Res.* 2000, 12, 59–69.
- [11] Park, K. S., Cho, S. Y., Kim, H., Paik, Y. K., *Int. J. Cancer* 2002, 97, 261–265.

- [12] Park, K. S., Kim, H., Kim, N. G., Cho, S. Y., et al., *Hepatology* 2002, 35, 1459–1466.
- [13] Kim, J., Kim, S. H., Lee, S. U., Ha, G. H., et al., *Electrophoresis* 2002, 23, 4142–4156.
- [14] Lim, S. O., Park, S. J., Kim, W., Park, S. G., et al., *Biochem. Biophys. Res. Commun.* 2002, 291, 1031–1037.
- [15] Tannapfel, A., Anhalt, K., Hausermann, P., Sommerer, F., et al., *J. Pathol.* 2003, 201, 238–249.
- [16] Takashima, M., Kuramitsu, Y., Yokoyama, Y., Iizuka, T., et al., *Proteomics* 2003, 3, 2487–2493.
- [17] van der Loop, F. T., Schaart, G., Timmer, E. D., Ramaekers, F. C., van Eys, G. J., *J. Cell Biol.* 1996, 134, 401–411.
- [18] van Eys, G. J., Voller, M. C., Timmer, E. D., Wehrens, X. H., et al., *Cell Struct. Funct.* 1997, 22, 65–72.
- [19] Wehrens, X. H., Mies, B., Gimona, M., Ramaekers, F. C., et al., *FEBS. Lett.* 1997, 405, 315–320.
- [20] Rensen, S. S., Thijssen, V. L., De Vries, C. J., Doevendans, P. A., et al., *Cardiovasc. Res.* 2002, 55, 850–863.
- [21] Blomhoff, R., Rasmussen, M., Nilsson, A., Norum, K. R., et al., *J. Biol. Chem.* 1985, 260, 13560–13565.
- [22] Wake, K., *Int. Rev. Cytol.* 1980, 66, 303–353.
- [23] Schurch, W., Seemayer, T. A., Gabbiani, G., *Am. J. Surg. Pathol.* 1998, 22, 141–147.
- [24] Ramadori, G., Veit, T., Schwogler, S., Dienes, H. P., et al., *Virchows Arch. B* 1990, 59, 349–357.
- [25] Ishikawa, R., Yamashiro, S., Matsumura, F., *J. Biol. Chem.* 1989, 264, 16764–16770.
- [26] Ishikawa, R., Yamashiro, S., Matsumura, F., *J. Biol. Chem.* 1989, 264, 7490–7497.
- [27] Bhattacharya, B., Prasad, G. L., Valverius, E. M., Salomon, D. S., Cooper, H. L., *Cancer Res.* 1990, 50, 2105–2112.
- [28] Mahadev, K., Raval, G., Bharadwaj, S., Willingham, M. C., et al., *Exp. Cell Res.* 2002, 279, 40–51.
- [29] Bharadwaj, S., Prasad, G. L., *Cancer Lett.* 2002, 183, 205–213.
- [30] Alaiya, A. A., Franzen, B., Fujioka, K., Moberger, B., et al., *Int. J. Cancer* 1997, 73, 678–683.
- [31] Ahrm, M., Best, C. J., Flaig, M. J., Gillespie, J. W., et al., *Mol. Carcinog.* 2002, 33, 9–15.
- [32] Jung, M. H., Kim, S. C., Jeon, G. A., Kim, S. H., et al., *Genomics* 2000, 69, 281–286.
- [33] Tada, A., Kato, H., Hasegawa, S., *Oncol. Rep.* 2000, 7, 1323–1326.
- [34] Taguchi, K., Takagi, Y., *Rinsho Byori* 2001, Suppl. 116, 117–124.
- [35] Adelman, R. C., Morris, H. P., Weinhouse, S., *Cancer Res.* 1967, 27, 2408–2413.
- [36] Kinoshita, M., Miyata, M., *Hepatology* 2002, 36, 433–438.
- [37] Kimura, T., *Hokkaido Igaku Zasshi* 1994, 69, 1232–1243.
- [38] Castaldo, G., Calcagno, G., Sibillo, R., Cuomo, R., et al., *Clin. Chem.* 2000, 46, 901–906.
- [39] Costanzo, P., Izzo, P., Lupo, A., Rippa, E., et al., *Ital. J. Biochem.* 1988, 37, 8–13.

Transplantation of Bone Marrow Cells Reduces CCl₄-Induced Liver Fibrosis in Mice

Isao Sakaida,¹ Shuji Terai,¹ Naoki Yamamoto,¹ Koji Aoyama,¹ Tsuyoshi Ishikawa,¹ Hiroshi Nishina,² and Kiwamu Okita¹

We investigated the effect of bone marrow cell (BMC) transplantation on established liver fibrosis. BMCs of green fluorescent protein (GFP) mice were transplanted into 4-week carbon tetrachloride (CCl₄)-treated C57BL6 mice through the tail vein, and the mice were treated for 4 more weeks with CCl₄ (total, 8 weeks). Sirius red and GFP staining clearly indicated migrated BMCs existing along with fibers, with strong expression of matrix metalloproteinase (MMP)-9 shown by anti-MMP-9 antibodies and *in situ* hybridization. Double fluorescent immunohistochemistry showed the expression of MMP-9 on the GFP-positive cell surface. Film *in situ* zymographic analysis revealed strong gelatinolytic activity in the periportal area coinciding with the location of MMP-9-positive BMCs. Four weeks after BMC transplantation, mice had significantly reduced liver fibrosis, as assessed by hydroxyproline content of the livers, compared to that of mice treated with CCl₄ alone. Subpopulation of Liv8-negative BMCs was responsible for this fibrolytic effect. **In conclusion**, mice with BMC transplants with continuous CCl₄ injection had reduced liver fibrosis and a significantly improved survival rate after BMC transplantation compared with mice treated with CCl₄ alone. This finding introduces a new concept for the therapy of liver fibrosis. *Supplementary material for this article can be found on the HEPATOLOGY website (<http://interscience.wiley.com/jpages/0270-9139/suppmat/index.html>). (HEPATOLOGY 2004;40: 1304–1311.)*

Recent reports have shown the capacity of the bone marrow cell (BMC) to differentiate into a variety of non-hematopoietic cell lineages.^{1–5} These results indicate that the BMC is an attractive cell source for regenerative medicine compared with tissue-specific stem cells.⁶ The capacity of the BMC to differentiate into hepatocytes and intestinal cells has been shown by Y-chromosome detection in autopsy analysis of human female

recipients of BMCs from male donors.^{7,8} Although Lagasse et al. reported that purified hematopoietic stem cells could differentiate into hepatocytes using a fumarylacacetate hydrazase-deficient model,⁵ Wagers et al. showed little evidence of plasticity in adult hematopoietic stem cells.⁹ Thus, although there is still controversy about which part of BMCs can differentiate into hepatocytes, the BMC seems to have the plasticity to differentiate into such cells. From the point of view of therapy, one of the targets of liver disease for BMC transplantation is liver cirrhosis with chronic liver failure. This is an unphysiological condition with excessive deposition of extracellular matrix and a relative lack of parenchymal cells (hepatocytes). Even if BMC transplantation is successful in supplying parenchymal cells, the fate of the extracellular matrix under these conditions is unknown. The present study clearly shows that transplanted BMCs reduce (degrade) carbon tetrachloride (CCl₄)-induced liver fibrosis with a significantly improved survival rate.

Abbreviations: BMC, bone marrow cell; CCl₄, carbon tetrachloride; GFP, green fluorescent protein; PBS, phosphate-buffered saline; IgG, immunoglobulin G; MMP, matrix metalloproteinase; NGS, normal goat serum; NRS, normal rabbit serum; DIG, digoxigenin.

From the ¹Department of Gastroenterology and Hepatology, School of Medicine, Yamaguchi University, Yamaguchi, Japan; and ²Department of Physiological Chemistry, Graduate School of Pharmaceutical Science, University of Tokyo, Tokyo, Japan.

Received September 30, 2003; accepted August 16, 2004.

Supported in part by grants-in-aid 16590597, 12670490, and 10470136 from the Ministry of Education, Culture, Sports, Science and Technology and by grants-in-aid for translational research from the Ministry of Health, Labor and Welfare of Japan.

Address reprint requests to: Isao Sakaida, M.D., Department of Gastroenterology and Hepatology, School of Medicine, Yamaguchi University, Minami-Kogushi 1-1-1 Ube, Yamaguchi-pref. 755-8505, Japan. E-mail: sakaida@yamaguchi-u.ac.jp; fax: (81) 836-22-2240.

Copyright © 2004 by the American Association for the Study of Liver Diseases.

Published online in Wiley InterScience (www.interscience.wiley.com).

DOI 10.1002/hep.20452

Materials and Methods

Mice. GFP-transgenic mice (TgN(β-act-EGFP)Osb) were kindly provided by Masaru Okabe (Genome Research Center, Osaka University, Osaka, Japan).¹⁰ C57BL6 female mice were purchased from Japan SLC