

37. Sansilvestri, P., A.A. Cardoso, P. Batard, B. Panterne, A. Hatzfeld and B. Lim *et al.*, 1995. Early CD34<sup>high</sup> cells can be separated into KITHigh cells in which transforming growth factor-beta (TGF-beta) downmodulates c-kit and KITlow cells in which anti-TGF-beta upmodulates c-kit. *Blood*, 86: 1729-1735.
38. Batard, P., M.N. Monier, N. Fortunel, K. Ducos, P. Sansilvestri-Morel, T. Phan, A. Hatzfeld and J.A. Hatzfeld, 2000. TGF-(beta)1 maintains hematopoietic immaturity by a reversible negative control of cell cycle and induces CD34 antigen up-modulation. *J. Cell Sci.*, 111: 1867-1875.
39. Bhatia, M., D. Bonnet, D. Wu, B. Murdoch, J. Wrana, L. Gallacher and J.E. Dick, 2001. Bone morphogenetic proteins regulate the developmental program of human hematopoietic stem cells. *J. Exp. Med.*, 189: 1139-48.
40. Zuckerman, K.S. and M.S. Wicha, 1983. Extracellular matrix production by the adherent cells of long-term murine bone marrow cultures. *Blood*, 61: 540-547.
41. Campbell, A., M.S. Wicha and M. Long, 1985. Extracellular matrix promotes the growth and differentiation of murine hematopoietic cells *in vitro*. *J. Clin. Invest.*, 75: 2085-2090.
42. Long, M.W., R. Briddell, A.W. Walter, E. Bruno and R. Hoffman, 1992. Human hematopoietic stem cell adherence to cytokines and matrix molecules. *J. Clin. Invest.*, 90: 251-255.
43. Adams, J.C. and F.M. Watt, 1993. Regulation of development and differentiation by the extracellular matrix. *Development*, 117: 1183-1198.
44. Long, M.W., 1992. Blood cell cytoadhesion molecules. *Exp. Hematol.*, 20: 288-301.
45. Verfaillie, C., R. Hurley, R. Bhatia and J.B. McCarthy, 1994. Role of bone marrow matrix in normal and abnormal hematopoiesis. *Crit. Rev. Oncol. Hematol.*, 16: 201-224.
46. Hynes, R.O., 1992. Integrins: versatility, modulation and signaling in cell adhesion. *Cell*, 69: 11-25.
47. Ruoslahti, E., 1991. Integrins. *J. Clin. Invest.*, 87: 1-5.
48. Kopan, R., J. Lee, M.H. Lin, A.J. Syder, J. Kesterson, N. Crutchfield, C.R. Li, W. Wu, J. Books and J.I. Gordon, 2002. Genetic mosaic analysis indicates that the bulb region of coat hair follicles contains a resident population of several active multipotent epithelial lineage progenitors. *Dev. Biol.*, 242: 44-57.
49. Marshman, E., C. Booth and C.S. Potten, 2002. The intestinal epithelial stem cell. *Bioessays*, 24: 91-8.
50. Nishimura, E.K., S.A. Jordan, H. Oshima, H. Yoshida, M. Osawa, M. Moriyama, I.J. Jackson, Y. Barrandon, Y. Miyachi and S. Nishikawa, 2002. Dominant role of the niche in melanocyte stem-cell fate determination. *Nature*, 416: 854-60.
51. Zhang, J., C. Niu, L. Ye, H. Huang, X. He, W.G. Tong, J. Ross, J. Haug, T. Johnson, J.Q. Feng, S. Harris, L.M. Wiedemann, Y. Mishina and L. Li, 2003. Identification of the haematopoietic stem cell niche and control of the niche size. *Nature*, 425: 836-41.
52. Calvi, L.M., G.B. Adams, K.W. Weibrecht, J.M. Weber, D.P. Olson, M.C. Knight, R.P. Martin, E. Schipani, P. Divieti, F.R. Bringhurst, L.A. Milner, H.M. Kronenberg and D.T. Scadden, 2003. Osteoblastic cells regulate the haematopoietic stem cell niche. *Nature*, 425: 841-6.
53. Yokota, T., K. Oritani, H. Mitsui, K. Aoyama, J. Ishikawa, H. Sugahara, I. Matsumura, S. Tsai, Y. Tomiyama, Y. Kanakura and Y. Matsuzawa, 1998. Growth-supporting activities of fibronectin on hematopoietic stem/progenitor cells *in vitro* and *in vivo*: structural requirement for fibronectin activities of CS1 and cell-binding domains. *Blood*, 91: 3263-3272.
54. Takakura, N., X.L. Huang, T. Naruse, I. Hamaguchi, D.J. Dumont, G.D. Yancopoulos and T. Suda, 1998. Critical role of the TIE2 endothelial cell receptor in the development of definitive hematopoiesis. *Immunity*, 9: 677-686.
55. Arai, F., A. Hirao, M. Ohmura, H. Sato, S. Matsuoka, K. Takubo, K. Ito, G.Y. Koh and T. Suda, 2004. Tie2/angiopoietin-1 signaling regulates hematopoietic stem cell quiescence in the bone marrow niche. *Cell*, 118: 149-61.
56. Varnum-Finney, B., L.E. Purton, M. Yu, C. Brashem-Stein, D. Flowers, S. Staats, K.A. Moore I. Le Roux, R. Mann, G. Gray, S. Artavanis-Tsakonas and I.D. Bernstein, 1998. The Notch ligand, Jagged-1, influences the development of primitive hematopoietic precursor cells. *Blood*, 91: 4084-4091.
57. Varnum-Finney, B., L. Xu, C. Brashem-Stein, C. Nourigat, D. Flowers, S. Bakkour, W.S. Pear and I.D. Bernstein, 2000. Pluripotent, cytokine-dependent, hematopoietic stem cells are immortalized by constitutive Notch1 signaling. *Nat. Med.*, 6: 1278-1281.
58. Karanu, F.N., B. Murdoch, L. Gallacher, D.M. Wu, M. Koremoto, S. Sakano *et al.*, 2000. The notch ligand jagged-1 represents a novel growth factor of human hematopoietic stem cells. *J. Exp. Med.*, 192: 1365-1372.
59. Karanu, F.N., B. Murdoch, T. Miyabayashi, M. Ohno, M. Koremoto, L. Gallacher *et al.*, 2001. Human homologues of Delta-1 and Delta-4 function as mitogenic regulators of primitive human hematopoietic cells. *Blood*, 97: 1960-1967.

60. Ohishi, K., B. Varnum-Finney and I.D. Bernstein, 2002. Delta-1 enhances marrow and thymus repopulating ability of human CD34(+)CD38(-) cord blood cells. *J. Clin. Invest.*, 110: 1165-1174.
61. Satoh, Y., I. Matsumura, H. Tanaka, S. Ezoe, H. Sugahara, M. Mizuki, H. Shibayama, E. Ishiko, J. Ishiko, K. Nakajima, Y. Karakura, 2004. Roles for c-Myc in self-renewal of hematopoietic stem cells. *J. Biol. Chem.*, 279: 24986-93.
62. Reya, T., M. O'Riordan, R. Okamura, E. Devaney, K. Willert, R. Nusse and R. Grosschedl, 2000. Wnt signaling regulates B lymphocyte proliferation through a LEF-1 dependent mechanism. *Immunity*, 13: 15-24.
63. Ivanova, N.B., J.T. Dimos, C. Schaniel, J.A. Hackney, K.A. Moore and I.R. Lemischka, 2002. A stem cell molecular signature. *Sciences*, 298: 601-4.
64. Reya, T., A.W. Duncan, L. Ailles, J. Domen, D.C. Scherer, K. Willert, L. Hintz, R. Nusse and I.L. Weissman, 2003. A role for Wnt signalling in self-renewal of haematopoietic stem cells. *Nature*, 423: 409-14.
65. Murdoch, B., K. Chadwick, M. Martin, F. Shojaei, K.V. Shah, L. Gallacher, R.T. Moon and M. Bhatia, 2003. Wnt-5A augments repopulating capacity and primitive hematopoietic development of human blood stem cells *in vivo*. *Proc. Natl. Acad. Sci. USA.*, 100: 3422-7.
66. Murone, M., A. Rosenthal and F.J. de Sauvage, 1999. Hedgehog signal transduction: from flies to vertebrates. *Exp. Cell Res.*, 253: 25-33.
67. Robbins, D.J., K.E. Nybakken, R. Kobayashi, J.C. Sisson, J.M. Bishop and P.P. Therond, 1997. Hedgehog elicits signal transduction by means of a large complex containing the kinesin-related protein costal2. *Cell*, 90: 225-234.
68. Bhardwaj, G., B. Murdoch and D. Wu, 2001. Sonic hedgehog induces the proliferation of primitive human hematopoietic cells via BMP regulation. *Natl. Immunol.*, 2: 178-180.
69. Melotti, P. and B. Calabretta, 1996. The transcription factors c-myc and GATA-2 act independently in the regulation of normal hematopoiesis. *Proc. Natl. Acad. Sci. USA.*, 93: 5313-5318.
70. Schmidt, M., V. Nazarov, L. Stevens, R. Watson and L. Wolff, 2000. Regulation of the resident chromosomal copy of c-myc by c-Myb is involved in myeloid leukemogenesis. *Mol. Cell. Biol.*, 20: 1970-1981.
71. Mucenski, M.L., K. McLain, A.B. Kier, S.H. Swerdlow, C.M. Schreiner and T.A. Miller *et al.*, 1991. A functional c-myc gene is required for normal murine fetal hepatic hematopoiesis. *Cell*, 65: 677-689.
72. Tsai, F.Y., G. Keller, F.C. Kuo, M. Weiss, J. Chen, M. Rosenblatt, F.W. Alt and S.H. Orkin, 1994. An early haematopoietic defect in mice lacking the transcription factor GATA-2. *Nature*, 371: 221-226.
73. Heyworth, C., K. Gale, M. Dexter, G. May and T. Enver, 1999. A GATA-2/estrogen receptor chimera functions as a ligand-dependent negative regulator of self-renewal. *Genes. Dev.*, 13: 1847-1860.
74. Ezoe, S., I. Matsumura, S. Nakata, K. Gale, K. Ishihara, N. Minegishi, T. Machii, Kitamura, T., Yamamoto, T. Enver and Y. Karakura, 2002. GATA-2/estrogen receptor chimera regulates cytokine-dependent growth of hematopoietic cells through accumulation of p21(WAF1) and p27(Kip1) proteins. *Blood*, 100: 3512-3520.
75. Kitajima, K., M. Masuhara, T. Era, T. Enver and T. Nakano, 2002. GATA-2 and GATA-2/ER display opposing activities in the development and differentiation of blood progenitors. *EMBO J.*, 21: 3060-3069.
76. Persons, D.A., J.A. Allay, E.R. Allay, R.A. Ashmun, D. Orlic and S.M. Jane, 1999. Enforced expression of the GATA-2 transcription factor blocks normal hematopoiesis. *Blood*, 93: 488-499.
77. Sauvageau, G., U. Thorsteinsdottir, C.J. Eaves, H.J. Lawrence, C. Largman, P.M. Lansdorp and R.K. Humphries, 1995. Overexpression of HOXB4 in hematopoietic cells causes the selective expansion of more primitive populations *in vitro* and *in vivo*. *Genes Dev.*, 9: 1753-1765.
78. Antonchuk, J., G. Sauvageau and R.K. Humphries, 2002. HOXB4-induced expansion of adult hematopoietic stem cells *ex vivo*. *Cell*, 109: 39-45.
79. Buske, C., M. Feuring-Buske, C. Abramovich, K. Spiekermann, C.J. Eaves, L. Coulombel, G. Sauvageau, D.E. Hogge and R.K. Humphries, 2002. Deregulated expression of HOXB4 enhances the primitive growth activity of human hematopoietic cells. *Blood*, 100: 862-868.
80. Krosi, J., P. Austin, N. Beslu, E. Kroon, R.K. Humphries and G. Sauvageau, 2003. *In vitro* expansion of hematopoietic stem cells by recombinant TAT-HOXB4 protein. *Natl. Med.*, 9: 1428-32.
81. Bjornsson, J.M., N. Larsson, A.C. Brun, M. Magnusson, E. Andersson and P. Lundstrom *et al.*, 2003. Reduced proliferative capacity of hematopoietic stem cells deficient in HOXB3 and HOXB4. *Mol. Cell. Biol.*, 23: 3872-3883.
82. Mahmoudi, T. and C.P. Verrijzer, 2001. Chromatin silencing and activation by Polycomb and trithorax group proteins. *Oncogene*, 20: 3055-3066.

83. Park, I.K., D. Qian, M. Kiel, M.W. Becker, M. Pihalja, I.L. Weissman, S.J. Morrison and M.F. Clarke, 2003. Bmi-1 is required for maintenance of adult self-renewing haematopoietic stem cells. *Nature*, 423: 302-305.
84. Jacobs, J.J., K. Kieboom, S. Marino, R.A. DePinho and M. van Lohuizen, 1999. The oncogene and Polycomb-group gene *bmi-1* regulates cell proliferation and senescence through the *ink4a* locus. *Nature*, 397: 164-168.
85. Wohlschlegel, J.A., B.T. Dwyer, S.K. Dhar, C. Cvetic, J.C. Walterm and A. Dutta, 2000. Inhibition of eukaryotic DNA replication by geminin binding to Cdt1. *Sci.*, 290: 2309-12.
86. Tada, S., A. Li, D. Maiorano, M. Mechali, J.J. Blow, 2001. Repression of origin assembly in metaphase depends on inhibition of RLF-B/Cdt1 by geminin. *Nat. Cell Biol.*, 3: 107-113.
87. Luo, L., X. Yang, Y. Takihara, H. Knoetgen and M. Kessel, 2004. The cell-cycle regulator geminin inhibits Hox function through direct and polycomb-mediated interactions. *Nature*, 427: 749-53.
88. Brugarolas, J., C. Chandrasekaran, J.I. Gordon, D. Beach, T. Jacks and G.J. Hannon, 1995. Radiation-induced cell cycle arrest compromised by p21 deficiency. *Nature*, 377: 552-557.
89. Deng, C., P. Zhang, J.W. Harper, S.J. Elledge and P. Leder, 1995. Mice lacking p21<sup>Cip1</sup>/WAF1 undergo normal development, but are defective in G1 checkpoint control. *Cell*, 82: 675-684.
90. Nakanishi, M., G.R. Adami, R.S. Robetorye, A. Noda, S.F. Venable and D. Dimitrov, 1995. Exit from G0 and entry into the cell cycle of cells expressing p21<sup>Sdi1</sup> antisense RNA. *Proc. Natl. Acad. Sci. USA.*, 92: 4352-4356.
91. Taniguchi, T., H. Endo, N. Chikatsu, K. Uchimar, S. Asano, T. Fujita, T. Nakahata and T. Motokura, 1999. Expression of p21<sup>Cip1</sup>/Waf1/Sdi1 and p27<sup>Kip1</sup> cyclin-dependent kinase inhibitors during human hematopoiesis. *Blood*, 93: 4167-4178.
92. Yaroslavskiy, B., S. Watkins, A.D. Donnenberg, T.J. Patton and R.A. Steinman, 1999. Subcellular and cell-cycle expression profiles of CDK-inhibitors in normal differentiating myeloid cells. *Blood*, 93: 2907-2917.
93. Cheng, T., N. Rodrigues, H. Shen, Y. Yang, D. Dombkowski, M. Sykes, D.T. Scadden, 2000. Hematopoietic stem cell quiescence maintained by p21<sup>Cip1</sup>/waf1. *Sciences*, 287: 1804-1808.
94. Nakayama, K., N. Ishida, M. Shirane, A. Inomata, T. Inoue, N. Shishido, I. Horii, D.Y. Loh and K. Nakayama, 1996. Mice lacking p27<sup>Kip1</sup> display increased body size, multiple organ hyperplasia, retinal dysplasia and pituitary tumors. *Cell*, 85: 707-720.
95. Kiyokawa, H., R.D. Kineman, K.O. Manova-Todorova, V.C. Soares, E.S. Hoffman, M. Ono, D. Khanam, A.C. Hayday, L.A. Frohman and A. Koff, 1996. Enhanced growth of mice lacking the cyclin-dependent kinase inhibitor function of p27<sup>Kip1</sup>. *Cell*, 85: 721-732.
96. Fero, M.L., M. Rivkin, M. Tasch, P. Porter, C.E. Carow, E. Firpo, K. Polyak, L.H. Tsai, V. Broudy, R.M. Perlmutter, K. Kaushansky and J.M. Roberts, 1996. A syndrome of multiorgan hyperplasia with features of gigantism, tumorigenesis and female sterility in p27<sup>Kip1</sup>-deficient mice. *Cell*, 85: 733-744.
97. Cheng, T., N. Rodrigues, D. Dombkowski, S. Stier and D.T. Scadden, 2000. Stem cell repopulation efficiency but not pool size is governed by p27<sup>Kip1</sup>. *Nat. Med.*, 6: 1235-1240.
98. Yuan, Y., H. Shen, D.S. Franklin, D.S., D.T. Scadden and T. Cheng, 2004. *In vivo* self-renewing divisions of haematopoietic stem cells are increased in the absence of the early G1-phase inhibitor, p18<sup>INK4C</sup>. *Nat. Cell Biol.*, 6: 436-42.
99. Furukawa, U., J. Kikuchi, M. Nakamura, S. Iwase, H. Yamada and M. Matsuda, 2000. Lineage-specific regulation of cell cycle control gene expression during haematopoietic cell differentiation. *Brit. J. Haemat.*, 110: 663-673.
100. Nakayama, K. and K. Nakayama, 1998. Cip/Kip cyclin-dependent kinase inhibitors: brakes of the cell cycle engine during development. *Bioessays*, 20: 1020-1029.
101. Serrano, M., H.W. Lee, L. Chin, C. Cordon-Cardo, D. Beach and R.A. DePinho, 1996. Role of the *INK4a* locus in tumor suppression and cell mortality. *Cell*, 85: 27-37.
102. Teofili, L., S. Rutella, P. Chiusolo, E.O. La Barbera, C. Rumi, F.O. Ranelletti, N. Maggiano, G. Leone and L.M. Larocca, 1998. Expression of p15<sup>INK4B</sup> in normal hematopoiesis. *Exp. Hematol.*, 26: 1133-1139.
103. Serrano, M., A.W. Lin, M.E. McCurrach, D. Beach and S.W. Lowe, 1997. Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16<sup>INK4a</sup>. *Cell*, 88: 593-602.
104. Malumbres, M., I. Perez De Castro, M.I. Hernandez, M. Jimenez, T. Corral and A. Pellicer, 2000. Cellular response to oncogenic ras involves induction of the Cdk4 and Cdk6 inhibitor p15<sup>INK4b</sup>. *Mol. Cell Biol.*, 20: 2915-2925.
105. Kamb, A., N.A. Gruis, J. Weaver-Feldhaus, Q. Liu, K. Harshman, S.V. Tavtigian, E. Stockert, R.S. Day, B.E. 3rd Johnson and M.H. Skolnick, 1994. A cell cycle regulator potentially involved in genesis of many tumor types. *Science*, 264: 436-440.

106. Ruas, M. and G. Peters, 1998. The p16INK4a/CDKN2A tumor suppressor and its relatives. *Biochim. Biophys. Acta.*, 11378: F115-F177.
107. Ogawa, S., A. Hangaishi, S. Miyawaki, S. Hirose, Y. Miura, K. Takeyama, N. Kamada, S. Ohtake, N. Uike and C. Shimazaki, 1995. Loss of the cyclin-dependent kinase 4-inhibitor (p16; MTS1) gene is frequent in and highly specific to lymphoid tumors in primary human hematopoietic malignancies. *Blood*, 86: 1548-1556.
108. Haidar, M.A., X.B. Cao, T. Manshouri, L.L. Chan, A. Glassman, H.M. Kantarjian, M.J. Keating, M.S. Beran and M. Albitar, 1995. p16INK4A and p15INK4B gene deletions in primary leukemias. *Blood*, 86: 311-315.
109. Gombart, A.F., R. Morosetti, C.W. Miller, J.W. Said and H.P. Koeffler, 1995. Deletions of the cyclin-dependent kinase inhibitor genes p16INK4A and p15INK4B in non-Hodgkin's lymphomas. *Blood*, 86: 1534-1539.
110. Hangaishi, A., S. Ogawa, N. Imamura, S. Miyawaki, Y. Miura, N. Uike, C. Shimazaki, N. Emi, K. Takeyama, S. Hirose, N. Kamada, Y. Kobayashi, Y. Takemoto, T. Kitani, K. Toyama, S. Ohtake, Y. Yazaki, R. Ueda and H. Hirai, 1996. Inactivation of multiple tumor-suppressor genes involved in negative regulation of the cell cycle, MTS1/p16INK4A/CDKN2, MTS2/p15INK4B, p53 and Rb genes in primary lymphoid malignancies. *Blood*, 87: 4949-4958.
111. Uchida, T., T. Kinoshita, H. Nagai, Y. Nakahara, H. Saito, T. Hotta and T. Murate, 1997. Hypermethylation of the p15INK4B gene in myelodysplastic syndromes. *Blood*, 90: 1403-1409.

厚生労働科学研究費補助金  
基礎研究成果の臨床応用推進研究事業  
Ex vivo 増幅臍帯血幹細胞を用いた  
トランスレーショナルリサーチ  
平成 14～16 年度 総合研究報告書

発行 平成 17 年 3 月  
発行者 中畑 龍俊 (主任研究者)  
事務局 〒650-0047 神戸市中央区港島南町2丁目2番  
先端医療センター再生医療研究部  
TEL : 078-304-5773  
印刷所 株式会社 トライス