

- mice with superoxide dismutase/catalase mimics. *Proc Natl Acad Sci USA* 2003;100:8526–31.
- [29] Lobo A, Launer LJ, Fratiglioni L, Andersen K, Di Carlo A, Breteler MM, et al. Prevalence of dementia and major subtypes in Europe: a collaborative study of population-based cohorts. *Neurol Dis Elderly Res Group Neurol* 2000;54:S4–9.
- [30] Lynch T, Cherny RA, Bush AI. Oxidative processes in Alzheimer's disease: the role of A β -metal interactions. *Exp Gerontol* 2000;35:445–51.
- [31] Lytras L, Cairns NJ, Jenner A, Jenner P, Halliwell B. An assessment of oxidative damage to proteins, lipids, and DNA in brain from patients with Alzheimer's disease. *J Neurochem* 1997;68:2061–9.
- [32] McCord JM, Fridovich I. Superoxide dismutase. Enzymatic function for erythrocuprein. *J Biol Chem* 1969;244:6049–55.
- [33] Mark RJ, Lovell MA, Markesberry WR, Uchida K, Mattson MP. A role for 4-hydroxyxonenal, an aldehydic product of lipid peroxidation, in disruption of iron homeostasis and neuronal death induced by amyloid β -peptide. *J Neurochem* 1997;68:255–64.
- [34] Markesberry WR. Oxidative stress hypothesis in Alzheimer's disease. *Free Radic Biol Med* 1997;23:134–47.
- [35] Matsuoka Y, Picciano M, La Francois J, Duff K. Fibrillar β -amyloid evokes oxidative damage in a transgenic mouse model of Alzheimer's disease. *Neuroscience* 2001;104:609–13.
- [36] Maurice T, Lockhart BP, Privat A. Amnesia induced by centrally administered β -amyloid peptides involves cholinergic dysfunction. *Brain Res* 1996;706:181–93.
- [37] Morris R. Developments of a water maze procedure for studying spatial learning in the rat. *J Neurosci Meth* 1984;11:47–60.
- [38] Niki E, Yamamoto Y, Komuro E, Sato K. Membrane damage due to lipid peroxidation. *Am J Clin Nutr* 1991;53:201S–5S.
- [39] Nitta A, Itoh A, Hasegawa T, Nabeshima T. β -Amyloid protein-induced Alzheimer's disease animal model. *Neurosci Lett* 1994;170:63–6.
- [40] Nitta A, Fukuta T, Hasegawa T, Nabeshima T. Continuous infusion of β -amyloid protein into cerebral ventricle induces learning impairment and neuronal and morphological degeneration. *Jpn J Pharmacol* 1997;73:51–7.
- [41] Oliver CN, Ahn BW, Moerman EJ, Goldstein S, Statman ER. Age-related changes in oxidized proteins. *J Biol Chem* 1987;262:5488–91.
- [42] Ratan RR, Murphy TH, Baraban JM. Macromolecular synthesis inhibitors prevent oxidative stress-induced apoptosis by shunting cysteine to glutathione. *J Neurosci* 1994;14:4385–92.
- [43] Reed DJ. Glutathione: toxicological implications. *Ann Rev Pharmacol Toxicol* 1990;30:603–31.
- [44] Stadtman ER. Protein oxidation and aging. *Science* 1992;257:1220–4.
- [45] Subramaniam R, Roediger F, Jorda B, Mattson MP, Keller JN, Waeg G, et al. The lipid peroxidation product, 4-hydroxy-2-trans-nonenal, alters the conformation of cortical synaptosomal membrane proteins. *J Neurochem* 1997;69:1161–9.
- [46] Tran MH, Yamada K, Olariu A, Mizuno M, Ren XH, Nabeshima T. Amyloid β -peptide induces nitric oxide production in rat hippocampus: association with cholinergic dysfunction and amelioration by inducible nitric oxide synthase inhibitors. *FASEB J* 2001;15:1407–9.
- [47] Tran MH, Yamada K, Nakajima A, Mizuno M, He J, Kamei H, et al. Tyrosine nitration of a synaptic protein synaptophysin contributes to amyloid β -peptide-induced cholinergic dysfunction. *Mol Psychiatry* 2003;8:407–12.
- [48] Woo JI, Lee JH, Yoo KY, Kim CY, Kim YI, Shin YS. Prevalence estimation of dementia in a rural area of Korea. *J Am Geriatr Soc* 1998;46:983–7.
- [49] Wortwein G, Stackman RW, Walsh TJ, Vitamin E. prevents the place learning deficit and the cholinergic hypofunction induced by AF64A. *Exp Neurol* 1994;125:15–21.
- [50] Yamada K, Tanaka T, Zou LB, Senzaki K, Yano K, Osada T, et al. Long-term deprivation of oestrogens by ovariectomy potentiates β -amyloid-induced working memory deficits in rats. *Br J Pharmacol* 1999;128:419–27.
- [51] Yamada K, Tanaka T, Han D, Senzaki K, Kameyama T, Nabeshima T. Protective effects of idebenone and alpha-tocopherol on β -amyloid-(1–42)-induced learning and memory deficits in rats: implication of oxidative stress in beta-amyloid-induced neurotoxicity in vivo. *Eur J Neurosci* 1999;11:83–90.
- [52] Yan JJ, Cho JY, Kim HS, Kim KL, Jung JS, Huh SO, et al. Protection against beta-amyloid peptide toxicity in vivo with long-term administration of ferulic acid. *Br J Pharmacol* 2001;133:89–96.
- [53] Yankner BA. Mechanisms of neuronal degeneration in Alzheimer's disease. *Neuron* 1996;250:279–82.
- [54] Yatin SM, Varadarajan S, Link CD, Butterfield DA. In vitro and in vivo oxidative stress associated with Alzheimer's amyloid β -peptide (1–42). *Neurobiol Aging* 1999;20:325–30.
- [55] Yatin SM, Varadarajan S, Butterfield DA. Vitamin E prevents Alzheimer's amyloid beta-peptide (1–42)-induced neuronal protein oxidation and reactive oxygen species production. *J Alzheimers Dis* 2000;2:123–31.