

PAPER

Differences in the relationship between lipid CHD risk factors and body composition in Caucasians and Japanese

N Iwao^{1,4}, S Iwao^{1,4}, DC Muller¹, M Koda², F Ando², H Shimokata², F Kobayashi³ and R Andres^{1*}

¹Gerontology Research Center/National Institutes for Health/National Institute on Aging, Baltimore, MD, USA; ²Department of Epidemiology, National Institute for Longevity Sciences, Gengo, Morioka, Obu, Aichi, Japan; and ³Department of Health and Psychosocial Medicine, Institute for Medical Science of Aging, Aichi Medical University School of Medicine, Yasago, Nagakute, Aichi-gun, Aichi, Japan

OBJECTIVES: To examine differences in the relationship between fat distribution and lipid coronary risk factors in Caucasian and Japanese population and further to determine whether the cut-points for body mass index (BMI) and waist circumference (WC) proposed by WHO and NHLBI are applicable to Japanese population as a predictor of a lipid risk factor abnormality or not.

RESEARCH METHODS AND PROCEDURES: Subjects were 895 participants of the Baltimore Longitudinal Study of Aging in the US (BLSA) and 1705 participants of the Longitudinal Study of Aging by the National Institutes for Longevity Science in Japan (NILS-LSA). Subjects were divided into four demographic groups as younger (age < 65 y) men and women, and older (age ≥ 65 y) men and women. Blood total cholesterol, triglycerides, LDL- and HDL-cholesterol and anthropometry were measured. Regression coefficients of BMI and WC on risk factors, sensitivity and specificity of the BMI and WC cut-points for blood lipid abnormality, and mean values of blood lipids at BMI or WC cut-points were computed in both populations.

RESULTS: Height, weight, WC and BMI were significantly greater in the BLSA than those in the NILS-LSA subjects. Total cholesterol, HDL- and LDL-cholesterol were significantly greater in the NILS-LSA than in the BLSA subjects. Sensitivities of BMI and WC cut-points were much lower in the NILS-LSA than in the BLSA subjects. Specificities of BMI and WC cut-points were higher in the NILS-LSA than in the BLSA subjects. Mean values of triglycerides, total cholesterol, HDL- and LDL-cholesterol at BMI = 25 were significantly greater in the NILS-LSA than in the BLSA subjects. At the WC cut-point (94 cm for men, 80 cm for women), mean values of all lipids were significantly greater in the NILS-LSA than in the BLSA subjects with the exception of triglycerides in younger women.

CONCLUSIONS: The Japanese subjects have smaller BMI and WC, worse total and LDL-cholesterol levels and better HDL-cholesterol levels compared to Caucasians. Sensitivities of BMI and WC for predicting lipid risk factor abnormality are much lower in Japanese. The cut-points for BMI and WC proposed by WHO and NHLBI may be too high for predicting an abnormality in triglycerides, total and LDL-cholesterol in Japanese. For detecting an abnormal HDL-cholesterol level, the BMI and WC cut-points may not be as beneficial for the Japanese population as for Caucasians.

International Journal of Obesity (2005) 29, 228–235. doi:10.1038/sj.ijo.0802615

Published online 30 November 2004

Keywords: racial difference; BMI; waist circumference; lipids; coronary risk factor

Introduction

The prediction and prevention of coronary heart disease (CHD) are of great importance, especially in industrial countries. The National Cholesterol Education Program (NCEP)¹ recommended that a fasting lipoprotein profile (total cholesterol, LDL-cholesterol, HDL-cholesterol and triglyceride) should be obtained once every 5 y in all adults aged 20 y or older. LDL-cholesterol was defined as the primary target of cholesterol-lowering therapy. Therefore, detecting an abnormality in blood lipids and initiating early treatment is essential to decrease the incidence of CHD.

*Correspondence: R Andres, National Institute on Aging, NIH, 5600 Nathan Shock Drive, Baltimore, MD 21224, USA.

E-mail: andres@grc.nia.nih.gov

Funding Source: Intramural Research Program, National Institute on Aging.

Presented in part at the Annual Meetings of the World Congress of Gerontology, 2001.

*The first two authors are equal contributors.

Received 14 October 2003; accepted 18 January 2004; published online 30 November 2004

It has been established that obesity, especially central distribution of body fat, is associated with many chronic diseases.²⁻⁶ Although imaging techniques such as magnetic resonance imaging (MRI) and computerized tomography (CT) have been the 'gold standard' methods for quantification of visceral fat, these methods are inconvenient and costly for routine clinical use. Therefore, several surrogates of intra-abdominal fat measurement have been examined.⁷⁻⁹ The body mass index (BMI) has long been used as a convenient, useful index of overweight and obesity. Studies have shown that waist circumference (WC) could be used to predict risk factor abnormalities as a surrogate of body fat distribution.⁹⁻¹¹ Han *et al*¹² indicated the linear relationship between WC and intra-abdominal fat. The World Health Organization (WHO)⁶ and the National Heart, Lung, and Blood Institute (NHLBI)⁵ issued comprehensive recommendations for classifying abnormalities in body weight and body fat distribution. Both reports recommended the BMI and the WC as measures of obesity and fat distribution, and concluded that risk of disease increases at a BMI of 25 kg/m² in both men and women; the WHO report noted increased risk at a WC of 94 cm for men and 80 cm for women.

It has been well known that obesity and fat distribution are influenced by complex factors including social, behavioral, cultural, physiological, metabolic and genetic factors. Differences in the relationships among BMI, WC and waist-hip ratio (WHR) in 19 populations, including Beijing Chinese, were reported by Molarius *et al*.¹³ They concluded that the

optimal screening cutoff point of WC and BMI may be population specific. Also, Hu *et al*¹⁴ showed that a rural Chinese population in developing countries has a different relationship between BMI and CHD from Western populations. However, NHLBI and WHO reports did not propose population-specific cut-points for either BMI or WC. Although it was shown that the cut-points proposed by WHO and NHLBI for BMI and WC were useful predictors for coronary risk factors in Caucasian younger and older men and women in our previous studies,^{15,16} it is still not clear whether or not these cut-points are applicable to relatively short stature and lighter weight populations in an industrial country such as Japan.

The purposes of the present study are (1) to examine the relationship between plasma lipid coronary risk factors and indices of body composition (BMI and WC) in Caucasian and Japanese populations; (2) to test whether recently proposed cut-points of BMI and WC can be applied to the younger and older Japanese population as well as to Caucasians.

Methods

Subjects

The subjects of this study consisted of 566 men and 329 women who participated in the Baltimore Longitudinal Study of Aging (BLSA) in USA, and 868 men and 837 women who participated in the National Longevity Sciences-Longitudinal Study of Aging (NILS-LSA) in Japan. Subjects of both

Table 1 Characteristics of BLSA and NILS subjects

	Younger (40-64 y)				Older (65-80 y)			
	BLSA		NILS		BLSA		NILS	
	N	Mean	N	Mean	N	Mean	N	Mean
Men								
Height (cm)	318	178.6	598	166.8**	248	174.3	270	160.9**
Weight (kg)	318	84.7	598	64.0**	248	77.1	270	57.5**
BMI (kg/m ²)	318	26.5	598	23.0**	248	25.3	270	22.2**
WC (cm)	318	93.8	598	81.9**	248	92.8	270	80.6**
Total cholesterol (mg/dl)	316	191.7	594	212.8**	248	188.4	268	211.1**
Triglycerides (mg/dl)	316	135.8	587	133.5	248	115.0	259	122.0
HDL-cholesterol (mg/dl)	316	41.1	596	57.9**	248	44.0	269	57.3**
LDL-cholesterol (mg/dl)	316	123.3	582	128.6*	248	121.3	257	130.5**
Women								
Height (cm)	183	163.6	620	153.7**	146	159.4	217	147.3**
Weight (kg)	183	65.9	620	53.4**	146	64.2	217	49.4**
BMI (kg/m ²)	183	24.6	620	22.6**	146	25.2	217	22.6**
WC (cm)	183	76.3	620	73.4**	146	79.9	217	75.6**
Total cholesterol (mg/dl)	180	188.5	615	221.9**	145	204.0	216	234.6**
Triglycerides (mg/dl)	180	98.6	596	98.0	145	112.8	211	118.8
HDL-cholesterol (mg/dl)	180	52.9	616	67.9**	145	55.4	217	64.4**
LDL-cholesterol (mg/dl)	179	115.8	595	135.1**	145	126.0	210	147.3**

**P<0.01, significant difference in mean values between BLSA and NILS-LSA subjects. *P<0.05, significant difference in mean values between BLSA and NILS-LSA subjects.

Table 2 Regression coefficient and trend analysis for BMI and WC

	Men				Women			
	Younger		Older		Younger		Older	
	BLSA	NILS	BLSA	NILS	BLSA	NILS	BLSA	NILS
(a) BMI								
Total cholesterol	1.33	1.39	-0.44	0.36	1.98	0.68	0.01	0.07
Triglycerides	7.14	8.74	8.05	4.47	4.04	5.25	3.13	4.47
HDL-cholesterol	-0.61	-1.75**	-1.33	-1.49	-0.40	-1.65**	-1.02	-1.72
LDL-cholesterol	0.51	1.39	-0.75	0.93	1.58	1.21	0.40	1.09
(b) WC								
Total cholesterol	0.43	0.42	-0.03	0.28	1.30	0.49*	-0.01	0.24
Triglycerides	2.71	3.65	2.44	1.77	2.53	2.40	1.46	2.08
HDL-cholesterol	-0.24	-0.74**	-0.45	-0.49	-0.30	-0.72**	-0.42	-0.74*
LDL-cholesterol	0.08	0.42	-0.16	0.39*	1.10	0.69	0.12	0.56

**P<0.01, significant difference in the coefficient between BLSA and NILS-LSA subjects. *P<0.05, significant difference in the coefficient between BLSA and NILS-LSA subjects.

cohorts were aged between 40 and 80 y old. The populations were dichotomized at age 65 y into younger and older age groups for each sex. The BLSA subjects were Caucasian and the NILS-LSA subjects were Japanese. Details of the selection process and procedures of the baseline examination in the BLSA and the NILS-LSA have been published previously.^{17,18} The subjects in the NILS-LSA were randomly selected from resident registrations in cooperation with the local government. The subjects in the BLSA were self-recruited, community-dwelling volunteers. Table 1 shows some descriptive statistics of the population.

Subjects who were being treated for hyperlipidemia that could influence the level of the risk factors were excluded. Measurements from women who were pregnant at the visit or who had had a baby less than 1 y prior to the visit were also excluded from these analyses.

Anthropometry

Height and weight were measured after an overnight fast with subjects wearing a light-weight hospital gown and without shoes. As an index of obesity, BMI was calculated as weight (kg) divided by the square of the height (m).

WC was used as the index of the body fat distribution. The waist was defined as the minimal abdominal circumference between the lower edge of the rib cage and the iliac crests in the BLSA, and as the circumference at the middle point between the lower edge of the rib cage and the iliac crests in the NILS-LSA. The circumferences were obtained with a flexible, metal tape measure, while maintaining close contact with skin and without compressing the underlying tissues. Subjects were in a standing position and breathing normally. The same small group of trained personnel made these measurements for the entire study in both the BLSA and the NILS-LSA.

Plasma lipids

After an overnight fast, an antecubital venous blood sample was drawn. The concentrations of plasma triglycerides and total cholesterol were determined by enzymatic method (Abbott Laboratories ABA-200 ATC Biochromatic Analyzer, Irving, TX 75015 for BLSA; Hitachi 7145 Automatic Analyzer, Hitachi Co., Ltd Japan for NILS-LSA). HDL-cholesterol was determined by dextran sulfate-magnesium precipitation procedure¹⁹ in the BLSA and enzymatic methods in NILS-LSA (Hitachi 7145 Automatic Analyzer, Hitachi Co., Ltd Japan). LDL-cholesterol concentrations were estimated by the Friedewald formula²⁰ in both BLSA and NILS-LSA.

Cut-points for abnormalities of BMI, WC and risk factors

Cut-points for BMI and WC abnormalities were determined according to NHLBI⁵ and WHO⁶ guidelines. The BMI and WC categories were dichotomized at the lower recommended cut-points due to the small number of subjects in the obese and high WC groups. In addition, the designation of the risk factor cut-points that define an abnormal level was derived from the 1994 recommendations of the National Cholesterol Education Program.²¹

BMI	25 kg/m ²
WC	80 cm for women, 94 cm for men
Total cholesterol	240 mg/dl
Triglyceride	200 mg/dl
HDL-cholesterol	35 mg/dl
LDL-cholesterol	160 mg/dl

Sensitivity and specificity of WC and BMI cut-points for risk factor abnormality

To examine the applicability of cut-points for WC and BMI proposed by WHO and NHLBI, the sensitivities and specificities of WC or BMI cut-points were calculated in the BLSA

and NILS-LSA subjects. By using cut-points both for risk factors and for BMI (or WC), subjects were divided into four groups as shown.

	Normal	Abnormal
Normal BMI (or WC)	A	B
Abnormal BMI (or WC)	C	D

Sensitivity is defined as D over the sum of B and D, and specificity is defined as A over the sum of A and C according to the general definition of these terms.¹³

Statistical methods

All data were analyzed using Statistical Analysis System (SAS) version 6. Standard methods were used to compute means and standard errors of the mean. *P*-values below 0.05 were regarded as indicating statistical significance. ANOVA was used to test for the presence of statistically significant differences in BMI, WC and risk factors between the BLSA and the NILS subjects. A test of whether or not the regression coefficients (slopes) of BMI (or WC) and the risk factors are consistent between the two populations was performed using general linear models. χ^2 analysis was performed for analyzing effects of population in the percentage of subjects with abnormal levels of BMI, WC and risk factors. Adjusted mean lipid values were computed using ANCOVA by setting the covariates (BMI or WC) to the appropriate cut-point value.

Results

Differences in BMI, WC and risk factors between the BLSA and NILS-LSA subjects

Height, weight, WC and BMI were significantly higher in the BLSA than those in the NILS-LSA subjects, regardless of the sex/age group (Table 1). The subjects in the NILS-LSA had significantly greater total and LDL-cholesterol in younger and older men and women. HDL-cholesterol was significantly greater in the NILS-LSA than that in the BLSA subjects for all demographic groups (younger and older men and women). There was no significant difference in triglycerides between BLSA and NILS-LSA.

Associations between BMI, WC and risk factors

The only lipid/anthropometric association that was consistently different between the two populations was that of HDL-cholesterol with both BMI and WC. In all cases, the regressions were negative, but the impact of the body composition variables was greater in the NILS population; in five of the eight regressions, these differences were statistically significant. These population differences were especially striking in the younger men and women.

There was also a consistent very large effect of both BMI and WC on the triglyceride levels, but there were no

Table 3 Percentage of subjects with an abnormal level in BMI, WC and lipid risk factors

	Men				Women			
	Younger		Older		Younger		Older	
	BLSA	NILS	BLSA	NILS	BLSA	NILS	BLSA	NILS
BMI	63.9	23.2**	48.8	17.9**	35.6	17.6**	44.1	21.8**
WC	49.4	5.2**	45.2	7.8**	31.7	19.0**	45.5	28.7**
Total cholesterol	8.2	20.5**	7.7	17.5**	12.2	30.9**	15.2	43.5**
Triglyceride	17.1	13.6	10.1	8.6	4.4	5.2	6.2	7.9
HDL-cholesterol	31.0	2.7**	23.4	2.2**	5.0	0.3**	6.2	0.0**
LDL-cholesterol	11.7	15.8*	10.1	11.2	11.7	20.8**	15.2	29.6**

***P*<0.01, significant difference between BLSA and NILS by χ^2 analysis.
**P*<0.05, significant difference between BLSA and NILS by χ^2 analysis.

significant differences in the slope between the two populations. The only other statistically significant slope differences were WC-total cholesterol in the younger women and WC-LDL-cholesterol in the older men, but these population differences were inconsistent in the other three demographic (age/sex) groups (Table 2).

Percentage of subjects with an abnormal risk factor status

Table 3 shows the percentages of subjects with an abnormal level of BMI, WC or risk factors for each age/sex group in the BLSA and NILS-LSA subjects. The percentage of subjects with an abnormal BMI as well as an abnormal WC was significantly greater in the BLSA than that in the NILS-LSA subjects in younger and older men and women. Consistent with these population differences, there was a significantly higher percentage of subjects with an abnormal HDL-cholesterol level in the BLSA than that in the NILS subjects in all age/sex groups. The percentage of subjects with an abnormal level of total and of LDL-cholesterol was significantly higher in NILS-LSA subjects than those in BLSA subjects with the exception of LDL-cholesterol in younger men. Abnormal triglyceride levels were similar in the two populations.

Sensitivity of BMI and WC

Table 4a shows the sensitivity of BMI cut-points proposed by WHO and NHLBI (BMI < 25 kg/m²) for predicting risk factor abnormality. Sensitivities of BMI were very much lower in the NILS-LSA than in the BLSA. Also, sensitivities of WC (Table 4b) were lower in the NILS-LSA than those in the BLSA subjects.

Specificity of BMI and WC

Specificity of BMI and WC is shown in Table 5a and b. Higher specificities of BMI were found in younger and older men

Table 4 Sensitivity of BMI and WC cut-point for lipid risk factor abnormality

	Men				Women			
	Younger		Older		Younger		Older	
	BLSA	NILS	BLSA	NILS	BLSA	NILS	BLSA	NILS
(a) BMI								
Total cholesterol	69.2	27.1**	63.2	17.0**	45.5	17.7**	31.8	20.7
Triglycerides	88.9	37.4**	84.0	29.2**	62.5	50.0	77.8	23.5**
HDL-cholesterol	78.4	12.5**	70.7	16.7**	75.0	—	75.0	—
LDL-cholesterol	70.3	33.0**	44.0	20.0**	42.9	16.4**	40.9	21.9
Mean	76.7	27.5	65.5	20.7	56.5	28.0	56.4	22.0
(b) WC								
Total cholesterol	50.0	8.2**	47.4	10.6**	45.5	24.1*	36.4	26.1
Triglycerides	75.9	14.5**	72.0	20.8**	62.5	59.4	66.7	41.2
HDL-cholesterol	65.0	6.3**	67.2	16.7**	75.0	50.0	75.0	—
LDL-cholesterol	46.0	9.6**	40.0	10.0**	42.9	23.4 (0.06)	40.9	31.6
Mean	59.2	9.6	56.7	14.5	56.5	39.2	54.7	33.0

** $P < 0.01$, significant difference in sensitivity between BLSA and NILS-LSA subjects. * $P < 0.05$, significant difference in sensitivity between BLSA and NILS-LSA subjects. —: there was no NILS-LSA subject in some categories that were used for calculation of sensitivity.

Table 5 Specificity of BMI and WC cut-point for lipid risk factor abnormality

	Men				Women			
	Younger		Older		Younger		Older	
	BLSA	NILS	BLSA	NILS	BLSA	NILS	BLSA	NILS
(a) BMI								
Total cholesterol	36.6	77.8**	52.4	81.9**	65.8	82.4**	53.7	76.3**
Triglycerides	41.2	79.2**	55.2	83.4**	65.7	84.2**	58.1	77.7**
HDL-cholesterol	42.5	76.6**	57.9	82.1**	66.3	82.3**	57.3	77.6**
LDL-cholesterol	36.9	78.7**	50.7	82.4**	65.4	82.0**	55.3	77.4**
Mean	39.3	78.1	54.0	82.5	65.8	82.7	56.1	77.3
(b) WC								
Total cholesterol	50.7	95.6**	55.0	92.8**	70.3	82.8**	52.9	67.8*
Triglycerides	56.1	96.2**	57.9	93.2**	69.8	83.0**	55.9	71.5**
HDL-cholesterol	58.0	94.8**	61.6	92.4**	70.4	80.8**	55.8	70.5**
LDL-cholesterol	50.5	95.5**	54.3	92.1**	69.8	81.8**	53.7	71.2**
Mean	53.8	95.5	57.2	92.6	70.0	82.1	54.5	70.3

** $P < 0.01$, significant difference in sensitivity between BLSA and NILS-LSA subjects. * $P < 0.05$, significant difference in sensitivity between BLSA and NILS-LSA subjects.

and women in the NILS-LSA subjects compared to the BLSA subjects. Specificities of WC were also higher in the NILS-LSA subjects than those in the BLSA subjects, especially in men.

Mean values of risk factors at cut-points of BMI and WC
At the cut-point of BMI, mean values of triglycerides, total and LDL-cholesterol were significantly greater (worse) in the NILS-LSA than those in the BLSA subjects regardless of demographic groups (Table 6a). Mean values of HDL-cholesterol in the NILS-LSA subjects were significantly higher (better) than in the BLSA subjects.

At the cut-points of WC, mean values of triglycerides, total and LDL-cholesterol were significantly greater (worse) in the NILS-LSA than those in the BLSA subjects in all groups with the exception of triglycerides in younger women (Table 6b). The mean values of HDL-cholesterol in younger and older men and women were greater (better) in the NILS-LSA than in the BLSA subjects.

Discussion

Our previous reports on a Caucasian population^{15,16} showed that BMI and WC independently were significant predictors of coronary risk factors in the blood pressure, glucose-

Table 6 Mean lipid values

	Men				Women			
	Younger		Older		Younger		Older	
	BLSA	NILS	BLSA	NILS	BLSA	NILS	BLSA	NILS
<i>(a) BMI (25 kg/m²)</i>								
Total cholesterol (mg/dl)	189.6	215.6**	188.5	212.1**	189.2	223.5**	204.0	234.8**
Triglyceride (mg/dl)	124.8	151.3**	112.4	134.7**	100.0	110.4*	112.1	129.1*
HDL-cholesterol (mg/dl)	42.0	54.3**	44.4	53.2**	52.7	63.9**	55.6	60.4**
LDL-cholesterol (mg/dl)	122.5	131.4**	121.5	133.1**	116.4	138.0**	125.9	149.9**
<i>(b) WC (94 cm for men, 80 cm for women)</i>								
Total cholesterol (mg/dl)	191.8	217.9**	188.3	214.9**	193.3	225.1**	204.0	235.7**
Triglyceride (mg/dl)	136.4	177.8**	117.9	145.7**	108.0	113.7	112.9	127.7*
HDL-cholesterol (mg/dl)	41.0	49.0**	43.4	50.8**	51.8	63.1**	55.4	61.2**
LDL-cholesterol (mg/dl)	123.3	133.7**	121.2	135.7**	119.9	139.6**	126.0	149.7**

***P*<0.01, significant difference between the BLSA and NILS-LSA subjects. **P*<0.05, significant difference between the BLSA and NILS-LSA subjects.

insulin and plasma lipid domains. Furthermore, the gradation of BMI into normal, overweight and obese zones according to NHLBI⁵ and WHO⁶ recommendations was supported. Also, the sex-specific WHO cut-points for WC that provided three zones (NHLBI standards provided two zones) were also found to be applicable to the risk factors in the four age-sex categories in Caucasians. However, it was pointed out that the BMI and/or WC in Asian populations are much lower than those in Caucasians.²² Therefore, the important questions that remain are whether BMI and/or WC associated with coronary risk factors in relatively short and light-weight Asian populations as well as in Caucasians, and whether the BMI and WC cut-points proposed by NHLBI and WHO are applicable to Asian populations as a predictor of risk factor abnormality.

In the present study, in order to examine these questions, the relationship between BMI, WC and lipid risk factors (regression) and the applicability (sensitivity, specificity) of the BMI and WC cut-points for these risk factors were compared between the BLSA (Caucasian) and the NILS-LSA subjects (Japanese).

Height, weight and BMI were lower in the NILS-LSA than in the BLSA subjects regardless of the age/sex groups. There was a methodological difference in the measurement of WC between BLSA and NILS-LSA (see the Methods section). Although the minimal circumferences between the lower edge of the rib cage and the iliac crests were measured as WC in the BLSA subjects, the circumference at the middle point between the lower edge of the rib cage and the iliac crests was measured in the NILS-LSA subjects. Therefore, the measured WC in the NILS-LSA subjects may not be minimal as was the BLSA measurement. Despite this, WC in the NILS-LSA is still much smaller than that in the BLSA. In addition, WC of the NILS-LSA subjects was as highly correlated to BMI and to lipids as WC of the BLSA subjects (data not shown). Therefore, we compared the data of WC between the BLSA and the NILS-LSA directly.

It is well known that greater BMI or WC results in higher lipid risk factor levels. However, Japanese subjects (NILS-LSA) had higher levels in total cholesterol and LDL-cholesterol with smaller weight, BMI and WC, compared to Caucasians (BLSA). This result may indicate that associations between BMI, WC and lipid levels in Japanese differ from those in Caucasians. In the present study, the relationship (slope of regression) between BMI, WC and total cholesterol, triglycerides and LDL-cholesterol was similar, with the exception of the relationship between WC and LDL-cholesterol in older men, and between WC and total cholesterol in younger women. However, relationships between BMI, WC and HDL-cholesterol in most demographic groups were different in the two populations. Although relationships between BMI, WC and risk factors (triglycerides, total and LDL-cholesterol) were similar in the two populations, the mean values of total and LDL-cholesterol, and the percentages of subjects with an abnormal level of total and LDL-cholesterol in the NILS-LSA were higher than the levels of these variables in the BLSA subjects. In addition, the mean values at the BMI and WC cut-points in triglycerides, total and LDL-cholesterol were also higher (worse) in the NILS-LSA than those in the BLSA subjects. The mean plasma lipids are different between Japanese and Caucasians, although Japanese and Caucasians have similar relationships between BMI, WC and triglycerides, total and LDL-cholesterol. Because Japanese have higher levels of plasma lipids (excepting HDL-cholesterol) with a smaller BMI and WC, a normal BMI and WC defined by WHO and NHLBI does not indicate a lower risk in triglycerides, total and LDL-cholesterol in the Japanese population. The BMI and WC cutoff points seem to be too high as a predictor of risk abnormality to detect an abnormality in triglycerides, total and LDL-cholesterol for the Japanese population. Present data of very low sensitivities of the BMI and WC cut-points in the NILS-LSA support these results. For example, 83% of older men in the NILS-LSA who had an abnormal total cholesterol level had a normal

BMI, and 89.4% older men in the NILS-LSA with an abnormal total cholesterol had a normal WC.

Thus, the cut-points for BMI and WC proposed by WHO and NHLBI may not be ideal or even useful predictors of risk abnormality in the Japanese population. If these cutoff points of abnormality for BMI and WC are defined at lower levels, they may become a useful index for the Japanese population because BMI and WC still correlate with lipid risk factors in the Japanese subjects (data not shown). Further examinations are needed for the selection of the specific cutoff points for BMI and WC in the Japanese population.

In the present study, different relationships between BMI, WC and HDL-cholesterol in the two populations were found in most demographic groups. Although the magnitudes of the decrease in HDL-cholesterol with the increases of BMI and WC were greater in the NILS-LSA than in the BLSA subjects, mean values of HDL-cholesterol at the BMI and WC cut-points were still greater (better) in the NILS-LSA subjects. In the present study, there was a difference in the lipid methodologies between the two populations (see the Methods section). However, accuracy of both methods has been certified by the Center for Disease Control and Prevention (CDC). In addition, mean HDL-cholesterol levels were very similar in the BLSA and NHANES III,²³ and also in NILS-LSA and the National Nutrition Survey (Japan).²⁴ Therefore, we do not believe that these significant differences in HDL-cholesterol between BLSA and NILS-LSA subjects were caused by the methodological difference in HDL-cholesterol measurement. Thus, we compared HDL-cholesterol levels between BLSA and NILS-LSA subjects directly. Our result shows that the Japanese population with a normal BMI or WC has a remarkably higher level of HDL-cholesterol. NCEP noted that a high HDL-cholesterol level appears to be protective against CHD, and a level of ≥ 60 mg/dl can even be called 'a negative risk factor'. The mean values of HDL-cholesterol at the BMI cut-point in the Japanese population were 54, 53, 64 and 60 mg/dl in younger men, older men, younger women and older women, respectively. And the mean values of HDL-cholesterol at the WC cut-point were 49, 51, 63 and 61 mg/dl in these groups. Mean HDL-cholesterol levels in Japanese women were close to being 'a negative risk factor', even if their BMI levels exceeded 25 kg/m² or their WC levels exceeded 80 cm.

From our results, it may seem that the BMI and WC cut-points proposed by WHO and NHLBI are at levels too low for detecting HDL-cholesterol abnormality in the NILS-LSA subjects. However, it should be taken into account that most Japanese have relatively small BMI and WC (percentage of subjects with an abnormal BMI and WC in the four age/sex groups are only 5–29). If cut-points of BMI and WC are set at higher levels, almost no Japanese will have an 'abnormal' WC or BMI. In this Japanese population, lowering the cut-points for BMI or WC will provide very low specificity and rising the cut-points will yield very low sensitivity. The only other solution would be to redefine the level of HDL-cholesterol abnormality for the Japanese population.

Accumulation of intra-abdominal fat has been shown to be associated with risk factor abnormalities.^{3,25,26} Although we did not quantify intra-abdominal fat using CT scans in the present study, the distribution of intra-abdominal fat might well be different in the two populations. This may underlie the fact that the BMI and WC cut-points do not have the same level of applicability for predicting risk abnormality in Caucasian and Japanese populations. Therefore, further study is needed to better understand the relation between abdominal fat distribution (CT scan) and risk factor abnormality; it should however be noted that Takami *et al*²⁷ reported that BMI and WHR are better predictors of metabolic abnormalities than abdominal fat measured by CT.

Also, the reason for the differences in blood lipids between the Japanese and Caucasian populations was beyond the scope of the present study. Diet, exercise, body composition or genetics might be expected to contribute to these differences.^{28–32}

In the present study, there was an interesting finding that correlations between BMI, WC and lipids (total and LDL-cholesterol) in older people were weak or not significant regardless of populations, although BMI and WC were still highly correlated with triglycerides and HDL-cholesterol in older people. We have previously reported the effects of age on the relationship between body composition and risk factors in Caucasians^{15,16,33} and these effects are also seen in the Japanese population.

In conclusion, the Japanese subjects have higher total and LDL-cholesterol levels with smaller mean BMI and WC measurements compared to Caucasians. Although mean values in triglycerides, total and LDL-cholesterol at the recommended cut-points of BMI and WC (BMI = 25 kg/m², WC = 94 cm for men 80 cm for women) are obviously high (worse) in the Japanese, the mean HDL-cholesterol level at the cut-point is higher (ie, better) in Japanese. Thus, these cut-points for BMI and WC proposed by WHO and NHLBI may be too high for predicting an abnormality in triglycerides, total and LDL-cholesterol in Japanese. Predicting an abnormal HDL-cholesterol level using the cut-points for BMI and WC may not be as beneficial for the Japanese population as for Caucasians.

It must be noted that direct analyses of the predictive power of the cut-points for BMI and WC on the development of CHD itself in Japanese and Caucasian populations would be instructive, since the variables examined are risk factors for CHD in the present study. We must also note that direct examinations of the relationship among the characteristics in plasma lipids, nutrition intake, and morbidity and mortality of CHD in the Japanese and Caucasians are desirable in future studies.

Acknowledgements

We thank Research Fellowships for Japanese Biomedical and Behavioral Researchers at NIH funded by the Japan Society for the Promotion of Science (JSPS).

References

- 1 National Cholesterol Education Program. Third report of the National Cholesterol Education Program (NCEP) Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III) final report. *Circulation* 2002; 106: 3143-3421.
- 2 Fujioka S, Matsuzawa Y, Tokunaga K, Tarui S. Contribution of intra-abdominal fat accumulation to the impairment of glucose and lipid metabolism in human obesity. *Metabolism* 1987; 36: 54-59.
- 3 Bjorntorp P. Metabolic implications of body fat distribution. *Diabetes Care* 1991; 14: 1132-1143.
- 4 Bjorntorp P. Abdominal fat distribution and disease: an overview of epidemiological data. *Ann Med* 1992; 24: 15-18.
- 5 NHLBI Obesity Education Initiative Expert Panel. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. *Obes Res* 1998; 6: 51S-209S.
- 6 WHO Consultation on Obesity. Obesity. In: WHO (ed). *Preventing and managing the global epidemic*. Division of Noncommunicable Diseases, Programme of Nutrition, Family and Reproductive Health, World Health Organization: Geneva, Switzerland; 1998.
- 7 Molarius A, Seidell J. Selection of anthropometric indicators for classification of abdominal fatness—a critical review. *Int J Obes Relat Metab Disord* 1998; 22: 719-727.
- 8 Poullet M, Despres J, Lemieux S, Moorjani S, Bouchard C, Tremblay A, Nadeau A, Lupien P. Waist circumference and abdominal sagittal diameter: best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. *Am J Cardiol* 1994; 73: 460-467.
- 9 Turcato E, Bosello O, Harris T, Zoico E, Bissoli L, Fracassi E, Zamboni M. Waist circumference and abdominal sagittal diameter as surrogates of body fat distribution in the elderly: their relation with cardiovascular risk factors. *Int J Obes Relat Metab Disord* 2000; 24: 1005-1010.
- 10 Lean M, Han T, Seidell J. Impairment of health and quality of life in people with large waist circumference. *Lancet* 1998; 351: 853-856.
- 11 VanItallie T. Waist circumference: a useful index in clinical care and health promotion. *Nutr Rev* 1998; 56: 300-313.
- 12 Han T, McNeill G, Seidell J, Lean M. Predicting intra-abdominal fatness from anthropometric measures: the influence of stature. *Int J Obes Relat Metab Disord* 1997; 21: 587-593.
- 13 Molarius A, Seidell J, Sans S, Tuomilehto J, Knulasmaa K. Varying sensitivity of waist action levels to identify subjects with overweight or obesity in 19 populations of the WHO MONICA project. *J Clin Epidemiol* 1999; 52: 1213-1224.
- 14 Hu F, Wang B, Chen C, Jin Y, Yang J, Stampfer M, Xu X. Body mass index and cardiovascular risk factors in a rural Chinese population. *Am J Epidemiol* 2000; 151: 88-97.
- 15 Iwao S, Iwao N, Muller D, Elahi D, Shimokata H, Andres R. Effects of aging on the relationship between multiple risk factors and waist circumference. *J Am Geriatr Soc* 2000; 48: 788-794.
- 16 Iwao N, Iwao S, Muller D, Elahi D, Shimokata H, Andres R. A test of recently proposed BMI standards with respect to old age. *Aging: Clin Exp Res* 2000; 12: 461-469.
- 17 Shock N, Greulich R, Andres R, Arenberg D, Costa PJ, Lakatta E, Tobin J. *Normal human aging*. U.S. Department of Health and Human Services, U.S. Government Printing Office, National Institutes of Health Publication: Washington, DC, 1984. pp 84-2450.
- 18 Shimokata H, Ando F, Niino N. A new comprehensive study on aging—the National Institute for Longevity Sciences, Longitudinal Study of Aging (NILS-LSA). *J Epidemiol* 2000; 10: S1-S9.
- 19 Warnick G, Benderson J, Albers J. Dextran sulfate-Mg²⁺ precipitation procedure for quantification of high-density-lipoprotein cholesterol. *Clin Chem* 1982; 28: 1379-1388.
- 20 Friedewald W, Levy R, Fredrickson D. Estimation of the concentration of low-density-lipoprotein cholesterol in plasma without the use of the preparation ultracentrifuge. *Clin Chem* 1972; 18: 499-502.
- 21 National Cholesterol Education Program. Second report of the expert panel on detection, evaluation, and treatment of high blood cholesterol in adults. *Circulation* 1994; 89: 1333-1432.
- 22 Molarius A, Seidell J, Sans S, Tuomilehto J, Kuulasmaa K. Waist and hip circumferences, and waist-hip ratio in 19 populations of the WHO MONICA Project. *Int J Obes Relat Metab Disord* 1999; 23: 116-125.
- 23 Brown C, Higgins M, Donato K, Rohde F, Garrison R, Obarzanek E, Ernst N, Horan M. Body mass index and the prevalence of hypertension and dyslipidemia. *Obes Res* 2000; 8: 605-619.
- 24 The Ministry of Health and Welfare. *National nutrition in 1998*. Daiichi Public: Tokyo, 1999 (in Japanese).
- 25 Nakamura T, Tokunaga K, Shimomura I, Nishida M, Yoshida S, Kotani K, Islam A, Keno Y, Kobatake T, Nagay Y, Fujioka S, Tarui S, Matsuzawa Y. Contribution of visceral fat accumulation to the development of coronary artery disease in non-obese men. *Atherosclerosis* 1994; 107: 239-246.
- 26 Despres J. Abdominal obesity as important component of insulin-resistance syndrome. *Nutrition* 1993; 9: 452-459.
- 27 Takami R, Takeda N, Hayashi M, Sasaki A, Kawachi S, Yoshino K, Takami K, Nakashima K, Akai A, Yamakita N, Yasuda K. Body fatness and fat distribution as predictors of metabolic abnormalities and early carotid atherosclerosis. *Diabetes Care* 2001; 24: 1248-1252.
- 28 Kumagi S, Tanaka H, Kitajima H, Kono S, Ogawa K, Yamauchi M, Morita N, Inoue M, Shindo M. Relationships of lipid and glucose metabolism with waist-hip ratio and physical fitness in obese men. *Int J Obes Relat Metab Disord* 1993; 17: 437-440.
- 29 Rodriguez B, Sharp D, Abbott R, Burchfiel C, Masaki K, Chyou P, Huang B, Yano K, Curb J. Fish intake may limit the increase in risk of coronary heart disease morbidity and mortality among heavy smokers. The Honolulu Heart Program. *Circulation* 1996; 94: 952-956.
- 30 Nakamura K, Shimai S, Kikuchi S, Tanaka M. Correlation between a liking for fat-rich foods and body fatness in adult Japanese: a gender difference. *Appetite* 2001; 36: 1-7.
- 31 Pott J, Simmons D. Sex and ethnic group differences in fat distribution in young United Kingdom South Asians and Europeans. *J Clin Epidemiol* 1994; 47: 837-841.
- 32 The Research Group on Serum Lipid Survey. Analysis of serum lipid levels in Japanese men and women according to body mass index. Increase in risk of atherosclerosis in postmenopausal women. *Atherosclerosis* 1999; 143: 55-73.
- 33 Iwao S, Iwao N, Muller D, Elahi D, Shimokata H, Andres R. Does waist circumference add to the predictive power of the body mass index for coronary risk? *Obes Res* 2001; 9: 658-695.

The Impact of Health Problems on Depression and Activities in Middle-Aged and Older Adults: Age and Social Interactions as Moderators

Yasuyuki Fukukawa,¹ Chiori Nakashima,¹ Satomi Tsuboi,¹ Naoakira Niino,¹
Fujiko Ando,¹ Shotaro Kosugi,² and Hiroshi Shimokata¹

¹Department of Epidemiology, National Institute for Longevity Sciences, Obu Aichi, Japan.

²School of Letters, Arts, and Sciences, Waseda University, Tokyo, Japan.

In this study, we compared the impact of health problems (HPs) on everyday activities and depressive symptoms between middle-aged and older adults. We also examined what type and source of social interactions moderate the noxious effects of HPs. Longitudinal analyses of data with 1,802 Japanese community-dwelling adults indicated that HPs were significantly related to (a) an increase in depressive symptoms among middle-aged adults and (b) a decline in everyday activities among older adults. The former was buffered by emotional family support, whereas the latter (b) was buffered by instrumental family support and, surprisingly, by negative interactions with family. In contrast, social interactions with other friends and acquaintances did not show any moderating effect.

MANY researchers have reported the relationship between health problems (HPs), such as disease or injury, and lower activities among older adults (Dargent-Molina, Hays, & Breart, 1996; Guccione et al., 1994; Mor et al., 1989). Yet, as aging per se is a normal phenomenon that leads to a limitation of activities (Dickerson & Fisher, 1993), the pathologic effects of disease on disability are sometimes confused with the results of normal aging (Pearson, 2000). However, the decline of physical function that is due to normal aging is relatively small and should not interfere with the ability of older adults to live an autonomous and high-quality life (Ferrucci et al., 2002). In fact, a noticeable percentage of persons who reach extreme longevity are still able to perform activities of daily living (ADLs; Franceschi et al., 2000). Consequently, we could expect older adults to live primarily free of disability if they do not suffer from HPs.

This assumption does not mean that the role age plays in the disablement process is negligible. For instance, Pohjasvaara, Erkinjuntti, Vataja, and Kaste (1997) examined whether ischemic strokes have an equivalent negative impact on the daily activities of younger versus older patients. They found that ADL functions deteriorated more significantly among the older patients than the younger patients after the stroke. The results suggest that older adults are more likely than younger adults to decrease their activities because of HPs. However, although most prior studies have statistically controlled for age when estimating the impact of HPs, the question of how age and HPs interact and influence everyday activities has rarely been addressed.

Psychosocial stress theory also considers HPs to be influential life events leading to a decline in psychological well-being such as depression (Holms & Rahe, 1967; Murrell, Norris, & Hutchins, 1984). Although little is known about the role of age in the stress process (Folkman, Lazarus, Pimley, & Novacek, 1987; Martin, Gruendahl, & Martin, 2001), theoretical and empirical findings suggest that HPs exert less impact

on well-being in older adults than in younger adults. For example, although HPs affect middle-aged adults' well-being by producing social role strain regarding work or parenting (Karasz & Ouellette, 1995), the number of such social roles decreases as people age (Aldwin, Sutton, Chiara, & Spiro, 1996). As Krause (1994) argued, stressors will be more strongly associated with well-being when they arise in important social roles. Thus, the fewer social roles of older adults may make dealing with HPs psychologically less challenging for them than it is for middle-aged adults. Indeed, Aldwin and colleagues (1996) found that older adults were not more likely to appraise HPs in terms of loss or threat, despite the fact that they were actually dealing with HPs.

Furthermore, according to the normative life events theory, events that are not expected for a particular period in life will have a greater impact on the well-being of individuals during that particular period than on individuals during other periods (Pearlin & Lieberman, 1979). Following the logic, the higher morbidity in older adults (Bowling & Grundy, 1997; Kriegsman, van Eijk, Penninx, Deeg, & Boeke, 1997) may attenuate the psychological impact of HPs on the age group, compared with younger groups. In fact, Hurwicz, Durham, Boyd-Davis, Gatz, and Bengtson (1992) found that, although over 30% of the older adults tended to report "ill health" as the event that had had the greatest impact on them, compared with 21% of the middle-aged and 16% of the younger adults, a significant association between the ill health and depressive symptoms was revealed only in the younger adults.

Taken as a whole, the primary purpose of the present study was to examine age differences in the impact of HPs on everyday activities and depressive symptoms among middle-aged and older adults. However, disability and depression are not merely dependent on a person's HPs but are also subject to psychosocial factors. This study also focused on one such factor, social interactions, and its role in the relationships between HPs and outcomes.

An extensive body of research has demonstrated that positive social interactions, such as social support, contribute to the improvement of a person's psychological and physical well-being (Cohen & Willis, 1985). However, there is a growing consensus that social support is a multidimensional concept, and the type and source of support should be considered when its effect is estimated. Some studies have found that social support from friends is more effective for psychological well-being than that from family (Dean, Kolody, & Wood, 1990; Lee & Ishii-Kuntz, 1987). Others assert that family support is more effective than support from friends (Chi & Chou, 2001; Yanagisawa et al., 2002). The findings from Felton and Berry (1992) are more complicated: Emotional support was more beneficial for psychological well-being when provided by nonkin, whereas instrumental support was more beneficial when provided by kin. Regarding physical health outcomes, some researchers have found support to be beneficial in improving physical function and activities (Duke, Leventhal, Brownlee, & Leventhal, 2002; Seeman et al., 1995), whereas Mendes de Leon, Gold, Glass, Kaplan, and George (2001) have indicated that instrumental support was associated with an increase of disability risk among older adults.

Although the previous findings are not altogether consistent, a careful review of the literature by Crohan and Antonucci (1989) revealed that, whereas friends tend to play their largest role in the arenas of socialization and the provision of day-to-day companionship, support provided by family becomes important when long-term sick care or daily living help is needed. This suggests that support from family would be more beneficial than support from other relationships, at least for people suffering from HPs.

The mixed findings on the effect of support by type may be partly derived from the analytical designs. Most previous studies have focused, on one hand, on the *direct* effect of social support: the effect of improving well-being regardless of whether stressors, such as HPs, are present or not. The *buffering* effect, on the other hand, posits that support is related to well-being for persons under stress, as it protects persons from the potentially pathologic influence of stress events (Cohen & Willis, 1985). Indeed, Penninx and associates (Kriegsman et al., 1997; Penninx et al., 1997, 1998) have reported that instrumental support buffered the incidence of mobility difficulties for older people with chronic lung disease, whereas emotional support buffered the increase of depressive symptoms for older people with cardiac disease and arthritis. As Cohen and Willis (1985) argued, these findings suggest that there must be a reasonable match between the type of available support and the consequences of HPs in order for buffering to occur.

The present study also investigated how negative interactions, that is, interference and criticism dimensions of social interactions, may moderate the relationship between HPs and outcomes. Although negative interactions are theoretically assumed to exacerbate the noxious association of stressors with well-being (Shinn, Lehman, & Wong, 1984), this negative buffering, or stress-amplification hypothesis (Okun, Melichar, & Hill, 1990), has not been well established. For example, Finch, Okun, Barrera, Zautra, and Reich (1989) failed to confirm the effects of negative interactions to aggravate the associations between HPs and psychological distress in older adults. However, because their study had a cross-sectional

analytical design using a relatively small sample ($N = 246$) and lacked attention to disability outcome, the results still have opened the door for further examination of the moderating effect of negative interactions on HPs and their consequences.

Hypotheses and Analytic Strategy

On the basis of the aforementioned grounds, this study examined the moderating effects of age and social interactions in the impact of HPs on everyday activities and depressive symptoms, using longitudinal data collected from Japanese middle-aged and older adults. In Japan, as in other Western countries, the population is aging rapidly, and estimating the influence of HPs on older adults' lives has become an important issue (Health and Welfare Statistics Association, 2002). This study would contribute to the available data for developing a cross-cultural view on the experience of HPs in later life.

The analysis was twofold. First, we addressed the question of whether the impact of HPs differed between middle-aged and older adults. We expected that the impact of HPs in decreasing everyday activities would be greater in older adults, compared with middle-aged adults. In contrast, compared with older adults, middle-aged adults would be more likely to increase their depressive symptoms because of HPs. To test the hypotheses, we conducted repeated measures analyses with a mixed procedure (SAS Institute Inc., 1996) to examine pattern differences of changes in activities and depression scores according to age group and HP experience.

Second, we examined what type and source of social interactions would moderate the negative consequences of HPs. To avoid problems associated with multicollinearity, we examined in separate analyses whether the effect of HPs is moderated by the level (high vs. low) of three types of social interactions (emotional support, instrumental support, and negative interactions), provided by the two interaction sources of family and others (friends or acquaintances). We expected that social interactions with family would more effectively buffer the negative consequences of HPs than interactions with others. Furthermore, we expected that emotional support would buffer the increase of depressive symptoms, whereas instrumental support would buffer the decline of everyday activities caused by HPs. We also predicted that negative interactions would amplify the noxious effects of HPs on activities and depressive symptoms.

METHODS

Participants

The data for this study are taken from the baseline (Wave 1, from November 1997 to April 2000) and the follow-up (Wave 2, from April 2000 to May 2002) surveys of the National Institute for Longevity Sciences—Longitudinal Study of Aging (NLS—LSA). The average follow-up interval was 2.1 years. The NLS—LSA participants were Japanese community-dwelling adults between 40 and 79 years of age, randomly recruited from areas around the institute. Details of the NLS—LSA have been described elsewhere (Shimokata, Ando, & Niino, 2000). The study sample consisted of 1,802 men and women who had completed examinations at both Wave 1 and Wave 2. The average age for the entire sample was 58.3 ± 10.6 years. For

the analyses in this article, we divided the sample into two groups according to their age at Wave 1 (older adults: 60–79 years old, $n = 833$; middle-aged adults: 40–59 years old, $n = 969$).

Measures

Health problems.—We identified HPs by using a life event checklist at Wave 2. The trained interviewer presented participants with the checklist of 43 items and asked them to report the occurrence of each event between Wave 1 and Wave 2 periods. Participants who reported the occurrence of “major injury or disease,” one of the listed items, were classified into the HP-present (HPP) group, whereas all others were classified into the HP-absent (HPA) group. Although information about the type of HP was collected from the HPP participants, no type-specific analyses were conducted in the study.

Everyday activities.—We measured everyday activities both at Wave 1 and Wave 2 by using the Tokyo Metropolitan Institute of Gerontology Index of Competence (TMIG-IC; Koyano, Shibata, Nakazato, Haga, & Suyama, 1987). This scale includes 13 items conceptually grouped into three categories: instrumental ADLs (using transportation, shopping, preparing meals, paying bills, and making bank deposits and withdrawals), intellectual or cultural activities (filling out forms, reading papers, reading books and magazines, and watching television), and social engagement activities (visiting a friend’s home, helping others, going to see someone in the hospital, and talking to others). Acceptable reliability ($\alpha = .86$) and validity (association with mortality) of the index have been reported (Koyano, Shibata, Nakazato, Haga, & Suyama, 1991). Participants indicate whether or not they can perform each activity with a “yes” (= 1) or a “no” (= 0), and then the responses are totaled and used as an index of everyday activities (a higher score indicates better performance). In the study, the internal reliability of the scale was .72.

Depressive symptoms.—We measured depressive symptoms at both Wave 1 and Wave 2 by using a Japanese version of the Center for Epidemiologic Studies Depression (CES-D) scale (Radloff, 1977; Shima, Shikano, Kitamura, & Asai, 1985). Participants indicated how often during the previous week they had experienced any of the 20 depressive symptoms described in the scale. Each item was rated on a 4-point scale ranging from 0 (rarely or none of the time) to 3 (most or all the time). Four positively worded items were reverse scored. The points were added together so that a higher score represented a higher level of depressive symptoms. For our sample, the internal reliability of the scale was .91.

Social interactions.—We measured social interactions at Wave 2, using a scale developed by Noguchi (1991). This scale comprises three subscales: emotional support, instrumental support, and negative interactions, each of which consists of four items. Participants indicated whether there was someone to provide specific interactions such as “listening to them when they have worries or problems” (emotional support), “caring for them when they are confined to bed for several weeks”

(instrumental support), or “being critical of them” (negative interactions). We rated the responses on a 4-point scale, ranging from 1 (none) to 4 (many), with higher scores indicating more social interactions. Each item was duplicated to refer to the two interaction sources of family and others (friends or acquaintances). Summing up item scores by type and source, we generated six social interaction measures (emotional support from family, emotional support from others, instrumental support from family, instrumental support from others, negative interactions with family, and negative interactions with others). The internal reliability of the measures ranged from .71 to .86. For analyses, participants with scores 1 *SD* below the age-specific mean of emotional support from family were classified into the low-emotional family support group and others were classified into the high-emotional family support group. We conducted the same operations for the other social interaction measures. However, for negative interaction measures, participants with scores 1 *SD* above the mean were classified into the high-negative interactions group and others were classified into the low-negative interactions group.

Control variables.—We used gender and annual family income at Wave 1 as control variables because of their significant associations with the outcome variables (data not shown). We measured annual family income by using a scale with 11 options (from 1 = income less than ¥1,500,000 to 11 = income greater than ¥20,000,000). In addition, in the analyses we considered the following information regarding participants’ health status at Wave 1. Subjective health was assessed by a single self-reported question (“How would you rate your health?”), with five options (from 1 = very bad to 5 = very good). The presence and history of seven diseases (stroke, hypertension, cardiovascular disease, diabetes, bronchitis, arthritis, and cancer) were totaled as an index of participants’ existing chronic conditions.

RESULTS

Sample Characteristics

Table 1 presents the results of bivariate tests of age differences in the study variables. Male participants represented just over half of the individuals in both age groups. The older group had a significantly lower income than the middle-aged group. The older group also felt less healthy and was in worse condition than the middle-aged group, although the differences were not very large. Those who reported experiencing HPs between the surveys made up 13.9% of the older group and 11.6% of the middle-aged group, which was not significantly different. Regarding the HP subtypes, participants who reported disease outstripped those who reported injury and those who reported both disease and injury or malfunctions from an unknown cause such as indefinite complaint (categorized as “other” in Table 1) in both age groups. The outcome variables did not differ between the age groups, except for the TMIG-IC score at Wave 2, which was slightly lower in the older group. Regarding social interactions, older adults reported fewer negative interactions with both family and others. Furthermore, older adults reported slightly less emotional and instrumental support from others than did middle-aged adults.

Table 1. Descriptive Information for Study Variables and Age Differences

Variable	Older	Middle-Aged
Gender, male (%)	53.4	51.0
Income (<i>M</i>)	5.1 (2.4)	7.3 (2.0)***
Subjective health (<i>M</i>)	3.1 (0.6)	3.3 (0.7)***
Chronic conditions (<i>M</i>)	0.8 (0.8)	0.3 (0.5)***
HPP (%)	13.9	11.6
Disease	7.7	5.6
Injury	5.0	4.6
Other	1.2	1.3
TMIG-IC (<i>M</i>)		
Wave 1	12.3 (1.1)	12.4 (1.0)
Wave 2	12.2 (1.2)	12.3 (1.1)*
CES-D (<i>M</i>)		
Wave 1	7.0 (6.7)	6.9 (6.2)
Wave 2	7.4 (6.7)	7.3 (6.7)
Social interactions, source-type (<i>M</i>)		
Family-emotional	12.5 (2.3)	12.4 (2.1)
Family-instrumental	12.1 (2.2)	12.0 (2.0)
Family-negative	8.6 (2.4)	9.6 (2.2)***
Other-emotional	11.2 (2.7)	11.6 (2.5)***
Other-instrumental	9.0 (2.8)	9.3 (2.7)*
Other-negative	7.0 (2.3)	8.3 (2.3)***

Notes: Entries in parentheses are standard deviations. HPP = health problem present; TMIG-IC = Tokyo Metropolitan Institute of Gerontology Index of Competence; CES-D = Center for Epidemiologic Studies Depression (scale).

* $p < .05$; *** $p < .001$.

Age and the Impact of HPs

We conducted repeated measures analyses to examine pattern differences of changes in the TMIG-IC and CES-D scores according to age (middle-aged vs. older) and HP (HPP vs. HPA) groups. The design was a split plot, involving the main effects of age, HP, and time (Wave 1 vs. Wave 2), and their two-way and three-way interaction terms. Gender, income, subjective health, chronic conditions, and the baseline outcome (TMIG-IC or CES-D) were used as covariates.

The analysis indicated that, among between-subject factors, gender (female), $F(1, 1741) = 10.88, p < .001$, subjective health, $F(1, 1741) = 9.45, p < .01$, and the baseline outcome, $F(1, 1741) = 5751.82, p < .001$, had significant positive associations with the TMIG-IC score. No significant main effects were found for age, HP, and Age \times HP interaction. Regarding within-subject factors, time, $F(1, 1743) = 13.32, p < .001$, and Age \times Time, $F(1, 1743) = 6.73, p < .01$, were revealed to be significant, suggesting participants decreased their activities during the surveys, regardless of age or HP effects, and the decline was more rapid in the older group. However, a significant three-way interaction among Age \times HP \times Time, $F(1, 1743) = 5.34, p < .05$, was also revealed, suggesting that the change in the TMIG-IC score depended on the participant's age and HP experience.

We performed the same analysis with the CES-D outcome. The results indicated that, among between-subject factors, chronic conditions, $F(1, 1737) = 8.02, p < .01$, and the baseline outcome, $F(1, 1737) = 5271.15, p < .001$, had significant positive associations with the CES-D score, whereas income,




Figure 1. Changes in everyday activities (Tokyo Metropolitan Institute of Gerontology Index of Competence, or TMIG-IC) over time relating to age and health problem (HP) experience. HPA = HP absent; HPP = HP present.

$F(1, 1737) = 4.23, p < .05$, and subjective health, $F(1, 1737) = 18.44, p < .001$, had significant inverse associations with the CES-D score. Furthermore, significant main effects with regard to age, $F(1, 1737) = 12.53, p < .001$, and HP, $F(1, 1737) = 3.93, p < .05$, emerged, suggesting that those in the middle-aged group and those in the HPP group had more depressive symptoms compared with those in the older group and those in the HPA group, respectively. However, these effects are qualified by a between-subject factor, Age \times HP interaction, $F(1, 1737) = 6.18, p < .05$, and within-subject factors, time, $F(1, 1701) = 8.90, p < .01$, and Age \times HP \times Time, $F(1, 1701) = 4.39, p < .05$. These results suggest that changes in depressive symptoms relating to HPs were not equivalent between the age groups.

Figure 1 and 2 illustrate the nature of the interactions by displaying the changes in the outcome scores over time by age and HP groups. We carried out further tests to examine simple main effects of time on each group by using the Tukey-Kramer adjusting method. After we controlled for gender, income, subjective health, chronic conditions, and the baseline outcome (TMIG-IC or CES-D), results revealed that the TMIG-IC score in older adults who had experienced HPs significantly decreased, $t(1743) = 3.64, p < .01$ (Figure 1), whereas the CES-D score in middle-aged adults who had experienced HPs significantly increased, $t(1701) = -3.14, p < .05$ (Figure 2), between the surveys. The HPA groups, in contrast, did not show any significant changes in their outcome scores, indicating stability of activities or mental health status over time irrespective of age group.

Moderating Effect of Social Interactions

Our second research goal was to determine the type and source of social interactions that moderate the noxious effects of HPs. On the basis of the results mentioned herein, we performed the analyses only on HPP participants. That is, for older adults with HPs, we carried out repeated measures analyses to determine whether the level (high vs. low) of each




Figure 2. Changes in depressive symptoms (Center for Epidemiologic Studies Depression, or CES-D, scale) over time relating to age and health problem (HP) experience. HPA = HP absent; HPP = HP present.




Figure 3. Changes in everyday activities (Tokyo Metropolitan Institute of Gerontology Index of Competence, or TMIG-IC) over time among older adults with health problems by the level of instrumental family support.

social interaction moderates (buffered or amplified) the change in TMIG-IC score over time. Similarly, for middle-aged adults with HPs, we examined whether the level of each social interaction moderates the change in CES-D score over time. The analyses tested the statistical significance of the main effect of each social interaction level and its interaction with time, controlling for gender, income, subjective health, chronic conditions, and the baseline outcome (TMIG-IC or CES-D).

The results were as follows. For older adults with HPs, significant main effects for levels of instrumental family support, $F(1, 104) = 11.09, p < .01$, emotional support from others, $F(1, 105) = 4.77, p < .05$, and negative interactions with family, $F(1, 104) = 4.45, p < .05$, emerged, with higher levels of these interactions increasing the TMIG-IC score. For middle-aged adults with HPs, significant main effects for levels of emotional family support, $F(1, 101) = 6.67, p < .05$, and instrumental family support, $F(1, 101) = 10.10, p < .01$, emerged, with higher levels of these interactions decreasing the CES-D score. No significant main effects were found for levels of other social interaction measures on both age groups, regardless of the outcome variable.

In addition to the main effects, three significant interactions emerged: Instrumental family support \times Time, $F(1, 109) = 7.65, p < .01$, and Negative interactions with family \times Time, $F(1, 109) = 5.55, p < .05$, in older adults, and Emotional family support \times Time in middle-aged adults, $F(1, 104) = 3.97, p < .05$. The effects of these interactions are illustrated in Figures 3, 4, and 5. Among older adults with HPs, there was a significant decrease in the TMIG-IC score over time only when instrumental family support level was low, $t(109) = 3.89, p < .01$ (Figure 3). Surprisingly, there was also a significant decrease of the TMIG-IC score over time when the level of negative interactions with family was low, $t(109) = 3.95, p < .001$ (Figure 4). Among middle-aged adults with HPs, there was a significant increase of the CES-D score over time only when the emotional family support level was low, $t(104) = -2.72, p < .05$ (Figure 5).

DISCUSSION

Age and the Impact of HPs

The three-way interaction of Age \times HP \times Time and the post hoc analyses indicated that the decline of everyday activities over time was statistically significant only for older adults who had experienced HPs between the surveys. The findings support our prediction that the impact of HPs in decreasing everyday activities would be greater in older adults, compared with middle-aged adults. As we noted earlier, aging per se is not so influential in limiting everyday activities of older adults (Dickerson & Fisher, 1993). However, although it is not inherently impaired, the reserve capacity to compensate for stress, metabolic derangement, and drug metabolism is increasingly limited with advancing age (Oskvig, 1999). Perhaps the age-related decline of reserve capacity in the human body allows HPs to have a greater impact on older adults, thus leading to limitations in the performance of everyday activities.

Another explanation for the difference is that the nature of HPs varies across age groups. Valderrama-Gama, Damian, Ruigomez, and Martin-Moreno (2002) argued that, whereas acute disease represents the main cause of HPs in young people, chronic conditions are more prevalent in older persons. Focusing on the type and, possibly, the severity of the HP would be the next step to developing and refining our findings.

In contrast, as we also expected, the increase of depressive symptoms over time was statistically significant only for middle-aged adults who had experienced HPs between the surveys. This is consistent with other studies (Aldwin et al., 1996; Hurwicz et al., 1992), including a Japanese one (Matsunaka, 2002), suggesting older adults are less emotionally reactive to HPs than younger adults. Thus, our results confirm the cross-cultural consistency in the psychological resilience of older adults with HPs. As we discussed earlier, because HPs threaten the salient social roles of middle-aged adults as parents or income earners (Karasz & Ouellette, 1995), they could become more critical for them and could lead to severe




Figure 4. Changes in everyday activities (Tokyo Metropolitan Institute of Gerontology Index of Competence, or TMIG-IC) over time among older adults with health problems by the level of negative interactions with family.

psychological distress. In contrast, as disengagement theory (Cumming & Henry, 1961) implies, aging is a process whereby people gradually withdraw from society and are relieved of annoying obligations and social roles. The decreased demands from society may allow older adults to feel at ease and less stressed even when they are suffering from HPs.

Our findings would also be compatible with the normative life events theory, which posits that events that are expected for a particular period in the life span will have less impact on well-being (Pearlin & Lieberman, 1979). We found that the older group felt less healthy and was in worse condition even at the study entry, compared with the middle-aged group (see Table 1). As Norris and Murrell (1988) pointed out, the prior experience of a stressor provides "inoculation" against strong emotional reaction to a similar stressor experienced in later days. Thus, relatively worse health status at baseline in older adults might make provision against newly experienced HPs and attenuate the influence on psychological well-being.

Moderating Effect of Social Interactions

We found that instrumental family support buffered the decline of everyday activities in older adults, whereas emotional family support buffered the increase of depressive symptoms in middle-aged adults. In contrast, social support provided by other relationships (friends or acquaintances) did not show any buffering effect in either age group. One of the implications of the findings is that, as we expected, social support is more effective at preventing negative consequences of HPs when it comes from family rather than from other relationships. According to a hierarchical-compensatory model of social support (Cantor, 1979), people select supportive ties from a hierarchy of relationships, with family members always selected before nonfamily members. Additionally, as in other Asian countries such as China (Chi & Chou, 2001) and Korea (Mo, 1999), Japanese society maintains cultural values and traditional norms of caregiving in which the family provides




Figure 5. Changes in depressive symptoms (Center for Epidemiologic Studies Depression, or CES-D, scale) over time among middle-aged adults with health problems by the level of emotional family support.

care for its older members (Asahara, Momose, & Murashima, 2002). These values and norms may also strengthen the central roles of family as the support provider, especially for people suffering from HPs. However, because we aggregated friends and acquaintances into one support source category of "other relationships," the effects of key network nonkin members might be masked (Barrera, Chassin, & Rogosch, 1993). Distinguishing "other relationships" across levels of closeness would then be needed to further assess the beneficial effect of nonkin support.

Regarding the type of social support, the results also supported our predictions: Emotional support buffered the increase of depressive symptoms, whereas instrumental support buffered the decline of everyday activities. The results are in line with those of other cross-sectional findings (Kriegsman et al., 1997; Penninx et al., 1997, 1998) and the theoretical suggestion that posits that support closely linked to the specific need elicited by a stressor will exert the buffering effect (Cohen & Willis, 1985). For example, emotional support might reduce depressive symptoms caused by HPs because this type of support enhances the person's self-esteem, whereas instrumental support attenuates limitations of activities because it helps the person seek medical care for the HPs. However, contrary to our findings, Mendes de Leon and colleagues (2001) found that older adults with more instrumental support showed a decline of ADLs over time. They examined how baseline instrumental support predicts lower ADL functions at the follow-up, with no regard to the presence or absence of HPs in the study population, whereas we examined whether older adults with HPs maintain their activity level by instrumental support measured at the follow-up. The inconsistency between the studies would then be due to the differences in analytical designs.

An unexpected finding in the study is that negative interactions with family did not amplify but rather buffered the noxious effect of HPs on older adults' everyday activities. A possible explanation is that social network members may, by being critical and demanding of an older person, induce him or

her to engage in beneficial health behaviors (Lewis & Rook, 1999). For example, a wife may be critical of her diabetic husband for not following an appropriate diet (Krause, 2001). Indeed, Krause, Goldenhar, Liang, Jay, and Maeda (1993) found that negative interactions predicted more frequent exercise among older Japanese adults. Perhaps negative interactions would potentially contribute to discouraging health-compromising behavior and promoting health-enhancing behavior.

Future Directions

Some limitations should be kept in mind when our findings are interpreted. First, Lawrence and Jette (1996) described the multistep pathway in the disablement process: Presence of disease leads to anatomic and structural abnormalities, which in turn lead to restrictions in basic physical and mental actions, which then lead to difficulty performing ADLs. Analyses using a 2-point assessment method, as in our study, are too simple to examine such a multistep disablement process. Second, we did not pursue gender differences in the effects of HPs to avoid the complicated interpretation of a four-way interaction (HP \times Age \times Gender \times Time). However, compared with men, women tend to report poorer physical health associated with chronic illness (O'Neill & Morrow, 2001) and more depressive symptoms (Nolen-Hoeksema, Larson, & Grayson, 1999). An elaborate analysis concerning these issues would modify and extend our findings.

Despite these limitations, the findings garnered from the present study add several specific points to the literature examining relationships between HPs and their negative consequences. Whereas most studies have restricted their sample to older adults or analyzed data by using age just as one of the confounders, this study made clear that the impact of HPs differs across age groups. Moreover, we highlight the importance of taking into account the type and source of social interactions when we are investigating how they moderate the impact of HPs. Although the results suggest that the relationships among HPs, everyday activities, and mental health are more complex than generally conceived, that is what makes it so challenging for researchers and professionals who are interested in stress and the development process in adulthood.

ACKNOWLEDGMENTS

This study was supported by a Grant-in-Aid (H14-choju-004) for Comprehensive Research on Aging and Health from the Ministry of Health, Labour and Welfare of Japan.

Address correspondence to Yasuyuki Fukukawa, Department of Epidemiology, National Institute for Longevity Sciences, 36-3 Gengo Morioka-cho, Obu Aichi, 474-8522, Japan. E-mail: fukukawa@nils.go.jp

REFERENCES

- Aldwin, C. M., Sutton, K. J., Chiara, G., & Spiro, A. (1996). Age differences in stress, coping, and appraisal: Findings from the Normative Aging Study. *Journal of Gerontology: Psychological Sciences*, *51B*, P179-P188.
- Asahara, K., Momose, Y., & Murashima, S. (2002). Family caregiving of the elderly and long-term care insurance in rural Japan. *International Journal of Nursing Practice*, *8*, 167-172.
- Barrera, M., Jr., Chassin, L., & Rogosch, F. (1993). Effects of social support and conflict on adolescent children of alcoholic and non-alcoholic fathers. *Journal of Personality and Social Psychology*, *64*, 602-612.
- Bowling, A., & Grundy, E. (1997). Activities of daily living: Changes in functional ability in three samples of elderly and very elderly people. *Age and Ageing*, *26*, 107-114.
- Cantor, M. H. (1979). Neighbors and friends: An overlooked resource in the informal support system. *Research on Aging*, *1*, 434-463.
- Chi, I., & Chou, K. L. (2001). Social support and depression among elderly Chinese people in Hong Kong. *International Journal of Aging and Human Development*, *52*, 231-252.
- Cohen, S., & Willis, T. A. (1985). Stress, social support, and the buffering hypothesis. *Psychological Bulletin*, *98*, 310-357.
- Crohan, S., & Antonucci, T. (1989). Friends as a source of social support in old age. In R. G. Adams & R. Blieszner (Eds.), *Older adult friendship: Structure and process* (pp. 129-146). Newbury Park, CA: Sage.
- Cumming, E., & Henry, W. E. (1961). *Growing old: The process of disengagement*. New York: Basic Books.
- Dargent-Molina, P., Hays, M., & Breart, G. (1996). Sensory impairments and physical disability in aged women living at home. *International Journal of Epidemiology*, *25*, 621-629.
- Dean, A., Kolody, B., & Wood, P. (1990). Effects of social support from various sources on depression in elderly persons. *Journal of Health and Social Behavior*, *31*, 148-161.
- Dickerson, A. E., & Fisher, A. G. (1993). Age differences in functional performance. *American Journal of Occupational Therapy*, *47*, 686-692.
- Duke, J., Leventhal, H., Brownlee, S., & Leventhal, E. A. (2002). Giving up and replacing activities in response to illness. *Journal of Gerontology: Psychological Sciences*, *57B*, P367-P376.
- Felton, B. J., & Berry, C. A. (1992). Do the sources of the urban elderly's social support determine its psychological consequences? *Psychology and Aging*, *7*, 89-97.
- Ferrucci, L., Cavazzini, C., Corsi, A., Bartali, B., Russo, C. R., Lauretani, F., et al. (2002). Biomarkers of frailty in older persons. *Journal of Endocrinological Investigation*, *25*, 10-15.
- Finch, J. F., Okun, M. A., Barrera, M., Jr., Zautra, A. J., & Reich, J. W. (1989). Positive and negative social ties among older adults: Measurement models and the prediction of psychological distress and well-being. *American Journal of Community Psychology*, *17*, 585-605.
- Folkman, S., Lazarus, R. S., Pimley, S., & Novacek, J. (1987). Age differences in stress and coping processes. *Psychology and Aging*, *2*, 171-184.
- Franceschi, C., Motta, L., Valensin, S., Rapisarda, R., Franzone, A., Berardelli, M., et al. (2000). Do men and women follow different trajectories to reach extreme longevity? Italian Multicenter Study on Centenarians (IMUSCE). *Aging: Clinical and Experimental Research*, *12*, 77-84.
- Guccione, A. A., Felson, D. T., Anderson, J. J., Anthony, J. M., Zhang, Y., Wilson, P. W., et al. (1994). The effects of specific medical conditions on the functional limitations of elders in the Framingham Study. *American Journal of Public Health*, *84*, 351-358.
- Health and Welfare Statistics Association. (2002). Kokumin eisei no doko [Health and welfare statistics in Japan]. *Journal of Health and Welfare Statistics*, *49*, 72-78.
- Holms, T. H., & Rahe, R. H. (1967). The social adjustment rating scale. *Journal of Psychosomatic Research*, *11*, 213-218.
- Hurwicz, M. L., Durham, C. C., Boyd-Davis, S. L., Gatz, M., & Bengtson, V. L. (1992). Salient life events in three-generation families. *Journal of Gerontology: Psychological Sciences*, *47*, P11-P13.
- Karasz, A., & Ouellette, S. C. (1995). Role strain and psychological well-being in women with systemic lupus erythematosus. *Women and Health*, *23*, 41-57.
- Koyano, W., Shibata, H., Nakazato, K., Haga, H., & Suyama, Y. (1987). Measurement of competence in the elderly living at home: Development of an index of competence. *Japanese Journal of Public Health*, *34*, 109-114.
- Koyano, W., Shibata, H., Nakazato, K., Haga, H., & Suyama, Y. (1991). Measurement of competence: Reliability and validity of the TMIG Index of competence. *Archives of Gerontology and Geriatrics*, *13*, 103-116.
- Krause, N. (1994). Stressors in salient social roles and well-being in later life. *Journal of Gerontology*, *49*, P137-P148.
- Krause, N. (2001). Social support. In R. H. Binstock, L. K. George, V. W. Marshall, A. M. O'Rand, & J. H. Schultz (Eds.), *Handbook of aging and the social sciences* (5th ed., pp. 272-294). San Diego, CA: Academic Press.

- Krause, N., Goldenhar, L., Liang, J., Jay, G., & Maeda, D. (1993). Stress and exercise among the Japanese elderly. *Social Science and Medicine*, 36, 1429-1441.
- Kriegsman, D. M., van Eijk, J. T., Penninx, B. W., Deeg, D. J., & Boeke, A. J. (1997). Does family support buffer the impact of specific chronic diseases on mobility in community-dwelling elderly? *Disability and Rehabilitation*, 19, 71-83.
- Lawrence, R. H., & Jette, A. M. (1996). Disentangling the disablement process. *Journal of Gerontology: Social Sciences*, 51B, S173-S182.
- Lee, G. R., & Ishii-Kuntz, M. (1987). Social interaction, loneliness, and emotional well-being among the elderly. *Research on Aging*, 9, 459-482.
- Lewis, M. A., & Rook, K. S. (1999). Social control in personal relationships: Impact on health behaviors and psychological distress. *Health Psychology*, 18, 63-71.
- Martin, M., Grüendahl, M., & Martin, P. (2001). Age differences in stress, social resources, and well-being in middle and older age. *Journal of Gerontology: Psychological Sciences*, 56B, P214-P222.
- Matsunaka, K. (2002). An investigation of personal factors on daily stress of people with visually impairment. *Shinrigaku Kenkyu*, 73, 340-345.
- Mendes de Leon, C. F., Gold, D. T., Glass, T. A., Kaplan, L., & George, L. K. (2001). Disability as a function of social networks and support in elderly African Americans and Whites: The Duke EPESE 1986-1992. *Journal of Gerontology: Social Sciences*, 56B, S179-S190.
- Mo, S. H. (1999). Population aging and changes in the family. In K. H. Cho, S. H. Hong, & K. W. Kim (Eds.), *Aging in Korea* (2nd ed., pp. 17-29). Seoul, Korea: Chung Ang Juk Sung Aptitude.
- Mor, V., Murphy, J., Masterson-Allen, S., Willey, C., Razmpour, A., Jackson, M. E., et al. (1989). Risk of functional decline among well elders. *Journal of Clinical Epidemiology*, 42, 895-904.
- Murrell, S. A., Norris, F. H., & Hutchins, G. L. (1984). Distribution and desirability of life events in older adults: Population and policy implications. *Journal of Community Psychology*, 12, 301-311.
- Noguchi, Y. (1991). Social support in elderly: The concept and measurement. *Social Gerontology*, 34, 37-48.
- Nolen-Hoeksema, S., Larson, J., & Grayson, C. (1999). Explaining the gender difference in depressive symptoms. *Journal of Personality and Social Psychology*, 77, 1061-1072.
- Norris, F. H., & Murrell, S. A. (1988). Prior experience as a moderator of disaster impact on anxiety symptoms in older adults. *American Journal of Community Psychology*, 16, 665-683.
- Okun, M. A., Melichar, J. F., & Hill, M. D. (1990). Negative daily events, positive and negative social ties, and psychological distress among older adults. *The Gerontologist*, 30, 193-199.
- O'Neill, E. S., & Morrow, L. L. (2001). The symptom experience of women with chronic illness. *Journal of Advanced Nursing*, 33, 257-268.
- Oskvig, R. M. (1999). Special problems in the elderly. *Chest*, 115, 158S-164S.
- Pearlin, L. I., & Lieberman, N. A. (1979). Social sources of emotional distress. *Research in Community and Mental Health*, 1, 217-248.
- Pearson, V. I. (2000). Assessment of function in older adults. In R. L. Kane & R. A. Kane (Eds.), *Assessing older persons: Measures, meaning, and practical applications* (pp. 17-48). New York: Oxford University Press.
- Penninx, B. W., van Tilburg, T., Boeke, A. J., Deeg, D. J., Kriegsman, D. M., & van Eijk, J. T. (1998). Effects of social support and personal coping resources on depressive symptoms: Different for various chronic diseases? *Health Psychology*, 17, 551-558.
- Penninx, B. W., van Tilburg, T., Erkinjuntti, T., Vataja, R., & Kaste, M. (1997). Comparison of stroke features and disability in daily life in patients with ischemic stroke aged 55 to 70 and 71 to 85 years. *Stroke*, 28, 729-735.
- Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1, 385-401.
- SAS Institute, Inc. (1996). The mixed procedure. In *SAS/STAT software: Changes and enhancements* (Release 6.11, pp. 531-656). Cary, NC: Author.
- Seeman, T. E., Berkman, L. F., Charpentier, P. A., Blazer, D. G., Albert, M. S., & Tinetti, M. E. (1995). Behavioral and psychosocial predictors of physical performance: MacArthur Studies of Successful Aging. *Journal of Gerontology: Medical Sciences*, 51A, M177-M183.
- Shima, S., Shikano, T., Kitamura, T., & Asai, M. (1985). New self-rating scales for depression. *Clinical Psychiatry*, 27, 717-723.
- Shimokata, H., Ando, F., & Niino, N. (2000). A new comprehensive study on aging—The National Institute for Longevity Sciences, Longitudinal Study of Aging (NILS-LSA). *Journal of Epidemiology*, 10, S1-S9.
- Shinn, M., Lehman, S., & Wong, N. W. (1984). Social interaction and social support. *Journal of Social Issues*, 40, 55-76.
- Valderrama-Gama, E., Damian, J., Ruigomez, A., & Martin-Moreno, J. M. (2002). Chronic disease, functional status, and self-ascribed causes of disabilities among noninstitutionalized older people in Spain. *Journal of Gerontology: Medical Sciences*, 57A, M716-M721.
- Yanagisawa, S., Baba, Y., Ito, C., Kobayashi, F., Kusagawa, Y., Kawai, F., et al. (2002). Effect of social support by family and friend on psychological QOL of the elderly. *Nippon Koshu Eisei Zasshi*, 49, 766-773.

Received December 20, 2002

Accepted September 4, 2003

Decision Editor: Margie E. Lachman, PhD

地域在住の中老年者の抑うつに関連する要因 ——その年齢差と性差——

国立長寿医療研究センター 坪井さとみ・福川康之・新野直明・安藤富士子・下方浩史

Age and gender differences in factors related to depressive symptoms among community-dwelling middle-aged and elderly people

Satomi Tsuboi, Yasuyuki Fukukawa, Naoakira Niino, Fujiko Ando, and Hiroshi Shimokata (Department of Epidemiology, the National Institute for Longevity Sciences, Gengo, Morioka-cho, Obu 474-8522)

This study was conducted to identify factors associated with depressive symptoms among males and females in two age groups; the middle-aged (40-59 years) and elderly (60-79 years). Subjects were 2 211 community-dwelling people (1 115 males and 1 096 females). Depressive symptoms were estimated by Center for Epidemiologic Studies Depression Scale (CES-D). Marital status, self-rated health, activities of daily living, age awareness, locus of control, and social support were assessed using self-administered questionnaires and interviews. In all groups, poor subjective health and external locus of control showed significant positive association with depressive symptoms. Age awareness was associated with depressive symptoms in female groups. Impairment of instrumental activities of daily living and lower social support were associated with depressive symptoms in the elderly groups. The findings suggested that there might be age and gender differences in the impact of factors associated with depressive symptoms.

Key words: middle-aged, elderly, gender, depressive symptoms, Center for Epidemiologic Studies Depression Scale.

老年期は、他の年齢層と比較して抑うつ状態の出現率が高いといわれている。多くの疫学調査でも、高齢者において抑うつ状態有症率の高いことが報告されている (Barusch, Rogers, & Abu-Bader, 1999; 新野, 1988)。老年期に抑うつ状態が多いのは、身体的健康及び経済的自立の損失、さらに定年退職や死別などによる社会や他者とのつながりの喪失が多くなることに関与するといわれるが (長谷川, 1987)、さらに詳しくその原因や構造を知るためには、老年期における抑うつの発生にかかわる要因について厳密に検討することが必要となる。

老年期の抑うつに関連する要因については、これまで様々な側面が論じられてきた。基本属性要因との関係では、性別 (川本・土井・山田・小国・岡山・鶴岡・佐藤・梶井, 1999; Zung, Broadhead, & Roth, 1993)、婚姻状態 (Pahkala, 1990) との関係があげられ、女性であることや、配偶者を持たないことが抑うつに関連するといわれている。身体的要因との関係では、うつ病と身体障害の関連 (滝沢・須賀・森田・内藤, 1988) や、日常生活の活動性が低い者に抑うつ状態の者が多い (井原, 1993; 村岡・生地・井原, 1996) ことが指摘されている。社会的要因との関係については、親密な友人関係や社会への参加度 (村岡他, 1996; Pahkala, 1990)、及びソーシャル・サポート (増地・岸, 2001) が、抑うつやうつ病と関連する

という報告がみられる。また、内的統制力の低さは老年期に特徴的な問題と考えられ、抑うつとの関連が指摘されている (Baum & Boxley, 1983)。

一方、老性自覚とは“加齢による心身の変化を主観的にどのように自覚しているか (中山・藤野, 1994, p. 22)”であるが、人は中年期の頃から、“もう若くはない”、“この頃、年をとったと思う”などの自覚体験が契機となり、新たに体験した自己と、それまで自分が抱いていた内的な自己とのギャップによる葛藤から、無力感や抑うつ感、心気症状を示す場合がある。特にいつまでも若くありつづけることに価値をおく社会においては、老性自覚と抑うつとの関連は強いといえよう。しかし、老性自覚体験が中年期から老年期の抑うつ関連要因の一つであることを検討した研究は少ない (Baum & Boxley, 1983)。Baum & Boxley (1983) は、高齢者の場合、年をとっていると自己を見なすことは抑うつに関係することを明らかにしたが、Linn & Hunter (1979) は、社会的地位や障害の程度などの変数を調整すると、年をとっていると感じることで、抑うつとの関係は見出せないとしている。

以上のように、老年期には様々な要因が抑うつと関連することが指摘されている。しかし、これまでの研究は高齢者に対象を限定した研究が多く、中年期と老年期の差異を検討するというような、広範な年齢層を含んだ詳細な検討は少ない。老年期の抑うつとの関連

が指摘されている変数の中には、老年期特有の変数、あるいは他の年齢においても同様の関連がみられる変数が混在している。また、抑うつに関連する要因は年齢や性別とも強く関連するものが多いため、各年齢や性別により、それぞれの要因が及ぼす影響の強さが異なる可能性がある。従って、本研究は、老年期の抑うつ関連要因を明らかにするために、中年期との比較、及び性差を検討することに焦点をおく。また、上記のように複数の要因が抑うつに影響すると指摘されているにもかかわらず、身体的要因や心理社会的要因などそれぞれの領域を別個に扱った研究が多く、各要因における相互間の関連を考慮した上で、総合的に抑うつとの関係を検討したものは少ない。そこで、これまでに関連が指摘されている要因を幅広くとらえ、これらの変数の相互関連を考慮し、中年期から老年期の男女の抑うつに関連する要因の特徴を探索的に示すことを第2の焦点とする。すなわち、本研究の目的は、地域在住の40歳から79歳の住民2267名に対し大規模な調査を行い、上記2点に焦点づけ、抑うつに関連する要因の年齢差及び性差を明らかにすることである。

方 法

調査対象

本研究の対象者は、国立長寿医療研究センターが行っている“老化に関する長期縦断疫学調査” (the National Institute for Longevity Sciences-Longitudinal Study of Aging: NILS-LSA) (Shimokata, Ando, & Niino, 2000) の第1回目調査 (1997年11月—2000年4月) に参加した地域住民2267名のうち、分析項目すべてについて有効回答が得られた者2211名 (男性1115名、女性1096名) である。

NILS-LSAは、医学・身体組成・運動機能・心理・栄養の各領域における様々な加齢変化を追跡し、正常な老化の進行過程を経時的に記録することを目的とした研究プロジェクトであり、本研究はその成果の一部である。対象者の選定については、センター近隣の愛知県大府市 (人口約7万人) 及び知多郡東浦町 (人口約4万人) を対象地域とし、自治体の協力を得て住民台帳から40—79歳の7855名を性、年代別に層化無作為抽出した。この地域は、名古屋市南部に位置し、大都市のベッドタウン及びトヨタグループを中心とした機械工業地帯として発展を遂げる一方、果樹栽培が盛んな地帯や田園地帯を多く残している。平成15年の住民基本台帳・外国人登録による調査では大府市の40—79歳の人口は、40代8945人 (26.3%)、50代11535人 (34.0%)、60代8721人 (25.7%)、70代4763人 (14.0%) である。また、平成12年の国勢調査より東浦町の40—79歳の人口は、40代5749人 (27.8%)、50代7240人 (35.0%)、60代4990人

(24.1%)、70代2703人 (13.1%) であり (カッコ内はいずれも40—79歳の全人口を100とした割合)、共に50代が多い構成になっている。しかし、我々の調査では性年齢別の比較を容易にするために、男女及び10歳刻みの年代で分けた八つの群に含まれる人数がほぼ同数になるように対象者の抽出、募集を行った。そのため、対象者の性年齢構成は地域を代表する訳ではないことに留意されたい。

抽出した7855名には、郵送にて調査の説明会への参加の意思についての問い合わせを行ったが、問い合わせ時にすでに65名が死亡ないし対象地域外へ転出していた。問い合わせ可能であった7790名のうち、3434名から回答が得られ、参加拒否の回答は881名であった。参加の意思のあった2553名のうち2510名が説明会に参加し、2267名が調査への参加に合意、インフォームドコンセントに署名した上で実際に調査に参加した。説明会への参加を呼びかけた者の29.1%が参加し、回答が得られた者からの参加率は66.0%であった。

対象者の学歴は、老年期女性のみ中卒者 (52.0%) が多かったが、中年期男女と老年期男性は、過半数 (55.7—85.6%) が高卒以上であった。職業に関しては、中年期男性では、技能・労務職 (30.9%)、事務・専門的技術職 (27.3%)、管理職 (21.8%) が多く、老年期男性では、無職 (60.2%) が多くを占め、有職者では、技能・労務職 (9.4%)、事務・専門的技術職 (6.9%)、農林漁業職 (6.0%) が多かった。中年期女性では、主婦またはパート職 (70.8%) が最も多く、有職者 (パート職含む) のうち、事務・専門技術職 (22.1%)、サービス・販売職 (15.0%) が多かった。老年期女性では、主婦またはパート職 (61.8%) が多く、無職が27.9%、農林漁業職が5.9%であった。

NILS-LSAについては、国立療養所中部病院倫理委員会の承認が得られている。

調査内容

抑うつの測定 抑うつ状態の評価には Center for Epidemiologic Studies Depression Scale (以下 CES-D; Radloff, 1977) の日本語版 (島・鹿野・北村・浅井, 1985) を用いた。この尺度は、米国国立精神衛生研究所において開発された疫学調査用の自己評価式抑うつ尺度で、20項目から成る。評定は4件法で、得点が高いほど抑うつ状態が強いことを示す (得点範囲0—60点)。ここでの抑うつ状態とは、抑うつ気分の他、悲観的な考え、被害的感情、精神運動抑制などを含む。

関連要因の測定 抑うつに関連する要因として、以下の変数を取り上げた。

1. 基本属性要因：配偶者の有無 (0=配偶者あり (内縁関係を含む)、1=配偶者なし (未婚、別居、離

Table 1
対象者の年齢及び性別の特性

		(n, %)	中年期群 (40—59 歳)		老年期群 (60—79 歳)		$\alpha^{a)}$				
			男性	女性	男性	女性					
人数		(n, %)	564	100.0	552	100.0	551	100.0	544	100.0	
基本属性要因											
配偶者	あり	(n, %)	522	92.6	502	90.9	513	93.1	377	69.3	
	なし	(n, %)	42	7.4	50	9.1	38	6.9	167	30.7	
身体的要因											
主観的健康感	非常に良い	(n, %)	17	3.0	26	4.7	17	3.1	8	1.5	
	良い	(n, %)	166	29.4	139	25.2	104	18.9	63	11.6	
	普通	(n, %)	336	59.6	344	62.3	351	63.7	398	73.2	
	悪い	(n, %)	44	7.8	42	7.6	75	13.6	69	12.7	
	非常に悪い	(n, %)	1	0.2	1	0.2	4	0.7	6	1.1	
活動能力	<i>M</i>		12.23		12.62		12.23		12.25		.57
	<i>SD</i>		1.17		0.78		1.22		1.19		
心理社会的要因											
老性自覚	なし	(n, %)	236	41.8	202	36.6	172	31.2	138	25.4	
	あり	(n, %)	328	58.2	350	63.4	379	68.8	406	74.6	
ローカス・オブ・コントロール	<i>M</i>		27.77		26.00		27.34		25.24		.74
	<i>SD</i>		4.67		4.75		4.70		4.93		
家族外サポート	<i>M</i>		28.84		30.39		29.59		30.92		.91
	<i>SD</i>		5.85		5.35		6.07		5.85		
抑うつ	<i>M</i>		6.88		7.18		7.39		8.37		.87
	<i>SD</i>		6.5		6.03		7.02		7.28		

^{a)} α = 信頼性係数。

婚、死別を含む)) の 2 分類とした。

2. 身体的要因：主観的健康感 (1 = 非常に良い - 5 = 非常に悪い、の 5 段階評定) と活動能力を尋ねた。活動能力の測定には、古谷野・柴田・中里・芳賀・須山 (1987) の老研式活動能力指標を用いた。この尺度は地域の中で自立した生活を営むために必要な能力である手段的日常生活動作能力 (Instrumental Activities of Daily Living: 以下 IADL)、機能的日常生活動作能力 (Functional Activities of Daily Living: 以下 FADL)、及び社会活動性を測定する。13 項目から成り、2 件法、13 点満点で、高得点ほど活動能力が高いことを示す。

3. 心理的要因：老性自覚の測定は、家族の死や病氣などの体験の有無を尋ねるライフイベント・チェックリスト中の 1 項目として、過去 2 年間における老いの自覚体験の有無を、1 = あり、0 = なしの 2 段階で尋ねた。過去 2 年間としたのは、NILS-LSA の測定が 2 年ごとに行われるためである。ローカス・オブ・コントロールの測定には、8 項目からなる尺度 (若林・蔭山・金井, 1995) を用いた。この尺度は 5 件法で回答を求め、高得点ほど内的統制、低得点ほど外的統制で

あることを示す (得点範囲 8—40 点)。ソーシャルサポートの測定には、家族以外の人々からのサポートを測定するソーシャルサポート尺度 (浦, 1992) を用いた。この尺度は 10 項目から成り、5 件法で、得点が高いほどサポートを受けていることを示す (得点範囲 10—50 点)。各尺度の信頼性係数は Cronbach α 係数を求めて検討した。各尺度の信頼性係数は Table 1 に示した。 α 係数は、活動能力でやや低かったが、これは中年期のデータがほとんど満点であったためと考えられ、尺度の測定する内容を知る重要度から、今後の分析に用いる必要があると判断した。

測定及び分析方法

老性自覚は、個別面接法によりその有無を測定した。その他の変数は、自記式調査票を用い、自宅で記入した上で面接時に持参するよう依頼し、面接時に再度未記入部分や不明点などを確認した。面接調査員はすべて調査方法について訓練を受けた臨床心理学専攻の大学院生である。

分析手順としては、まず、対象者を中年期 (40—59 歳)、老年期 (60—79 歳) の二つの年齢コホート (世

代)、及び性別のそれぞれ2群に分割し、各要因における年齢差・性差を検討した。カテゴリー変数の比率の差の検定には年齢・性別に χ^2 検定を用いた。連続変数については年齢×性の2要因分散分析を行った。

次にCES-Dを基準変数とし、関連要因の各変数を説明変数として、 F 値の変動が、変数追加では.05、変数削除では.10の確率で起きた場合に変数の追加と削除を行うステップワイズ法による重回帰分析を行い、関連要因相互の影響を取り除いた上で年齢及び性別の4群における抑うつに関連する要因について検討した。統計処理は、すべて統計プログラムパッケージSAS (Ver. 6.12, 1997)を用いた。

結 果

対象者の特徴

分析対象者の年齢及び性別による各特性の値をTable 1に示す。配偶者の有無については、 χ^2 検定より年齢・性の頻度の偏りが共に有意で($\chi^2(1)=52.18, p<.001$; $\chi^2(1)=75.75, p<.001$)、老年期及び女性で配偶者なし(未婚、離婚、死別を含む)の割合が高かった。主観的健康感、 χ^2 検定の結果年齢・性の頻度の偏りが共に有意で($\chi^2(4)=68.21, p<.001$; $\chi^2(4)=12.36, p<.05$)、特に老年期で“悪い”、“非常に悪い”者が多かった。老性自覚についても、 χ^2 検定より、年齢・性の頻度の偏りが共に有意で($\chi^2(1)=29.53, p<.001$; $\chi^2(1)=7.66, p<.01$)、老年期及び女性で“あり”の者が多かった。また、活動能力は、年齢×性の2要因分散分析を行ったところ、年齢の主効果及び性の主効果が共に有意であり、($F(1, 2207)=15.35, p<.001$, 老年期<中年期; $F(1, 2207)=19.74, p<.001$, 男性<女性)、両条件の交互作用に有意差がみられた($F(1, 2207)=15.31, p<.001$)。そこで、男性群、女性群それぞれにおける年齢要因の単純主効果を検定したところ、女性群のみ年齢の効果が有意であり($F(1, 1094)=37.06, p<.001$)、中年期の活動能力が高かったが、男性では年齢差がみられなかった。ローカス・オブ・コントロールは、2要因分散分析の結果、年齢の主効果及び性の主効果が共に有意であり、($F(1, 2207)=8.65, p<.01$, 老年期<中年期; $F(1, 2207)=91.38, p<.001$, 女性<男性)、交互作用は見出されなかった。家族以外からのソーシャルサポートは、2要因分散分析の結果年齢の主効果及び性の主効果が共に有意で($F(1, 2207)=6.76, p<.01$; $F(1, 2207)=34.22, p<.001$)、老年期及び女性で高く交互作用は見出されなかった。抑うつについては、2要因分散分析の結果、年齢の主効果及び性の主効果が共に有意であり($F(1, 2207)=8.78, p<.01$; $F(1, 2207)=4.92, p<.05$)、老年期及び女性で高く、交互作用はみられなかった。以上より、すべての要因で年齢・性

差がみられたので、以後の分析は、年齢及び性別の4群で行った。

抑うつの関連要因

分析に用いた基準変数及び説明変数の各変数間の相関をピアソンの相関係数により検討したところ、CES-D得点と正の相関がみられた変数は、配偶者(なし)、主観的健康感(悪い)、老性自覚(あり)であった。負の相関がみられた変数は、活動能力、ローカス・オブ・コントロール、家族以外からのソーシャルサポートであった。CES-D得点とすべての変数間で統計的に有意な相関が示されたが、いずれも相関係数は低かった。また、説明変数間の相互相関についても、弱い関連しかみられなかった。次に、CES-D得点を基準変数とする各群別の重回帰分析によって、抑うつの関連要因について検討した。説明変数として投入した変数は、今回の研究に用いた基本属性要因、身体的要因、心理社会的要因のすべてである。Table 2に示すように、回帰式全体の決定係数(R^2)は、中年期男性群=.211、中年期女性群=.149、老年期男性群=.276、老年期女性群=.184であり、いずれの年齢群においても有意であった(中年男性群: $F=29.81, p<.001$, 中年女性群: $F=19.11, p<.001$, 老年期男性群 $F=34.64, p<.001$, 老年期女性群 $F=20.12, p<.001$)。またすべての説明変数間の相関は低く、トレランス値も.78-.99であったため、変数間の多重共線性の問題はないものと判断した。

各投入変数の標準偏回帰係数を観察すると、いずれの群に対してもCES-D得点との有意な関連を示したのは、主観的健康感とローカス・オブ・コントロールであった。つまり、主観的健康感が不良の者、外的統制の者は、年齢・性別にかかわらず抑うつが高いという関係が認められた。(主観的健康感: $\beta=.207-.263$, ローカス・オブ・コントロール: $\beta=-.284-.188$)。配偶者の有無については、中年期群、老年期男性群でCES-D得点に影響し(中年期男性群: $\beta=.100, p<.01$, 中年期女性群: $\beta=.081, p<.05$, 老年期男性群: $\beta=.152, p<.001$)、配偶者なしの者は抑うつが高いことが示されたが、老年期女性群のみ抑うつとの有意な関係が認められなかった。活動能力については、老年期の男女群共にCES-D得点に有意な効果を示し(老年期男性群: $\beta=-.134, p<.001$, 老年期女性群: $\beta=-.109, p<.01$)、社会生活や日常生活に支障がある高齢者ほど抑うつを訴えることが多いことが示された。しかし、中年期群では有意な効果はなく、活動能力の程度はCES-D得点に影響しなかった。老性自覚については、女性群のみでCES-D得点に影響し(中年期女性群: $\beta=.083, p<.05$, 老年期女性群: $\beta=.078, p<.05$)、老性自覚を持つ女性ほど、抑うつが高くなることが示されたが、男性群では