

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
General anesthetic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Hypogenic sadative agent	Male	7	2.6	8	2.4	13	4.4	30	11.2	4	9.3	62	5.1
	Female	6	2.0	11	3.9	31	10.8	49	17.8	5	14.7	102	8.7
	Total	13	2.3	19	3.1	44	7.5	79	14.6	9	11.7	164	6.9
Antiepileptic agent	Male	0	0.0	0	0.0	2	0.7	2	0.7	0	0.0	4	0.3
	Female	2	0.7	2	0.7	4	1.4	3	1.1	0	0.0	11	0.9
	Total	2	0.4	2	0.3	6	1.0	5	0.9	0	0.0	15	0.6
Antipyretic analgesic agent	Male	31	11.7	20	6.0	24	8.1	41	15.4	9	20.9	125	10.4
	Female	61	20.7	34	11.9	46	16.1	52	18.9	5	14.7	198	16.9
	Total	92	16.4	54	8.8	70	12.0	93	17.2	14	18.2	323	13.6
Stimulant drug, antihypnotic drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antiparkinson drugs	Male	0	0.0	1	0.3	1	0.3	1	0.4	0	0.0	3	0.2
	Female	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Total	1	0.2	1	0.2	1	0.2	1	0.2	0	0.0	4	0.2
Antipsychotic drug	Male	5	1.9	6	1.8	12	4.0	9	3.4	0	0.0	32	2.7
	Female	5	1.7	4	1.4	15	5.2	17	6.2	3	8.8	44	3.7
	Total	10	1.8	10	1.6	27	4.6	26	4.8	3	3.9	76	3.2
Drug for common cold	Male	23	8.6	23	6.9	24	8.1	14	5.2	3	7.0	87	7.2
	Female	37	12.6	20	7.0	24	8.4	13	4.7	3	8.8	97	8.3
	Total	60	10.7	43	7.0	48	8.2	27	5.0	6	7.8	184	7.7
Other agent acting on CNS	Male	0	0.0	0	0.0	1	0.3	3	1.1	0	0.0	4	0.3
	Female	0	0.0	1	0.4	0	0.0	1	0.4	0	0.0	2	0.2
	Total	0	0.0	1	0.2	1	0.2	4	0.7	0	0.0	6	0.3
Local anesthetic	Male	1	0.4	1	0.3	0	0.0	0	0.0	0	0.0	2	0.2
	Female	0	0.0	0	0.0	2	0.7	0	0.0	0	0.0	2	0.2
	Total	1	0.2	1	0.2	2	0.3	0	0.0	0	0.0	4	0.2
Muscle relaxant	Male	0	0.0	0	0.0	0	0.0	2	0.7	1	2.3	3	0.2
	Female	0	0.0	0	0.0	1	0.3	1	0.4	0	0.0	2	0.2
	Total	0	0.0	0	0.0	1	0.2	3	0.6	1	1.3	5	0.2

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Agent for autonomic nervous system	Male	0	0.0	1	0.3	0	0.0	3	1.1	0	0.0	4	0.3
	Female	1	0.3	0	0.0	0	0.0	3	1.1	0	0.0	4	0.3
	Total	1	0.2	1	0.2	0	0.0	6	1.1	0	0.0	8	0.3
Antispasmodic agent	Male	5	1.9	3	0.9	5	1.7	5	1.9	1	2.3	19	1.6
	Female	4	1.4	3	1.1	6	2.1	5	1.8	3	8.8	21	1.8
	Total	9	1.6	6	1.0	11	1.9	10	1.8	4	5.2	40	1.7
Diaphoretic, antiperspirants	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent for PNS	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Ophthalmic solution (others)	Male	51	19.2	61	18.4	64	21.5	60	22.5	13	30.2	249	20.7
	Female	48	16.3	48	16.8	73	25.5	97	35.3	14	41.2	280	23.9
	Total	99	17.7	109	17.7	137	23.5	157	29.0	27	35.1	529	22.2
Ophthalmic solution (for cataract)	Male	0	0.0	5	1.5	14	4.7	38	14.2	7	16.3	64	5.3
	Female	0	0.0	2	0.7	28	9.8	66	24.0	8	23.5	104	8.9
	Total	0	0.0	7	1.1	42	7.2	104	19.2	15	19.5	168	7.1
Ophthalmic solution (pilocarpine)	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Total	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0	2	0.1
Ophthalmic solution (β blocker)	Male	0	0.0	1	0.3	5	1.7	10	3.7	2	4.7	18	1.5
	Female	0	0.0	2	0.7	8	2.8	6	2.2	1	2.9	17	1.4
	Total	0	0.0	3	0.5	13	2.2	16	3.0	3	3.9	35	1.5
Ophthalmic solution (epinephrine)	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Ophthalmic solution (for glaucoma)	Male	0	0.0	2	0.6	5	1.7	7	2.6	1	2.3	15	1.2
	Female	0	0.0	1	0.4	4	1.4	5	1.8	1	2.9	11	0.9
	Total	0	0.0	3	0.5	9	1.5	12	2.2	2	2.6	26	1.1
Ophthalmic solution (steroid)	Male	2	0.8	4	1.2	8	2.7	12	4.5	4	9.3	30	2.5
	Female	3	1.0	3	1.1	5	1.7	18	6.5	5	14.7	34	2.9
	Total	5	0.9	7	1.1	13	2.2	30	5.5	9	11.7	64	2.7

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Ophthalmic solution (unknown)	Male	0	0.0	1	0.3	3	1.0	11	4.1	1	2.3	16	1.3
	Female	3	1.0	3	1.1	4	1.4	8	2.9	3	8.8	21	1.8
	Total	3	0.5	4	0.6	7	1.2	19	3.5	4	5.2	37	1.6
Agent for ENT use	Male	4	1.5	1	0.3	3	1.0	5	1.9	0	0.0	13	1.1
	Female	0	0.0	1	0.4	1	0.3	1	0.4	1	2.9	4	0.3
	Total	4	0.7	2	0.3	4	0.7	6	1.1	1	1.3	17	0.7
Anti-vertigeneous drug	Male	0	0.0	1	0.3	3	1.0	5	1.9	0	0.0	9	0.7
	Female	0	0.0	5	1.8	3	1.0	12	4.4	3	8.8	23	2.0
	Total	0	0.0	6	1.0	6	1.0	17	3.1	3	3.9	32	1.3
Other agent acting on sensory organ	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Digitalis agent	Male	0	0.0	0	0.0	2	0.7	5	1.9	1	2.3	8	0.7
	Female	1	0.3	2	0.7	3	1.0	1	0.4	0	0.0	7	0.6
	Total	1	0.2	2	0.3	5	0.9	6	1.1	1	1.3	15	0.6
Xanthine agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other inotropic agent	Male	0	0.0	0	0.0	1	0.3	1	0.4	0	0.0	2	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	1	2.9	1	0.1
	Total	0	0.0	0	0.0	1	0.2	1	0.2	1	1.3	3	0.1
β blocker	Male	3	1.1	10	3.0	8	2.7	13	4.9	1	2.3	35	2.9
	Female	3	1.0	7	2.5	4	1.4	8	2.9	1	2.9	23	2.0
	Total	6	1.1	17	2.8	12	2.1	21	3.9	2	2.6	58	2.4
Agents for membrane-stabilizing activity	Male	0	0.0	1	0.3	3	1.0	4	1.5	0	0.0	8	0.7
	Female	0	0.0	0	0.0	1	0.3	5	1.8	0	0.0	6	0.5
	Total	0	0.0	1	0.2	4	0.7	9	1.7	0	0.0	14	0.6
Other antiarrhythmic agent	Male	0	0.0	1	0.3	2	0.7	2	0.7	0	0.0	5	0.4
	Female	0	0.0	0	0.0	3	1.0	1	0.4	0	0.0	4	0.3
	Total	0	0.0	1	0.2	5	0.9	3	0.6	0	0.0	9	0.4
Thyazide diuretic	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	1	0.3	0	0.0	1	0.3	4	1.5	0	0.0	6	0.5
	Total	1	0.2	0	0.0	2	0.3	4	0.7	0	0.0	7	0.3

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Loop diuretic	Male	0	0.0	0	0.0	2	0.7	4	1.5	1	2.3	7	0.6
	Female	1	0.3	1	0.4	4	1.4	3	1.1	0	0.0	9	0.8
	Total	1	0.2	1	0.2	6	1.0	7	1.3	1	1.3	16	0.7
Potassium sparing diuretic	Male	0	0.0	0	0.0	1	0.3	2	0.7	1	2.3	4	0.3
	Female	0	0.0	0	0.0	4	1.4	5	1.8	1	2.9	10	0.9
	Total	0	0.0	0	0.0	5	0.9	7	1.3	2	2.6	14	0.6
Carbonate dehydratase inhibitor	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0
Other diuretic	Male	0	0.0	0	0.0	0	0.0	2	0.7	0	0.0	2	0.2
	Female	0	0.0	0	0.0	3	1.0	4	1.5	1	2.9	8	0.7
	Total	0	0.0	0	0.0	3	0.5	6	1.1	1	1.3	10	0.4
ACE inhibitor	Male	4	1.5	22	6.6	27	9.1	41	15.4	6	14.0	100	8.3
	Female	4	1.4	11	3.9	26	9.1	38	13.8	5	14.7	84	7.2
	Total	8	1.4	33	5.4	53	9.1	79	14.6	11	14.3	184	7.7
Ca antagonist	Male	8	3.0	41	12.4	63	21.2	74	27.7	10	23.3	196	16.3
	Female	8	2.7	25	8.8	44	15.4	93	33.8	12	35.3	182	15.5
	Total	16	2.9	66	10.7	107	18.4	167	30.8	22	28.6	378	15.9
Other antihypertensive agent	Male	1	0.4	6	1.8	9	3.0	13	4.9	1	2.3	30	2.5
	Female	0	0.0	3	1.1	9	3.1	7	2.5	2	5.9	21	1.8
	Total	1	0.2	9	1.5	18	3.1	20	3.7	3	3.9	51	2.1
Vascular reinforcing agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Vasoconstrictor	Male	0	0.0	1	0.3	0	0.0	1	0.4	0	0.0	2	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0	2	0.1
Vasodilator	Male	1	0.4	4	1.2	15	5.1	23	8.6	5	11.6	48	4.0
	Female	1	0.3	1	0.4	8	2.8	25	9.1	4	11.8	39	3.3
	Total	2	0.4	5	0.8	23	3.9	48	8.9	9	11.7	87	3.7
Antihyperlipidemic agent	Male	5	1.9	19	5.7	20	6.7	29	10.9	3	7.0	76	6.3
	Female	2	0.7	24	8.4	44	15.4	76	27.6	5	14.7	151	12.9
	Total	7	1.3	43	7.0	64	11.0	105	19.4	8	10.4	227	9.5

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other agent on cardiovascular system	Male	1	0.4	6	1.8	9	3.0	20	7.5	3	7.0	39	3.2
	Female	1	0.3	1	0.4	8	2.8	17	6.2	7	20.6	34	2.9
	Total	2	0.4	7	1.1	17	2.9	37	6.8	10	13.0	73	3.1
Respiratory stimulator	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antitussive agent	Male	3	1.1	0	0.0	1	0.3	3	1.1	0	0.0	7	0.6
	Female	7	2.4	6	2.1	3	1.0	7	2.5	0	0.0	23	2.0
	Total	10	1.8	6	1.0	4	0.7	10	1.8	0	0.0	30	1.3
Expectorant	Male	3	1.1	2	0.6	6	2.0	6	2.2	3	7.0	20	1.7
	Female	9	3.1	3	1.1	9	3.1	8	2.9	1	2.9	30	2.6
	Total	12	2.1	5	0.8	15	2.6	14	2.6	4	5.2	50	2.1
Antitussive agent and expectorant	Male	3	1.1	3	0.9	1	0.3	2	0.7	0	0.0	9	0.7
	Female	7	2.4	5	1.8	2	0.7	4	1.5	1	2.9	19	1.6
	Total	10	1.8	8	1.3	3	0.5	6	1.1	1	1.3	28	1.2
Bronchodilator	Male	4	1.5	3	0.9	3	1.0	11	4.1	3	7.0	24	2.0
	Female	6	2.0	2	0.7	3	1.0	5	1.8	0	0.0	16	1.4
	Total	10	1.8	5	0.8	6	1.0	16	3.0	3	3.9	40	1.7
Gargle	Male	1	0.4	0	0.0	2	0.7	0	0.0	0	0.0	3	0.2
	Female	1	0.3	1	0.4	0	0.0	1	0.4	0	0.0	3	0.3
	Total	2	0.4	1	0.2	2	0.3	1	0.2	0	0.0	6	0.3
Other agent acting on respiratory system	Male	2	0.8	1	0.3	1	0.3	4	1.5	0	0.0	8	0.7
	Female	2	0.7	1	0.4	1	0.3	2	0.7	0	0.0	6	0.5
	Total	4	0.7	2	0.3	2	0.3	6	1.1	0	0.0	14	0.6
Antidiarrheal drug	Male	4	1.5	5	1.5	8	2.7	7	2.6	4	9.3	28	2.3
	Female	6	2.0	1	0.4	6	2.1	15	5.5	0	0.0	28	2.4
	Total	10	1.8	6	1.0	14	2.4	22	4.1	4	5.2	56	2.4
Antiulcerative agent	Male	16	6.0	23	6.9	37	12.5	67	25.1	11	25.6	154	12.8
	Female	16	5.4	19	6.7	42	14.7	66	24.0	5	14.7	148	12.6
	Total	32	5.7	42	6.8	79	13.6	133	24.5	16	20.8	302	12.7
Stomachic	Male	26	9.8	26	7.9	22	7.4	18	6.7	3	7.0	95	7.9
	Female	14	4.8	21	7.4	18	6.3	22	8.0	4	11.8	79	6.7
	Total	40	7.1	47	7.6	40	6.9	40	7.4	7	9.1	174	7.3

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Antacid	Male	1	0.4	2	0.6	1	0.3	1	0.4	0	0.0	5	0.4
	Female	0	0.0	0	0.0	0	0.0	4	1.5	0	0.0	4	0.3
	Total	1	0.2	2	0.3	1	0.2	5	0.9	0	0.0	9	0.4
Laxative	Male	2	0.8	6	1.8	5	1.7	28	10.5	1	2.3	42	3.5
	Female	13	4.4	20	7.0	16	5.6	29	10.5	1	2.9	79	6.7
	Total	15	2.7	26	4.2	21	3.6	57	10.5	2	2.6	121	5.1
Cholagogue	Male	0	0.0	3	0.9	8	2.7	4	1.5	1	2.3	16	1.3
	Female	0	0.0	1	0.4	0	0.0	2	0.7	0	0.0	3	0.3
	Total	0	0.0	4	0.6	8	1.4	6	1.1	1	1.3	19	0.8
Combined digestive drug	Male	1	0.4	3	0.9	1	0.3	2	0.7	0	0.0	7	0.6
	Female	3	1.0	0	0.0	0	0.0	3	1.1	0	0.0	6	0.5
	Total	4	0.7	3	0.5	1	0.2	5	0.9	0	0.0	13	0.5
Other agent acting on alimentary system	Male	11	4.1	4	1.2	14	4.7	13	4.9	3	7.0	45	3.7
	Female	7	2.4	7	2.5	7	2.4	25	9.1	5	14.7	51	4.3
	Total	18	3.2	11	1.8	21	3.6	38	7.0	8	10.4	96	4.0
Pituitary hormone drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	1	2.3	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	1	1.3	1	0.0
Hormone drug of salivary gland	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Thyroid parathyroid hormone drug	Male	1	0.4	1	0.3	1	0.3	1	0.4	0	0.0	4	0.3
	Female	1	0.3	6	2.1	2	0.7	6	2.2	0	0.0	15	1.3
	Total	2	0.4	7	1.1	3	0.5	7	1.3	0	0.0	19	0.8
Anabolic steroid drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Adrenocortical hormone drug	Male	6	2.3	8	2.4	4	1.3	7	2.6	1	2.3	26	2.2
	Female	7	2.4	9	3.2	5	1.7	4	1.5	0	0.0	25	2.1
	Total	13	2.3	17	2.8	9	1.5	11	2.0	1	1.3	51	2.1
Androgen drug	Male	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	1	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Estrogen or progesterone drug	Male	0	0.0	0	0.0	0	0.0	5	1.9	0	0.0	5	0.4
	Female	3	1.0	5	1.8	2	0.7	1	0.4	0	0.0	11	0.9
	Total	3	0.5	5	0.8	2	0.3	6	1.1	0	0.0	16	0.7
Mixed hormone drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Total	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0
Other hormone drug	Male	1	0.4	1	0.3	9	3.0	3	1.1	1	2.3	15	1.2
	Female	1	0.3	3	1.1	5	1.7	10	3.6	1	2.9	20	1.7
	Total	2	0.4	4	0.6	14	2.4	13	2.4	2	2.6	35	1.5
Agent acting on urinary system	Male	0	0.0	0	0.0	0	0.0	0	0.0	1	2.3	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	1	1.3	1	0.0
Agent acting on genital system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Uterine constrictor	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Contraceptive	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Total	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0
Drugs for hemorrhoids	Male	1	0.4	2	0.6	1	0.3	0	0.0	1	2.3	5	0.4
	Female	0	0.0	2	0.7	1	0.3	2	0.7	0	0.0	5	0.4
	Total	1	0.2	4	0.6	2	0.3	2	0.4	1	1.3	10	0.4
Other agent acting on urogenital or anus	Male	0	0.0	3	0.9	10	3.4	21	7.9	3	7.0	37	3.1
	Female	2	0.7	0	0.0	1	0.3	5	1.8	1	2.9	9	0.8
	Total	2	0.4	3	0.5	11	1.9	26	4.8	4	5.2	46	1.9
Integumentary microbiosite	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Protective agent for wound	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Pyostatic	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.0
Analgenic (external application)	Male	3	1.1	4	1.2	4	1.3	14	5.2	2	4.7	27	2.2
	Female	9	3.1	6	2.1	6	2.1	16	5.8	1	2.9	38	3.2
	Total	12	2.1	10	1.6	10	1.7	30	5.5	3	3.9	65	2.7
Integumentary Parasiticide	Male	1	0.4	2	0.6	3	1.0	0	0.0	0	0.0	6	0.5
	Female	1	0.3	0	0.0	1	0.3	0	0.0	0	0.0	2	0.2
	Total	2	0.4	2	0.3	4	0.7	0	0.0	0	0.0	8	0.3
Emollient	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Hair tonic	Male	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	1	0.0
Bath liquid	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other surface-acting agent	Male	0	0.0	2	0.6	1	0.3	0	0.0	0	0.0	3	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	2	0.3	1	0.2	0	0.0	0	0.0	3	0.1
Dental local anesthetic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for devitalization of the pulp	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Analgesic or sedative for dental use	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for mummification of the pulp	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Agent for pulp capping	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics for dental use	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Total	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0
Other agent for dental use	Male	0	0.0	1	0.3	1	0.3	1	0.4	0	0.0	3	0.2
	Female	0	0.0	1	0.4	2	0.7	1	0.4	0	0.0	4	0.3
	Total	0	0.0	2	0.3	3	0.5	2	0.4	0	0.0	7	0.3
Vitamin A or D	Male	0	0.0	1	0.3	2	0.7	2	0.7	1	2.3	6	0.5
	Female	1	0.3	7	2.5	12	4.2	30	10.9	4	11.8	54	4.6
	Total	1	0.2	8	1.3	14	2.4	32	5.9	5	6.5	60	2.5
Vitamin B1	Male	2	0.8	1	0.3	2	0.7	7	2.6	0	0.0	12	1.0
	Female	0	0.0	1	0.4	4	1.4	7	2.5	1	2.9	13	1.1
	Total	2	0.4	2	0.3	6	1.0	14	2.6	1	1.3	25	1.1
Vitamin B complex	Male	6	2.3	7	2.1	12	4.0	18	6.7	4	9.3	47	3.9
	Female	4	1.4	8	2.8	14	4.9	33	12.0	2	5.9	61	5.2
	Total	10	1.8	15	2.4	26	4.5	51	9.4	6	7.8	108	4.5
Vitamin C	Male	1	0.4	3	0.9	1	0.3	2	0.7	0	0.0	7	0.6
	Female	7	2.4	5	1.8	6	2.1	5	1.8	0	0.0	23	2.0
	Total	8	1.4	8	1.3	7	1.2	7	1.3	0	0.0	30	1.3
Vitamin E	Male	0	0.0	2	0.6	1	0.3	3	1.1	3	7.0	9	0.7
	Female	2	0.7	8	2.8	4	1.4	7	2.5	1	2.9	22	1.9
	Total	2	0.4	10	1.6	5	0.9	10	1.8	4	5.2	31	1.3
Vitamin K	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	1	0.4	0	0.0	5	1.8	0	0.0	6	0.5
	Total	0	0.0	1	0.2	0	0.0	6	1.1	0	0.0	7	0.3
Mixed vitamin	Male	5	1.9	4	1.2	7	2.4	17	6.4	4	9.3	37	3.1
	Female	11	3.7	7	2.5	17	5.9	26	9.5	4	11.8	65	5.5
	Total	16	2.9	11	1.8	24	4.1	43	7.9	8	10.4	102	4.3
Other vitamin	Male	9	3.4	10	3.0	14	4.7	10	3.7	1	2.3	44	3.7
	Female	9	3.1	7	2.5	9	3.1	9	3.3	0	0.0	34	2.9
	Total	18	3.2	17	2.8	23	3.9	19	3.5	1	1.3	78	3.3

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Calcium	Male	0	0.0	2	0.6	3	1.0	3	1.1	2	4.7	10	0.8
	Female	2	0.7	5	1.8	9	3.1	15	5.5	2	5.9	33	2.8
	Total	2	0.4	7	1.1	12	2.1	18	3.3	4	5.2	43	1.8
Mineral preparation	Male	0	0.0	1	0.3	0	0.0	1	0.4	0	0.0	2	0.2
	Female	4	1.4	2	0.7	2	0.7	6	2.2	2	5.9	16	1.4
	Total	4	0.7	3	0.5	2	0.3	7	1.3	2	2.6	18	0.8
Saccharide preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	1	0.1
	Total	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	1	0.0
Organic acid preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
protein or amino acid preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	1	0.3	1	0.4	0	0.0	0	0.0	0	0.0	2	0.2
	Total	1	0.2	1	0.2	0	0.0	0	0.0	0	0.0	2	0.1
Hormone drugs made from internal organ	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0
Drugs for infant	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other tonic	Male	10	3.8	6	1.8	7	2.4	6	2.2	3	7.0	32	2.7
	Female	3	1.0	4	1.4	10	3.5	8	2.9	2	5.9	27	2.3
	Total	13	2.3	10	1.6	17	2.9	14	2.6	5	6.5	59	2.5
Blood substitute	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Total	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	2	0.1
Styptic	Male	1	0.4	1	0.3	2	0.7	3	1.1	0	0.0	7	0.6
	Female	1	0.3	0	0.0	4	1.4	1	0.4	2	5.9	8	0.7
	Total	2	0.4	1	0.2	6	1.0	4	0.7	2	2.6	15	0.6
Anticoagulant	Male	0	0.0	2	0.6	2	0.7	8	3.0	0	0.0	12	1.0
	Female	0	0.0	2	0.7	3	1.0	1	0.4	0	0.0	6	0.5
	Total	0	0.0	4	0.6	5	0.9	9	1.7	0	0.0	18	0.8

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other blood substitute	Male	0	0.0	4	1.2	20	6.7	38	14.2	3	7.0	65	5.4
	Female	0	0.0	5	1.8	6	2.1	22	8.0	3	8.8	36	3.1
	Total	0	0.0	9	1.5	26	4.5	60	11.1	6	7.8	101	4.2
Agent for hemodialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for peritoneal dialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent for dialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for liver disease	Male	0	0.0	4	1.2	5	1.7	2	0.7	0	0.0	11	0.9
	Female	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Total	0	0.0	4	0.6	5	0.9	3	0.6	0	0.0	12	0.5
Antidote	Male	0	0.0	0	0.0	0	0.0	2	0.7	0	0.0	2	0.2
	Female	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Total	0	0.0	0	0.0	1	0.2	2	0.4	0	0.0	3	0.1
Agent for habitual intoxication	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Drugs for gout	Male	6	2.3	10	3.0	16	5.4	20	7.5	5	11.6	57	4.7
	Female	0	0.0	1	0.4	6	2.1	5	1.8	1	2.9	13	1.1
	Total	6	1.1	11	1.8	22	3.8	25	4.6	6	7.8	70	2.9
Enzyme preparation	Male	6	2.3	3	0.9	1	0.3	2	0.7	0	0.0	12	1.0
	Female	1	0.3	1	0.4	2	0.7	5	1.8	0	0.0	9	0.8
	Total	7	1.3	4	0.6	3	0.5	7	1.3	0	0.0	21	0.9
Antidiabetics	Male	2	0.8	14	4.2	22	7.4	27	10.1	2	4.7	67	5.6
	Female	1	0.3	11	3.9	12	4.2	13	4.7	1	2.9	38	3.2
	Total	3	0.5	25	4.1	34	5.8	40	7.4	3	3.9	105	4.4
Other agent on metabolic system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Unclassified agent on metabolic system	Male	1	0.4	1	0.3	2	0.7	12	4.5	0	0.0	16	1.3
	Female	1	0.3	8	2.8	5	1.7	16	5.8	4	11.8	34	2.9
	Total	2	0.4	9	1.5	7	1.2	28	5.2	4	5.2	50	2.1
Chlorophyll agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pigment agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other cell activation agent	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.0
Alkylating agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antimetabolite	Male	1	0.4	0	0.0	0	0.0	0	0.0	1	2.3	2	0.2
	Female	0	0.0	1	0.4	0	0.0	1	0.4	0	0.0	2	0.2
	Total	1	0.2	1	0.2	0	0.0	1	0.2	1	1.3	4	0.2
Antibiotics for antitumor agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Plant component for antitumor agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other antitumor agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	1	2.3	1	0.1
	Female	1	0.3	0	0.0	2	0.7	1	0.4	0	0.0	4	0.3
	Total	1	0.2	0	0.0	2	0.3	1	0.2	1	1.3	5	0.2
Antihistaminic drug	Male	5	1.9	6	1.8	7	2.4	2	0.7	0	0.0	20	1.7
	Female	5	1.7	6	2.1	5	1.7	2	0.7	1	2.9	19	1.6
	Total	10	1.8	12	1.9	12	2.1	4	0.7	1	1.3	39	1.6
Agent for stimulus therapy	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	0	0.0	1	0.4	1	0.3	1	0.4	0	0.0	3	0.3
	Total	0	0.0	1	0.2	2	0.3	1	0.2	0	0.0	4	0.2

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Agent for nonspecific immunogenicity	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other antiallergic drug	Male	16	6.0	11	3.3	14	4.7	13	4.9	3	7.0	57	4.7
	Female	15	5.1	11	3.9	15	5.2	14	5.1	2	5.9	57	4.9
	Total	31	5.5	22	3.6	29	5.0	27	5.0	5	6.5	114	4.8
Crude drug	Male	8	3.0	11	3.3	9	3.0	15	5.6	1	2.3	44	3.7
	Female	11	3.7	9	3.2	20	7.0	13	4.7	2	5.9	55	4.7
	Total	19	3.4	20	3.2	29	5.0	28	5.2	3	3.9	99	4.2
Herb medicine	Male	6	2.3	9	2.7	10	3.4	10	3.7	3	7.0	38	3.2
	Female	11	3.7	8	2.8	12	4.2	12	4.4	1	2.9	44	3.7
	Total	17	3.0	17	2.8	22	3.8	22	4.1	4	5.2	82	3.4
Other drugs by a herb medicine	Male	0	0.0	1	0.3	2	0.7	0	0.0	0	0.0	3	0.2
	Female	1	0.3	1	0.4	0	0.0	1	0.4	0	0.0	3	0.3
	Total	1	0.2	2	0.3	2	0.3	1	0.2	0	0.0	6	0.3
Antibiotics (G(+))	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(-))	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(+), G(-))	Male	6	2.3	9	2.7	7	2.4	1	0.4	0	0.0	23	1.9
	Female	8	2.7	6	2.1	9	3.1	4	1.5	0	0.0	27	2.3
	Total	14	2.5	15	2.4	16	2.7	5	0.9	0	0.0	50	2.1
Antibiotics (G(+), G(-), Mycoplasma)	Male	1	0.4	1	0.3	2	0.7	2	0.7	2	4.7	8	0.7
	Female	5	1.7	3	1.1	3	1.0	3	1.1	0	0.0	14	1.2
	Total	6	1.1	4	0.6	5	0.9	5	0.9	2	2.6	22	0.9
Antibiotics (Rickettsia, Chlamidia)	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	1	0.3	1	0.4	0	0.0	0	0.0	0	0.0	2	0.2
	Total	1	0.2	1	0.2	0	0.0	0	0.0	0	0.0	2	0.1
Antibiotics against acid-fast bacterium	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Antibiotics against fungus	Male	1	0.4	0	0.0	0	0.0	1	0.4	0	0.0	2	0.2
	Female	0	0.0	1	0.4	1	0.3	0	0.0	0	0.0	2	0.2
	Total	1	0.2	1	0.2	1	0.2	1	0.2	0	0.0	4	0.2
Other antibiotics	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Sulfa drug	Male	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Female	1	0.3	1	0.4	0	0.0	1	0.4	0	0.0	3	0.3
	Total	2	0.4	1	0.2	0	0.0	1	0.2	0	0.0	4	0.2
Antituberculous agent	Male	1	0.4	0	0.0	1	0.3	0	0.0	0	0.0	2	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	1	0.2	0	0.0	1	0.2	0	0.0	0	0.0	2	0.1
Drugs for leprosy	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Synthetic antibiotics	Male	5	1.9	1	0.3	5	1.7	0	0.0	0	0.0	11	0.9
	Female	2	0.7	2	0.7	1	0.3	4	1.5	0	0.0	9	0.8
	Total	7	1.3	3	0.5	6	1.0	4	0.7	0	0.0	20	0.8
Antiviral agent	Male	2	0.8	0	0.0	1	0.3	0	0.0	0	0.0	3	0.2
	Female	0	0.0	1	0.4	1	0.3	0	0.0	0	0.0	2	0.2
	Total	2	0.4	1	0.2	2	0.3	0	0.0	0	0.0	5	0.2
Other agent for chemotherapy	Male	5	1.9	3	0.9	3	1.0	1	0.4	1	2.3	13	1.1
	Female	0	0.0	1	0.4	0	0.0	2	0.7	0	0.0	3	0.3
	Total	5	0.9	4	0.6	3	0.5	3	0.6	1	1.3	16	0.7
Vaccine	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Poison or toxoid	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antitoxic serum or leptospira serum	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Blood supplements	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for biological test	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Mixed biological agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other biological agent	Male	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0
Protozoicide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Parasiticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other drug against parasitic disease	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Diluent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Ointment base	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0
Solution	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Corrective, tinction	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Emulsifier	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other dispensing drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
X-ray contrast medium	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for functional examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other test	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antiseptic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pesticide for prevention of epidemics	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Insecticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Insecticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Rodenticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other drug used for public health	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Reagent for general examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for blood analysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for chemical analysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for immunologic test	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for microbiological examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for pathological examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Radiopharmaceutical for diagnosis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other reagent for diagnosis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Strap	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other drug not for treatment	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Opiate alkaloid agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Coca alkaloid agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other alkaloid agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Synthetic narcotic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Unclassifiable drug	Male	1	0.4	1	0.3	3	1.0	1	0.4	0	0.0	6	0.5
	Female	3	1.0	0	0.0	3	1.0	6	2.2	0	0.0	12	1.0
	Total	4	0.7	1	0.2	6	1.0	7	1.3	0	0.0	18	0.8
No use of drug	Male	103	38.7	125	37.8	76	25.6	46	17.2	6	14.0	356	29.6
	Female	99	33.7	101	35.4	61	21.3	26	9.5	2	5.9	289	24.6
	Total	202	36.1	226	36.7	137	23.5	72	13.3	8	10.4	645	27.1
Total	Male	266	100.0	331	100.0	297	100.0	267	100.0	43	100.0	1204	100.0
	Female	294	100.0	285	100.0	286	100.0	275	100.0	34	100.0	1174	100.0
	Total	560	100.0	616	100.0	583	100.0	542	100.0	77	100.0	2378	100.0

IV. Nutritional Examinations

IV. Nutritional Examinations

Three days dietary record (3DR)

Foods and nutrition intakes were assessed by the three-consecutive-day weighing dietary records. On this method, foods and dietary supplements were weighed separately with scales before or after cooked. If the subjects did not weigh foods, they asked to record food size by portion. The subjects were asked to take the photos before and after eating foods using disposable camera. The dietary records were checked and coded by nutritionists. The average intakes of food and nutrition per day were calculated according to the 5th edition Standard Tables of Foods Consumption edited by the Science and Technology Agency in Japan and others. The substituted food composition table for Japanese foods were developed by National Cancer Center for the assessment of fatty acid intakes (Sasaki et al., *J Epidemiol*, 9: 190-207, 1999).

Definition and categories of dietary supplement in NILS-LSA

Dietary supplement were defined as any products intended for ingestion as a supplement to the diet from unnatural food forms such as capsules, tablets, or powders, but not from functional foods or modified foods. It contains vitamins, minerals, herbs, botanicals, and other substances such as enzymes, organ tissues, metabolites, concentrates, and constituent extracts of these substances.

Dietary supplement was grouped into seven major categories according to primary nutrient contents. It contains vitamin, mineral, fat, protein, dietary fiber, others, and drug (which contains some nutrients, for example a remedy for the cold which containing vitamin C.)

Dietary supplement frequency questionnaire (DSFQ)

Dietary supplement user was defined as a subject who used one or more dietary supplement one or more times in previous one year.

Three days dietary record

1) Food intake (3DR)

Cereals

Potatoes and starches

Sugars and sweeteners

Pulses

Nuts and seeds