

2004 00637 A

厚生労働科学研究費補助金  
エイズ対策研究事業

新作用機序の抗 HIV-1 薬剤の開発に関する研究

平成 16 年度 総括研究報告書

主任研究者 岡田 誠 治

平成 17 (2005) 年 3 月

## 目 次

I. 総括研究報告		
新作用機序の抗 HIV-1 薬剤の開発に関する研究	-----	1
岡田 誠 治		
II. 研究成果の刊行に関する一覧表	-----	5
III. 研究成果の刊行物・別刷	-----	7

## 新作用機序の抗 HIV-1 薬剤開発に関する研究

主任研究者 岡田 誠治 熊本大学エイズ学研究センター予防開発分野 教授

研究要旨 HIV-1 アクセサリー蛋白 Nef がマクロファージのチロシンキナーゼ Hck と会合することにより M-CSF の受容体経路を阻害するという新しい事実を見いだした。更に、この系を用いて Nef を標的とする新たな薬剤のスクリーニング系を開発した。また、免疫不全マウスにヒト臍帯血を移植することにより 4 週に渡ってヒトの免疫系が構築される系と臍帯血中の造血幹細胞を移植することにより長期に渡ってヒトの免疫系が構築される系を確立した。

分担研究者：鈴 伸也

熊本大学エイズ学研究センター予防開発分野 講師

### A. 研究目的

エイズの原因ウイルスであるヒト免疫不全ウイルス (HIV-1) は、年々その感染者が増加している。近年 HAART 療法のような有効な薬剤治療法が開発され、HIV-1 感染者の治療状況は大幅に改善された。しかし、現在の治療法では HIV-1 の完全な排除は不可能であり、薬剤の長期服用による副作用や薬剤耐性の出現など多くの問題が生じている。そのため、新たな作用機序を持つ薬剤開発は急務である。

HIV-1 アクセサリー蛋白 Nef はエイズ発症の病原性因子として良く知られている。近年、Nef と宿主細胞内チロシンリン酸化酵素 Hck との会合が発症に大きな役割を果たす事が明らかにされた。Hck は単球・マクロファージ特異的酵素である事から、Nef が Hck との会合を通して単球・マクロファージに機能変化をもたらす事が発症の引き金と予想されるが、その分子機序は不明である。この分子機序を解明し、その作用点を選択的に標的とする薬剤を見出すことにより新たな作用機序を持つ抗 HIV-1 薬を開発し、薬剤耐性ウイルスの克服に貢献する事が本研究の大きな目的である。先ず、Nef-Hck 会合により単球・マクロファージ内のシグナル伝達機構がどのように攪乱されるかを明らかにすると共に薬剤の *in vitro* 評

価系を確立する。特に、Hck はサイトカインシグナルの下流に於いても重要な働きを果たす分子である事から、Nef-Hck 会合のサイトカイン受容体経路に及ぼす影響に焦点を当て研究を行う。同時に、ヒト造血系・免疫系を構築したマウスを作成して、HIV-1 を感染させることにより、抗 HIV-1 薬の *in vivo* 評価系を確立する。

### B. 研究方法

Nef-Hck 会合のサイトカインシグナルに及ぼす影響を、特に、単球・マクロファージの増殖・分化・活性化を司る最も主要なサイトカインである M-CSF に着目して解析を行った。細胞株として TF-1-*fms* を用いた。TF-1-*fms* はヒト単球系の細胞であり、M-CSF に依存して増殖し、M-CSF とホルボールエステル TPA の共存在下でマクロファージへ分化する特徴を持つ。この TF-1-*fms* にエストロゲン受容体と融合させた Nef 蛋白質を発現させ、安定細胞株 (TF-1-*fms*-Nef-ER) を樹立した。この細胞株を用いて、合成エストロゲン 4-HT で Nef 活性化を誘導する事で Nef の M-CSF/M-CSF 受容体系に及ぼす影響を解析した。具体的には、M-CSF 反応性の TF-1-*fms* 細胞の増殖能を flow cytometry 法を用いて解析した。同時に、M-CSF/TPA によるマクロファージ分化に及ぼす Nef の影響を解析した。更に、M-CSF によるシグナル経路の活性化に及ぼす影響を種々のシグナル分子に対する抗体を用いた免疫沈降・Western

Blot 法により解析した。

ヒト造血系・免疫系を構築したマウスを作製する為に、NOD/Scid/ $\beta_2$  microglobulin 欠損マウス (Jackson 研究所) 及び NOD/Scid/Jak-3 欠損マウスへヒト臍帯血由来造血幹細胞移植を行った。NOD/Scid/Jak-3 欠損マウスは、NOD/Scid マウスに Jak-3 欠損マウス (理化学研究所 RCAI 斉藤隆博士から供与) をバッククロスして作成した。純化したヒト臍帯血中 CD34 陽性細胞を 2.5 Gy 放射線照射したマウスへ移植し、経時的に末梢血や種々のマウス臓器中におけるヒト血液細胞の出現を flow cytometry 法等により解析した。また、ヒト末梢血及び臍帯血単核球を放射線照射した NOD/Scid/ $\beta_2$  microglobulin 欠損マウスに移植し、短期間ヒトの免疫系が構築可能かどうかを検討し、更に作成されたマウスに HIV-1 実験室株 (JRFL) の感染を試みた。

#### (倫理面への配慮)

ヒト由来試料 (末梢血・臍帯血等) を用いた研究は、熊本大学大学院医学薬学研究部等倫理委員会の承認を受け、規則に従い実施している。また、免疫不全マウスの作成及び移植実験等の動物実験は、毎年熊本大学本荘地区動物実験委員会の承認を得た上で熊本大学動物実験指針に従い実施している。

#### 1) 研究対象者に対する人権擁護上の配慮

研究に用いる臍帯血・末梢血は、他の研究目的には使用しない。臍帯血・末梢血は匿名処理を行うため、個人情報流出することはない。また、同意書に署名後も試料採取・使用までの期間に同意を撤回することを可能としている。

#### 2) 研究方法による研究対象者に対する利益・不利益

本研究により、直接提供者が医学上の利益・不利益を得ることはない。

#### 3) 危険性の排除

臍帯血は臍帯を切り離した後で、臍帯・胎盤に残った血液を採取するため、新生児と母体への影響はない。また、臍帯血の採取は母子共に安全な分娩のみに限るとし、臍帯血採取によって分娩時の危険性が増す可能性を排除している。

末梢血は、医師が問診した上で健康に問題ないと判断した場合に限り、医師が採血している。採

血に伴う身体への危険性はあるが、これは通常の診療行為を越えるものではない。一回の採取量は 10-100 ml であり、採血量は、本人の了解のもとに決定している。

#### 4) インフォームドコンセントに係わる状況

臍帯血採取に関しては、協力医療機関の医療スタッフ (医師) が本研究の趣旨を説明し、臍帯血提供の同意を得られた方のみ同意書に署名していただいている。この際、説明を行った医療スタッフ名を明記し、同意書は協力医療機関において厳重に保管している。

末梢血採取に関しては、熊本大学エイズ学研究センターのスタッフ (医師) が本研究の趣旨を説明し、末梢血提供の同意を得られた方のみ同意書に署名していただいている。この際、説明を行った医師名を明記し、同意書はエイズ学研究センターにおいて厳重に保管している。

#### 5) 実験動物に対する動物愛護上の配慮

動物実験は、熊本大学動物実験指針を遵守し、極力動物の苦痛軽減に配慮して行っている。動物実験における実験処置に対する倫理基準では、カテゴリ B (動物に対してほとんど不快感を与えないと思われる実験) レベルの実験であり、解析時には頸椎脱臼により安楽死させている。

### C. 研究結果

Nef 活性化で TF-1-*fms*-Nef-ER 細胞の M-CSF 依存性増殖が著しく抑制された。この増殖抑制は早期アポトーシスを伴っていた。また、細胞形態の観察と付着細胞の出現率の測定から M-CSF/TPA によるマクロファージへの分化も大きく障害される事が明らかになった。これらの現象の原因として、Nef が、M-CSF によって引き起こされる最も早期の細胞内変化である M-CSF 受容体の活性化 (自己チロシンリン酸化) を強く障害する事が明らかとなった。この為、M-CSF 受容体活性化に引き続き起こる、受容体との種々のシグナル分子との会合も障害されていた。更に重要な事に、Nef によって活性化された Hck が恒常的に M-CSF 受容体と会合していた。Hck は通常 M-CSF によって活性化された M-CSF 受容体とのみ会合する分子である。従って以上の結果は、Nef が Hck を活性化し Hck と M-CSF 受容体の非生理的な会合をもたらす事で、M-CSF 受容体の活性化を障害し、その結果、M-CSF の生物活性を抑制したと考えられた。

また、上記の実験結果を元に Nef-Hck の会合を阻害する新規作用機序の抗 HIV-1 薬のスクリーニング法を確立した。この方法は、TF-1-*fms*-Nef-ER 細胞に候補となる薬剤を加えて培養し、4-HT を添加して Nef の活性化を誘導した際に、TF-1-*fms*-Nef-ER 細胞のアポトーシスを阻害するか否かで判定する方法である（特許出願中）。本法により、短期間に大量の低分子物質のスクリーニングが可能となった。

臍帯血単核球を移植した NOD/Scid/ $\beta_2$  microglobulin 欠損マウスでは、移植後 2-6 週に渡りヒト T 細胞の構築が認められた。また、HIV-1 実験室株 JRFL を感染させたところ、感染 2 週後にマウス末梢血において ELISA 法により p24 が同定されたことから、マウス体内で HIV-1 の感染・増殖が起きていることが示唆された。T 細胞の構築と HIV-1 の感染の程度は、末梢血単核球を移植した場合と同等であった。

一方、臍帯血由来 CD34 陽性ヒト造血幹細胞を移植した NOD/Scid/ $\beta_2$  microglobulin 欠損マウスと NOD/Scid/Jak-3 欠損マウスでは、移植後 20 週以上に渡りヒト由来の細胞による造血系の構築が確認された。マウス骨髄と脾臓において、ヒト由来の B 細胞と CD14 陽性の単球系細胞の安定的な生着が確認された。更に NOD/Scid/Jak-3 欠損マウスでは、T 細胞の出現も確認された。以上の結果より、長期に渡りヒト単球系の細胞を構築するマウスの作製が可能となった。

#### D. 考察

Nef-Hck の会合が Hck-M-CSF 受容体の恒常的会合という非生理的な変化をもたらす事で、M-CSF 受容体の活性化を障害し、結果として Nef が M-CSF の生物活性を抑制するという全く新しい事実を明らかにした。最近他のグループにより、単球・マクロファージに作用する、もう一つの主要なサイトカインである GM-CSF 受容体系には Nef-Hck は障害を与えない事が報告されている。これらの結果は、Nef-Hck 会合がサイトカイン系を異常制御する事で、単球・マクロファージに機能的な偏りを生じさせ、その事がエイズ発症に繋がる一因である可能性を示唆する。今後、その機能的偏りを規定する分子群の同定がエイズ発症の新たな機構を明らかにすると予想される。また、Nef-Hck 会合が M-CSF 系のみを阻害するという特異性は、本研究計画で目指す、高い選択性の新規作用機序の抗 HIV-1 薬開発が可能とするものである。

本年度の研究で新規作用機序の抗 HIV-1 薬剤スクリーニングのための、極めて簡便で且つ多検体をスクリーニング出来る *in vitro* 培養系は十分に確立された。現在、熊本大学薬学部等の協力で様々な骨格の低分子化合物の入手が出来ており、今後これら化合物の中からリード化合物の同定を行っていく。又、継続して、Nef-Hck 会合がサイトカイン受容体経路に及ぼす影響を更に詳細に明らかにしていく。

また、ヒトの造血・免疫系を構築したマウスは、最近国内外のいくつかの施設で報告されているが、造血幹細胞を移植したマウスにおいては、エイズのモデルとして使われているという報告は今のところ得られていない。今後これらのマウスに HIV-1 の感染を試みて、マウスモデルの作成を試みる予定である。このマウスは、上記研究で得られた研究成果を *in vivo* の系で確認するのみでなく、今後薬剤やワクチン等の治療法の開発や HIV-1 の病態解明に非常に有用なツールとなりうると考えられる。

#### E. 結論

HIV-1 Nef が、マクロファージのチロシンキナーゼ Hck と会合することにより M-CSF 受容体経路を阻害することによりマクロファージの機能を攪乱することを見いだした。そして、新規作用機序の抗 HIV-1 薬剤をスクリーニングする為の *in vitro* 培養系を確立し、当初の計画通り進める事が出来た。我々が見出した、Nef によるサイトカインシグナル伝達の攪乱作用 (Nef が Hck との結合・活性化を通して、単球・マクロファージ特異的サイトカイン M-CSF とその受容体の経路に障害を与える) は新しい知見であり、薬剤開発に最も適した分子標的のひとつであると考えられた。今後、より詳細な作用機構の解明を進めると同時に、薬剤のスクリーニングを行う。一方、*in vivo* 評価系の確立を目指したヒト造血幹細胞の移植による免疫不全マウスでのヒト単球系細胞の構築は、既に NOD/Scid/ $\beta_2$  microglobulin 欠損マウスと NOD/Scid/Jak-3 欠損マウスを用いることにより研究基盤を確立する事ができた。これらの *in vitro*、*in vivo* の系を有機的に駆使して、Nef をターゲットとした新規薬剤の開発とその評価系の確立を目指す予定である。

## F. 健康危機情報

該当なし

## G. 研究発表

### 1. 論文発表

1) Suzu. S., Harada H, Matsumoto, T., and Okada, S. HIV-1 Nef interferes with M-CSF receptor signaling through Hck activation and inhibits M-CSF bioactivities. *Blood*, in press.

### 2. 学会発表

1) Suzu. S., Harada H, and Okada, S. HIV-1 Nef Protein Interferes with M-CSF Receptor Signaling through Hck Activation and Inhibits M-CSF Bioactivities. 12<sup>th</sup> Conference on Retroviruses and Opportunistic Infections. 22-25 Feb. 2005, Boston, USA

2) Okada, S., and Suzu. S. HIV-1 Nef interferes with M-CSF receptor signaling through Hck activation and inhibits M-CSF bioactivities. 46<sup>th</sup> Annual Meeting of the American Society of Hematology. 4-7 Dec. 2004. San Diego USA.

3) 鈴木 伸也, 岡田誠治. M-CSF による細胞増殖と分化に必要な細胞内シグナルの比較解析. 第66回日本血液学会総会、2004年9月17-19日、京都

4) 松本貴博, 鈴木伸也, 岡田誠治. HIV-1 Nef は単球・マクロファージ特異的サイトカインである M-CSF とそのレセプター経路を阻害する. 第52回日本ウイルス学会学術集会、2004年11月21-23日、横浜

5) 後藤優美, 原田英樹, 鈴木伸也, 岡田誠治. ヒト臍帯血単核球を用いた HIV-1 感染マウスモデルの開発. 第18回日本エイズ学会学術集会、2004年12月9-11日、静岡 (優秀演題)

## H. 知的所有権の出願・取得状況 (予定を含む)

特許出願 2004-319531

発明者：岡田誠治、鈴木 伸也

抗 HIV-1 薬剤のスクリーニング系及びスクリーニング方法

## 研究成果の刊行に関する一覧表

### 書籍

著者名	論文タイトル名	書籍全体の編集者名	書籍名	出版社名	出版地	出版年	ページ

### 雑誌

発表者名	論文タイトル	発表誌名	巻号	ページ	出版年
鈴 伸也 岡田 誠治	HIV-1 Nef interferes with M-CSF Receptor signaling through Hck and Inhibits M-CSF bioactivities.	<i>Blood</i>		印刷中	





## HIV-1 Nef interferes with M-CSF receptor signaling through Hck activation and inhibits M-CSF bioactivities

Shinya Suzu, Hideki Harada, Takahiro Matsumoto, and Seiji Okada

HIV-1 Nef protein is a major determinant of the pathogenicity of the virus. It has been shown that Nef activates Hck, a member of Src family kinase, in monocytes/macrophages and that the interaction is critical for AIDS-like disease progression in a mouse model. However, it was unclear how the molecular interaction in monocytes/macrophages leads to disease progression. Here we show for the first time that Nef interferes with the macrophage-colony-stimulating factor (M-CSF)/M-CSF receptor pathway. In this

study, we introduced a conditionally active Nef into myeloid leukemia TF-1-*fms* cells and analyzed their responsiveness to M-CSF. We found that Nef-activated Hck constitutively associated with the M-CSF receptor complex. The formation of the molecular complex should occur under physiologic conditions, that is, on M-CSF stimulation. Because of aberrant molecular association, the tyrosine-phosphorylation/activation of the receptor in response to M-CSF was markedly diminished in Nef-active cells. Consequently,

Nef activation caused the inhibition of M-CSF-mediated proliferation of TF-1-*fms* cells and macrophage differentiation of the cells induced by M-CSF and 12-*O*-tetradecanoylphorbol 13-acetate. These results indicate that HIV-1 Nef interferes with M-CSF receptor signaling through Hck activation and thereby inhibits M-CSF functions in monocytes/macrophages. (Blood. 2005;105:000-000)

© 2005 by The American Society of Hematology

### Introduction

Nef is an accessory protein of HIV-1, a causative virus for AIDS. A number of reports, including studies of HIV-1-infected patients and of animal models, have demonstrated that the Nef protein is a major determinant of the pathogenicity of HIV-1.<sup>1-4</sup> Transgenic mice expressing the complete coding sequences of HIV-1 under the regulatory sequences of human CD4 gene developed severe AIDS-like abnormalities: loss of CD4<sup>+</sup> T cells, thymus atrophy, failure to thrive, diarrhea, wasting, premature death, interstitial pneumonitis, and tubulo-interstitial nephritis.<sup>4</sup> Using this mouse model and the introduction of mutation into selected HIV-1 gene(s), Hanna et al<sup>4</sup> clearly demonstrated that Nef harbored a major disease determinant. Therefore, much attention has been given to Nef to explain its contribution to HIV-1 pathogenesis and to investigate it as a target for antiviral drug development.

CD4<sup>+</sup> T cells and monocytes/macrophages are the principal target cells for HIV-1, and the functions of Nef in CD4<sup>+</sup> T cells are generally accepted as accounting for many aspects of viral pathogenesis (reviewed in Fackler and Baur<sup>5</sup> and Peterlin and Trono<sup>6</sup>). For example, Nef has been shown to cause the down-regulation of cell surface molecules such as cell surface receptor CD4<sup>7</sup> and major histocompatibility complex (MHC) class I in CD4<sup>+</sup> T cells.<sup>8</sup> The down-regulation of MHC class I is considered to diminish the recognition of HIV-1-infected cells by cytotoxic T cells. In contrast, the contribution of monocytes/macrophages to viral pathogenesis is less well understood. Several lines of evidence support the idea that monocytes/macrophages and CD4<sup>+</sup> T cells are important for the development and progression of AIDS. Recent studies have demonstrated that Nef induces the production of CC

chemokines (macrophage inflammatory proteins-1 $\alpha$  and -1 $\beta$ ) and soluble forms of CD23 and intracellular adhesion molecule-1 by macrophages.<sup>9,10</sup> These Nef-induced factors from macrophages might stimulate the chemotaxis and activation of resting CD4<sup>+</sup> T cells, thereby promoting the permissiveness of CD4<sup>+</sup> T cells to HIV-1 infection.<sup>9,10</sup> This Nef function is likely to be mediated by the activation of NF- $\kappa$ B transcription factor.<sup>10</sup>

Another important feature of Nef is the binding at high affinity to myeloid lineage-specific Src family kinase Hck to activate its kinase activity.<sup>11,12</sup> The proline-rich (PxxP) motif in Nef binds to the Src homology 3 (SH3) domain of Hck.<sup>11</sup> Interestingly, Hck was found to bind preferentially and with higher affinity to Nef than other Src kinases.<sup>11</sup> The pathologic relevance of the molecular interaction *in vivo* was revealed by studies of HIV-1 transgenic mice. The mutation in the SH3-binding motif of Nef abolished the development of AIDS-like disease in the HIV-1 transgenic mice.<sup>13</sup> Moreover, the breeding of transgenic mice expressing the complete coding sequences of HIV-1 on a *hck*<sup>-/-</sup> background resulted in the delay of disease development.<sup>13</sup> These studies suggest that the modulation of macrophage functions by Nef is important for disease development because the molecular interaction between Nef and Hck occurs in cells of macrophage lineage but not in CD4<sup>+</sup> T cells. However, it is unclear how the molecular interaction in monocytes/macrophages contributes to HIV-1 pathogenesis. To answer the question, it is necessary to identify intracellular signaling pathways that would be perturbed by the Nef-Hck interaction.

From the Division of Hematopoiesis, Center for AIDS Research, Kumamoto University, Japan.

Submitted June 3, 2004; accepted December 21, 2004. Prepublished online as *Blood* First Edition Paper, December 30, 2004; DOI 10.1182/blood-2004-06-2084.

Supported in part by a grant from Uehara Memorial Foundation and by Health and Labour Sciences Research Grants from the Ministry of Health, Labour and Welfare of Japan.

Reprints: Seiji Okada, Division of Hematopoiesis, Center for AIDS Research, Kumamoto University, Honjo 2-2-1, Kumamoto 860-0811, Japan; e-mail: okadas@kaiju.medic.kumamoto-u.ac.jp.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2005 by The American Society of Hematology

Macrophage-colony-stimulating factor (M-CSF) is a primary cytokine for monocytes/macrophages and regulates their development and various effector functions (reviewed in Roth and Stanley<sup>14</sup>). The importance of M-CSF in macrophage development *in vivo* has been clearly proven using a naturally occurring osteopetrotic (*op/op*) mouse.<sup>15</sup> The M-CSF biologic effects are mediated by a unique receptor encoded by the proto-oncogene *c-fms*, a receptor tyrosine kinase (reviewed in Bourette and Rohrschneider<sup>16</sup>). The binding of M-CSF induces the dimerization of the receptor and the activation of its kinase activity.<sup>16</sup> This leads to the autophosphorylation of specific tyrosine residues in the cytoplasmic domain and the subsequent interactions of the phosphorylated residues with other proteins, each initiating signaling along specific pathways.<sup>16</sup> The Src family is one of the well-documented partners for the activated M-CSF receptor. M-CSF stimulation results in the activation of Src kinases, including Hck; simultaneously, these kinases associate with the ligand-activated M-CSF receptor.<sup>17-21</sup> We and other investigators<sup>17-21</sup> have demonstrated that the activation of Src kinases is required for biologic effects of M-CSF such as mitogenic signal and differentiation-inducing signal. This finding prompted us to examine whether the molecular interaction between HIV-1 Nef and Hck would affect M-CSF receptor signaling and thereby modulate M-CSF bioactivities. In this paper, we demonstrate for the first time that Nef interferes with M-CSF-induced activation of its receptor, the earliest signaling event induced by the ligand, through Hck activation and thereby inhibits M-CSF functions on monocytes/macrophages.

## Materials and methods

### Cells and culture conditions

TF-1-*fms* cells were established by introducing the human M-CSF receptor (*c-fms*) gene into a human myeloid leukemia cell line, TF-1, as described previously.<sup>22</sup> Original TF-1 cells were growth factor-dependent cells,<sup>23</sup> and the proliferation of TF-1-*fms* cells was dependent on M-CSF.<sup>22</sup> In this study, we established a TF-1-*fms* clone expressing a fusion protein composed of Nef and murine estrogen receptor hormone-binding domain (Nef-ER) by introducing pEBB-Nef-ER-IRES-puro plasmid<sup>24</sup> using Effectene reagent (Qiagen, Valencia, CA). The *Nef* gene was derived from NL4-3 strain of HIV-1,<sup>24</sup> and its product has been shown to activate the kinase activity of Hck.<sup>11,12</sup> The transfection was performed according to the manufacturer's recommendations. Transfected cells were selected in medium containing 1.5  $\mu$ g/mL puromycin (Sigma, St Louis, MO), followed by limiting dilution to establish stable cell lines. The stable TF-1-*fms* clones were analyzed for Nef-ER expression by Western blotting using -Nef rabbit antiserum.<sup>25</sup> We also established a TF-1-*fms* clone expressing enhanced green fluorescence protein (EGFP). This was achieved with a retrovirus vector (MSCV2.2/IRES-EGFP),<sup>26</sup> as described previously.<sup>27</sup> In brief, the amphotropic packaging PT67 cells (Clontech, Palo Alto, CA) were transfected with MSCV2.2/IRES-EGFP vector using LipofectAMINE2000 reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's recommendations. Then, TF-1-*fms* cells were cocultured with the transfected PT67 cells in the presence of 8  $\mu$ g/mL Polybrene (Sigma). The stable TF-1-*fms* clones were obtained by limiting dilution and were analyzed for EGFP expression by flow cytometry. Parental TF-1-*fms* cells and EGFP-expressing cells were maintained with RPMI 1640 (Sigma)-10% fetal calf serum (FCS) in the presence of recombinant human M-CSF (100 ng/mL)<sup>22</sup> and 200  $\mu$ g/mL G418 (Life Technologies, Grand Island, NY). Nef-ER-expressing cells were maintained with RPMI 1640-10% FCS in the presence of M-CSF, G418, and puromycin. To activate the Nef-ER protein, an estrogen analog, 4-hydroxytamoxifen (4-HT; Sigma), was added to the cultures.<sup>24</sup> Unless otherwise stated, 4-HT was used at a final concentration of 1  $\mu$ M. To induce the differentiation of TF-1-*fms*-Nef-ER cells into

macrophages, cells were cultured with M-CSF and 100 ng/mL 12-*O*-tetradecanoylphorbol 13-acetate (TPA; Sigma).

### Cell count and flow cytometry

Viable cell counts were obtained by enumerating cells that excluded trypan blue dye. Adherent cells were harvested by trypsinization.

To detect apoptotic cells, the flow cytometric analysis of propidium iodide (PI)-stained nuclei was performed as described previously.<sup>28</sup> In brief, cells were washed with phosphate-buffered saline (PBS) and then incubated in hypotonic lysing buffer (0.1% sodium citrate, 0.3% Nonidet P-40, and 50  $\mu$ g/mL PI) at 4°C for 30 minutes. Alternatively, cells were stained with annexin V conjugated to phycoerythrin (PE) (PharMingen, San Jose, CA) and vital dye 7-aminoactinomycin D (7-AAD; PharMingen). Cultured cells were washed with PBS, resuspended in binding buffer (10 mM HEPES [N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid], 140 mM NaCl, and 2.5 mM CaCl<sub>2</sub>), and incubated with annexin V-PE and 7-AAD (1.25  $\mu$ g/mL) at room temperature for 15 minutes. Flow cytometry analyses were performed with FACSCalibur using Cell Quest Software (Becton Dickinson, Mountain View, CA).

In this study, we analyzed the expression level of M-CSF receptor on TF-1-*fms*-Nef-ER cells by flow cytometry using Flag-tagged M-CSF protein. The Flag (DYKDDDDK) sequence was introduced into the C-terminus of mature M-CSF polypeptides<sup>22</sup> by polymerase chain reaction (PCR) amplification with an upstream primer (5'-TGCCGGGACCCAGCTGCCCGTATGA-3') and a downstream primer (5'-CTCACTTGTGTCATCGTCTTTGTAGTCTCGGCTAGAGCA-3'). Codons specifying the Flag sequence are underscored. The PCR product was subcloned into an expression vector, pEF-BOS.<sup>29</sup> The resultant plasmid was transfected into COS7 cells using LipofectAMINE2000 reagent (Invitrogen),<sup>30</sup> and the culture supernatant was used as a source of Flag-tagged M-CSF. For analysis, cells were incubated with the Flag-tagged M-CSF, followed by biotin-labeled anti-Flag M2 antibody (10  $\mu$ g/mL; Sigma) and PE-labeled streptavidin (5  $\mu$ g/mL; PharMingen).<sup>31</sup>

### Immunoprecipitation and Western blotting

Immunoprecipitation and Western blotting were performed essentially as described previously.<sup>17</sup> Cells were growth factor depleted for 14 hours in RPMI 1640-10% FCS and then stimulated with M-CSF (100 ng/mL) for the indicated periods. For the activation of Nef, 4-HT (1  $\mu$ M) was added to the culture at the initiation of M-CSF deprivation. Then cells were solubilized with Nonidet P-40 lysis buffer (1% Nonidet P-40, 50 mM Tris HCl, 150 mM NaCl) containing protease inhibitors (1 mM EDTA [ethylenediaminetetraacetic acid], 1 mM phenylmethylsulfonyl fluoride, 1  $\mu$ g/mL aprotinin, 1  $\mu$ g/mL leupeptin, and 1  $\mu$ g/mL pepstatin) and phosphatase inhibitors (1 mM Na<sub>3</sub>VO<sub>4</sub> and 1 mM NaF). Cell lysates were centrifuged, and the resultant supernatants were subjected to Western blotting or immunoprecipitation. Antibodies (purchased from Santa Cruz Biotechnology, Santa Cruz, CA) used for Western blotting were as follows: anti-M-CSF receptor rabbit immunoglobulin G (IgG) (C-20), antiphosphotyrosine mouse IgG (PY99), -ERK rabbit IgG (K-23), antiphosphorylated ERK mouse IgG (E-4), and anti-Hck rabbit IgG (N-30). Anti-Hck mouse IgG (clone 18; Transduction Laboratories, Lexington, KY) was also used. Antibodies used for immunoprecipitation were as follows: anti-M-CSF receptor rat IgG (12-2D6; Zymed, South San Francisco, CA), anti-Hck rabbit IgG (Santa Cruz Biotechnology), and antiphosphotyrosine mouse IgG (PY99; Santa Cruz Biotechnology). Immune complexes were precipitated with Protein A/G PLUS-Agarose (Santa Cruz Biotechnology). Cleared cell lysates and immunoprecipitates were resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) under reducing conditions, and proteins were transferred to a nylon membrane (Hybond-P; Amersham, Buckinghamshire, United Kingdom). Detection was performed using the Enhanced Chemiluminescence Western Blotting Detection System (Amersham) with horseradish peroxidase-coupled anti-immunoglobulin (Amersham).

AQ: 2

AQ: 3

**Expression of M-CSF receptor, Nef, and Hck in 293T cells**

Expression plasmid for human M-CSF receptor was prepared as described previously.<sup>22</sup> The plasmids for Nef and human Hck were provided by E. T. Sawai<sup>32</sup> and Y. Murakami,<sup>33</sup> respectively. The *Nef* gene was derived from the SF2 strain of HIV-1,<sup>32</sup> and its product has been shown to activate the kinase activity of Hck.<sup>34</sup> The human embryonic kidney cell line 293T was obtained from the American Type Culture Collection (Manassas, VA) and was maintained with Dulbecco modified Eagle medium (DMEM; Sigma)–10% FCS. Cells were grown on a 6-well tissue culture plate and were transfected with plasmids for M-CSF receptor (0.4 μg), Nef (0.8 μg) and Hck (0.4 μg), using LipofectAMINE2000 reagent (Invitrogen). After transfection, the cells were cultured in DMEM–10% FCS for 36 hours and then cultured in DMEM–0.1% FCS for an additional 12 hours. In a selected experiment, cells were stimulated with M-CSF for 2 minutes. Cells were solubilized with Nonidet P-40 lysis buffer, and the cleared cell lysates were subjected to immunoprecipitation with anti-M-CSF receptor rat IgG (Zymed) or anti-Hck rabbit IgG (Santa Cruz Biotechnology). Immunoprecipitates were analyzed by Western blotting with antiphosphotyrosine mouse IgG (Santa Cruz Biotechnology), anti-M-CSF receptor rabbit IgG (Santa Cruz Biotechnology), or anti-Hck mouse IgG (Transduction Laboratories).

AQ: 4


**Results**

**Inhibition of M-CSF-mediated cell proliferation and differentiation by Nef**

We initially attempted to determine whether Nef would affect the biologic effects of M-CSF. To achieve this, we used the human myeloid leukemia cell line TF-1-*fms*, the proliferation of which was dependent on M-CSF.<sup>22</sup> We established the TF-1-*fms* clone expressing Nef-murine estrogen receptor hormone-binding domain fusion protein (TF-1-*fms*-Nef-ER) (Figure 1A). The Nef-ER fusion protein could be seen as a 60-kDa band on Western blot (Figure 1B). In this system, Nef was basally inactive, but its function could be inducibly activated by the estrogen analog 4-HT.<sup>24</sup> As shown in Figure 1C, we found that the proliferation of TF-1-*fms*-Nef-ER cells mediated by M-CSF was markedly inhibited on the addition of 4-HT (ie, the activation of Nef). The inhibitory effect was dependent on the concentrations of 4-HT added to the cultures and was specific because 4-HT never affected the proliferation rate of the parental TF-1-*fms* cells (Figure 1C). Furthermore, we found that Nef activation was associated with the appearance of apoptotic subdiploid cells in the culture of TF-1-*fms*-Nef-ER cells (Figure 1D). In parallel, Nef activation increased the percentage of annexin V–positive but 7-AAD–negative cells in the culture of TF-1-*fms*-Nef-ER cells (Figure 1E). These findings indicated that Nef induced apoptotic cell death in TF-1-*fms*-Nef-ER cells.

We next attempted to determine whether Nef also could inhibit another biologic activity of M-CSF. The phorbol ester TPA is well known to induce macrophage differentiation of various myeloid leukemia cell lines.<sup>23</sup> Recently, we found that, in the presence of TPA, M-CSF markedly stimulates the differentiation of TF-1-*fms* cells into mature macrophages but that it does not stimulate the proliferation of the cells (S. Suzu et al, unpublished results). This was based on several observations. First, the combination of TPA and M-CSF caused more drastic morphologic changes in TF-1-*fms* cells than did treatment with TPA alone. Second, culture in the presence of TPA and M-CSF contained more adherent cells than in the presence of TPA alone. Third, the phagocytic activity of cells treated with TPA and M-CSF was significantly higher than that of cells treated with TPA alone. Using this culture system, we

AQ: 6



**Figure 1. TF-1-*fms* cells expressing Nef-ER show impaired proliferation and apoptotic cell death on Nef activation.** (A) Schematic diagram of the Nef-ER-IRES-puro construct. ER indicates estrogen receptor hormone-binding domain; IRES, internal ribosomal entry sequence; puro, puromycin resistance gene. (B) Total cell lysates from the parental TF-1-*fms* cells (parent) or the TF-1-*fms* clone stably expressing Nef-ER (Nef-ER) were analyzed for the expression of Nef-ER by Western blotting with anti-Nef rabbit antiserum. (C) Parental TF-1-*fms* (□) or TF-1-*fms*-Nef-ER cells (▨) were seeded at a density of  $1 \times 10^4$  cells/mL in the presence of M-CSF (100 ng/mL) and increasing concentrations of 4-HT. Cells were cultured for 2 days (top row) or 4 days (bottom row), and viable cells were enumerated. Error bars from triplicate assays are shown. Results are representative of 3 independent experiments. (D) TF-1-*fms*-Nef-ER cells were cultured with M-CSF in the absence (top) or the presence of 1 μM 4-HT (bottom) for 48 hours. Apoptotic subdiploid cells was detected by flow cytometry. (E) Parental TF-1-*fms* (top row) or TF-1-*fms*-Nef-ER cells (bottom row) were cultured with M-CSF in the absence (left column) or the presence of 1 μM 4-HT (right column) for 24 hours. Cells were analyzed for the presence of apoptotic cells by staining with annexin V-PE, and 7-AAD (lower right quadrant).

AQ: 7

AQ: 8

AQ: 8


examined the effect of Nef on the macrophage differentiation-inducing activity of M-CSF. As shown, the morphologic changes in TF-1-*fms*-Nef-ER cells induced by TPA and M-CSF were markedly inhibited on the activation of Nef (Figure 2A). Furthermore, the number of cells adhering to the dishes in the 4-HT-containing culture was lower than that in the 4-HT-free culture (Figure 2B). The inhibitory effect was dependent on the concentration of 4-HT added to the cultures (Figure 2C).

F2

**Inhibitory effect of Nef on M-CSF bioactivity is not mediated by soluble factors or down-regulation of M-CSF receptor**

The inhibitory effects of Nef on M-CSF-induced cell proliferation (Figure 1) and macrophage differentiation (Figure 2) suggested that Nef affected signaling pathways transduced by M-CSF. However, because Nef has been shown to induce the expression of a number of soluble factors (chemokines and cytokines) in monocytes/macrophages,<sup>9,10,35-37</sup> the inhibitory effects of Nef observed in our culture systems might be mediated through the secretion of inhibitory molecule(s) by Nef rather than through a direct effect on the M-CSF receptor signaling pathways. To address this issue, we conducted a coculture experiment with parental TF-1-*fms* cells and TF-1-*fms*-Nef-ER cells. Parental cells were engineered to express EGFP protein, allowing us to distinguish parental cells from Nef-ER cells in the cocultures through flow cytometric analysis. As shown in Figure 3A, the relative cell number of Nef-ER cells was


F3




**Figure 2. Nef activation causes inhibition in macrophage differentiation.** (A) Morphologies of TF-1-*fms*-Nef-ER cells cultured for 24 hours with M-CSF (100 ng/mL), M-CSF/TPA (100 ng/mL), or M-CSF/TPA/4-HT (1  $\mu$ M), as indicated in each case. (B) TF-1-*fms*-Nef-ER cells were seeded into 6-well culture plates at a density of  $2 \times 10^5$  cells/well. Cells were cultured in the presence of M-CSF, M-CSF/TPA, or M-CSF/TPA/4-HT for 12 hours or 24 hours. After culture, total cells in the wells (□) and cells that adhered to the dishes (■) were enumerated. (C) TF-1-*fms*-Nef-ER cells were seeded as in panel B. Cells were cultured for 12 hours in the presence of M-CSF, TPA, and increasing concentrations of 4-HT. Cells adhering to the dishes (●) or remaining in suspension (○) were enumerated. (B-C) Error bars from triplicate assays are shown. Results are representative of 3 independent experiments.

comparable to that of the parental cells in the absence of 4-HT (left panel) but was markedly low in the presence of 4-HT (right panel). In fact, the absolute cell number of Nef-ER cells in 4-HT-containing cocultures was reduced to approximately 10% when compared with that in 4-HT-free cocultures (Figure 3B). However, and of importance, the absolute cell numbers of the parental cells remained unchanged between 4-HT-containing cocultures and 4-HT-free cocultures (Figure 3B). These results suggested that soluble factors from Nef-active cells, if any, were not involved in the proliferation inhibitory effect of Nef observed in our culture system.

Another well-known function of Nef is the down-regulation of cell surface molecules such as CD4 and MHC class I.<sup>7,8</sup> Therefore, we next examined whether Nef activation caused the down-regulation of the receptor for M-CSF in TF-1-*fms*-Nef-ER cells. As shown by the culture in Figure 4A, on which Western blotting analysis was performed, we did not find any change in the expression level of M-CSF receptor, even when TF-1-*fms*-Nef-ER cells were cultured for up to 2 days in the presence of 4-HT. Flow cytometric analysis with the Flag-tagged M-CSF also revealed that



**Figure 3. Secretion of soluble factors is not involved in the impaired proliferation of TF-1-*fms*-Nef-ER cells induced by Nef activation.** Parental TF-1-*fms* cells, but not TF-1-*fms*-Nef-ER cells, were engineered to express EGFP protein by the retroviral infection system. The cell density of the TF-1-*fms*-EGFP (parent-EGFP) and TF-1-*fms*-Nef-ER (Nef-ER) cells was adjusted to  $1 \times 10^4$  cells/mL, and the cells were cultured after an equal volume of the cell suspensions was combined. These cultures were incubated for 3 days with M-CSF in the presence or the absence of 4-HT. (A) After the cocultures, cells were subjected to flow cytometric analysis to determine the relative cell numbers. (B) Absolute cell numbers of the parental cells and Nef-ER cells were calculated from the relative cell numbers (A) and the total cell numbers in the cocultures. Error bars from triplicate assays are shown. These results are representative of 3 independent experiments.



**Figure 4. Down-regulation of M-CSF receptor is not involved in the impaired proliferation of TF-1-*fms*-Nef-ER cells induced by Nef activation.** (A) Parental TF-1-*fms* and TF-1-*fms*-Nef-ER cells were cultured with 4-HT for the indicated periods, and the levels of M-CSF receptor proteins were analyzed by Western blotting. Total cell lysates from TF-1 cells, but not from TF-1-*fms* cells, in which the M-CSF receptor gene had not been introduced, were included in the analysis as a negative control (*fms*-negative). (B) TF-1-*fms*-Nef-ER cells were left untreated (top) or were treated with 4-HT for 24 hours (bottom), and the level of cell surface M-CSF receptor expression was analyzed by flow cytometry with Flag-tagged M-CSF (solid lines). Profiles of cells incubated with a Flag-tagged protein,<sup>30</sup> unrelated to M-CSF, are also shown as a control (broken lines).


the level of cell surface expression of M-CSF receptor in TF-1-*fms*-Nef-ER cells treated for 24 hours with 4-HT was comparable with that in Nef-inactive cells (Figure 4B). Taken together with the result of the coculture experiment (Figure 3), these results strongly suggested that Nef affected intrinsic M-CSF receptor signaling pathways and thereby inhibited M-CSF activities.

**Mechanism by which Nef inhibits M-CSF receptor signaling**

Because M-CSF receptor is a receptor tyrosine kinase,<sup>14,16</sup> we initially compared the M-CSF-induced protein tyrosine phosphorylation in Nef-active TF-1-*fms*-Nef-ER cells with that in Nef-inactive cells. On M-CSF stimulation, a number of signaling molecules, such as proteins, with molecular weights of 150 kDa to 160 kDa, 100 kDa, and 60 kDa were shown to be rapidly tyrosine phosphorylated in Nef-inactive cells (Figure 5A, open arrowheads). However, the tyrosine phosphorylation of these proteins in response to M-CSF stimulation was markedly diminished in Nef-active cells (Figure 5A). Yet the blot also showed that the effect of Nef was not an overall reduction in tyrosine phosphorylation. The proteins, seen as a closely spaced doublet at approximately 55 kDa, were phosphorylated only on M-CSF stimulation in Nef-inactive cells but were phosphorylated before stimulation in Nef-active cells (Figure 5A, asterisks). Moreover, the tyrosine phosphorylation of protein, seen at 40 to 45 kDa, was elevated in Nef-active cells (Figure 5A, filled arrowhead). Judging by its molecular weight, the latter protein (40-45 kDa, filled arrowhead) seemed to be extracellular signal-related protein kinase (ERK).<sup>17</sup> This was confirmed by further analyses in which total cell lysates were immunoprecipitated with antiphosphotyrosine antibody and then were probed with -ERK antibody or in which total cell lysates were directly probed with phosphorylated ERK-specific antibody (Figure 5B). Using a similar approach, we identified the protein seen at 150 to 160 kDa, the tyrosine phosphorylation of which was induced by M-CSF stimulation but was diminished in Nef-active cells (Figure 5A, top open arrowhead) as M-CSF receptor (Figure 5C, top and middle panels). The blot shown in the middle panel also revealed that the 60-kDa protein (Figure 5A, bottom open arrowhead) was a component of M-CSF receptor complex, and its association with the complex was markedly diminished in Nef-active cells (Figure 5C, open arrowhead). These results indicated

F4

AQ: 7




**Figure 5.** Nef activation causes perturbations in M-CSF receptor signaling in TF-1-*fms*-Nef-ER cells. (A-C) TF-1-*fms*-Nef-ER cells were deprived of M-CSF for 14 hours and restimulated with M-CSF for the indicated periods. For the activation of Nef, 4-HT was added to the culture at the initiation of M-CSF deprivation. Total cell lysates or immunoprecipitates were analyzed by Western blotting. (A) Total cell lysates from untreated or 4-HT-treated cells were analyzed with antiphosphotyrosine (pTyr) antibody. See "Mechanism by which Nef inhibits M-CSF receptor signaling" for explanations of the symbols in detail. (B) Immunoprecipitates with -pTyr antibody were analyzed with -ERK antibody (top). Alternatively, total cell lysates were analyzed with antibody specific for phosphorylated ERK (middle). The blot shown in the bottom panel, in which the total cell lysates were analyzed with -ERK antibody, verified that comparable amounts of proteins were loaded in the top panel. (C) Immunoprecipitates with -pTyr antibody were analyzed with anti-M-CSF receptor (*c-fms*) antibody (top). Alternatively, the immunoprecipitates with anti-M-CSF receptor antibody were analyzed with -pTyr antibody (middle). The blot was exposed to autoradiography film for 10 seconds (short) or 1 minute (long). See "Mechanism by which Nef inhibits M-CSF receptor signaling" for explanations of the symbols in detail. (D) TF-1-*fms*-Nef-ER cells were cultured with or without 4-HT for 14 hours under M-CSF-free conditions. Total cell lysates were prepared and subjected to immunoprecipitation with anti-M-CSF receptor (*c-fms*) antibody or anti-Hck antibody. Anti-M-CSF receptor immunoprecipitates were analyzed with anti-Hck antibody or anti-M-CSF receptor antibody. Anti-Hck immunoprecipitates were analyzed with anti-pTyr antibody, anti-M-CSF receptor antibody, or anti-Hck antibody.

that Nef activation caused inhibition in the tyrosine phosphorylation/activation of M-CSF receptor and the molecular association with the activated receptor, which was the earliest signaling event induced by M-CSF.

We next examined whether the activation of Hck by Nef was involved in the inhibitory effect of Nef on M-CSF receptor signaling. We focused attention on the proteins seen as a closely spaced doublet at approximately 55 kDa. The pattern in tyrosine phosphorylation was different between Nef-inactive and Nef-active cells—the proteins were tyrosine phosphorylated in response to M-CSF stimulation in Nef-inactive cells but were phosphorylated before stimulation in Nef-active cells (Figure 5A, asterisks). Moreover, and of interest, the proteins were recruited to the complex of M-CSF receptor only after M-CSF stimulation in Nef-inactive cells but preexisted in the complex in Nef-active cells irrespective of M-CSF stimulation (Figure 5C, asterisks). We confirmed that the proteins were Hck. When total cell lysates were prepared from cells that were cultured without M-CSF, immunoprecipitated with anti-M-CSF receptor antibody, and analyzed for the

presence of Hck protein in the M-CSF receptor complex, we found that Hck was present in the complex in Nef-active cells but not in Nef-inactive cells (Figure 5D). Analyses of anti-Hck immunoprecipitates further revealed the tyrosine phosphorylation of Hck and its association with M-CSF receptor complex in Nef-active cells (Figure 5D). Given that Src family kinases, including Hck, should associate with M-CSF receptor only if the receptor is activated by its ligand,<sup>18-20</sup> the constitutive association of Hck with M-CSF receptor complex that occurred in Nef-active TF-1-*fms*-Nef-ER cells was an aberrant interaction.

Finally, we performed transient expression experiments with 293T cells and confirmed that Nef induced the activation of Hck, the aberrant molecular interaction between Hck and M-CSF receptor, and the impaired M-CSF receptor activation. Tyrosine phosphorylation of Hck in 293T cells was enhanced by cotransfection with Nef (Figure 6A, top panel), possibly reflecting the finding that Nef stimulated the autophosphorylation of Hck.<sup>12,33,34,38</sup> The association between Hck and M-CSF receptor, in the absence of M-CSF stimulation, was stimulated by cotransfection with Nef (Figure 6A, middle panel; Figure 6B, upper panel). The tyrosine phosphorylation of M-CSF receptor in response to M-CSF was impaired in 293T cells coexpressing M-CSF receptor, Nef, and Hck (Figure 6C, upper panel). Such impairment was not observed in 293T cells coexpressing M-CSF receptor and Nef or M-CSF receptor and Hck (Figure 6C, upper panel). In summary, our data strongly suggested that HIV-1 Nef interfered with the activation of M-CSF receptor in response to its ligand by means of the activation of Hck and the subsequent induction of an aberrant molecular association between Hck and M-CSF receptor, thereby inhibiting M-CSF bioactivities.



**Figure 6.** Nef causes tyrosine phosphorylation of Hck, constitutive association of Hck with M-CSF receptor complex, and inhibition in tyrosine phosphorylation of M-CSF receptor in 293T cells. (A-C) 293T cells were transfected with M-CSF receptor (*c-fms*) expression plasmid, alone or in combination with Nef or Hck expression plasmids. Total cell lysates were prepared and subjected to immunoprecipitation/Western blotting analyses. (C) Transfected 293T cells were treated in the absence or presence of M-CSF for 2 minutes before lysis. (A) Immunoprecipitates with anti-Hck antibody were analyzed with antiphosphotyrosine (pTyr) antibody (top) or anti-M-CSF receptor antibody (middle). The blot shown in the bottom panel, in which the immunoprecipitates were analyzed with anti-Hck antibody, is a loading control experiment for upper panels. (B) Immunoprecipitates with anti-M-CSF receptor antibody were analyzed with anti-Hck antibody (top). The blot shown in the bottom panel, in which the immunoprecipitates were analyzed with anti-M-CSF receptor antibody, is a loading control experiment for the top panel. (C) Immunoprecipitates with anti-M-CSF receptor antibody were analyzed with anti-pTyr antibody (top). The blot shown in the bottom panel, in which the immunoprecipitates were analyzed with anti-M-CSF receptor antibody, is a loading control experiment for the top panel.

## Discussion

The present study identified for the first time a dysregulated signaling event in monocytes/macrophages induced by the interaction between HIV-1 Nef and the cellular kinase Hck. We have demonstrated that Nef activates Hck and recruits the activated Hck to the M-CSF receptor complex. By means of the induction of the unphysiologic signaling event, Nef interferes with M-CSF receptor signaling pathways, thereby inhibiting M-CSF bioactivities and possibly causing dysregulated functions of HIV-1-infected monocytes/macrophages.

Since Hanna et al<sup>13</sup> reported the importance of the interaction between Nef and Hck for the development of AIDS-like disease through studies with HIV-1 transgenic mice and Hck knockout mice, efforts have been made to identify the signaling molecules for which activation is affected by Nef-Hck interaction in monocytes/macrophages. Signal transducer and activator of transcription (STAT) family molecules were considered possible candidates for the Nef-Hck complex. A recent study reported that Nef induced the activation of STAT3 in a myeloid cell line.<sup>34</sup> However, it was not established whether the Nef-Hck interaction directly led to STAT3 activation.<sup>34</sup> Similarly, another recent study reported that Nef activated STAT-1 $\alpha$  and -1 $\beta$  in human macrophages derived from peripheral blood monocytes, but the activation occurred through the secretion of unidentified soluble factors.<sup>36</sup> Thus, the STAT family molecules may not be direct targets for the Nef-Hck complex. The secretion of soluble factors, such as cytokines and chemokines, from monocytes/macrophages is a well-established function of Nef,<sup>9,10,35-37</sup> and macrophage inflammatory proteins 1 $\alpha$  and 1 $\beta$  are well-known examples, as reported by Swingle et al.<sup>9</sup> Interestingly, the possibility that the Nef-Hck interaction did not participate in the secretion of these chemokines was raised by the authors.<sup>9</sup> After all, in spite of its importance in disease development, the role of the Nef-Hck interaction in the modulation of monocyte/macrophage signaling remains to be determined.

Our present study strongly suggests that M-CSF receptor is a target for the Nef-Hck complex. We demonstrated that Nef caused the inhibition of the activation of M-CSF receptor in response to M-CSF and the molecular association with the activated receptor, which were the earliest signaling events for M-CSF receptor pathways (Figures 5A, C, 6C). As an inevitable consequence, Nef caused the inhibition of biologic activities of M-CSF, such as cell proliferation (Figure 1) and macrophage differentiation (Figure 2) of TF-1-*fms* cells. We observed a similar effect of Nef in primary myeloid cells. When cord blood CD34<sup>+</sup> cells were retrovirally transduced with the *Nef* gene and then cultured with M-CSF, we found a marked decrease in the proportion of Nef-positive cells during the culture with M-CSF (data not shown). Although further analyses are necessary, the finding raises the possibility that Nef does inhibit M-CSF receptor signaling in primary myeloid cells. The proliferation inhibitory effect of Nef was not mediated by the secretion of inhibitory molecules (Figure 3) or by the down-regulation of M-CSF receptor (Figure 4), further supporting the conclusion that Nef affected intrinsic M-CSF receptor signaling. The most likely mechanism by which Nef interferes with M-CSF receptor signaling is the unphysiologic behavior of Hck. The Hck proteins seem to be constitutively phosphorylated in Nef-active TF-1-*fms*-Nef-ER cells (Figure 5A, C). The induction of tyrosine phosphorylation of Hck by Nef was confirmed by transient expression experiments in 293T cells (Figure 6A). More important, Hck constitutively formed the complex with M-CSF receptor in

active Nef-expressing cells (Figures 5D, 6A, B). In contrast, Hck associates with M-CSF receptor only if the receptor is activated by its ligand (Figure 5C, D).<sup>18-20</sup> It is reasonable to expect that the aberrant constitutive association of Nef-activated Hck to M-CSF receptor complex would inhibit the conformational change of the receptor that would be induced by its ligand and would be required for initiating signaling cascades. Therefore, we concluded that Nef interfered with M-CSF receptor signaling through the activation of Hck and its recruitment to M-CSF receptor complex.

Given that Nef activates the kinase activity of Hck<sup>11,12</sup> and that the activation of Src kinases, including Hck, is required for biologic effects of M-CSF such as mitogenic signal and differentiation-inducing signal,<sup>17-21</sup> one may speculate that Nef mimics the effect of M-CSF on monocytes/macrophages. However, this is not the case. Nef caused the inhibition of biologic activities of M-CSF such as cell proliferation (Figure 1) and macrophage differentiation (Figure 2). In addition, the activation of Nef did not cause M-CSF-independent proliferation of TF-1-*fms*-Nef-ER cells (data not shown). The reason Nef-activated Hck does not mimic M-CSF functions might be explained by the difference in the mode of Hck activation. The activity of Src family kinases, including Hck, is suppressed by 2 intramolecular interactions, SH2-tail and SH3-linker.<sup>39,40</sup> Nef has been shown to activate Hck by binding to its SH3 domain, thus preventing the SH3-linker interaction.<sup>12,38</sup> In contrast, the ligand-activated M-CSF receptor has been shown to associate with the SH2 domain but not with the SH3 domain of Src kinases.<sup>19</sup> It is reasonable to expect that the different modes of Hck activation might generate distinct output signals. That Nef-activated Hck formed an unphysiologic molecular association with M-CSF receptor complex supported this idea.

Nef activation did not necessarily cause the overall reduction of M-CSF receptor pathways. Tyrosine phosphorylation/activation of ERK in response to M-CSF was enhanced by Nef activation (Figure 5B). This finding raises the possibility that Nef-activated Hck inhibits the process of "full activation" of M-CSF receptor. The activation process of M-CSF receptor is considered to involve the ligand-induced dimerization and subsequent autophosphorylation of specific tyrosine residues in their cytoplasmic region.<sup>16</sup> Phosphorylated tyrosine residues then serve as docking sites for SH2 domain-containing molecules, each initiating signaling cascades along specific pathways.<sup>16</sup> Future studies in which we determine whether Nef-activated Hck directly binds to M-CSF receptor and identify site(s) within the cytoplasmic region of the receptor to which Nef-activated Hck binds will clarify the mechanism for the selective inhibition of Nef on the M-CSF receptor pathways.

Our finding that Nef inhibits the process of macrophage differentiation mediated by M-CSF (Figure 2) explains the mechanism for dysfunction, such as defective phagocytosis of opportunistic pathogens, of monocytes/macrophages in HIV-1-infected persons. It is well documented that the phagocytosis of pathogens such as *Mycobacterium avium* complex, *Pneumocystis carinii*, *Toxoplasma gondii*, and *Candida albicans* by peripheral blood monocytes, tissue macrophages, and monocyte-derived macrophages is impaired by in vitro and in vivo HIV-1 infection (reviewed in Kedzierska et al<sup>41</sup>). M-CSF is an important cytokine for macrophage differentiation<sup>15</sup> and has been shown to stimulate various effector functions of macrophages.<sup>14</sup> In particular, a number of in vitro and in vivo studies have shown that M-CSF activates antimicrobial activities of macrophages.<sup>42-45</sup> Moreover, the clinical efficacy of M-CSF has been proven in patients with invasive fungal infections.<sup>46,47</sup>

We showed that the percentage of apoptotic cells in M-CSF-containing culture of Nef-active TF-1-*fms*-Nef-ER cells was significantly higher than that of Nef-inactive cells (Figure 1D, E). In contrast, the degree of apoptotic cell death induced by M-CSF withdrawal in Nef-active TF-1-*fms*-Nef-ER cells was similar to that in Nef-inactive cells (data not shown). These results indicate that Nef does not actively induce apoptotic cell death in the cells but inhibits the signaling from M-CSF. On the other hand, a recent study has revealed that *Nef* does not inhibit but that it mimics signaling initiated by granulocyte/macrophage-CSF (GM-CSF),<sup>34</sup> another cytokine that stimulates the proliferation and differentiation of monocytes/macrophages. That HIV-1 Nef selectively inhibits the M-CSF/M-CSF receptor system may address how monocytes/macrophages contribute to the development and progression of AIDS *in vivo*. Monocytes/macrophages are considered to be involved in the disease progression process by their secretion of soluble factors<sup>9,10</sup>; the profile of gene expression of M-CSF-derived macrophages differs from that of GM-CSF-derived macrophages.<sup>48</sup> It is possible that the selective inhibition of Nef on cytokine signaling results in the modification of macrophage

functions, such as cytokine/chemokine production, in HIV-1-infected persons. This may create an immunologic environment favorable for HIV-1 virus production. Alternatively, this may lead to abnormal tissue distribution of cells of the lineage, thereby facilitating the transfer of HIV-1 virus to bystander CD4<sup>+</sup> T cells. Our findings provide a novel clue to understanding how monocytes/macrophages contribute to the development and progression of AIDS and to clarifying the role of the Nef-Hck interaction in the disease process.

## Acknowledgments

We thank K. Tabata and S. Okamura for research support. M-CSF was kindly provided by Morinaga Milk Industry. Nef antiserum and pEBB-Nef-ER-IRES-puro were obtained from the National Institutes of Health AIDS Research and Reference Reagent Program (Division of AIDS, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD).

## References

- Kestler HW, Ringler DJ, Mori K, et al. Importance of the *nef* gene for maintenance of high virus loads and for development of AIDS. *Cell*. 1991; 65:651-662.
- Kirchhoff F, Greenough TC, Brettler DB, Sullivan JL, Desrosiers RC. Brief report: absence of intact *nef* sequences in a long-term survivor with non-progressive HIV-1 infection. *N Engl J Med*. 1995; 332:228-232.
- Deacon NJ, Tsykin A, Solomon A, et al. Genomic structure of an attenuated quasi species of HIV-1 from a blood transfusion donor and recipients. *Science*. 1995;270:988-991.
- Hanna Z, Kay DG, Rebai N, Guimond A, Jothy S, Jolicoeur P. Nef harbors a major determinant of pathogenicity for an AIDS-like disease induced by HIV-1 in transgenic mice. *Cell*. 1998;95:163-175.
- Fackler OT, Baur AS. Live and let die: Nef functions beyond HIV replication. *Immunity*. 2002;16: 493-497.
- Peterlin BM, Trono D. Hide, shield and strike back: how HIV-infected cells avoid immune eradication. *Nat Rev Immunol*. 2003;3:97-107.
- Garcia JV, Miller AD. Serine phosphorylation-independent downregulation of cell-surface CD4 by nef. *Nature*. 1991;350:508-511.
- Schwartz O, Marechal V, Le Gall S, Lemonnier F, Heard JM. Endocytosis of major histocompatibility complex class I molecules is induced by the HIV-1 Nef protein. *Nat Med*. 1996;2:338-342.
- Swingler S, Mann A, Jacque J-M, et al. HIV-1 Nef mediates lymphocyte chemotaxis and activation by infected macrophages. *Nat Med*. 1999;5:997-1003.
- Swingler S, Brichacek B, Jacque J-M, Ulich C, Zhou J, Stevenson M. HIV-1 Nef intersects the macrophage CD40L signalling pathway to promote resting-cell infection. *Nature*. 2003;424:213-219.
- Saksela K, Cheng G, Baltimore D. Proline-rich (PxxP) motifs in HIV-1 Nef bind to SH3 domains of a subset of Src kinases and are required for the enhanced growth of Nef<sup>+</sup> viruses but not for down-regulation of CD4. *EMBO J*. 1995;14:484-491.
- Moarefi I, LaFevre-Bernt M, Sicheri F, et al. Activation of the Src-family tyrosine kinase Hck by SH3 domain displacement. *Nature*. 1997;385: 650-653.
- Hanna Z, Weng X, Kay DG, Poudrier J, Lowell C, Jolicoeur P. The pathogenicity of human immunodeficiency virus (HIV) type 1 Nef in CD4C/HIV transgenic mice is abolished by mutation of its SH3-binding domain, and disease development is delayed in the absence of Hck. *J Virol*. 2001; 75:9378-9392.
- Roth P, Stanley ER. The biology of CSF-1 and its receptor. *Curr Top Microbiol Immunol*. 1992;181, 141-167.
- Yoshida H, Hayashi S-I, Kunisada T, et al. The murine mutation osteopetrosis is in the coding region of the macrophage colony stimulating factor gene. *Nature*. 1990;345:442-444.
- Bourette RP, Rohrschneider LR. Early events in M-CSF receptor signaling. *Growth Factors*. 2000; 17:155-166.
- Suzu S, Tanaka-Douzon M, Nomaguchi K et al. p56<sup>lck</sup> as a cytokine-inducible inhibitor of cell proliferation and signal transduction. *EMBO J*. 2000;19:5114-5122.
- Courtneidge SA, Dhand R, Pliat D, Twamley GM, Waterfield MD, Rousset MF. Activation of Src family kinases by colony stimulating factor-1, and their association with its receptor. *EMBO J*. 1993; 12:943-950.
- Alonso G, Koegl M, Mazurenko N, Courtneidge SA. Sequence requirements for binding of Src family kinases to activated growth factor receptors. *J Biol Chem*. 1995;270:9840-9848.
- Barone MV, Courtneidge SA. Myc but not Fos rescue of PDGF signalling block caused by kinase-inactive Src. *Nature* 1995;378:509-512.
- Marks DC, Csar XF, Wilson NJ, et al. Expression of a Y559F mutant CSF-1 receptor in M1 myeloid cells: a role for Src kinases in CSF-1 receptor-mediated differentiation. *Mol Cell Biol Res Commun*. 1999;1:144-152.
- Suzu S, Kimura F, Ota J, et al. Biologic activity of proteoglycan macrophage colony-stimulating factor. *J Immunol*. 1997;159:1860-1867.
- Kitamura T, Tange T, Terasawa T, et al. Establishment and characterization of a unique human cell line that proliferates dependently on GM-CSF, IL-3, or erythropoietin. *J Cell Physiol*. 1989;140: 323-334.
- Walk SF, Alexander M, Maier B, Hammarskjold M-L, Rekosch DM, Ravichandran KS. Design and use of an inducibly activated immunodeficiency virus type 1 Nef to study immune modulation. *J Virol*. 2001;75:834-843.
- Shugars DC, Smith MS, Glueck DH, Nantermet PV, Seillier-Moisewitsch F, Swanstrom R. Analysis of human immunodeficiency virus type 1 nef gene sequences present *in vivo*. *J Virol*. 1993;67: 4639-4650.
- Kume A, Xu R, Ueda Y, Urabe M, Ozawa K. Long-term tracking of murine hematopoietic cells transduced with a bicistronic retrovirus containing CD24 and EGFP genes. *Gene Ther*. 2000;7: 1193-1199.
- Hartatik T, Okada S, Okabe S, Arima M, Hatano M, Tokuhisa T. Binding of BAZF and Bcl6 to STAT6-binding DNA sequences. *Biochem Biophys Res Commun*. 2001;284:26-32.
- Okada S, Zhang H, Hatano M, Tokuhisa T. A physiologic role of Bcl-x<sub>L</sub> induced in activated macrophages. *J Immunol*. 1998;160:2590-2596.
- Mizushima S, Nagata S. pEF-BOS, a powerful mammalian expression vector [abstract]. *Nucleic Acids Res*. 1990;18:5322.
- Suzu S, Hayashi Y, Harumi T, et al. Molecular cloning of a novel immunoglobulin superfamily gene preferentially expressed by brain and testis. *Biochem Biophys Res Commun*. 2002;296:1215-1221.
- Schneider P, MacKay F, Steiner V, et al. BAFF, a novel ligand of the tumor necrosis factor family, stimulates B cell growth. *J Exp Med*. 1999;189: 1747-1756.
- Sawai ET, Baur A, Struble H, Peterlin BM, Levy JA, Cheng-Mayer C. Human immunodeficiency virus type 1 Nef associates with a cellular serine kinase in T lymphocytes. *Proc Natl Acad Sci U S A*. 1994;91:1539-1543.
- Murakami Y, Fukazawa H, Kobatake T, et al. A mammalian two-hybrid screening system for inhibitors of interaction between HIV Nef and the cellular tyrosine kinase Hck. *Antiviral Res*. 2002; 55:161-168.
- Briggs SD, Scholtz B, Jacque J-M, Swingler S, Stevenson M, Smithgall TE. HIV-1 Nef promotes survival of myeloid cells by a Stat3-dependent pathway. *J Biol Chem*. 2001;276:25605-25611.
- De SK, Venkateshan CNS, Seth P, Gajdusek DC, Gibbs CJ. Adenovirus-mediated human immunodeficiency virus-1 Nef expression in human monocytes/macrophages and effect of Nef on downmodulation of Fcγ receptors and expression of monokines. *Blood*. 1998;91:2108-2117.
- Federico M, Percario Z, Olivetta E, et al. HIV-1 Nef activates STAT1 in human monocytes/macrophages through the release of soluble factors. *Blood*. 2001;98:2752-2761.

37. Olivetta E, Percario Z, Fiorucci G, et al. HIV-1 Nef induces the release of inflammatory factors from human monocytes/macrophages: involvement of Nef endocytotic signals and NF- $\kappa$ B activation. *J Immunol*. 2003;170:1716-1727.
38. Lerner EC, Smithgall TE. SH3-dependent stimulation of Src-family kinase autophosphorylation without tail release from the SH2 domain *in vivo*. *Nat Struct Biol*. 2002;9:365-369.
39. Xu W, Harrison SC, Eck MJ. Three-dimensional structure of the tyrosine kinase c-Src. *Nature*. 1997;385:595-602.
40. Sicheri F, Moarefi I, Kuriyan J. Crystal structure of the Src family tyrosine kinase Hck. *Nature*. 1997;385:602-609.
41. Kedzierska K, Azzam R, Ellery P, Mak J, Jaworowski A, Crowe SM. Defective phagocytosis by human monocyte/macrophages following HIV-1 infection: underlying mechanisms and modulation by adjunctive cytokine therapy. *J Clin Virol*. 2003;26:247-263.
42. Rollides E, Lyman CA, Mertins SD, et al. Ex vivo effects of macrophage colony-stimulating factor on human monocyte activity against fungal and bacterial pathogens. *Cytokine*. 1996;8:42-48.
43. Brummer E, Stevens DA. Macrophage colony-stimulating factor induction of enhanced macrophage anticytotoxic activity: synergy with fluconazole for killing. *J Infect Dis*. 1994;170:173-179.
44. Gonzalez CE, Lyman CA, Lee S, et al. Recombinant human macrophage colony-stimulating factor augments pulmonary host defences against *Aspergillus fumigatus*. *Cytokine*. 2001;15:87-95.
45. Rollides E, Lyman CA, Sein T, Petraitiene R, Walsh TJ. Macrophage colony-stimulating factor enhances phagocytosis and oxidative burst of mononuclear phagocytes against *Penicillium marneffei* conidia. *FEMS Immunol Med Microbiol*. 2003;15:19-26.
46. Nemunaitis J, Meyers JD, Buckner CD, et al. Phase I trial of recombinant human macrophage colony-stimulating factor in patients with invasive fungal infections. *Blood*. 1991;78:907-913.
47. Nemunaitis J, Shannon-Dorcy K, Appelbaum FR, et al. Long-term follow-up of patients with invasive fungal disease who received adjunctive therapy with recombinant human macrophage colony-stimulating factor. *Blood*. 1993;82:1422-1427.
48. Hashimoto S-I, Suzuki T, Dong H-Y, Yamazaki N, Matsushima K. Serial analysis of gene expression in human monocytes and macrophages. *Blood*. 1999;94:837-844.