

Figure 4. Kaplan-Meier curves for disease-free survival according to Akt1/Akt2 expression in 56 patients with HCC. Patients with low Akt2 expression had a significantly worse prognosis than those with high Akt2 expression ($p < 0.001$). Such a relationship was not observed with Akt1 expression.




Figure 5. Kaplan-Meier curves for overall survival rate according to Akt1/Akt2 expression in 56 patients with HCC. Patients with high levels of Akt2 protein had a significantly better survival than those with low level Akt2 expression ($p = 0.038$). Expression levels of Akt1 were not significant.

Table III. Univariate and multivariate analysis (Cox proportional hazards model).

Categories	Univariate	Multivariate		
	p-value	p-value	Risk ratio	95% Confidence interval
Akt2 (high:low)	0.001	0.012	6.351	1.276-7.017
Differentiation (well/mod.:poor/undiff.)	0.028	0.048	3.904	0.223-0.994
Portal vein invasion (yes:no)	0.006	0.420	0.651	0.235-1.828
No. of tumors (solitary:multible)	0.081	0.288	1.130	0.259-1.193

Well/mod., well/moderately-differentiated; poor/undiff., poorly-/undifferentiated.

prognosis of patients with HCC. Indeed, both disease-free survival rate and overall survival rate were significantly better in patients with high Akt2 expression than in those with low Akt2 expression (Figs. 4 and 5). The significance of

Akt2 protein as a prognostic factor was also confirmed by multivariate analysis (Table III). These findings may be explained by the histopathological results, which revealed that prognostic factors such as cell differentiation, portal vein

involvement and number of tumors were significantly less frequent in patients with high Akt2 expression compared to those with low Akt2 expression (Table II). In addition, the risk ratio of Akt2 was 6.35, higher than that of histopathological differentiation, another significant prognostic factor by multivariate analysis (Table III). These results indicate that Akt2 protein expression is a useful prognostic indicator in patients with HCC.

Akt is known to play an important role in various aspects of cancer progression, such as anti-apoptosis, angiogenesis, tumor invasion and metastasis (23-27). In this study, however, histopathological differentiation, and presence of portal tumor thrombus and multiple intrahepatic tumors were inversely related with Akt2 protein expression (Table II), suggesting that Akt2 may be a positive regulator of tumor differentiation in HCC. The exact causes for these controversial findings remain unknown at present. Akt2 may be involved in the early stages of hepatocarcinogenesis, as it was overexpressed in well-differentiated HCC, which is an early stage tumor, and may transform into a more undifferentiated type (33). For example, Akt2 is induced in colonic polyp-tissues at a very early stage in colon carcinogenesis (34). One possible mechanism for Akt2 overexpression in early HCC might be hypoxia, decreased tissue blood flow in well-differentiated HCC, which results from increased cellularity due to proliferating tumor cells (35). In response to hypoxia, Akt can be activated and induce expression of vascular endothelial factor (VEGF) and cyclooxygenase-2 (COX-2) via the transcription factors hypoxia-inducible factor or nuclear factor κ B (NF- κ B) (36,37). Both VEGF and COX-2 play important roles in angiogenesis, one of the representative features of more advanced, dedifferentiated HCC (38). Indeed, the expression of COX-2 and VEGF in HCC tissues decreases according to tumor dedifferentiation (39,40), a feature similar to that of Akt2 protein expression. Based on these findings, further studies are essential to elucidate the cause-effect relationship between Akt2 expression and malignant features of HCC.

In non-tumor tissues, Akt2 protein expression was more highly expressed in liver cirrhosis than in normal liver or chronic hepatitis (Table I), suggesting the relationship with progression of the underlying chronic liver disease. Akt, a direct target of PI-3 kinase, is activated following cell stimulation by various growth factors (9) and is considered to regulate intracellular processes including gene transcription of cell cycle-related proteins (41-43). Molecules involved in the G1-S phase of the cell cycle, such as cyclin D1, CDC25A, or inhibitors of the cell cycle, such as p21Cip1/WAF1 and P27KIP1, are all molecules acting down-stream of Akt. Overexpression of Akt2 can transform NIH 3T3 cells (44). HBx protein, a constitutive protein of hepatitis B virus, also interacts with Akt/PKB and stimulates its kinase activity, which in turn blocks apoptosis by transforming growth factor (TGF)- β inhibition, caspase 3 inactivation or H-Ras activation (45-47). In addition, HBx, collaborating with other oncogenes, such as H-Ras, can transform immortalized rodent cells through activation of Akt (47). Therefore, it is possible that enhanced Akt activation may induce unrestricted cell proliferation leading to carcinogenesis in chronic liver disease.

In contrast to Akt2 protein expression, Akt1 protein was expressed only faintly in all tumor and non-tumor tissues, and there was no relationship with histopathological characteristics of HCC. Organ- and tissue-specific expression of Akt protein has been demonstrated. Indeed, Akt2 protein has been demonstrated to be a predominant isoform in ovarian cancer, colon cancer and pancreatic cancer, whereas Akt1 protein was predominantly detected in gastric cancer (10,11,29,34). Akt mRNA was expressed in both normal and tumor tissues of the colon, breast, prostate and lung (48). In this study, the expression of Akt2 protein was higher in both non-cancer and cancer tissues of the liver compared with that of Akt1 protein (Figs. 1 and 2). The reason for the difference between Akt1 and Akt2 protein expression is not yet well understood, and the degree of Akt immunostaining is not always correlated with its kinase activity. However, these findings suggest that Akt2 may play a more specific role than Akt1 in the regulation of carcinogenesis and progression of chronic liver disease.

In conclusion, we have demonstrated in the present study that expression of Akt2 in liver tissue was closely associated with both hepatocarcinogenesis and the post-operative prognosis of HCC.

References

1. Parkin DM, Pisani P and Ferlay J: Global cancer statistics. *CA. Cancer J Clin* 49: 33-64, 1999.
2. El-Serag HB and Mason AC: Rising incidence of hepatocellular carcinoma in the United States. *N Engl J Med* 340: 745-750, 1999.
3. Wu CC, Ho WL, Yeh DC, Huang CR, Liu TJ and Peng FK: Hepatic resection of hepatocellular carcinoma in cirrhotic livers: is it unjustified in impaired liver function? *Surgery* 120: 34-39, 1996.
4. De La Coste A, Romagnolo B, Billuart P, *et al*: Somatic mutations of the beta-catenin gene are frequent in mouse and human hepatocellular carcinomas. *Proc Natl Acad Sci USA* 95: 8847-8851, 1998.
5. Murakami Y, Hayashi K, Hirohashi S and Sekiya T: Aberrations of the tumor suppressor p53 and retinoblastoma genes in human hepatocellular carcinomas. *Cancer Res* 51: 5520-5525, 1991.
6. Nishida N, Fukuda Y, Komeda T, *et al*: Amplification and overexpression of the cyclin D1 gene in aggressive human hepatocellular carcinoma. *Cancer Res* 54: 3107-3110, 1994.
7. Ito Y, Matsuura N, Sakon M, *et al*: Expression and prognostic roles of the G1-S modulators in hepatocellular carcinoma: p27 independently predicts the recurrence. *Hepatology* 30: 90-99, 1999.
8. Franke TF, Yang SI, Chan, *et al*: The protein kinase encoded by the Akt proto-oncogene is a target of the PDGF-activated phosphatidylinositol 3-kinase. *Cell* 81: 727-736, 1995.
9. Testa JR and Bellacosa A: Membrane translocation and activation of the Akt kinase in growth factor-stimulated hematopoietic cells. *Leuk Res* 21: 1027-1031, 1997.
10. Staal SP: Molecular cloning of the akt oncogene and its human homologues AKT1 and AKT2: amplification of AKT1 in a primary human gastric adenocarcinoma. *Proc Natl Acad Sci USA* 84: 5034-5037, 1987.
11. Cheng JQ, Godwin AK, Bellacosa A, *et al*: AKT2, a putative oncogene encoding a member of a subfamily of protein-serine/threonine kinases, is amplified in human ovarian carcinomas. *Proc Natl Acad Sci USA* 89: 9267-9271, 1992.
12. Nakatani K, Sakaue H, Thompson DA, Weigel RJ and Roth RA: Identification of a human Akt3 (protein kinase B gamma) which contains the regulatory serine phosphorylation site. *Biochem Biophys Res Commun* 257: 906-910, 1999.
13. Staal SP, Huebner K, Croce CM, Parsa NZ and Testa JR: The AKT1 proto-oncogene maps to human chromosome 14, band q32. *Genomics* 2: 96-98, 1988.
14. Altomare DA, Guo K, Cheng JQ, Sonoda G, Walsh K and Testa JR: Cloning, chromosomal localization and expression analysis of the mouse Akt2 oncogene. *Oncogene* 11: 1055-1060, 1995.

15. Franke TF, Kaplan DR, Cantley LC and Toker A: Direct regulation of the Akt proto-oncogene product by phosphatidylinositol-3,4-bisphosphate. *Science* 275: 665-668, 1997.
16. Klippel A, Kavanaugh WM, Pot D and Williams LT: A specific product of phosphatidylinositol 3-kinase directly activates the protein kinase Akt through its pleckstrin homology domain. *Mol Cell Biol* 17: 338-344, 1997.
17. Alessi DR, Andjelkovic M, Caudwell B, Cron P, Morrice N, Cohen P and Hemmings BA: Mechanism of activation of protein kinase B by insulin and IGF-1. *EMBO J* 15: 6541-6551, 1996.
18. Brownawell AM, Kops GJ, Macara IG and Burgering BM: Inhibition of nuclear import by protein kinase B (Akt) regulates the subcellular distribution and activity of the forkhead transcription factor AFX. *Mol Cell Biol* 21: 3534-3546, 2001.
19. Ueki K, Yamamoto-Honda R, Kaburagi Y, *et al*: Potential role of protein kinase B in insulin-induced glucose transport, glycogen synthesis, and protein synthesis. *J Biol Chem* 273: 5315-5322, 1998.
20. Eapen AK, Henry MK, Quelle DE and Quelle FW: Dna damage-induced G(1) arrest in hematopoietic cells is overridden following phosphatidylinositol 3-kinase-dependent activation of cyclin-dependent kinase 2. *Mol Cell Biol* 21: 6113-6121, 2001.
21. Kennedy SG, Kandel ES, Cross TK and Hay N: Akt/protein kinase B inhibits cell death by preventing the release of cytochrome c from mitochondria. *Mol Cell Biol* 19: 5800-5810, 1999.
22. Cardone MH, Roy N, Stennicke HR, *et al*: Regulation of cell death protease caspase-9 by phosphorylation. *Science* 282: 1318-1321, 1998.
23. Fulton D, Gratton JP, McCabe TJ, *et al*: Regulation of endothelium-derived nitric oxide production by the protein kinase Akt. *Nature* 399: 597-601, 1999.
24. Dimmeler S, Fleming I, Fisslthaler B, Hermann C, Busse R and Zeiher AM: Activation of nitric oxide synthase in endothelial cells by Akt-dependent phosphorylation. *Nature* 399: 601-605, 1999.
25. Kang SS, Kwon T, Kwon DY and Do SI: Akt protein kinase enhances human telomerase activity through phosphorylation of telomerase reverse transcriptase subunit. *J Biol Chem* 274: 13085-13090, 1999.
26. Park BK, Zeng X and Glazer RI: Akt1 induces extracellular matrix invasion and matrix metalloproteinase-2 activity in mouse mammary epithelial cells. *Cancer Res* 61: 7647-7653, 2001.
27. Ringel MD, Hayre N, Saito J, *et al*: Overexpression and over-activation of Akt in thyroid carcinoma. *Cancer Res* 61: 6105-6111, 2001.
28. Yuan ZQ, Sun M, Feldman RI, *et al*: Frequent activation of AKT2 and induction of apoptosis by inhibition of phosphoinositide-3-OH kinase/Akt pathway in human ovarian cancer. *Oncogene* 19: 2324-2330, 2000.
29. Cheng JQ, Ruggeri B, Klein WM, Sonoda G, Altomare DA, Watson DK and Testa JR: Amplification of AKT2 in human pancreatic cells and inhibition of AKT2 expression and tumorigenicity by antisense RNA. *Proc Natl Acad Sci USA* 93: 3636-3641, 1996.
30. Sun M, Wang G, Paciga JE, *et al*: AKT1/PKBalpha kinase is frequently elevated in human cancers and its constitutive activation is required for oncogenic transformation in NIH3T3 cells. *Am J Pathol* 159: 431-437, 2001.
31. Burgering BM and Coffey PJ: Protein kinase B (c-Akt) in phosphatidylinositol-3-OH kinase signal transduction. *Nature* 376: 599-602, 1995.
32. Takemasa I, Yamamoto H, Sekimoto M, *et al*: Overexpression of CDC25B phosphatase as a novel marker of poor prognosis of human colorectal carcinoma. *Cancer Res* 60: 3043-3050, 2000.
33. Sugihara S, Nakashima O, Kojiro M, Majima Y, Tanaka M and Tanikawa K: The morphologic transition in hepatocellular carcinoma. A comparison of the individual histologic features disclosed by ultrasound-guided fine-needle biopsy with those of autopsy. *Cancer* 70: 1488-1492, 1992.
34. Roy HK, Olusola BF, Clemens DL, Karolski WJ, Ratashak A, Lynch HT and Smyrk TC: AKT proto-oncogene overexpression is an early event during sporadic colon carcinogenesis. *Carcinogenesis* 23: 201-205, 2002.
35. Kim KR, Moon HE and Kim KW: Hypoxia-induced angiogenesis in human hepatocellular carcinoma. *J Mol Med* 80: 703-714, 2002.
36. Mazure NM, Chen EY, Laderoute KR and Giaccia AJ: Induction of vascular endothelial growth factor by hypoxia is modulated by a phosphatidylinositol 3-kinase/Akt signaling pathway in Ha-ras-transformed cells through a hypoxia inducible factor-1 transcriptional element. *Blood* 90: 3322-3331, 1997.
37. Ji YS, Xu Q and Schmedtje JF Jr: Hypoxia induces high-mobility-group protein I(Y) and transcription of the cyclooxygenase-2 gene in human vascular endothelium. *Circ Res* 83: 295-304, 1998.
38. Kohno H, Nagasue N and Rahman MA: COX-2- a target for preventing hepatic carcinoma? *Expert Opin Ther Targets* 6: 483-489, 2002.
39. Kondo M, Yamamoto H, Nagano H, *et al*: Increased expression of COX-2 in nontumor liver tissue is associated with shorter disease-free survival in patients with hepatocellular carcinoma. *Clin Cancer Res* 5: 4005-4012, 1999.
40. Yamaguchi R, Yano H, Iemura A, Ogasawara S, Haramaki M and Kojiro M: Expression of vascular endothelial growth factor in human hepatocellular carcinoma. *Hepatology* 28: 68-77, 1998.
41. Fuhrmann G, Leisser C, Rosenberger G, *et al*: Cdc25A phosphatase suppresses apoptosis induced by serum deprivation. *Oncogene* 20: 4542-4553, 2001.
42. Zhou BP, Liao Y, Xia W, Spohn B, Lee MH and Hung MC: Cytoplasmic localization of p21Cip1/WAF1 by Akt-induced phosphorylation in HER-2/neu-overexpressing cells. *Nat Cell Biol* 3: 245-252, 2001.
43. Liang J, Zubovitz J, Petrucelli T, *et al*: PKB/Akt phosphorylates p27, impairs nuclear import of p27 and opposes p27-mediated G1 arrest. *Nat Med* 8: 1153-1160, 2002.
44. Cheng JQ, Altomare DA, Klein MA, Lee WC, Kruh GD, Lissy NA and Testa JR: Transforming activity and mitosis-related expression of the AKT2 oncogene: evidence suggesting a link between cell cycle regulation and oncogenesis. *Oncogene* 14: 2793-2801, 1997.
45. Shih WL, Kuo ML, Chuang SE, Cheng AL and Doong SL: Hepatitis B virus X protein inhibits transforming growth factor-beta-induced apoptosis through the activation of phosphatidylinositol 3-kinase pathway. *J Biol Chem* 275: 25858-25864, 2000.
46. Lee YI, Kang-Park S, Do SI and Lee YI: The hepatitis B virus-X protein activates a phosphatidylinositol 3-kinase-dependent survival signaling cascade. *J Biol Chem* 276: 16969-16977, 2001.
47. Kim YC, Song KS, Yoon G, Nam MJ and Ryu WS: Activated ras oncogene collaborates with HBx gene of hepatitis B virus to transform cells by suppressing HBx-mediated apoptosis. *Oncogene* 20: 16-23, 2001.
48. Zinda MJ, Johnson MA, Paul JD, *et al*: AKT-1, -2 and -3 are expressed in both normal and tumor tissues of the lung, breast, prostate, and colon. *Clin Cancer Res* 7: 2475-2479, 2001.

Clinical value of alterations in *p73* gene, related to *p53* at 1p36, in human hepatocellular carcinoma

TARO AOKI, ATSUSHI MIYAMOTO, SHIGERU MARUBASHI, HIROAKI NAGANO, KOJI UMESHITA, YOSHIYUKI FUJIWARA, KEIZO DONO, SHOJI NAKAMORI, MASATO SAKON and MORITO MONDEN

Department of Surgery and Clinical Oncology, Graduate School of Medicine, Osaka University, Osaka, Japan

Received September 22, 2003; Accepted November 4, 2003

Abstract. A novel gene, *p73*, encoding a protein with significant homology to *p53* and showing functional similarities to *p53*, was identified at chromosome 1p36, at which tumor suppressor gene of hepatocellular carcinoma (HCC) is supposed to be. Involvement of *p73* in hepatocarcinogenesis is controversial and clinical value of *p73* alterations remains obscure. We investigated allelic status of *p73* in 63 patients with HCC. Loss of heterozygosity (LOH) in *p73* was analyzed by PCR-RFLP analysis. The results were compared with LOH on chromosome 1p surrounding *p73* locus, mutations of *p53* and *p73*, and clinicopathologic characteristics. LOH on *p73* was observed in 33% of informative tumors. LOH in *p73* was not always observed between the regions with LOH on chromosome 1p examined despite the significant association of LOH in *p73* with LOH on chromosome 1p. No mutations were detected in *p73*. Tumors with LOH in *p73* were more frequently detected in liver without cirrhosis than that with cirrhosis. There was no significant statistic association between the presence of LOH in *p73* and six different clinicopathologic characteristics such as age, sex, histological type, T stage, tumor diameter, and virus status. Disease-free survival rates of the patients with LOH in *p73* were significantly poorer than those without LOH in *p73*. Multivariate analysis indicated that presence of LOH in *p73* was independent prognostic factor in patients with HCC. These findings suggested that *p73* might play some role in tumor progression of HCC even though *p73* should not be considered a candidate gene on chromosome 1p of HCC and does not function as a tumor suppressor gene like *p53*.

Identifying the patients with LOH of *p73* in tumors could be useful to predict early recurrence and to stratify the patients who need adjuvant therapy after operation.

Introduction

Hepatocellular carcinoma (HCC) is one of the most common human cancers in the world and it carries a poor prognosis (1). Various studies have confirmed its association with persistent hepatitis B or C viruses infection and other liver-damaging factors such as alcoholic cirrhosis and exposure to mycotoxins (1,2). Although each of these etiologic agents appears to predispose a hepatocyte to development of HCC, either directly or indirectly, the molecular mechanisms involved in hepatocarcinogenesis remain obscure. Tumorigenesis and progression of malignancy, however, are considered to be the multiple processes involving genetic alterations of several types, such as chromosomal deletions, chromosomal translocations, point mutations, and gene amplification (3). These changes can lead to activation of oncogenes or inactivation of tumor suppressor genes. There is evidence for widespread involvement of putative suppressor genes such as *p53* and *Rb* genes in development and progression of HCC (4,5), while a few alterations have been found in known oncogenes (6-9).

Somatic inactivation of tumor suppressor genes is usually achieved by intragenic mutations in one allele of the gene and by the loss of a chromosomal region spanning the second allele (3,10). Chromosomal analysis using microsatellite polymorphic markers has revealed loss of heterozygosity (LOH) of specific chromosomal regions in various types of cancers, e.g., colon, breast, and lung cancer (3,11-13). The mapping of regions with a high frequency of LOH has been critical for identifying tumor suppressor genes (3). In HCC, previous studies mainly relying upon either restriction fragment length polymorphism (RFLP) markers or microsatellite markers restricted to specific chromosome arms, have also defined a number of LOH on various chromosome arms (14-19). Especially on chromosome 1p, frequent LOHs in HCC were reported, suggesting the presence of tumor suppressor genes in this chromosomal regions (20,21). However, the gene responsible for this chromosomal region has not been identified yet.

Recently, a novel putative tumor suppressor gene, *p73*, mapped to the 1p36 region (22), in which frequent deletions were reported in patients with neuroblastoma (23,24) as well

Correspondence to: Dr Shoji Nakamori, Department of Surgery and Clinical Oncology, Graduate School of Medicine, Osaka University, E2, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan
E-mail: nakamori@surg2.med.osaka-u.ac.jp

Abbreviations: HCC, human hepatocellular carcinoma; PCR, polymerase chain reaction; LOH, loss of heterozygosity; RFLP, restriction length polymorphism; RT-PCR, reverse transcription-PCR; SSCP, single strand conformation polymorphism; DFS, disease-free survival

Key words: *p73*, chromosome 1p, loss of heterozygosity, *p53*, hepatocellular carcinoma, polymerase chain reaction

as HCC. This gene encodes a protein with remarkable sequence similarity to the DNA-binding, transactivation, and oligomerization domains of *p53*, which is widely accepted to be involved in tumorigenesis and progression of a variety of malignant tumors including HCC (25,26). It also activates transcription of *p53* responsive genes and inhibit cell growth in a *p53*-like manner, by inducing apoptosis (27).

Despite several reports that demonstrated *p73* alterations in HCC, the involvement in hepatocarcinogenesis is still controversial due to the lack of the number of patients examined (28,29). Although increased expression of this gene in HCC is reported to be associated with poor patient prognosis in only one immunohistochemical study (30), clinical value of *p73* gene alterations remains obscure. In the present study, we examine LOH on *p73* including chromosome 1p region surrounding *p73* locus and mutations of *p73*, with reference to the *p53* mutation status, in 63 patients with HCC in order to elucidate the potential involvement of *p73* gene alterations in the development and progression of HCC.

Patients and methods

Patients and tissue preparation. Tumor and corresponding non-tumorous liver tissues from 63 HCC patients who underwent curative surgery at the Department of Surgery and Clinical Oncology, Graduate School of Medicine, Osaka University between June 1994 and August 1998, were collected with informed consent. This study included only patients from whom adequate tumor tissue was available for pathologic examination and molecular genetic analysis. Each tissue specimen was divided into two pieces at surgery. One for molecular analysis was immediately frozen by liquid nitrogen at the time of surgery and stored at -80°C until examination. Another was processed for pathologic examination. All 63 tumors were macroscopically or microscopically examined to determine location, size, extent and mode of cancer invasion, and metastasis to lymph nodes or distant organs according to the International Union Against Cancer tumor-nodes-metastasis (TNM) classification (31). Mean age of the patients was 61.5 years (range, 35-74 years). Fifty-two patients were men and 11 were women. Thirty-eight patients had HCV infection, 19 had HBV infection, and 6 had no HCV/HBV infection. Thirty-four patients had liver cirrhosis. Mean tumor diameter was 4.1 cm (range, 0.8-19.0 cm). Histologically, 9 tumors were classified as well-differentiated adenocarcinoma, 24 were moderately differentiated adenocarcinoma, and 30 were poorly differentiated adenocarcinoma. According to the TNM classification (31), 14 patients were T1N0M0 (stage I), 29 were T2N0M0 (stage II), 11 were T3N0M0 (stage III-A), and 9 were T4N0M0 (stage IV-A) (Table I). Patient follow-up was organized in Department of Surgery and Clinical Oncology, Graduate School of Medicine, Osaka University. All patients were followed until recurrence of disease or the end of the observation period (April 30, 2002). Mean follow-up period was 36.9±13.4 months.

A pair of tumor and the corresponding non-tumorous liver tissue was prepared for DNA and RNA analysis. Genomic DNA samples and total RNA were isolated using SepaGene

Table I. Summary of clinicopathologic characteristics of 63 patients.

Age (mean)	35-74 years (61.5)
Sex (Male/Female)	52/11
Histological type (well/mod/poor)	9/24/30
T stage (1/2/3/4)	14/29/11/9
Tumor diameter (mean)	0.8-19.0 cm (4.1)
HCV infection (+/-)	38/25
HBV infection (+/-)	19/44
Liver cirrhosis (+/-) ^a	34/26

^aThree patients unknown.

(SankoJyunyaku, Tokyo, Japan) and TRIzol (Life Technologies, Grand island, NY) according to the manufacturer's protocols, respectively.

Microsatellite polymorphism analysis. Genetic alterations on chromosome 1p in HCC were studied by microsatellite polymorphism analysis, as previously described (21,32). The CA-repeat microsatellite markers, D1S243, D1S228, D1S513, and D1S211 (33), which are located near the chromosomal region of *p73* were used. Primer sets labeled with fluorescent dye for PCR from Human MAPPAIRS (Research Genetics, Huntsville, AL) were used. PCR was performed in a 20 µl reaction mixture containing 0.05 µg of genomic DNA template, 1X PCR buffer, 1.5 mM MgCl₂, 0.8 mM deoxy-nucleotide triphosphates, 0.2 mM of each primers, and one unit of AmpliTaq Gold (Perkin-Elmer, Foster city, CA) using Perkin-Elmer Cetus 9600 thermocycler (Perkin-Elmer Cetus). Reactions were carried for 30 cycles. Each cycle consisted of 45 sec at 95°C for denaturation, 1 min at 57°C for primer annealing, and 1 min at 72°C for primer extension. The first denaturation step was lengthened to 12 min for activation of AmpliTaq. The last extension step was lengthened to 7 min. Each PCR product (5 µl), mixed with 2 µl of blue dextran (80% formamide; 10 mM EDTA; 0.6% blue dextran), was heated at 95°C for 20 min and, then, electrophoresed in 6% denaturing polyacrylamide gel, containing 7.7 M Urea at 1500 V for 2 h. The image of fluorescent-labeled PCR products was scanned and acquired by a FMBIO II Multi-View (Takara, Shiga, Japan). When there was an absence, or a >50% reduction in intensity of one allele in tumor specimens comparing to the allele in normal liver, tumor was scored as exhibiting LOH, as previously described (21,32).

Examination of LOH at the *p73* locus by PCR-RFLP analysis. PCR-RFLP analysis with restriction enzyme *SpyI* (Boehringer Mannheim, Mannheim, Germany) was used to distinguish two distinct alleles, representing tightly linked polymorphisms in *p73* exon 2, as previously described (22). An allelic polymorphism consisting of a double nucleotide substitution (G→A) and (C→T) at position 4 and 14 of exon 2. Nested PCR

was performed under the following conditions. First PCR was performed with primer sets (CACCTGCTCCAGGG ATGC, AAAATAGAAGCGTCAGTC) (22) in a 25 μ l reaction mixture containing 0.05 μ g of genomic DNA template, 1X PCR buffer, 1.5 mM MgCl₂, 0.8 mM deoxynucleotide triphosphates, 0.2 mM of each primers, and one unit of AmpliTaq Gold using Perkin-Elmer Cetus 9600 thermocycler. Reactions were carried for 35 cycles. Each cycle consisted of 30 sec at 95°C for denaturation, and 30 sec at 58°C for primer annealing. First denaturation step was lengthened to 9 min for activation of AmpliTaq. The last primer extension step was consisted of 10 min at 68°C. Second PCR step was performed with primer sets (CAGGCC CACTTGCCTGCC, CTGTCCCAAGCTGATGA) (22) under the same condition of first PCR step using 0.5 μ l of first PCR product. After second PCR, a specific 482-bp fragment was obtained. The PCR product (5 μ l) was then digested with S_{ty}I overnight. S_{ty}I digestion of PCR products yields two smaller size fragments from AT allele-derived amplicons of 376 and 106 bp, respectively, whereas the GC allele-derived amplicons remain uncut, as previously described (22). The digestion results were analyzed on a 2% agarose gel.

PCR-SSCP analysis for the mutation screening of the p53 and p73. PCR-SSCP analysis was performed to screen the mutation of p53 in exon 5-8 and mutation of p73 in exon 1-13. Regarding PCR-SSCP analysis of the p53, primer pairs for p53 were: (TTCAACTCTGTCTCCTTCCT, CAGCCC TGTCGTCTCTCCAG) for exon 5, (GCCTCTGATTCC TCACTGAT, TTAACCCCTCCTCCCAGAGA) for exon 6, (CTTGCCACAGGTCTCCCAA, TGTGCAGGGTGGCAA GTGGC) for exon 7, (TTCCTTACTGCCTCTTGCTT, CGC TTCTGTCTCTGCTT) for exon 8 (Nippongene, Tokyo, Japan). PCR was performed in a 20 μ l reaction mixture containing 0.05 μ g of genomic DNA template in the presence of a [³²P]-dCTP, 1X PCR buffer, 1.5 mM MgCl₂, 0.8 mM deoxynucleotide triphosphates, 0.2 mM of each primers, and one unit of AmpliTaq Gold using Perkin-Elmer Cetus 9600 thermocycler. Reactions were carried for 35 cycles. Each cycle consisted of 30 sec at 95°C for denaturation, 30 sec at 64°C for primer annealing. The first denaturation step was lengthened to 12 min for activation of AmpliTaq. The last primer extension step was consisted of 10 min at 72°C, then, followed by electrophoretic separation on 5% non-denaturing polyacrylamide gel with 5% glycerol at room temperature. To screen mutation of p73 gene, RT-PCR-SSCP analysis was performed as previously described (34). Reverse transcription (RT) was performed with 1 μ g of total RNA and Reverse Transcription System (Promega, Madison, WI), according to the manufacturer's protocols. p73 cDNA was amplified into 8 overlapping fragment (S1-AS1, S2-AS2, ..., S7-AS7, S8-AS8 primer sets) (34). The gel was stained by SYBR Green II RNA Gel Stain (FMC, Rockland, ME) and the image of PCR products was scanned and acquired by a FMBIO II Multi-View.

Statistical analysis. The clinicopathological characteristics of patients in relation to LOH in p73 were analyzed by using Mann-Whitney U test or χ^2 test. The characteristics were also




Figure 1. Representative result of LOH analysis in p73 by PCR-RFLP analysis. Ethidium bromide staining comparing PCR-RFLP products with tumor (T) and matched normal liver (N) DNA samples. *Revealing LOH in T. **Homozygous samples. Lane M; size marker (50 bp DNA ladder).




Figure 2. Schematic representation of partial deletions on chromosome 1p and p73 in 19 HCC patients. Patient numbers are shown above, and the loci on the left. Closed circles, LOH; open circles, retained heterozygosity; blank space, uninformative.

analyzed by multivariate analysis. Disease-free survival (DFS) curves were plotted according to the Kaplan-Meier method and were analyzed using the log-rank test. Statistical analysis was performed using Stat-View software version 5.0 (ABACUS, Berkeley, CA). A value of $p < 0.05$ was considered statistically significant.

Results

Twenty-four (38.1%) of 63 patients examined were GC/AT heterozygous for the exon 2 polymorphism and were informative for LOH analysis, while 35 patients (55.6%) were GC/GC homozygous and 4 (6.3%) were AT/AT homozygous. In 24 informative patients, LOH in p73 was observed in 8 patients (33.3%) (Fig. 1).

All of 63 patients were informative for at least one microsatellite marker examined on chromosome 1p. More than one LOH on chromosome 1p were observed in 16 (25.4%) of 63 patients. The deletion mapping in 19 cases showing LOH on chromosome 1p or in p73 is summarized in Fig. 2. Although the association of frequency of LOH in p73 with LOH on chromosome 1p was statistically significant ($p < 0.0005$; Table II), LOH in p73 gene was not always observed between the regions with LOH on chromosome 1p examined (patient no. 9, 27, 29, 31, 33, 38, 48, 50, 52 in Fig. 2).

To examine the possibility that p73 function as a tumor suppressor, mutation in p73 was screened using RT-PCR-

Table II. Association of LOH in *p73* with clinicopathologic variables.

Variables	LOH in <i>p73</i>		p-value
	(+) (n=8)	(-) (n=16)	
Age (year)	61.0	61.8	0.85
Sex (male/female)	7/1	13/3	0.70
Histological type (well-mod/poor)	6/2	9/7	0.37
T stage (T1-2/T3-4)	6/2	10/6	0.54
Tumor diameter (≥ 2 cm/ < 2 cm)	1/7	6/10	0.20
LOH on chromosome 1p (+/-)	5/3	0/16	0.0004
HCV infection (+/-)	5/3	12/4	0.65
HBV infection (+/-)	2/6	4/12	>0.99
Liver cirrhosis (+/-)	1/7	11/5	0.027


Table III. Relationship between LOH in *p73* and *p53* mutation.

	<i>p53</i> mutation	
	(-) (%)	(+) (%)
LOH in <i>p73</i> (-)	12 (75)	4 (25)
(+)	7 (87.5)	1 (12.5)
NI	32	7
Total	51	12

NI, not informative.

SSCP analysis. Several SSCP shifts were detected in a few exons in the pair of the tumor and normal liver DNA, suggesting that those should reflect sequence polymorphisms as previously described (22,34,35) (data not shown). Mutation screening in *p53* was also performed using PCR-SSCP analysis to examine the relationship between LOH on *p73* and *p53* mutations. Mutation in *p53* was observed in 12 (19.1%) among the 63 tumors. There were no mutations on *p53* in 7 of 8 patients with LOH in *p73*. Although LOH on *p73* was more frequent in tumors without *p53* mutations, statistical difference was not significant (Table III).

To study whether the alterations of *p73* are involved in hepatocarcinogenesis and progression, the relation between LOH in *p73* and clinicopathologic characteristics was examined. Although tumors with LOH in *p73* were more frequently detected in liver without cirrhosis than that with cirrhosis ($p=0.027$), there was no significant statistical association between the presence of LOH on *p73* and 6 different clinicopathologic features such as age, sex, histological type, T stage, tumor diameter, and virus status of 24 HCC patients informative for LOH on *p73* (Table II). Furthermore, the

Figure 3. Comparison of disease-free survival in LOH (+) group vs. LOH (-) group. There was a significant difference in disease-free survival between the two groups ($p=0.0001$).

presence of LOH on chromosome 1p was significantly associated to LOH on *p73*. There were no significant statistical difference in DFS rate between the presence or the absence of LOH on chromosome 1p in all 63 patients (data not shown). However, DFS curve of the 8 patients with LOH on *p73* was significantly poorer than that of the 16 patients without LOH in *p73* ($p=0.0001$) (Fig. 3). Multivariate analysis using the Cox proportional hazard model indicated that presence of LOH in *p73*, tumor diameter and liver cirrhosis were independent prognostic factors in patients with HCC in spite of small number of patients (Table IV).

Discussion

p73 has recently been identified on chromosome 1p as a first family member of the *p53* tumor-suppressor gene (22). *p73* is predicted to encode a protein with significant amino acid sequence similarity to *p53* (22), and, like *p53*, can activate the transcription of *p53*-responsive genes, such as *p21waf1/cip1*, to inhibit cell growth in a *p53*-like manner by inducing apoptosis when over produced (27). Because of the close similarity between *p73* and *p53* in both structure and function, it was suggested that *p73* might also be a tumor suppressor (22). Furthermore, *p73* was shown to be located at the chromosomal localization 1p36, which is a region frequently deleted in a variety cancers including HCC (11,12,20,21). Up to this point, however, numerous studies have not discovered any coding region mutations of *p73*, an important feature of a tumor suppressor, in tumors including HCC (22,34-36). It is still controversial whether *p73* should be considered a tumor suppressor gene in hepatocarcinogenesis.

Several genetic changes which support the presence of tumor suppressor genes in HCC have been detected as loss of heterozygosity (37) and a genetic model of multistage carcinogenesis and progression has been proposed (38,39). In the present study, we first investigated whether or not *p73* may play a role as a tumor suppressor gene in HCC. Allelotyping analysis to detect LOH on *p73* and on chromosome 1p surrounding *p73* locus was performed on 63 surgically treated

Table IV. Multivariate analysis for disease recurrence.

Variable	Odds ratio (95% CI)	p-value
Age (year) ≥60 vs. <60	6.60 (0.66-65.80)	0.11
Sex Male vs. female	2.47 (0.37-16.55)	0.35
Histological type Well-mod vs. poor	1.68 (0.37-7.70)	0.51
T stage T1-2 vs. T3-4	2.80 (0.68-11.60)	0.16
Tumor diameter >2 cm vs. ≤2 cm	0.06 (0.004-0.832)	0.04
LOH in p73 (-) vs. (+)	15.00 (1.61-139.42)	0.02
LOH on chromosome 1p (-) vs. (+)	0.55 (0.06-5.29)	0.60
HCV infection (-) vs. (+)	0.67 (0.12-3.73)	0.64
HBV infection (-) vs. (+)	12.74 (0.87-186.83)	0.06
Liver cirrhosis (-) vs. (+)	0.04 (0.003-0.53)	0.01

95% CI, 95% confidence interval.

HCC tumors and corresponding non-cancerous liver tissues. LOH in *p73* was observed in 33% of tumors examined, a frequency higher than that in previous reports (28,29) and there was a significant correlation between the patients with LOH in *p73* and the patients with LOH on chromosome 1p ($p=0.0004$). Deletion map, however, showed that LOH in *p73* was not always detected between the regions with LOH on chromosome 1p examined. Although LOH on chromosome 1p have occurred in tumors at an early stage or with a well-differentiated HCC as well as in tumors at more advanced stages, and even in tumors <2 cm (21), LOH on *p73* was not associated with these features.

It is widely accepted that loss or inactivation of a tumor suppressor gene, *p53* contribute to the development of 50% of all human cancers (25,26). In HCC, *p53* mutations are frequently observed in moderately and poorly differentiated tumors (38), and, consequently, is to be associated with the grade of cancer cell atypia that advanced with tumor growth (39). However, frequency of *p53* aberration in HCC (4,5,38) were <30% as examined in the present study, and were lower than other malignant tumors (25,26), suggesting the presence of other suppressor genes except *p53*. Because of the remarkable sequence similarity to the DNA-binding transactivation, and oligomerization domains of *p53* (22), *p73* has been considered as a candidate suppressor gene in

HCC. Although the present study showed that most of the patients with LOH in *p73* had no mutation in *p53* (87.5%), we did not detect any coding region mutations of *p73* in 63 tumors as previously reported in other types of tumors including HCC (22,28,34-36). These findings suggested that *p73* might not be a tumor suppressor gene on chromosome 1p. Otherwise, *p73* unlike *p53* may function in a different manner from Knudson's two-hit theory (40), while mutation type of *p53* is loss of one allele and subtle structural change of the remaining allele (41,42). Further studies are needed to clarify the precise mechanism of *p73* on hepatocarcinogenesis.

The present study indicated LOH in *p73* as an indicator of early recurrence for patients with HCC treated by curative surgical resection. Patients with tumors which had LOH on *p73* had a statistically significant shorter disease-free survival than those without LOH on *p73*. Although cirrhotic status of liver is also one of important prognostic indicators for the recurrence of HCC after surgery (2), LOH on *p73* was frequently observed in patients without cirrhosis. After comparing *p73* LOH status with other factors known to be of prognostic value in patients with surgically curative resection for HCC in a multivariate model, we identified LOH on *p73* as a new and independent marker for assessing the early recurrence of patients after curative resection. The previous study also identified the increased expression of *p73* protein as an indicator of poor prognosis for patients with HCC treated by surgical tumor resection (30). However, the molecular mechanisms not only of the increased expression of *p73* in HCC but also of influence on poor prognosis remain obscure. Since overexpressed *p73* protein is likely to be wild-type (30) also shown in the present study, LOH of *p73* may influence overexpression of *p73* protein which could antagonize *p53* function increasing proliferation in tumor with a poor prognosis (30). Further work will have to clarify the relationship between LOH of *p73* and overexpression of *p73*.

In conclusion, we demonstrated that *p73* might play some role in tumor progression of HCC even though *p73* should not be considered a candidate gene on chromosome 1p of HCC and it does not function as a tumor suppressor gene like *p53*. Identifying the patients with LOH of *p73* in tumors could be useful to predict early recurrence and to stratify the patients who need adjuvant therapy after operation.

Acknowledgements

This study was supported in part by Grants-in-Aid for the Second Term Comprehensive 10-year Strategy for Cancer Control and Cancer Research from the Ministry of Health and Welfare, Japan, and Grants-in-Aid for Scientific Research on Priority Areas, and Basic Research from the Ministry of Education, Science, Sports, and Culture, Japan.

References

- Okuda K: Hepatocellular carcinoma: recent progress. *Hepatology* 15: 948-963, 1992.
- Okada S, Shimada K, Yamamoto J, *et al*: Predictive factors for postoperative recurrence of hepatocellular carcinoma. *Gastroenterology* 106: 1618-1624, 1994.
- Fearon E and Vogelstein B: A genetic model for colorectal tumorigenesis. *Cell* 61: 759-767, 1990.

4. Murakami Y, Hayashi K, Hirohashi S and Sekiya T: Aberrations of the tumor suppressor p53 and retinoblastoma genes in human hepatocellular carcinomas. *Cancer Res* 51: 5520-5525, 1991.
5. Hayashi H, Sugio K, Matsumata T, Adachi E, Takenaka K and Sugimachi K: The clinical significance of p53 gene mutation in hepatocellular carcinomas from Japan. *Hepatology* 22: 1702-1707, 1995.
6. Lee H, Rajagopalan M and Vyas G: A lack of direct role of hepatitis B virus in the activation of ras and c-myc oncogenes in human hepatocellular carcinogenesis. *Hepatology* 8: 1116-1120, 1988.
7. Takada S and Koike K: Activated N-ras gene was found in human hepatoma tissue but only in a small fraction of the tumor cells. *Oncogene* 4: 189-193, 1989.
8. Tsuda H, Hirohashi S, Shimosato Y, Ino Y, Yoshida T and Terada M: Low incidence of point mutation of c-Ki-ras and N-ras oncogenes in human hepatocellular carcinoma. *Jpn J Cancer Res* 80: 196-199, 1989.
9. Tada M, Omata M and Ohto M: Analysis of ras gene mutations in human hepatic malignant tumors by polymerase chain reaction and direct sequencing. *Cancer Res* 50: 1121-1124, 1990.
10. Knudson A: Hereditary cancer, oncogenes, and antioncogenes. *Cancer Res* 45: 1437-1443, 1985.
11. Leister I, Weith A, Bruderlein S, *et al.*: Human colorectal cancer: high frequency of deletions at chromosome 1p35. *Cancer Res* 50: 7232-7235, 1990.
12. Genuardi M, Tsihira H, Anderson D and Saunders G: Distal deletion of chromosome 1p in ductal carcinoma of the breast. *Am J Hum Genet* 45: 73-82, 1989.
13. Brauch H, Johnson B, Hovis J, *et al.*: Molecular analysis of chromosome 3 in small-cell and non-small-cell carcinoma of the lung. *N Engl J Med* 317: 1109-1113, 1987.
14. Nishida N, Fukuda Y, Kokuryu H, *et al.*: Accumulation of allelic loss on arms of chromosomes 13q, 16q and 17p in the advanced stages of human hepatocellular carcinoma. *Int J Cancer* 51: 862-868, 1992.
15. Walker G, Hayward N, Falvey S and Cooksley W: Loss of somatic heterozygosity in hepatocellular carcinoma. *Cancer Res* 51: 4367-4370, 1991.
16. Nagai H, Pineau P, Tjollais P, Buendia M and Dejea A: Comprehensive allelotyping of human hepatocellular carcinoma. *Oncogene* 14: 2927-2933, 1997.
17. Konishi M, Kikuchi-Yanoshita R, Tanaka K, *et al.*: Genetic changes and histopathological grades in human hepatocellular carcinomas. *Jpn J Cancer Res* 84: 893-899, 1993.
18. Emi M, Fujiwara Y, Nakajima T, *et al.*: Frequent loss of heterozygosity for loci on chromosome 8p in hepatocellular carcinoma, colorectal cancer, and lung cancer. *Cancer Res* 52: 5368-5372, 1992.
19. Takahashi K, Kudo K, Ishibashi H, Hirata Y and Niho Y: Frequent loss of heterozygosity on chromosome 22 in hepatocellular carcinoma. *Hepatology* 17: 794-799, 1993.
20. Yeh S, Chen P, Chen H, Lai M, Wang C and Chen D: Frequent genetic alterations at the distal region of chromosome 1p in human hepatocellular carcinoma. *Cancer Res* 54: 4188-4192, 1994.
21. Kuroki T, Fujiwara Y, Tsuchiya E, *et al.*: Accumulation of genetic changes during development and progression of hepatocellular carcinoma: loss of heterozygosity on chromosome arm 1p occurs at an early stage of hepatocarcinogenesis. *Genes Chromosomes Cancer* 13: 163-167, 1995.
22. Kaghad M, Bonnet H, Yang A, *et al.*: Monoallelically expressed gene related to p53 at 1p36, a region frequently deleted in neuroblastoma and other human cancers. *Cell* 90: 809-819, 1997.
23. Caron H, van Sluis P, van Hoesve M, *et al.*: Allelic loss of chromosome 1p36 in neuroblastoma is of preferential maternal origin and correlates with N-myc amplification. *Nat Genet* 4: 187-190, 1993.
24. Cheng J, Hiemstra J, Schneider S, *et al.*: Preferential amplification of the paternal allele of the N-myc gene in human neuroblastomas. *Nat Genet* 4: 191-194, 1993.
25. Hollstein M, Sidransky D, Vogelstein B and Harris C: p53 mutations in human cancers. *Science* 253: 49-53, 1991.
26. Levine A, Momand J and Finlay C: The p53 tumor suppressor gene. *Nature* 351: 453-456, 1991.
27. Jost C, Marin M and Kaelin W Jr: p73 is a human p53-related protein that can induce apoptosis. *Nature* 389: 191-194, 1997.
28. Mihara M, Nimura Y, Ichimiya S, *et al.*: Absence of mutation of the p73 gene localized at chromosome 1p36.3 in hepatocellular carcinoma. *Br J Cancer* 79: 164-167, 1999.
29. Herath NI, Kew MC, Whitehall VLJ, *et al.*: p73 is up-regulated in a subset of hepatocellular carcinomas. *Hepatology* 31: 601-695, 2000.
30. Tannapfel A, Wasner M, Krause K, *et al.*: Expression of p73 and its relation to histopathology and prognosis in hepatocellular carcinoma. *J Natl Cancer Inst* 91: 1154-1158, 1999.
31. Sobin L and Wittekind C: In: TNM classification of Malignant Tumors. 5th edition. Wiley-Liss, New York, pp74-77, 1997.
32. Tamura S, Nakamori S, Kuroki T, *et al.*: Association of cumulative allelic losses with tumor aggressiveness in hepatocellular carcinoma. *J Hepatol* 27: 669-676, 1997.
33. Gyapay G, Morissette J, Vignal A, *et al.*: The 1993-94 Genethon human genetic linkage map. *Nat Genet* 7: 246-339, 1994.
34. Nomoto S, Haruki N, Kondo M, *et al.*: Search for mutations and examination of allelic expression imbalance of the p73 gene at 1p36.33 in human lung cancers. *Cancer Res* 58: 1380-1383, 1998.
35. Mai M, Yokomizo A, Qian C, *et al.*: Activation of p73 silent allele in lung cancer. *Cancer Res* 58: 2347-2349, 1998.
36. Takahashi H, Ichimiya S, Nimura Y, Watanabe M, Furusato M and Wakui S: Mutation, allelotyping, and transcription analyses of the p73 gene in prostatic carcinoma. *Cancer Res* 58: 2076-2077, 1998.
37. Ding SF and Habib NA: Loss of heterozygosity in liver tumors. *J Hepatol* 22: 230-238, 1995.
38. Oda T, Tsuda H, Scarpa A, Sakamoto M and Hirohashi S: p53 gene mutation spectrum in hepatocellular carcinoma. *Cancer Res* 52: 6358-6364, 1992.
39. Teramoto T, Satonaka K, Kitazawa S, Fujimori T, Hayashi K and Maeda S: p53 gene abnormalities are closely related to hepatoviral infections and occur at a late stage of hepatocarcinogenesis. *Cancer Res* 54: 231-235, 1994.
40. Knudson A: Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 68: 820-823, 1971.
41. Baker S, Fearon E, Nigro J, *et al.*: Chromosome 17 deletions and p53 gene mutations in colorectal carcinomas. *Science* 244: 217-221, 1989.
42. Nigro J, Baker S, Preisinger A, *et al.*: Mutations in the p53 gene occur in diverse human tumor types. *Nature* 342: 705-708, 1989.

Featured Article

Molecular Prediction of Response to 5-Fluorouracil and Interferon- α Combination Chemotherapy in Advanced Hepatocellular Carcinoma

Yukinori Kurokawa,^{1,2} Ryo Matoba,²
Hiroaki Nagano,¹ Masato Sakon,¹
Ichiro Takemasa,¹ Shoji Nakamori,¹ Keizo Dono,¹
Koji Umeshita,¹ Noriko Ueno,² Shin Ishii,³
Kikuya Kato,² and Morito Monden¹

¹Department of Surgery and Clinical Oncology, Graduate School of Medicine, Osaka University, Osaka, Japan; and ²Taisho Laboratory of Functional Genomics and ³Laboratory of Theoretical Life Science, Nara Institute of Science and Technology, Nara, Japan

ABSTRACT

Purpose: The prognosis of hepatocellular carcinoma (HCC) is very poor, particularly in patients with tumors that have invaded the major branches of the portal vein. Combination chemotherapy with intra-arterial 5-fluorouracil and subcutaneous interferon- α has shown promising results for such advanced HCC, but it is important to develop the ability to accurately predict chemotherapeutic responses.

Experimental Design: We analyzed the expression of 3,080 genes using a polymerase chain reaction-based array in 20 HCC patients who were treated with combination chemotherapy after reduction surgery. After unsupervised analyses, a supervised classification method for predicting chemotherapeutic responses was constructed. To minimize the number of predictive genes, we used a random permutation test to select only significant ($P < 0.01$) genes. A leave-one-out cross-validation confirmed the gene selection. We also prepared an additional 11 cases for validation of predictive performance.

Results: Hierarchical clustering analysis and principal component analysis with all 3,080 genes revealed distinct

gene expression patterns in responders (those with complete response or partial response) and nonresponders (those with stable disease or progressive disease) to the combination chemotherapy. Using a weighted-voting classification method with either all genes or only significant genes as assessed by permutation testing, the objective responses to treatment were correctly predicted in 17 of 20 cases (accuracy, 85%; positive predictive value, 100%; negative predictive value, 80%). Moreover, patients in the validation dataset could be classified into two distinct prognostic groups using 63 predictive genes.

Conclusions: Molecular analysis of 63 genes can predict the response of patients with advanced HCC and major portal vein tumor thrombi to combination chemotherapy with 5-fluorouracil and interferon- α .

INTRODUCTION

Hepatocellular carcinoma (HCC), the predominant histologic subtype of primary liver cancer, is one of the major causes of death from malignancy worldwide. Although recent progress in both diagnostic and surgical techniques has resulted in considerable improvement in the morbidity and mortality rates, the overall outcome remains far from satisfactory. The prognosis is miserable, particularly in patients with tumors that have invaded the major branch of the portal vein, and the median survival time is several months (1–3). Indeed, the 1-year survival rate was <50% even in resectable cases (3, 4). We and others have recently developed a new treatment regimen for such advanced HCC patients: combination chemotherapy of 5-fluorouracil (5-FU) and interferon (IFN)- α (5, 6). In follow-up studies, about 40% to 50% of patients showed promising responses to the therapy and lived >2 years. However, for patients who did not respond to this treatment, survival was too short to receive another type of treatment. Furthermore, this chemotherapy may have multiple adverse effects, including leukopenia, thrombocytopenia, and depression. Therefore, accurate prediction of chemosensitivity is desirable, not only so that nonresponding patients do not lose a limited chance to take advantage of other possible treatments, but also to eliminate suffering of these patients due to debilitating side effects. Unfortunately, there are currently no useful indicators to distinguish between patients who are likely to respond to this combination chemotherapy and patients who are not.

In this study, genes possessing the ability to predict patient responses to 5-FU and IFN- α combination chemotherapy were selected by gene expression profile analysis using adaptor-tagged competitive polymerase chain reaction (ATAC-PCR) technology, a polymerase chain reaction (PCR)-based array system (7). The

Received 2/6/04; revised 6/9/04; accepted 6/17/04.

Grant support: Grant-in-Aid for the Development of Innovative Technology from the Ministry of Education, Culture, Sports, Science and Technology, Japan; Grants-in-Aid for the Second Term Comprehensive 10-Year Strategy for Cancer Control and Cancer Research from the Ministry of Health and Welfare, Japan; and Grants-in-Aid for Scientific Research on Priority Areas, and Basic Research from the Ministry of Education, Science, Sports, and Culture, Japan.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Morito Monden, Department of Surgery and Clinical Oncology, Graduate School of Medicine, Osaka University, 2-2, Yamadaoka, Suita city, Osaka 565-0871, Japan. Phone: 81-6-6879-3251; Fax: 81-6-6879-3259; E-mail: monden@surg2.med.osaka-u.ac.jp.

©2004 American Association for Cancer Research.

performance of this prediction method was estimated with both a complete cross-validation and an external validation dataset.

MATERIALS AND METHODS

Tissues and Patients. Between August 1998 and January 2003, 20 HCC patients with multiple tumors spreading to bilateral lobes with tumor thrombi in the major branches of the portal vein underwent palliative surgery to reopen the portal flow and recover liver function. After obtaining informed consent, we collected HCC tissue from the main resected tumors and isolated the total RNA from these samples for PCR-based array experiments. All 20 patients had visible tumors in the remaining liver. The patients were then subjected to a treatment regimen of combination chemotherapy with 5-FU and IFN- α as described previously (5). Briefly, patients received continuous arterial infusion of 5-FU (450–500 mg/d) for the initial 2 weeks and subcutaneous IFN- α injection (5 million IU) 3 times per week for 4 weeks. The sizes of the remaining tumors in the liver, which are thought to be intrahepatic metastases from the resected primary tumor, were measured before and after treatment by either abdominal computed tomography scan or ultrasonography. The chemotherapeutic response was clinically evaluated according to the Eastern Cooperative Oncology Group (ECOG) criteria (8). In this study, responders were defined as patients with complete response (CR) or partial response (PR); nonresponders were defined as patients with stable disease or progressive disease (PD).

Furthermore, as an independent validation dataset, we collected liver tissue specimens and clinical data from 11 HCC patients who underwent hepatic resection of all visible tumors between October 1998 and October 2002. Although these patients had tumor(s) with major portal vein tumor thrombi, the area of spreading tumor(s) was limited to two segments of the liver. The treatment regimen of combination chemotherapy and the method of follow-up in the validation dataset were the same as the procedures used in the original dataset. All aspects of our study protocol were approved by the ethics committee of Osaka University Medical School.

Polymerase Chain Reaction-Based Array System. To select genes expressed in liver tissues, we constructed three cDNA libraries: one from a mixture of HCC and nontumorous livers, one from normal livers, and one from metastatic liver cancers, as described previously (9). We designed PCR primers for ATAC-PCR reactions for a total of 2,666 genes from these expressed sequence tag collections. In total, we prepared 3,080 primers for ATAC-PCR; this total includes an additional 414 genes established in previous literature. The specificity of this gene selection provides an advantage over more universal gene sets, such as those selected from the UniGene database, which include genes not expressed in liver tissues. The ATAC-PCR experimental procedure was performed as described previously (10). The complete list of genes and detailed protocols for the ATAC-PCR experimental procedure are available on our web site.⁴

Analysis of Polymerase Chain Reaction-Based Array Data. The relative expression level of each gene was calculated by calibration using a standard mixture of normal

liver tissues, as described previously (11–15). After conversion to a logarithmic scale (base 2), the data matrix was normalized to a median of 0 by standardizing each sample.

As a supervised classification method, we adopted a weighted-voting (WV) algorithm generally used in gene expression profiling (16–19). Briefly, we calculated the signal to noise (S2N) ratio, $S_i = (\mu_R - \mu_N)/(\sigma_R + \sigma_N)$, where μ and σ represent the mean and SD of expression for responders and nonresponders, respectively. The magnitude of the gene vote (v_i) reflects the deviation of the test sample X_i value from the average of the two groups: $v_i = S_i \times (X_i - (\mu_R + \mu_N)/2)$. We summed the v_i values to obtain the total votes for a “good signature” group (V_G) and a “poor signature” group (V_P). The prediction strength is $(V_G - |V_P|)/(V_G + |V_P|)$, and we adopted a threshold of 0. If the strength is a positive number, the test sample belongs to the good signature group. If the strength is a negative number, the test sample belongs to the poor signature group. This model was evaluated by leave-one-out cross-validation (16–21), wherein one sample was randomly withheld, and the model was regenerated with the remaining samples and then used to predict the group of the withheld sample. This process was repeated until every sample was tested, and the cumulative accuracy was recorded. When an external validation dataset was used to evaluate this WV model, we used the S2N ratio (S_i) obtained from the original dataset. In calculating the weighted vote (v_i), the mean expression level of the entire validation dataset was substituted for the average of the two groups $((\mu_R + \mu_N)/2)$ in the original dataset.

Permutation testing, which involves randomly permuting group labels to determine gene-group correlations, was used to determine statistical significance (16–19). The original score, $S_o = |\mu_R - \mu_N|$, of each gene was calculated without permuting the labels (responder or nonresponder). The labels of all of the samples were then randomly permuted, and the scores were recalculated between two groups consisting of the new members. Repetition of this permutation 50,000 times provides a score number (N_o) larger than the original score (S_o). For each gene, the permutation P value was determined by $P = N_o/50,000$. To estimate the false discovery rate, which is the percentage of genes identified by chance (22), we calculated the Q value of every gene using downloaded software.⁵ The Q value provides a measure of each feature’s significance, automatically taking into account the problem of multiple testing (23).

Hierarchical cluster analysis using an unweighted pair group method using arithmetic mean method with Pearson’s correlation and principal component analysis (PCA) was performed using GeneMaths 2.0 software. Other statistical analyses were performed using StatView 5.0J software. Correlations between the responses to our combination chemotherapy and clinicopathological parameters were evaluated by the χ^2 test with the Yates correction. Overall and disease-free survival rates were calculated using the Kaplan-Meier method. Differences in survival curves were estimated by the log-rank test.

⁴ http://love2.aist-nara.ac.jp/laboratory/index_frame.html.

⁵ <http://faculty.washington.edu/~jstorey/qvalue>.

Table 1 Characteristics of 20 patients

Case no.	Age (y)/sex	HBs Ag	HCV Ab	Child grade	Cirrhosis	AFP*	Edmondson classification	Main tumor size (cm)	Cycles of chemotherapy	Response†
1	74/M	-	+	A	Present	Low	3	3.0	3	CR
2	67/F	-	+	A	Present	High	3	9.2	3	CR
3	49/F	+	-	A	Absent	High	4	6.8	3	PD
4	47/M	+	-	A	Present	High	3	12.0	3	PD
5	51/F	-	-	B	Absent	High	3	15.0	3	PR
6	56/M	-	-	B	Present	Low	2	0.8	3	PD
7	69/M	-	+	A	Absent	High	3	8.0	3	CR
8	66/M	+	+	A	Present	High	3	4.0	3	CR
9	65/M	-	+	B	Present	High	3	1.3	2	PD
10	53/M	+	-	A	Present	High	3	17.0	3	PD
11	52/M	+	-	B	Absent	High	3	18.0	3	CR
12	66/M	-	-	B	Present	High	3	3.5	1	PD
13	39/M	+	-	B	Present	High	3	4.5	1	PD
14	56/M	-	+	A	Present	High	3	5.5	3	SD
15	67/M	-	+	A	Present	High	3	3.2	2	PD
16	47/M	+	-	A	Absent	High	3	13.0	3	PD
17	70/M	-	+	A	Absent	Low	3	9.4	3	CR
18	70/M	-	+	A	Absent	High	3	9.8	3	CR
19	71/M	-	+	A	Absent	High	3	9.0	3	PD
20	29/M	+	-	A	Absent	High	3	18.0	2	PD

Abbreviations: HBs Ag, hepatitis B surface antigen; HCV Ab, hepatitis C virus antibody; SD, stable disease.

* Divided into either low (≤ 50 ng/mL) or high (> 50 ng/mL).

† Assessment based on ECOG criteria.

RESULTS

Patient Characteristics. The characteristics of the 20 HCC patients in the original dataset are shown in Table 1. All patients had multiple tumors spreading to bilateral lobes with tumor thrombi in the major branches of the portal vein and were treated with 5-FU and IFN- α combination chemotherapy after reduction surgery. Fifteen of the 20 patients were treated with three cycles of the above-mentioned regimen. The remaining five patients received only one or two cycles of 5-FU due to extensive progression of the tumors. According to the ECOG criteria of objective response, 8 patients (40%) were classified as responders, demonstrating either CR or PR for at least 4 weeks. The remaining 12 patients (60%) were classified as nonresponders, exhibiting either stable disease or PD. The clinicopathological characteristics of responders (CR and PR) and nonresponders (stable disease and PD) were compared by the χ^2 test (Table 2). There were no significant differences ($P < 0.05$) in any factor between the two response groups.

Unsupervised Analyses. We performed a hierarchical cluster analysis of the samples using all 3,080 genes. When the clinical samples were sorted on the basis of similarity in gene expression, the samples could be separated according to the response to 5-FU and IFN- α combination chemotherapy, *i.e.*, responders and nonresponders, with only a few exceptions (Fig. 1). Next, we applied PCA, a statistical method for reducing the number of data dimensions, to more simply present the relationships between the samples. On displaying the expression patterns of all 3,080 genes in three-dimensional space, we observed that most responders and nonresponders were located separately, indicating distinct gene expression patterns (Fig. 2).

Supervised Analyses. To construct a molecular prediction system, we used a supervised approach, a WV algorithm using S2N correlation ratios. The ability of this method to

Table 2 Clinicopathological characteristics between responders and nonresponders

Characteristics	Responders (n = 8)	Nonresponders (n = 12)	P
Age (y)			>0.05
<65	2	8	
≥ 65	6	4	
Sex			>0.05
Male	6	11	
Female	2	1	
HBs Ag			>0.05
Negative	6	6	
Positive	2	6	
HCV Ab			>0.05
Negative	2	8	
Positive	6	4	
Child grade			>0.05
A	6	8	
B	2	4	
Liver cirrhosis			>0.05
Absent	5	4	
Present	3	8	
AFP (ng/mL)			>0.05
≤ 50	2	1	
> 50	6	11	
Histological grade			>0.05
Moderate	0	1	
Poor	8	10	
Undiff	0	1	
Tumor size (cm)			>0.05
≤ 5	2	5	
> 5	6	7	

Abbreviations: Moderate, moderately differentiated hepatocellular carcinoma; Poor, poorly differentiated hepatocellular carcinoma; Undiff, undifferentiated carcinoma.




Fig. 1 Hierarchical clustering of 3,080 genes in 20 HCC cases. Each *column* represents a single gene, and each *row* represents a patient sample. The tissue type of each sample is designated by a *bar* at the *right*, and *green* and *red* bars indicate responders and nonresponders, respectively. The *color scale* at the *bottom* indicates the relative expression levels in terms of SDs from the mean.

predict patient responses to 5-FU and IFN- α combination chemotherapy was evaluated by leave-one-out cross-validation. In this first cross-validation analysis, we used all 3,080 genes to avoid any selection bias (Fig. 3A). The prediction accuracy of

this WV method was 85.0%, and its 95% confidence interval ranged from 66.0% to 95.9%. The positive and negative predictive values were 100.0% and 80.0%, respectively.

These results demonstrate that this method provides a




Fig. 2 PCA in 20 HCC cases. The variation is reduced to three-dimensional space; three components represent 37.0% of the total variance. Each *sphere* represents a single sample: *green* for a responder and *red* for a nonresponder.

○ responders ● non-responders




Fig. 3 Flow chart for selecting samples and gene set for analysis. A, the first approach, which used the original dataset and all genes. B, the second approach, which used the original dataset and only significant genes ($P < 0.01$). C, the third approach, which used the validation dataset and 63 significant genes selected with the original dataset.

valuable prediction of the chemotherapeutic response. To determine whether all 3,080 genes are necessary for prediction, we compared the prediction accuracy of a small number of genes to the prediction accuracy of all 3,080 genes. When the predictive genes were defined as genes with significant P values ($P < 0.01$) by the random permutation tests (Fig. 3B), the complete cross-validation without information leakage showed an identical performance with the initial cross-validation using all 3,080 genes. We therefore selected only 63 genes with significant P values between the 8 responder patients and the 12 nonresponder patients (Table 3). The median of their Q values, which is a measure in terms of the false discovery rate, was $<25\%$. The expression patterns of these 63 genes exhibited distinct profiles between the two groups (Fig. 4).

To further evaluate our prediction system, we prepared an independent dataset, which consisted of 11 HCC patients with major portal vein tumor thrombi. Although these patients had no visible tumors after resection before combination chemotherapy, the risk of recurrence within the early postoperative period due to intrahepatic micrometastasis was very high (3, 4, 24). Moreover, in these advanced HCC cases with major portal vein tumor thrombi, the survival time was strongly correlated to the chemotherapeutic response. In fact, the median survival times of responders and nonresponders in the original dataset were 28 and 7 months, respectively, and the difference was statistically significant ($P = 0.002$). We therefore performed overall survival analysis and disease-free survival analysis instead of estimation to determine the prediction accuracy of objective response. Our prediction method using 63 genes classified patients into either a good signature group (those who were predicted to have good responses to this chemotherapy) or a poor signature group (those who were predicted to have poor responses; Fig. 3C). The overall survival rates were significantly different between these two predicted groups ($P = 0.001$; Fig. 5A). In addition, the good signature group had a distinctly better prognosis for disease-free survival than the poor signature group ($P = 0.002$; Fig. 5B).

DISCUSSION

HCC remains a major problem worldwide and appears to be increasing in developed Western countries (25). HCC is notorious for its poor prognosis because the overall 5-year survival rate after resection has remained as poor as 35% to 50%, even if curative surgery is performed (26–28). Moreover, once the tumor invades the major branch of the portal vein, the prognosis is extremely poor (1–4). Various complications including gastrointestinal bleeding, ascites, and hemorrhage from esophageal varices are provoked by stenosis of the portal vein. Until now, there has been no standard treatment for advanced HCC with portal tumor thrombi. However, we and others have recently developed a new combination chemotherapy regimen using intra-arterial 5-FU and subcutaneous IFN- α (5, 6), which has proven to be effective in about half of such advanced HCC patients. In this study, we have demonstrated that a molecular prediction method using expression data of 63 genes was able to distinguish patients who responded to this therapy from those who did not respond.

DNA microarray technology allows parallel expression analysis of thousands of genes to address complex questions in tumor biology. Many trials predicting the prognosis of various human malignancies have been reported using DNA microarrays (29–31). Moreover, a novel prediction method using gene expression profiling has recently been reported for the treatment of breast cancer patients with the taxanes (docetaxel and paclitaxel; ref. 21). Although DNA microarrays have contributed to such expression profiling studies to some degree, DNA microarrays can detect only a fraction of the changes in gene expression detectable by reverse transcription-PCR (32). We therefore performed a high-throughput quantitative PCR based on ATAC-PCR (7) to analyze the genetic differences in HCC. This assay requires smaller amounts of RNA than DNA microarray analysis. PCR-based analysis of selected genes costs much less than DNA microarrays, which are likely to require at least several hundred spotted genes for diagnosis. The aforementioned ben-

Table 3 The list of 63 predictive genes

P	Up/down	UniGene ID	Gene symbol	Gene definition
0.0001	Down	Hs.74088	<i>EGR3</i>	Early growth response 3
0.0002	Up	Hs.432818	<i>MFAP3</i>	Microfibrillar-associated protein 3
0.0002	Down	-	<i>EST</i>	DNC15 <i>Homo sapiens</i> cDNA
0.0003	Up	Hs.151738	<i>MMP9*</i>	Matrix metalloproteinase 9
0.0005	Down	Hs.257697	<i>PDCD4</i>	Programmed cell death 4
0.0006	Up	Hs.337534	<i>EST</i>	<i>H. sapiens</i> clone 25061 mRNA sequence
0.0009	Down	Hs.293189	<i>FLJ37853</i>	<i>H. sapiens</i> cDNA FLJ37853 fis
0.0010	Down	Hs.6838	<i>ARHE</i>	Ras homolog gene family, member E
0.0010	Down	Hs.79078	<i>MAD2L1</i>	MAD2 mitotic arrest deficient-like 1
0.0011	Down	Hs.158244	<i>NMNAT2</i>	Nicotinamide nucleotide adenylyltransferase 2
0.0012	Down	Hs.175483	<i>FLJ22016</i>	<i>H. sapiens</i> cDNA: FLJ22016 fis
0.0012	Down	Hs.511400	<i>SDNI</i>	Staphylococcal nuclease domain containing 1
0.0012	Up	Hs.301242	<i>MIDORI</i>	Myocytic induction/differentiation originator
0.0013	Down	Hs.471660	<i>FLJ36229</i>	<i>H. sapiens</i> cDNA FLJ36229 fis
0.0014	Down	Hs.81874	<i>MGST2</i>	Microsomal glutathione S-transferase 2
0.0015	Down	Hs.414156	<i>CTPS2</i>	CTP synthase II
0.0015	Up	Hs.118633	<i>OASL†</i>	2'-5'-oligoadenylate synthetase-like
0.0016	Down	Hs.406266	<i>HK2*</i>	Hexokinase 2
0.0017	Down	Hs.75309	<i>EEF2</i>	Eukaryotic translation elongation factor 2
0.0019	Down	Hs.121150	<i>Taxilin</i>	Taxilin
0.0019	Down	Hs.67201	<i>NT5C</i>	5', 3'-nucleotidase, cytosolic
0.0021	Down	Hs.417369	<i>CLNSIA</i>	Chloride channel, nucleotide-sensitive, 1A
0.0023	Down	Hs.13350	<i>UNC5B</i>	Unc-5 homolog B
0.0024	Down	Hs.345908	<i>TSC</i>	Hypothetical protein FLJ20607
0.0025	Down	Hs.2007	<i>TNFSF6*</i>	Tumor necrosis factor (ligand) superfamily, member 6
0.0025	Up	Hs.440899	<i>KIAA1691</i>	KIAA1691 protein
0.0025	Down	Hs.143873	<i>S100A10</i>	S100 calcium binding protein A10
0.0026	Down	Hs.272630	<i>ATP5VID</i>	ATPase, H+ transporting, lysosomal 34 kDa, V1 subunit D
0.0027	Down	Hs.413226	<i>ASB4</i>	Ankyrin repeat and SOCS box-containing 4
0.0027	Down	Hs.226391	<i>AGR2</i>	Anterior gradient 2 homolog
0.0027	Down	Hs.3610	<i>KIAA0205</i>	KIAA0205 gene product
0.0029	Down	Hs.335118	<i>JJAZ1</i>	Joined to JAZF1
0.0031	Down	Hs.12840	<i>LOC285859</i>	Hypothetical protein LOC285859
0.0031	Down	Hs.82324	<i>KIAA0157</i>	KIAA0157 protein
0.0031	Up	Hs.421825	<i>PPA2</i>	Inorganic pyrophosphatase 2
0.0036	Up	Hs.143773	<i>eIF3k</i>	Eukaryotic translation initiation factor 3 subunit k
0.0036	Up	Hs.380877	<i>SGKL</i>	Serum/glucocorticoid regulated kinase-like
0.0038	Down	Hs.387183	<i>BLMH</i>	Bleomycin hydrolase
0.0038	Up	Hs.114648	<i>ERG-1</i>	Estrogen-regulated gene 1
0.0042	Down	Hs.34012	<i>BRCA2</i>	Breast cancer 2, early onset
0.0048	Down	Hs.512628	<i>PPP2R2A</i>	Protein phosphatase 2, regulatory subunit B, α isoform
0.0051	Up	Hs.377975	<i>IGKC</i>	Immunoglobulin κ constant
0.0053	Down	Hs.435765	<i>ENPEP†</i>	Glutamyl aminopeptidase
0.0055	Down	Hs.356076	<i>BIRC4†</i>	Baculoviral IAP repeat-containing 4
0.0055	Down	Hs.220745	<i>FLJ10244</i>	Ral-A exchange factor RalGPS2
0.0060	Down	Hs.376531	<i>LOC283167</i>	<i>H. sapiens</i> LOC283167
0.0061	Down	Hs.128410	<i>GLS</i>	Glutaminase
0.0061	Down	Hs.356876	<i>GPR125</i>	G protein-coupled receptor 125
0.0065	Down	Hs.250696	<i>KDELR3</i>	KDEL endoplasmic reticulum protein retention receptor 3
0.0066	Down	Hs.10949	<i>ERO1L</i>	ERO1-like
0.0068	Down	Hs.380446	<i>LENG8</i>	Leukocyte receptor cluster member 8
0.0070	Up	Hs.22129	<i>DJ1042K10.2</i>	Hypothetical protein DJ1042K10.2
0.0070	Down	Hs.93177	<i>IFNB1</i>	Interferon β 1, fibroblast
0.0072	Down	Hs.389461	<i>SMAP-1</i>	Smooth muscle cell-associated protein 1
0.0074	Up	Hs.512654	<i>SET07</i>	PR/SET domain containing protein 07
0.0080	Down	Hs.432574	<i>POLR2H</i>	Polymerase II polypeptide H
0.0080	Down	Hs.183986	<i>PVRL2</i>	Poliovirus receptor-related 2
0.0088	Down	Hs.406351	<i>EST</i>	<i>H. sapiens</i> clone IMAGE:120162 mRNA sequence
0.0089	Down	Hs.2533	<i>ALDH9A1</i>	Aldehyde dehydrogenase 9 family, member A1
0.0090	Down	Hs.282410	<i>CALM1†</i>	Calmodulin 1
0.0091	Down	Hs.37288	<i>NR1D2</i>	Nuclear receptor subfamily 1, group D, member 2
0.0096	Down	Hs.11355	<i>TMPO</i>	Thymopoietin
0.0098	Up	Hs.299208	<i>COQ3</i>	Coenzyme Q3 homolog, methyltransferase

NOTE. P values were calculated by random permutation test. Up-regulation and down-regulation were defined as expression in responders compared with nonresponders.

* Genes previously reported to be associated with sensitivity to 5-FU.

† Genes previously reported to be associated with sensitivity to IFN.

responder *non-responder*




Fig. 4 The expression pattern of the 63 predictive genes in 20 HCC cases. Each column represents a patient sample, and each row represents a single gene.




Fig. 5 Overall survival curves (A) and disease-free survival curves (B) calculated using the Kaplan-Meier method for the validation dataset. Differences in survival curves were estimated by the log-rank test.

efits and the strength of this system for cancer research, established in previous reports on HCC, breast, and colorectal cancers make this technique a powerful means to obtain a better understanding of the molecular characteristics of cancers (11–15).

In the unsupervised analyses with all examined genes, the difference of expression patterns between responders (CR and PR) and nonresponders (stable disease and PD) was clear. Of course, it was controversial how to divide two response groups. However, the response to this regimen with 5-FU and IFN- α was usually typical; that is, PR or stable disease cases were rare, as shown in Table 1. Clinically, radiologically, and biochemically, the PR case in this study revealed a near CR, and the stable disease case was similar to PD cases. We therefore classified the PR patient as a responder and the stable disease patient as a nonresponder due to the conventional grouping method (33, 34). In supervised analyses, we selected 63 significant genes by a random permutation test to minimize the number of predictive genes. These 63 predictive genes included some genes that have previously been reported to be associated with sensitivity to 5-FU or IFN. For example, a gene related to 5-FU sensitivity is tumor necrosis factor ligand superfamily member 6 (TNFSF6), also called FAS (CD95) ligand, which plays a major role in the induction of apoptosis in response to a variety of extracellular signals, including anticancer drugs. In hepatoma cells or human breast cancer, up-regulation of

TNFSF6 was associated with poor sensitivity to 5-FU chemotherapy (35, 36). Matrix metalloproteinase 9 (MMP9), a member of the secreted zinc metalloprotease group, is also inhibited by 5-FU treatment (37). Hexokinase 2 (HK2) plays an important role in intracellular glucose metabolism and is strongly associated with the metabolism of 2'-fluoro-2'-deoxy-D-mannose, which has been correlated to 5-FU response (38). A gene involved in sensitivity to IFN- α is OASL, a member of the 2'-5'-oligoadenylate synthetase gene families. OASL plays an important role in the antiviral effects of IFN and is involved in apoptosis and control of cellular growth (39, 40). Baculoviral IAP repeat-containing 4 (BIRC4; XIAP), encodes a protein that significantly inhibits apoptosis. IFN inhibited BIRC4 function by inducing XIAP associated factor-1 (XAF-1; ref. 41). Glutamyl aminopeptidase is a differentiation-related kidney glycoprotein of 160 kDa (gp160), and the absence of glutamyl aminopeptidase expression was predictive of IFN- α sensitivity in renal cell carcinoma (42). Calmodulin-1 (CALM1) plays a role in growth and the cell cycle and in signal transduction. Calmodulin inhibitors affect not only the uptake and processing of IFN inducers but also the release of IFN from induced cells (43). Our selection approach to these predictive genes was confirmed with both a complete cross-validation and an external validation dataset.

In our univariate analysis with clinicopathological factors, there was no significant indicator that distinguished responders and nonresponders. Patt *et al.* (6) reported that low levels (≤ 50 ng/mL) of serum α -fetoprotein (AFP) were a predictor of response to 5-FU and IFN- α combination chemotherapy, but the serum AFP level was not a significant indicator of chemotherapeutic response in our study. Because we treated only extremely advanced HCC patients with major portal vein tumor thrombi with this combination therapy, only 3 of the 20 patients showed low levels of serum AFP. It may be too early to make conclusions regarding this factor, but we believe that another appropriate method will be required to more accurately predict the response to this treatment.

Indeed, this combination chemotherapy of 5-FU and IFN- α is promising for advanced HCC, but we need to consider the side effects. IFN- α and/or 5-FU sometimes induce severe adverse effects, including myelosuppression, fever, and depression (44). In particular, adverse myelosuppression is an important factor in HCC cases not only because thrombocytopenia and/or leukopenia are frequently present before chemotherapy, but also because treatment often has to be discontinued due to these side effects. Moreover, the survival of patients with such extremely advanced HCC was too short to allow a secondary treatment (1–4). Therefore, accurate prediction of sensitivity to the first-line chemotherapy is necessary for advanced HCC patients so that they do not lose their limited chance to receive treatment and the patients who will not respond to the therapy can be protected from debilitating side effects.

These results illustrate the potential of biological technology to advance diagnostic methods, allowing physicians to plan beyond empirical results toward a more molecularly well-defined, personalized therapy. Particularly for patients with poor prognosis, such as advanced HCC patients, accurate prediction of chemotherapeutic responses before treatment is necessary. Because our molecular prediction system involves only a small

number of predictive genes and a simplified algorithm that does not require either statistical software or specialists, this system should easily lead to future clinical application.

ACKNOWLEDGMENTS

We thank Chiyuri Maruyama, Keiko Ikenaga, Mihoko Yoshino, and Satoko Kinjo for expert technical assistance as well as Dr. Hideo Ota for helpful suggestions.

REFERENCES

- Raoul JL, Guyader D, Bretagne JF, et al. Randomized controlled trial for hepatocellular carcinoma with portal vein thrombosis: intra-arterial iodine-131-iodized oil versus medical support. *J Nucl Med* 1994;35:1782-7.
- Lee HS, Kim JS, Choi JJ, et al. The safety and efficacy of transcatheter arterial chemoembolization in the treatment of patients with hepatocellular carcinoma and main portal vein obstruction. A prospective controlled study. *Cancer (Phila)* 1997;79:2087-94.
- Tanaka A, Morimoto T, Yamaoka Y. Implications of surgical treatment for advanced hepatocellular carcinoma with tumor thrombi in the portal vein. *Hepatogastroenterology* 1996;43:637-43.
- Ikai I, Yamaoka Y, Yamamoto Y, et al. Surgical intervention for patients with stage IV-A hepatocellular carcinoma without lymph node metastasis: proposal as a standard therapy. *Ann Surg* 1998;227:433-9.
- Sakon M, Nagano H, Dono K, et al. Combined intraarterial 5-fluorouracil and subcutaneous interferon-alpha therapy for advanced hepatocellular carcinoma with tumor thrombi in the major portal branches. *Cancer (Phila)* 2002;94:435-42.
- Patt YZ, Yoffe B, Chamsangavej C, et al. Low serum alpha-fetoprotein level in patients with hepatocellular carcinoma as a predictor of response to 5-FU and interferon-alpha-2b. *Cancer (Phila)* 1993;72:2574-82.
- Kato K. Adaptor-tagged competitive PCR: a novel method for measuring relative gene expression. *Nucleic Acids Res* 1997;25:4694-6.
- Oken MM, Creech RH, Tormey DC, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol* 1982;5:649-55.
- Matoba R, Kato K, Saito S, et al. Gene expression in mouse cerebellum during its development. *Gene (Amst)* 2000;241:125-31.
- Matoba R, Kato K, Kurooka C, et al. Correlation between gene functions and developmental expression patterns in the mouse cerebellum. *Eur J Neurosci* 2000;12:1357-71.
- Kurokawa Y, Matoba R, Takemasa I, et al. Molecular features of non-B, non-C hepatocellular carcinoma: a PCR-array gene expression profiling study. *J Hepatol* 2003;39:1007-15.
- Kurokawa Y, Matoba R, Nakamori S, et al. PCR-array gene expression profiling of hepatocellular carcinoma. *J Exp Clin Cancer Res* 2004;23:135-41.
- Kurokawa Y, Matoba R, Takemasa I, et al. Molecular-based prediction of early recurrence in hepatocellular carcinoma. *J Hepatol* 2004;41:284-91.
- Iwao K, Matoba R, Ueno N, et al. Molecular classification of primary breast tumors possessing distinct prognostic properties. *Hum Mol Genet* 2002;11:199-206.
- Muro S, Takemasa I, Oba S, et al. Identification of expressed genes linked to malignancy of human colorectal carcinoma by parametric clustering of quantitative expression data. *Genome Biol* <http://genome-biology.com/2003/4/3/R21>.
- Golub TR, Slonim DK, Tamayo P, et al. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science (Wash DC)* 1999;286:531-7.
- MacDonald TJ, Brown KM, LaFleur B, et al. Expression profiling of medulloblastoma: PDGFRA and the RAS/MAPK pathway as therapeutic targets for metastatic disease. *Nat Genet* 2001;29:143-52.
- Pomeroy SL, Tamayo P, Gaasenbeek M, et al. Prediction of central nervous system embryonal tumor outcome based on gene expression. *Nature (Lond)* 2002;415:436-42.
- Armstrong SA, Staunton JE, Silverman LB, et al. MLL translocations specify a distinct gene expression profile that distinguishes a unique leukemia. *Nat Genet* 2002;30:41-7.
- Ramaswamy S, Ross KN, Lander ES, Golub TR. A molecular signature of metastasis in primary solid tumors. *Nat Genet* 2003;33:49-54.
- Chang JC, Wooten EC, Tsimelzon A, et al. Gene expression profiling for the prediction of therapeutic response to docetaxel in patients with breast cancer. *Lancet* 2003;362:362-9.
- Tusher VG, Tibshirani R, Chu G. Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci USA* 2001;98:5116-21.
- Storey JD, Tibshirani R. Statistical significance for genome-wide studies. *Proc Natl Acad Sci USA* 2003;100:9440-5.
- Sakon M, Umeshita K, Nagano H, et al. Clinical significance of hepatic resection in hepatocellular carcinoma. *Arch Surg* 2000;135:1456-9.
- Befeler AS, Di Bisceglie AM. Hepatocellular carcinoma: diagnosis and treatment. *Gastroenterology* 2002;122:1609-19.
- Lai EC, Fan ST, Lo CM, et al. Hepatic resection for hepatocellular carcinoma. An audit of 343 patients. *Ann Surg* 1995;221:291-8.
- Poon RT, Fan ST, Lo CM, et al. Improving survival results after resection of hepatocellular carcinoma: a prospective study of 377 patients over 10 years. *Ann Surg* 2001;234:63-70.
- Ono T, Yamanoi A, Nazmy EI, et al. Adjuvant chemotherapy after resection of hepatocellular carcinoma causes deterioration of long-term prognosis in cirrhotic patients: metaanalysis of three randomized controlled trials. *Cancer (Phila)* 2001;91:2378-85.
- Beer DG, Kardias SL, Huang CC, et al. Gene-expression profiles predict survival of patients with lung adenocarcinoma. *Nat Med* 2002;8:816-24.
- Rosenwald A, Wright G, Chan WC, et al. The use of molecular profiling to predict survival after chemotherapy for diffuse large-B-cell lymphoma. *N Engl J Med* 2002;346:1937-47.
- van de Vijver MJ, He YD, van't Veer LJ, et al. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med* 2002;347:1999-2009.
- Holland MJ. Transcript abundance in yeast varies over six orders of magnitude. *J Biol Chem* 2002;277:14363-6.
- Rowinsky EK, Jiroutek M, Bonomi P, Johnson D, Baker SD. Paclitaxel steady-state plasma concentration as a determinant of disease outcome and toxicity in lung cancer patients treated with paclitaxel and cisplatin. *Clin Cancer Res* 1999;5:767-74.
- Cohen V, Panet-Raymond V, Sabbaghian N, et al. Methylene-tetrahydrofolate reductase polymorphism in advanced colorectal cancer: a novel genomic predictor of clinical response to fluoropyrimidine-based chemotherapy. *Clin Cancer Res* 2003;9:1611-5.
- Eichhorst ST, Muller M, Li-Weber M, et al. A novel AP-1 element in the CD95 ligand promoter is required for induction of apoptosis in hepatocellular carcinoma cells upon treatment with anticancer drugs. *Mol Cell Biol* 2000;20:7826-37.
- Sjostrom J, Blomqvist C, von Boguslawski K, et al. The predictive value of bcl-2, bax, bcl-xL, bag-1, fas, and fasL for chemotherapy response in advanced breast cancer. *Clin Cancer Res* 2002;8:811-6.
- Ragoowansi R, Khan U, Brown RA, McGrouther DA. Reduction in matrix metalloproteinase production by tendon and synovial fibroblasts after a single exposure to 5-fluorouracil. *Br J Plast Surg* 2001;54:283-7.
- McSheehy PM, Leach MO, Judson IR, Griffiths JR. Metabolites of 2'-fluoro-2'-deoxy-D-glucose detected by ¹⁹F magnetic resonance spectroscopy in vivo predict response of murine RIF-1 tumors to 5-fluorouracil. *Cancer Res* 2000;60:2122-7.

39. Hartmann R, Olsen HS, Widder S, Jorgensen R, Justesen J. p59OASL, a 2'-5' oligoadenylate synthetase like protein: a novel human gene related to the 2'-5' oligoadenylate synthetase family. *Nucleic Acids Res* 1998;26:4121-8.
40. Eskildsen S, Justesen J, Schierup MH, Hartmann R. Characterization of the 2'-5'-oligoadenylate synthetase ubiquitin-like family. *Nucleic Acids Res* 2003;31:3166-73.
41. Chawla-Sarkar M, Lindner DJ, Liu YF, et al. Apoptosis and interferons: role of interferon-stimulated genes as mediators of apoptosis. *Apoptosis* 2003;8:237-49.
42. Nanus DM, Pfeffer LM, Bander NH, Bahri S, Albino AP. Antiproliferative and antitumor effects of alpha-interferon in renal cell carcinomas: correlation with the expression of a kidney-associated differentiation glycoprotein. *Cancer Res* 1990;50:4190-4.
43. Lin HY, Davis PJ, Davis FB, Chadha KC, Thacore HR. Effect of calmodulin antagonists on the interferon system: induction and action of interferons. *J Interferon Res* 1990;10:39-46.
44. Sachs E, Di Bisceglie AM, Dusheiko GM, et al. Treatment of hepatocellular carcinoma with recombinant leucocyte interferon: a pilot study. *Br J Cancer* 1985;52:105-9.

Molecular-based prediction of early recurrence in hepatocellular carcinoma[☆]

Yukinori Kurokawa^{1,2}, Ryo Matoba², Ichiro Takemasa¹, Hiroaki Nagano¹,
Keizo Dono¹, Shoji Nakamori¹, Koji Umeshita¹, Masato Sakon¹, Noriko Ueno²,
Shigeyuki Oba³, Shin Ishii³, Kikuya Kato², Morito Monden^{1,*}

¹Department of Surgery and Clinical Oncology, Graduate School of Medicine, Osaka University, 2-2, Yamadaoka, Suita city, Osaka 565-0871, Japan

²Taisho Laboratory of Functional Genomics, Nara Institute of Science and Technology, Nara 630-0101, Japan

³Laboratory of Theoretical Life Science, Nara Institute of Science and Technology, Nara 630-0101, Japan

See Editorial, pages 336–339

Background/Aims: Hepatocellular carcinoma (HCC) has a very poor prognosis, due to the high incidence of tumor recurrence. As the current morphological indicators are often insufficient for therapeutic decisions, we sought to identify additional biologic indicators for early recurrence.

Methods: We analyzed gene expression using a PCR-based array of 3072 genes in 100 HCC patients. Informative genes predicting early intrahepatic recurrence were selected by random permutation testing, and a weighted voting prediction method was constructed. Following estimation of prediction accuracy, a multivariate Cox analysis was performed.

Results: By permutation testing, we selected 92 genes demonstrated distinct expression patterns differing significantly between recurrence cases and recurrence-free cases. Our prediction method, using the 20 top-ranked genes, correctly predicted the early intrahepatic recurrence for 29 of 40 cases within the validation group, and the odds ratio was 6.8 (95% CI 1.7–27.5, $P = 0.010$). The 2-year recurrence rates in the patients with the good signature and those with the poor signature were 29.4 and 73.9%, respectively. Multivariate Cox analysis revealed that molecular-signature was an independent indicator for recurrence (hazard ratio 3.82, 95% CI 1.44–10.10, $P = 0.007$).

Conclusions: Our molecular-based prediction method using 20 genes is clinically useful to predict early recurrence of HCC.

© 2004 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.

Keywords: Hepatocellular carcinoma; Permutation testing; Multivariate Cox analysis

1. Introduction

Hepatocellular carcinoma (HCC), the predominant histological subtype of primary liver cancer, is one of the major causes of death from malignancy throughout the world. HCC is notorious for poor prognosis, due to its invasiveness and frequent association with cirrhosis. Although recent progress in diagnostic procedures and

surgical techniques have resulted in considerably improved morbidity and mortality rates, the overall five-year survival rate following resection has remained as poor as 35–50% due to high recurrence rate in the remnant liver following resection [1–3]. Classification of primary HCC tumors according to clinicopathological findings has been proposed and applied to assess the risk of recurrence following resection [4,5]. Although the UICC TNM staging has been the most prevalent method of estimating the early prognosis of HCC patients after surgery, it is not yet accepted as the definitive staging system for HCC [6,7]. This uncertainty inherent in the conventional classification methods may be attributed to incomplete exploration of the biological differences of tumors in patients belonging to identical

Received 6 January 2004; received in revised form 23 April 2004; accepted 28 April 2004; available online 26 May 2004

* Supplementary data associated with this article can be found at doi: 10.1016/j.jhep.2004.04.031

* Corresponding author. Tel.: +81-6-6879-3251; fax: +81-6-6879-3259.
E-mail address: monden@surg2.med.osaka-u.ac.jp (M. Monden).