

- genome by *in vitro* processing analysis. *Proc. Natl. Acad. Sci. USA* **88**:5547–5551.
17. Hope, R. G., and J. McLauchlan. 2000. Sequence motifs required for lipid droplet association and protein stability are unique to the hepatitis C virus core protein. *J. Gen. Virol.* **81**:1913–1925.
  18. Hope, R. G., D. J. Murphy, and J. McLauchlan. 2002. The domains required to direct core proteins of hepatitis C virus and GB virus-B to lipid droplets share common features with plant oleosin proteins. *J. Biol. Chem.* **277**:4261–4270.
  19. Hüseyin, P., H. Langen, J. Mous, and H. Jacobsen. 1996. Hepatitis C virus core protein: carboxy-terminal boundaries of two processed species suggest cleavage by a signal peptide peptidase. *Virology* **224**:93–104.
  20. Isoyama, T., S. Kuge, and A. Nomoto. 2002. The core protein of hepatitis C virus is imported into the nucleus by transport receptor Kap123p but inhibits Kap121p-dependent nuclear import of yeast API-like transcription factor in yeast cells. *J. Biol. Chem.* **277**:39634–39641.
  21. Lai, M. M., and C. F. Ware. 2000. Hepatitis C virus core protein: possible roles in viral pathogenesis. *Curr. Top. Microbiol. Immunol.* **242**:117–134.
  22. Leung, S. W., M. T. Harreman, M. R. Hodel, A. E. Hodel, and A. H. Corbett. 2003. Dissection of the karyopherin alpha nuclear localization signal (NLS)-binding groove: functional requirements for NLS binding. *J. Biol. Chem.* **278**:41947–41953.
  23. Liu, Q., C. Tackney, R. A. Bhat, A. M. Prince, and P. Zhang. 1997. Regulated processing of hepatitis C virus core protein is linked to subcellular localization. *J. Virol.* **71**:657–662.
  24. Lo, S. Y., F. Masiarz, S. B. Hwang, M. M. Lai, and J. H. Ou. 1995. Differential subcellular localization of hepatitis C virus core gene products. *Virology* **213**:455–461.
  25. Lo, S. Y., M. Selby, M. Tong, and J. H. Ou. 1994. Comparative studies of the core gene products of two different hepatitis C virus isolates: two alternative forms determined by a single amino acid substitution. *Virology* **199**:124–131.
  26. Lu, W., A. Strohecker, and J. H. Ou. 2001. Post-translational modification of the hepatitis C virus core protein by tissue transglutaminase. *J. Biol. Chem.* **276**:47993–47999.
  27. Machida, K., K. Tsukiyama-Kohara, E. Seike, S. Tone, F. Shibasaki, M. Shimizu, H. Takahashi, Y. Hayashi, N. Funata, C. Taya, H. Yonekawa, and M. Kohara. 2001. Inhibition of cytochrome c release in Fas-mediated signaling pathway in transgenic mice induced to express hepatitis C viral proteins. *J. Biol. Chem.* **276**:12140–12146.
  28. Matsumoto, M., S. B. Hwang, K. S. Jeng, N. Zhu, and M. M. Lai. 1996. Homotypic interaction and multimerization of hepatitis C virus core protein. *Virology* **218**:43–51.
  29. McLauchlan, J. 2000. Properties of the hepatitis C virus core protein: a structural protein that modulates cellular processes. *J. Viral Hepatitis* **7**:2–14.
  30. McLauchlan, J., M. K. Lemberg, G. Hope, and B. Martoglio. 2002. Intramembrane proteolysis promotes trafficking of hepatitis C virus core protein to lipid droplets. *EMBO J.* **21**:3980–3988.
  31. Mihara, K. 2000. Targeting and insertion of nuclear-encoded preproteins into the mitochondrial outer membrane. *Bioessays* **22**:364–371.
  32. Moradpour, D., C. Englert, T. Wakita, and J. R. Wands. 1996. Characterization of cell lines allowing tightly regulated expression of hepatitis C virus core protein. *Virology* **222**:51–63.
  33. Moriishi, K., T. Okabayashi, K. Nakai, K. Moriya, K. Koike, S. Murata, T. Chiba, K. Tanaka, R. Suzuki, T. Suzuki, T. Miyamura, and Y. Matsuura. 2003. Proteasome activator PA28gamma-dependent nuclear retention and degradation of hepatitis C virus core protein. *J. Virol.* **77**:10237–10249.
  34. Moriya, K., H. Fujie, Y. Shintani, H. Yotsuyanagi, T. Tsutsumi, K. Ishibashi, Y. Matsuura, S. Kimura, T. Miyamura, and K. Koike. 1998. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat. Med.* **4**:1065–1067.
  35. Moriya, K., K. Nakagawa, T. Santa, Y. Shintani, H. Fujie, H. Miyoshi, T. Tsutsumi, T. Miyazawa, K. Ishibashi, T. Horie, K. Imai, T. Todoroki, S. Kimura, and K. Koike. 2001. Oxidative stress in the absence of inflammation in a mouse model for hepatitis C virus-associated hepatocarcinogenesis. *Cancer Res.* **61**:4365–4370.
  36. Moriya, K., H. Yotsuyanagi, Y. Shintani, H. Fujie, K. Ishibashi, Y. Matsuura, T. Miyamura, and K. Koike. 1997. Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J. Gen. Virol.* **78**:1527–1531.
  37. Neupert, W. 1997. Protein import into mitochondria. *Annu. Rev. Biochem.* **66**:863–917.
  38. Okamoto, K., K. Moriishi, T. Miyamura, and Y. Matsuura. 2004. Intramembrane proteolysis and endoplasmic reticulum retention of hepatitis C virus core protein. *J. Virol.* **78**:6370–6380.
  39. Okuda, M., K. Li, M. R. Beard, L. A. Showalter, F. Scholle, S. M. Lemon, and S. A. Weinman. 2002. Mitochondrial injury, oxidative stress, and antioxidant gene expression are induced by hepatitis C virus core protein. *Gastroenterology* **122**:366–375.
  40. Perlemuter, G., A. Sabile, P. Letteron, G. Vona, A. Topilco, Y. Chretien, K. Koike, D. Pessayre, J. Chapman, G. Barba, and C. Bréchet. 2002. Hepatitis C virus core protein inhibits microsomal triglyceride transfer protein activity and very low density lipoprotein secretion: a model of viral-related steatosis. *FASEB J.* **16**:185–194.
  41. Raiston, R., K. Thudium, K. Berger, C. Kuo, B. Gervase, J. Hall, M. Selby, G. Kuo, M. Houghton, and Q. L. Choo. 1993. Characterization of hepatitis C virus envelope glycoprotein complexes expressed by recombinant vaccinia viruses. *J. Virol.* **67**:6753–6761.
  42. Ray, R. B., L. M. Lagging, K. Meyer, and R. Ray. 1996. Hepatitis C virus core protein cooperates with ras and transforms primary rat embryo fibroblasts to tumorigenic phenotype. *J. Virol.* **70**:4438–4443.
  43. Roth, J., D. J. Taatjes, and M. J. Warhol. 1989. Prevention of non-specific interactions of gold-labeled reagents on tissue sections. *Histochemistry* **92**:47–56.
  44. Sabile, A., G. Perlemuter, F. Bono, K. Kohara, F. Demaugre, M. Kohara, Y. Matsuura, T. Miyamura, C. Brechot, and G. Barba. 1999. Hepatitis C virus core protein binds to apolipoprotein AII and its secretion is modulated by fibrates. *Hepatology* **30**:1064–1076.
  45. Santolini, E., G. Migliaccio, and N. La Monica. 1994. Biosynthesis and biochemical properties of the hepatitis C virus core protein. *J. Virol.* **68**:3631–3641.
  46. Selby, M. J., Q. L. Choo, K. Berger, G. Kuo, E. Glazer, M. Eckart, C. Lee, D. Chien, C. Kuo, and M. Houghton. 1993. Expression, identification and subcellular localization of the proteins encoded by the hepatitis C viral genome. *J. Gen. Virol.* **74**:1103–1113.
  47. Shimoike, T., S. Mimori, H. Tani, Y. Matsuura, and T. Miyamura. 1999. Interaction of hepatitis C virus core protein with viral sense RNA and suppression of its translation. *J. Virol.* **73**:9718–9725.
  48. Suzuki, R., Y. Matsuura, T. Suzuki, A. Ando, J. Chiba, S. Harada, I. Saito, and T. Miyamura. 1995. Nuclear localization of the truncated hepatitis C virus core protein with its hydrophobic C terminus deleted. *J. Gen. Virol.* **76**:53–61.
  49. Suzuki, R., K. Tamura, J. Li, K. Ishii, Y. Matsuura, T. Miyamura, and T. Suzuki. 2001. Ubiquitin-mediated degradation of hepatitis C virus core protein is regulated by processing at its carboxyl terminus. *Virology* **280**:301–309.
  50. Tanaka, Y., T. Shimoike, K. Ishii, R. Suzuki, T. Suzuki, H. Ushijima, Y. Matsuura, and T. Miyamura. 2000. Selective binding of hepatitis C virus core protein to synthetic oligonucleotides corresponding to the 5' untranslated region of the viral genome. *Virology* **270**:229–236.
  51. Tellinghuisen, T. L., and C. M. Rice. 2002. Interaction between hepatitis C virus proteins and host cell factors. *Curr. Opin. Microbiol.* **5**:419–427.
  52. Thomson, M., and T. J. Liang. 2000. Molecular biology of hepatitis C virus. p. 1–23. *In* T. J. Liang and J. H. Hoofnagle (ed.), *Hepatitis C*. Academic Press, San Diego, Calif.
  53. Weihofen, A., K. Binns, M. K. Lemberg, K. Ashman, and B. Martoglio. 2002. Identification of signal peptide peptidase, a presenilin-type aspartic protease. *Science* **296**:2215–2218.
  54. Weis, K., I. W. Mattaj, and A. I. Lamond. 1995. Identification of hSRP1 alpha as a functional receptor for nuclear localization sequences. *Science* **268**:1049–1053.
  55. Yanagi, M., R. H. Purcell, S. U. Emerson, and J. Bukh. 1997. Transcripts from a single full-length cDNA clone of hepatitis C virus are infectious when directly transfected into the liver of a chimpanzee. *Proc. Natl. Acad. Sci. USA* **94**:8738–8743.
  56. Yasui, K., T. Wakita, K. Tsukiyama-Kohara, S. I. Funahashi, M. Ichikawa, T. Kajita, D. Moradpour, J. R. Wands, and M. Kohara. 1998. The native form and maturation process of hepatitis C virus core protein. *J. Virol.* **72**:6048–6055.

## Identification of Basal Promoter and Enhancer Elements in an Untranslated Region of the TT Virus Genome

Tetsuro Suzuki,<sup>1\*</sup> Ryosuke Suzuki,<sup>1</sup> Jin Li,<sup>1</sup> Minako Hijikata,<sup>2</sup> Mami Matsuda,<sup>1</sup> Tiang-Cheng Li,<sup>1</sup> Yoshiharu Matsuura,<sup>3</sup> Shunji Mishiro,<sup>4</sup> and Tatsuo Miyamura<sup>1</sup>

Department of Virology II, National Institute of Infectious Diseases,<sup>1</sup> and Department of Respiratory Diseases, Research Institute, International Medical Center of Japan,<sup>2</sup> Shinjuku-ku, and Department of Medical Sciences, Toshiba General Hospital, Shinagawa-ku,<sup>4</sup> Tokyo, and Research Center for Emerging Infectious Diseases, Research Institute for Microbial Diseases, Osaka University, Suita-shi, Osaka,<sup>3</sup> Japan

Received 2 April 2004/Accepted 19 May 2004

**The regulation of TT virus (TTV) gene expression was characterized. Transient-transfection assays using reporter constructs revealed that a 113-nucleotide (nt) sequence within the untranslated region, proximal to the transcription initiation site and containing a TATA box motif, has a basal promoter activity. This sequence is well conserved among different TTV genotypes. Upstream stimulating factor bound to a consensus binding motif within this region and positively regulates TTV transcription. Furthermore, a 488-nt region upstream of the basal promoter exhibited enhancer activity, presumably in a cell type-specific manner. This study illustrates some of the mechanisms involved in the transcriptional regulation of TTV.**

TT virus (TTV), which was discovered in a patient with acute hepatitis, is an unenveloped, single-stranded, circular DNA virus, with a genome of approximately 3.8 kb (6). TTV is thought to be a new member of the *Circoviridae* family of viruses, and it was recently proposed that the virus be named Torque Teno virus (6). The TTV genome includes an untranslated region (UTR) of approximately 1.2 kb and a coding region of approximately 2.6 kb, including two major open reading frames which are sandwiched by the TATA box and polyadenylation signal motifs (11, 13, 15). Analyses of TTV transcripts have revealed three spliced mRNA species of 3.0, 1.2, and 1.0 kb with common 5' and 3' termini (9, 14). However, the molecular mechanisms controlling TTV transcription are still unknown. In this study, the basal promoter and enhancer of a TTV isolate, SANBAN of genogroup 3 (5, 18), were identified and functionally characterized.

First, we determined the transcription initiation sites of the TTV genome by 5' rapid amplification of cDNA end (5'-RACE) analysis (Marathon cDNA amplification kit; Clontech) using poly(A)-rich RNA from a human hepatocellular carcinoma cell line, HepG2, transfected with a cloned TTV genome. The 5'-RACE PCR products were cloned and sequenced. We observed two potential transcription initiation sites, which map at nucleotides (nt) 121 and 110 (numbered according to the sequence deposited in DDBJ/GenBank/EMBL databases under accession number AB025946). Although transcription may be initiated at both sites, the upper site was designated position +1 in this study.

The UTR of the TTV genome contains a TATA box element between positions -40 and -35, as well as a number of putative transcription factor-binding motifs (Fig. 1A). Despite

considerable genetic diversity throughout the whole genome, the UTR sequence was relatively conserved among the different TTV genotypes, presumably reflecting its functional constraints (15, 16). Thus, we analyzed transcriptional regulation of the UTR sequence.

To characterize TTV promoter activity, a firefly luciferase reporter plasmid, p(-890/+115), was constructed by subcloning the TTV sequence from positions -890 to +115, which was amplified by PCR using appropriate primers with restriction sites at the 5' ends, into the promoterless pGL3-Basic (Promega). Eleven different cell lines were transfected with p(-890/+115), along with a *Renilla* luciferase expression vector, pRL-TK, as an internal standard for determining transfection efficiency. Luciferase activities in cell lysates prepared after 16 h of transfection were determined (2). It is of interest that the 1.0-kb fragment demonstrated a pronounced promoter activity in all the hepatocellular carcinoma cell lines tested. Human (Huh7, HepG2, and FLC4 [1, 2]) and mouse (Hepa1-clc7) hepatocellular carcinoma cells were tested (Fig. 1B). This fragment demonstrated the greatest activity in Huh7 cells (~10-fold greater than in other cells). We observed substantial promoter activity in GL37 (African green monkey kidney) and CHO (Chinese hamster ovary) cells, whereas limited activity was observed in Caco2 (human colon carcinoma), MOLT4 (human T-cell leukemia), CV1 (African green monkey kidney fibroblast), 3T3 Swiss (mouse fibroblast), and CMT93 (mouse rectal carcinoma) cells. These results indicate that the UTR of the TTV genome functions as a promoter in a cell type-specific manner.

To assess basal, proximal promoter activity in the UTR, a series of 5' deletions fused to the luciferase gene were constructed and transfected into Huh7 and HepG2 cells (Fig. 2). A deletion extending to nt -601 [p(-601/+115)] enhanced promoter activity in both cell lines (by ~1.5-fold), while deletion of another 274 nt [p(-327/+115)] decreased promoter activity by more than 80%, suggesting that there is a negative regula-

\* Corresponding author. Mailing address: Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan. Phone: (81) 3-5285-1111. Fax: (81) 3-5285-1161. E-mail: tesuzuki@nih.go.jp.




FIG. 1. Functional analysis of the TTV promoter in the viral UTR. (A) Schematic representation of the 1.0-kb UTR sequence. The TATA box element and putative transcription factor-binding sites are shown. Transcription factor-binding sites were identified using the TRASFAC database and a search program (<http://motif.genome.ad.jp/>). The transcription initiation site (+1) is indicated and corresponds to nt 121 (AB025946). The numbers at the bottom of vertical dotted lines indicate the start points of the full-length promoter construct and deletion mutants of the promoter constructs used in Fig. 1B and 2. (B) Cell type specificity of the TTV promoter activities. Cells were transfected with p(-890/+115) (+) or promoter-less pGL3-Basic (-) together with pRL-TK (*Renilla* luciferase). Cell extracts were prepared 16 h after transfection, and luciferase activities in the extracts were determined using a dual-luciferase reporter assay system (Promega) with the Lumat LB9501 luminometer (Berthold). All values were normalized to *Renilla* luciferase activities and are shown as means ± standard deviations (error bars) of three independent samples.

tory element between nt -890 and -601. A deletion extending to nt -113 [p(-113/+115)] resulted in a slight to moderate reduction in activity, but promoter activity still remained greater than that observed with p(+15/+113), in which the

TATA box and the transcription start sites were deleted. These findings suggest that the 113 nt immediately upstream of the transcription initiation site contains a basal promoter region critical for TTV gene expression.




FIG. 2. Deletion analysis of the TTV promoter in Huh7 and HepG2 cells. The structures of the luciferase reporter constructs containing various lengths of the TTV UTR sequence are shown to the left. A series of DNA fragments with 5' deletions of the TTV promoter were amplified by PCR using the full-length TTV DNA of SANBAN isolate as a template with the same reverse primer and various forward primers. The fragments were cloned into pGL3-Basic at XhoI and HindIII sites. The indicated constructs were transfected into Huh7 cells or HepG2 cells. Relative luciferase activity in each transfectant was determined as described in the legend to Fig. 1B. Results are shown as means ± standard deviations (error bars) of three independent samples.







FIG. 3. Basal promoter activity of TTV regulated by USF. (A) Alignment of the putative basal promoter regions from TTV isolates SANBAN (DDBJ/GenBank/EMBL accession number AB025946), TA278 (AB017911), JA20 (AF122914), TUS01 (AB017613), IR1031 (AB038619), S039 (AB038620), TJN01 (AB028668), and TYM9 (AB050448). The transcription initiation site is numbered +1. The TATA box and positions of putative binding sites for USF and SP1 are indicated. Nucleotides that are identical to those in the SANBAN isolate (-) are indicated. (B) Effect of deleting DNA from the basal promoter region on the TTV basal promoter activity. A series of DNA fragments with 5' or internal deletions of the basal promoter region were amplified by PCR and cloned into pGL3-Basic at XhoI and HindIII sites. The indicated constructs were transfected into Huh7 cells (gray bars) or HepG2 cells (white bars). Relative luciferase activity in each transfectant was determined as described in the legend




FIG. 4. Enhancer activity of the 488-nt fragment ( $-601/-114$ ) of the TTV UTR. The 488-nt ( $-614/-114$ ) and 363-nt ( $-476/-114$ ) fragments were PCR amplified using primers with 5' overhangs containing BglIII (sense) and BamHI (antisense) sites. The fragments were then cloned into pGL3-Promoter at the BamHI site. The indicated constructs (left panel) were transfected into Huh7 cells or HepG2 cells. Relative luciferase activity in each transfectant was determined as described in the legend to Fig. 1B. Results are shown as means  $\pm$  standard deviations (error bars) of three independent samples. SV40, simian virus 40.

Computer-assisted analysis of this basal promoter region identified potential binding sites recognized by upstream stimulating factor (USF) and Sp1, which are conserved among TTV genotypes (Fig. 3A). To determine whether these sequences contribute to TTV promoter activity, 5' or internal deletion mutations were introduced into p( $-113/+115$ ), which was then examined for dual luciferase activity. Deletions, those found in p( $-92/+115$ ) and pBPd1, reduced promoter activity by 70 to 85%, suggesting that the USF-binding sequence is crucial for TTV promoter activity (Fig. 3B). Deletion of a Sp1-binding sequence (pBPd4) also conferred a decrease in promoter activity, although their effects were relatively moderate, suggesting that the Sp1-binding motif and/or its encompassing sequence may play a role in regulating TTV promoter activity by maintaining the structural integrity of the transcriptional machinery.

In genes where USF regulates transcription, cotransfection of USF expression vectors with reporter genes stimulates reporter activities. To further investigate the effect of USF on TTV promoter activity, we cotransfected USF1 or USF2 expression vectors (pCMV-USF1 and pCMV-USF2) (7) with p( $-113/+115$ ) into HepG2 cells. The cotransfection significantly increased promoter activity (by threefold), suggesting that USF proteins regulate TTV transcription (Fig. 3C).

USF is a family of basic-helix-loop-helix-leucine zipper transcription factors, initially identified by their ability to bind to

the 5'-CACGTG-3' sequence within the adenovirus major late promoter (3, 4, 10). USF1 and USF2 have been subsequently shown to bind to the promoters of various cellular and viral genes. To determine whether the TTV basal promoter was capable of USF binding, gel mobility shift assays were performed on an end-labeled oligonucleotide (nt  $-113$  to  $-84$ ) containing the putative USF-binding motif (Fig. 3D). DNA-protein-binding complexes were observed in nuclear extracts from cells transfected with pCMV-USF1 (Fig. 3D, lane 5), pCMV-USF2 (lane 8), and the empty vector (lane 2). An excess of unlabeled homologous probe competed with the protein binding (lanes 3, 6, and 10), whereas a mutated USF sequence failed to compete (lanes 4, 7, and 10). The addition of anti-USF antibodies to the binding reaction mixture supershifted the DNA-protein complexes (lanes 12 and 14). The combined data demonstrate that USF binds to its binding motif within the TTV basal promoter to up-regulate viral transcription.

On the basis of the results of the luciferase assays using 5' deletions of the TTV UTR (Fig. 2), the positive regulatory element appears to be located immediately upstream of the basal promoter. To ascertain whether the 488-bp fragment between nt  $-601$  and  $-113$  functions as the enhancer region, this fragment or a 5' deletion of this fragment was placed downstream of the polyadenylation signal in pGL3-Promoter (Promega), driven by the simian virus 40 promoter, in either

to Fig. 1B. Results are shown as a percentage of the activity in cells transfected with p( $-113/+115$ ); values are shown as means  $\pm$  standard deviations (error bars) ( $n = 3$  per group). (C) Effects of USF overexpression on basal promoter activity. HepG2 cells were cotransfected with each reporter construct with pCMV-USF1 (USF1), pCMV-USF2 (USF2), or empty pC<sub>0</sub> vector (C<sub>0</sub>). Luciferase activity was determined 48 h after transfection. For each reporter construct, relative luciferase activity is presented as a percentage of the activity in pC<sub>0</sub>-transfected cells. (D) Binding of USF proteins to the region from nt  $-113$  to  $-84$  in the TTV basal promoter. The electrophoretic mobility shift assays were performed as described previously (19). A double-stranded oligonucleotide corresponding to the TTV sequence from nt  $-113$  to  $-84$  was used as a probe. Nuclear extracts from the cells transiently transfected with pCMV-USF1 (USF1; lanes 5, 6, 7, 11, and 12), pCMV-USF2 (USF2; lanes 8, 9, 10, 13, and 14), or pC<sub>0</sub> (C<sub>0</sub>; lanes 2 to 4) or no extract (lane 1) were mixed with <sup>32</sup>P-labeled probe for the binding reaction mixtures. Competitors, unlabeled probe (USF), and a mutant with the USF-binding motif (USFmut) were added at a 25-fold molar excess. The sense sequence (5'-TCACTAAC CAATGACCCGTCTCGCCCAAC [the mutated nucleotides are underlined]) and complementary sequence of the mutant with the USF-binding motif were added. A supershift experiment was also performed by incubating antibody against USF1 (lane 12) or USF2 (lane 14) (+) with the nuclear extracts before the probe was added. The positions of specific binding complexes (arrows) and supershifted complexes (arrowheads) are indicated.

the sense or antisense orientation. Luciferase activity of constructs containing the 488-bp fragment [pEN(-601/-114) and pEN(-114/-601)] led to 50- and 10-fold stimulation in Huh7 and HepG2 cells, respectively. The 5' deletion extending to nt -476 [pEN(-476/-114) and pEN(-114/-476)] reduced enhancer activity (Fig. 4). No enhancement was observed by transfection of MOLT4 cells with pEN(-601/-114) and pEN(-114/-601) (data not shown). These results demonstrate that the 488-bp region upstream of the basal promoter contains an enhancer element, suggesting cell-specific transcription of the TTV genome. It is noteworthy that the enhancer element is conserved among TTV genotypes. For example, 72% homology has been observed between clones SANBAN and TA278, and the database search has revealed at least 20 potential transcription factor-binding sites within this element, including CREB and CRB, which are activated upon cyclic AMP signaling-dependent phosphorylation (12, 17).

While the manuscript was being prepared, Kamada et al. reported the promoter and enhancer activities in the UTR of TTV, clone VT416 whose genome is 98% similar to that of TA278, and its cell tropism (8). However, they did not identify transcription factors that bind to the region and regulate TTV transcription. In summary, the findings reported by Kamada et al. and the findings of our present study emphasize the important role of the UTR as a basal promoter and enhancer within the UTR. Other areas of interest for further study include the identification of additional factors involved in tissue-specific TTV transcription and determining the significance of polymorphism of the regulatory elements.

We are grateful to M. Sawadogo (University of Texas M. D. Anderson Cancer Center, Houston, Tex.) for providing pCMV-USF1, pCMV-USF2, and pC<sub>1</sub>. We thank M. Yahata and S. Yoshizaki for technical assistance and T. Mizoguchi for preparation of the manuscript.

This work was supported in part by grants-in-aid from the Ministry of Health, Labor and Welfare of Japan.

#### REFERENCES

- Aizaki, H., S. Nagamori, M. Matsuda, H. Kawakami, O. Hashimoto, H. Ishiko, M. Kawada, T. Matsuura, S. Hasumura, Y. Matsuura, T. Suzuki, and T. Miyamura. 2003. Production and release of infectious hepatitis C virus from human liver cell cultures in the three-dimensional radial-flow bioreactor. *Virology* 314:16-25.
- Aoki, Y., H. Aizaki, T. Shimoike, H. Tani, K. Ishii, I. Saito, Y. Matsuura, and T. Miyamura. 1998. A human liver cell line exhibits efficient translation of HCV RNAs produced by a recombinant adenovirus expressing T7 RNA polymerase. *Virology* 250:140-150.
- Atchley, W. R., and W. M. Fitch. 1997. A natural classification of the basic helix-loop-helix class of transcription factors. *Proc. Natl. Acad. Sci. USA* 94:5172-5176.
- Gregor, P. D., M. Sawadogo, and R. G. Roeder. 1990. The adenovirus major late transcription factor USF is a member of the helix-loop-helix group of regulatory proteins and binds to DNA as a dimer. *Genes Dev.* 4:1730-1740.
- Hijikata, M., K. Takahashi, and S. Mishiuro. 1999. Complete circular DNA genome of a TT virus variant (isolate name SANBAN) and 44 partial ORF2 sequences implicating a great degree of diversity beyond genotypes. *Virology* 260:17-22.
- Hino, S. 2002. TTV, a new human virus with single stranded circular DNA genome. *Rev. Med. Virol.* 12:151-158.
- Ismail, P. M., T. Lu, and M. Sawadogo. 1999. Loss of USF transcriptional activity in breast cancer cell lines. *Oncogene* 18:5582-5591.
- Kamada, K., T. Kamahora, P. Kobat, and S. Hino. 2003. Transcriptional regulation of TT virus: promoter and enhancer regions in the 1.2-kb non-coding region. *Virology* 321:341-348.
- Kamahora, T., S. Hino, and H. Miyata. 2000. Three spliced mRNAs of TT virus transcribed from a plasmid containing the entire genome in COS1 cells. *J. Virol.* 74:9980-9986.
- Lin, Q., X. Luo, and M. Sawadogo. 1994. Archaic structure of the gene encoding transcription factor USF. *J. Biol. Chem.* 269:23894-23903.
- Miyata, H., H. Tsunoda, A. Kazi, A. Yamada, M. A. Khan, J. Murakami, T. Kamahora, K. Shiraki, and S. Hino. 1999. Identification of a novel GC-rich 113-nucleotide region to complete the circular, single-stranded DNA genome of TT virus, the first human circovirus. *J. Virol.* 73:3582-3586.
- Montminy, M. 1997. Transcriptional regulation by cyclic AMP. *Annu. Rev. Biochem.* 66:807-822.
- Mushahwar, I. K., J. C. Erker, A. S. Muerhoff, T. P. Leary, J. N. Simons, L. G. Birkenmeyer, M. L. Chalmers, T. J. Pilot-Matias, and S. M. Dexai. 1999. Molecular and biophysical characterization of TT virus: evidence for a new virus family infecting humans. *Proc. Natl. Acad. Sci. USA* 96:3177-3182.
- Okamoto, H., T. Nishizawa, A. Tawara, M. Takahashi, J. Kishimoto, T. Sai, and Y. Sugai. 2000. TT virus mRNAs detected in the bone marrow cells from an infected individual. *Biochem. Biophys. Res. Commun.* 279:700-707.
- Okamoto, H., T. Nishizawa, and M. Ukita. 1999. A novel unenveloped DNA virus (TT virus) associated with acute and chronic non-A to G hepatitis. *Intervirology* 42:196-204.
- Prescott, L. E., D. M. MacDonald, F. Davidson, J. Mokili, D. I. Pritchard, D. E. Arnot, E. M. Riley, B. M. Greenwood, S. Hamid, A. A. Saeed, M. O. McClure, D. B. Smith, and P. Simmonds. 1999. Sequence diversity of TT virus in geographically dispersed human populations. *J. Gen. Virol.* 80:1751-1758.
- Sassone-Corsi, P. 1998. Coupling gene expression to cAMP signalling: role of CREB and CREM. *Int. J. Biochem. Cell. Biol.* 30:27-38.
- Takahashi, K., M. Hijikata, E. I. Samokhvalov, and S. Mishiuro. 2000. Full or near full-length nucleotide sequences of TT virus variants (types SANBAN and YONBAN) and the TT virus-like mini virus. *Intervirology* 43:119-123.
- Tsutsumi, T., T. Suzuki, T. Shimoike, R. Suzuki, K. Moriya, Y. Shintani, H. Fujie, Y. Matsuura, K. Koike, and T. Miyamura. 2002. Interaction of hepatitis C virus core protein with retinoid X receptor alpha modulates its transcriptional activity. *Hepatology* 35:937-946.