



ACADEMIC  
PRESS

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

NeuroImage

NeuroImage 19 (2003) 1674–1685

[www.elsevier.com/locate/ynimg](http://www.elsevier.com/locate/ynimg)

## Context-dependent cortical activation in response to financial reward and penalty: an event-related fMRI study

Yuko Akitsuki,<sup>a,b,\*</sup> Motoaki Sugiura,<sup>b</sup> Jobu Watanabe,<sup>c,d</sup> Keiichi Yamashita,<sup>e</sup> Yuko Sassa,<sup>d,f</sup>  
Shuichi Awata,<sup>a</sup> Hiroo Matsuoka,<sup>a</sup> Yasuhiro Maeda,<sup>g</sup> Yoshihiko Matsue,<sup>g</sup>  
Hiroshi Fukuda,<sup>c</sup> and Ryuta Kawashima<sup>b</sup>

<sup>a</sup> Department of Psychiatry, Tohoku University Graduate School of Medicine, Sendai, Japan

<sup>b</sup> NICHe, Tohoku University, Sendai, Japan

<sup>c</sup> IDAC, Tohoku University, Sendai, Japan

<sup>d</sup> LBC Research Center, Tohoku University 21st Century Center of Excellence Program in Humanities, Sendai, Japan

<sup>e</sup> Department of Neurosurgery Tottori University School of Medicine, Yonago, Japan

<sup>f</sup> Graduate School of International Cultural Studies, Tohoku University, Tohoku, Japan

<sup>g</sup> Kansei Fukushi Research Center, Tohoku Fukushi University, Sendai, Japan

Received 20 November 2002; revised 8 March 2003; accepted 24 April 2003

### Abstract

An event-related fMRI technique was used to assess neural responses to financial reward and penalty during a simple gambling task. We attempted to determine whether brain activities are dependent on the unique context of an event sequence. Thirty-six healthy volunteers participated in the study. The task was to guess the color of the suit of a card on each trial and to respond by pressing a button. Every correct response (“win”) and incorrect response (“loss”) was associated with financial reward and penalty, respectively. The magnitude of reward or penalty in each trial did not change; however, the subjects’ self-reported emotional arousal was significantly higher for the events of “the fourth win of four wins in a row” and “the fourth loss of four losses in a row.” We also found that the bilateral anterior cingulate and medial prefrontal cortices were specifically activated when the subjects experienced “the fourth win of four wins in a row” and “the fourth loss of four losses in a row.” When the subjects experienced “a win following four losses in a row” or “a loss following four wins in a row,” the right dorsolateral prefrontal cortex was specifically activated. Our data indicate that there exist brain activities associated with the event-sequence context in which abstract reward or penalty is received. These context-dependent activities appear to be crucial for adapting oneself to new circumstances and may account for clinical symptoms of various mental illnesses in which dysfunction of these regions has been reported.

© 2003 Elsevier Inc. All rights reserved.

### Introduction

Receiving rewards or penalties is not a simply passive behavior. Animals should predict potential rewards/penalties, distinguish their nature, and respond appropriately by changing their attentional or arousal level for successful adaptive behavior. Single-unit studies of nonhuman primates revealed the brain structures that are related to re-

wards and penalties. In the monkey, midbrain dopamine neurons were reported to be activated by the occurrence of rewards after their presentation and visual or auditory stimuli that predict rewards (Schultz, 1986; Romo and Schultz, 1990; Schultz and Romo, 1990; Ljungberg et al., 1992). Dopamine neurons also encode an error in the temporal prediction of rewards; the dopamine activity is enhanced by surprising rewards and it is depressed by the omission of predicted rewards (Ljungberg et al., 1992; Mirenowicz and Schultz, 1994). Other neurons in the ventral striatum (Apicella et al., 1991; Shidara et al., 1998) and orbitofrontal cortex (Niki et al., 1972) also respond to the delivery of

\* Corresponding author. NICHe, Tohoku University, Aoba 10, Aramaki, Aobaku, Sendai, Miyagi, 980-8579, Japan.

E-mail address: [akitsuki@idac.tohoku.ac.jp](mailto:akitsuki@idac.tohoku.ac.jp) (Y. Akitsuki).

rewards. Furthermore, neurons within the orbitofrontal cortex not only discriminate rewards and penalties (Thorpe et al., 1983), but also respond to reward expectancy (Tremblay and Schultz, 1999, 2000). Reward expectation neurons are also found in the striatum (Shidara et al., 1998; Schlutz et al., 1992). Neurons in the lateral prefrontal cortex were reported to be sensitive to the presence or absence of expected rewards (Watanabe, 1990, 1992) and play an important role in coding the discrepancy between the expectancy of a specific reward and the response outcome (Watanabe, 1996).

On the other hand, imaging studies revealed that human brain activities are associated with the cognition of abstract rewards and penalties. In particular, the “gambling task” has been recently used as a laboratory model for investigating brain activities in response to abstract types of reward and punishment (e.g., monetary reward or penalty) in normal subjects. A previous positron emission tomography (PET) study revealed that the human orbitofrontal cortex can be activated with monetary reward (Thut et al., 1997). A functional magnetic resonance imaging (fMRI) study supports this finding, demonstrating that an increase in the activity of the medial orbitofrontal cortex is related to the subjects’ receipt of monetary reward and deactivation following punishments (O’Doherty et al., 2001). The authors recorded the converse pattern of activation in the lateral orbitofrontal cortex (activation following punishments, deactivation following rewards). Moreover, in these areas, the authors found a correlation between the magnitude of brain activation and that of rewards and punishments received.

However, neural responses related to human reward systems under different psychological contexts are studied less extensively. On this issue, some groups focused on the context of anticipation of abstract rewards and punishments. They demonstrated that whereas anticipation of increasing rewards elicited nucleus accumbens activation (Breiter et al., 2001; Knutson et al., 2001b), the anticipation of increasing punishment did not activate the nucleus accumbens (Knutson et al., 2001a). Nucleus accumbens activation correlated with individual differences in self-reported happiness elicited by the reward cues (Knutson et al., 2001a). On the other hand, Elliott and colleagues (2000) focused on the psychological context of a winning or losing streak. They demonstrated that brain activities in the bilateral globus pallidus, thalamus, and subgenual cingulate are associated with rewards in the context of winning streak. They also showed that responses to both rewards occurring as part of a winning streak and penalties as part of a losing streak are observed in the caudate, insula, and ventral prefrontal cortex. This is consistent with the findings that the medial orbitofrontal and ventromedial prefrontal cortices are involved in the representation of rewards and risk (Bechara et al., 1994, 1996, 1997; Rolls et al., 1994).

Taking these results into account, a prior emotional experience with streaks of rewards/penalties appears to shape the subjective “feeling” of an incoming expected reward/

penalty or an unexpected one for that matter. Moreover, there seems to be dissociable neural responses that are dependent on the psychological context in which abstract rewards and penalties are received in the human brain. Therefore, based on the study by Elliott et al., we would like to determine the relationship between subjective feelings and the context of winning/losing streaks and how these context are expressed in the human brain. In this experiment, an event-related fMRI technique was used to assess neural responses to financial reward and penalty during a gambling task whose paradigm was based on the study by Elliott and Critchley (Critchley et al., 2000). Our aim is to determine whether subjective feelings and brain activities are dependent on the unique context of an event sequence, that is, four wins in a row, four losses in a row, a win following four losses in a row, or a loss following four wins in a row. On the basis of previous studies, we hypothesize that:

1. The medial prefrontal cortex is activated by risk-taking and emotionally salient events, such as the context of a winning or losing streak.
2. The lateral prefrontal cortex is activated by the discrepancy between the expectancy and the outcome of rewards/penalties, such as the context of switching from a winning or losing streak to the opposite event.

## Methods

### Subjects

Thirty-six healthy volunteers (19 men and 17 women), whose mean age was 20.8 years (SD = 2.98), participated in this study. All were native Japanese speakers and right-handed, as assessed by the Edinburgh Handedness Inventory. Subjects with a history of psychiatric, neurological, or other serious physical illnesses; drug or alcohol abuse; or second-degree relatives with a history of major psychiatric disorders were excluded.


Approval to carry out this experiment was obtained from Ethics Committee of Tohoku Fukushi University. In accordance with the Helsinki Declaration of Human Rights (1975), written informed consent was obtained from all participants after complete and detailed description of the study to the subjects.

### Experimental paradigm

Brain activity associated with the psychological context of reward/penalty in a simple gambling task was examined, using an event-related fMRI technique. The experimental paradigm we used here was based on the previous studies done by Elliott and Critchley (Elliott et al., 2000; Critchley et al., 2000).

Subjects were presented with cards as visual stimuli on a

### Gambling task



### Cumulative reward score




Fig. 1. Gambling task (top). During the first part, subjects were instructed to guess the color of the suit and respond by pressing a button. (red, button 1; black, button 2). During the second part, the correct color and a "win!" or "loss!" message were presented. HRF, hemodynamic response function. Cumulative reward score (bottom). Because the outcome feedback ("win" or "loss") sequence was predetermined regardless of the subjects' actual choices, the cumulative reward score was also predetermined as shown here.

small screen on a head coil. The task was to guess the color of the suit (red or black) of a card on each trial and respond by pressing a button. (Fig. 1) Each trial consists of two parts and each part lasted 2.4 s, so that the duration of the entire trial was 4.8 s. In the first part, a card was presented for 2.4 s, and during this period, subjects were instructed to press button 1 if they thought the suit color would be red or

press button 2, if black. In the first part, the cumulative "reward" score was also presented. Subjects began this task with a 0 yen "stake." For the subsequent 2.4 s (the second part), the correct color and a "win!" or "loss!" message were presented. Every correct response (win) in the second part was associated with an increment of 1000 yen in the cumulative reward score for the subsequent first part of the next

trial, whereas every incorrect response (loss) was associated with a decrement of 1000 yen in the score. In fact, this outcome feedback (win or loss) sequence was predetermined according to a pseudorandom sequence, regardless of the subjects' actual choices. Therefore, the cumulative reward score was also predetermined (Fig. 1).

To examine the neural response related to the context in which rewards and penalties are experienced in this task, artificial repetition of win or loss was designed, that is, "four wins in a row" and "four losses in a row." There were 215 trials in each run. The total number of win versus loss, "winning four trials in a row" versus "losing four trials in a row" and "a win subsequent to losing four trials in a row" versus "a loss subsequent to winning four trials in a row" was counterbalanced across the task.

Immediately after the fMRI measurements, the subjects were asked to indicate how happy or unhappy they found themselves at winning or losing events during the task performance using an 11-point rating scale (i.e., "5" indicates "happiest," "0" indicates "neutral," and "−5" indexes "unhappiest").

#### fMRI measurement

T1-weighted structural images were acquired for each volunteer, using a 1.5-T Siemens Vision plus scanner (Siemens Magnetron Vision, Erlanger, Germany; repetition time 9.7 ms, echo time 4 ms, flip angle 12°, FOV = 250 mm, pixel size 1.25 × 0.98 mm, matrix 175 × 256). To measure neural responses, gradient echo, echoplanar T2\*-weighted images with blood oxygenation level-dependent (BOLD) contrast were acquired (repetition time 4000 ms, echo time 60 ms, flip angle 90°, FOV = 256 × 256 × 256 mm<sup>3</sup>, voxel size 4 × 4 × 3 mm<sup>3</sup>, matrix 64 × 64, slice thickness 3mm, interslice gap 0.99 mm). Thirty-four axial slices were oriented over the whole brain.

To allow for T1 equilibration effects, two preliminary scans were acquired and subsequently discarded. Thereafter, scans were acquired continuously every 4.0 s, so each block of five behavioral trials corresponded to six scans. This temporal asynchrony is significant for avoiding a systematic bias in sampling over peristimulus time.

#### Data analysis

Data were analyzed on a work station (Sun Microsystems), using MATLAB (Mathworks Inc., Natick, MA) and statistical parametric mapping (SPM99; Wellcome Department of Cognitive Neurology, London, UK). The initial two scans were discarded for the nonequilibrium state of magnetization.

Before statistical analysis, a series of spatial transformation stages were required. First, slice time adjustment was performed to correct for differences in acquisition time among 34 slices of each scan. Second, to correct for artifacts caused by small head movements, images from each subject




Fig. 3. Brain areas that were significantly activated when subjects experience four wins in a row and four losses in a row. Depicted here are areas of the brain that yielded significantly higher activity in W4 than in W1 (a), and in L4 than in L1 (b) and rendered onto the mean anatomically standardized T1 image of all subjects. Both activations were significant at  $P < 0.05$  (corrected).

Fig. 4. Brain areas that were significantly activated when subjects experience a win following four losses in a row and a loss following four wins in a row. Depicted here are areas of the brain that yielded significantly higher activity in SW than in W1 (a) and in SL than in L1 (b). Both activations were significant at  $P < 0.05$  (corrected). Frontal views are shown in the top panel and lateral views from the right are shown in the lower panel. SW, sudden win; SL, sudden loss.

were realigned to the first image and resliced using a sinc interpolation. Third, a mean image created from the realigned volumes was coregistered with the structural T1 volume and the structural volumes were spatially normalized to a standard template in the space of Talairach and Tournoux (1988) using nonlinear basis functions. Fourth, the derived spatial transformation was applied to the realigned T2\* volumes, which were then spatially smoothed with a three-dimensional isotropic Gaussian kernel (12 mm full-width at half-maximum). This improves the signal-to-noise ratio and accommodates for residual variations in functional neuroanatomy that usually persist between subjects after spatial normalization.

A key aspect of rewarding situation is how reward (or penalty)-related responses are modulated by the psychological context. We, therefore, hypothesized that neural responses depend on the context of winning or losing streak and presumed the following 10 conditions for each subject:

1. W1: the win other than W2, W3, W4, and SW
2. W2: the second win of two wins in a row
3. W3: the third win of four wins in a row
4. W4: the fourth win of four wins in a row
5. SW: a win following four losses in a row; SW stands for sudden win
6. L1: the loss other than L2, L3, L4, and SL
7. L2: the second loss of two losses in a row
8. L3: the third loss of three losses in a row
9. L4: the fourth loss of four losses in a row
10. SL: a loss following four wins in a row; SL stands for sudden loss

For both W4 and SW, which were designed as a special context, a subtraction method was used to contrast the brain activity associated with W1 that seems less meaningful or less exciting in terms of psychological context. On the other hand, for both L4 and SL, a subtraction method was used to contrast the brain activity associated with L1. We also compared the brain activities between W1 and L1, W4 and L4, and all wins (including W1, W2, W3, W4, and SW) and all losses (including L1, L2, L3, L4, and SL).

In addition, we performed the correlation analysis between the subjects' arousal and the fMRI signals. In this analysis, neural activities were examined for parametric modulation by the scores of arousal rating scale.

fMRI data were analyzed using the general linear model. For the statistical analysis on the intrasubject basis, the time series of images were correlated with the SPM99 built-in "Canonical HRF" that approximates activation patterns. The onset of the expected response was positioned at the beginning of the second part of each trial (i.e., the point at which the correct color of suit and a "win!" or "loss!" message were presented (see Fig. 1). Condition effects at each voxel were estimated according to this general linear model and regionally specific effects were compared using linear contrasts. Global changes were adjusted by propor-




Fig. 2. Mean subjects' ratings. Mean subjects' ratings of emotional experience in each winning and losing event. Each subject rated each event individually (W1, W2, W3, W4, SW, L1, L2, L3, L4, and SL). SW stands for sudden win. SL stands for sudden loss. Two way ANOVA and post hoc Tukey's tests were performed. Subjects' ratings showed significant differences in W1 versus W4 and L1 versus L4 ( $P < 0.000001$ ), but not in W1 versus SW and L1 versus SL.

tional scaling and low-frequency confounding effects were removed using an appropriate high-pass filter. Voxel values for each contrast yielded a statistical parametric map of the  $t$  statistic (SPM  $t$ ), which was subsequently transformed to the unit normal distribution (SPM  $Z$ ).

Intersubject maps were then produced by performing one-sample  $t$  tests to identify voxels that survive a voxel-wise multiple comparison ( $P < 0.05$ , corrected). At this step, we used masking procedures: The resulting SPM $\{T\}$  of W4 - W1, L4 - L1, SW - W1, and SL - L1 were masked with the statistically thresholded ( $P < 0.05$  corrected) mask of W4, L4, SW, and SL, respectively.

Finally, the resulting activation maps were created and displayed by projection onto the mean anatomically standardized T1 image of all subjects to identify cortical structures.

## Results

### Subjects' ratings

Two-way ANOVA (analysis of variance) and post hoc Tukey's tests were performed. Examination of the subjects' self-reported ratings indicated that W4 was a significantly more pleasant/happier event than W1 (mean  $\pm$  SD, W1,  $1.58 \pm 0.87$ ; W4,  $4.5 \pm 0.78$ ,  $P < 0.000001$ ), and L4 was a significantly more unpleasant/unhappier event than L1 (mean  $\pm$  SD, L1,  $-1.28 \pm 1.03$ ; L4,  $-4.17 \pm 1.36$ ,  $P < 0.000001$ ) (Fig. 2).

However, there was no significant difference between W1 and SW (mean  $\pm$  SD, L1,  $1.58 \pm 0.87$ ; SW,  $2.11 \pm 1.58$ ,  $P > 0.05$ ) or between L1 and SL (mean  $\pm$  SD, L1,  $-1.28 \pm 1.03$ ; SL,  $-1.58 \pm 1.40$ ,  $P > 0.05$ ).

Table 1  
Brain activities for four wins in a row and four losses in a row

	L/R	Region	Number of voxels ( <i>k</i> )	Talairach coordinates (mm)			<i>T</i> score
				<i>x</i>	<i>y</i>	<i>z</i>	
<b>W4 – W1</b>							
	R	Medial aspect of superior frontal gyrus	81 <sup>a</sup>	8	22	58	8.11
	R	Medial aspect of superior frontal gyrus		12	58	22	7.16
	L	Medial aspect of superior frontal gyrus	908 <sup>b</sup>	–2	58	10	8.21
	L	Medial aspect of superior frontal gyrus		–2	40	46	6.85
<b>L4 – L1</b>							
	R	Medial aspect of superior frontal gyrus	1229	8	56	22	8.78
	R	Medial aspect of superior frontal gyrus	32	10	18	60	7.7
	L	Medial aspect of superior frontal gyrus	111 <sup>c</sup>	–8	22	58	6.55
	L	Medial aspect of superior frontal gyrus		–4	50	20	8.35
	R	Middle temporal gyrus	22	60	–12	–12	5.29

Note. Maximally activated voxels in brain areas where significantly higher activities were found in W4 than in W1 and in L4 than in L1 ( $P < 0.05$ , corrected). Stereotaxic coordinates are derived from the human atlas of Talairach and Tournoux (1998) and refer to medial–lateral position (*x*) relative to the midline (positive = right), anterior–posterior position (*y*) relative to the anterior commissure (positive = anterior), and superior–inferior position (*z*) relative to the commissural line (positive = superior). L, left; R, right.

<sup>a</sup> This amount is the total number of activated voxels for the right medial aspect of the superior frontal gyrus in W4 – W1.

<sup>b</sup> This amount is the total number of activated voxels for the left medial aspect of the superior frontal gyrus in W4 – W1.

<sup>c</sup> This amount is the total number of activated voxels for the left medial aspect of the superior frontal gyrus in L4 – L1.

### fMRI data

Figure 3a and Table 1 show the significantly activated regions in the subtraction analysis of W4 – W1. In the event of the fourth win of four wins in a row (W4), significant activation was observed in the medial aspect of the superior frontal gyrus and the anterior cingulate gyrus, in comparison with the W1 event. There were two activation peaks in the cluster of the right superior frontal gyrus and in that of the left superior frontal gyrus.

Figure 3b and Table 1 illustrate the significantly activated regions in the subtraction analysis of L4 – L1. In the event of the fourth loss of four losses in a row (W4), significant activation was found in the medial aspect of the frontal gyrus, the anterior cingulate gyrus, and the right middle temporal gyrus. There were two activation peaks in the cluster of the left medial aspect of the superior frontal gyrus. The activation pattern of W4 – W1 highly overlapped with that of L4 – L1 (Figs. 3a and b).

Table 2 shows the significantly activated regions in the subtraction analysis of W1 – L1. In the event of W1, significant activation was observed in the bilateral ventral striatum, bilateral lingual gyrus, bilateral inferior temporal gyrus, and bilateral posterior lobe of the cerebellum, compared with the L1 event. However, no significant activation was found in L1 – W1, W4 – L4, and L4 – W4.

Table 3 shows the comparison between neural activation for all wins and that for all losses. In the events of win, the bilateral ventral striatum, bilateral posterior lobe of the cerebellum, and left lateral occipital cortex were significantly activated in comparison with the events of loss. No significant activation was found in the subtraction analysis of all wins – all losses.

Table 4 shows the brain regions that correlate with the subjects' arousal. The medial aspect of the superior frontal gyrus and the anterior cingulate gyrus significantly correlate with the subjects' arousal in the events of both win and loss. This activation pattern appears to be very similar to those of W4 – W1 and that of L4 – L1 (Figs. 3a, b).

Figure 4a and Table 5 show the significantly activated regions in the subtraction analysis of SW – L1. In the event of a win after four losses in a row (SW), significant activation was observed in the right middle frontal gyrus, in comparison with the W1 event. There were two activation peaks in the cluster of the right middle frontal gyrus.

Figure 4b and Table 5 illustrate the significantly activated regions in the subtraction analysis of SL – L1. Compared with L1, in the event of a loss after four wins in a row (SL), significant activation was found in the right middle frontal gyrus. There were two activation peaks in the cluster of the right middle frontal gyrus. The activation pattern of SW versus W1 appears to be very similar to that of SL versus L1. (Figs. 4a and b)

### Discussion

In this study, using a simple gambling task, we demonstrated brain activities associated with the event-sequence context in which abstract reward or penalty is received.

The most significant findings in our study are as follows:

1. The magnitude of each reward or penalty was the same across the whole events; however, psychological data demonstrated that subjective emotional experiences are different because of the context in which

Table 2  
Brain activities for W1 versus L1 and W4 versus L4

	L/R	Region	Number of voxels ( <i>k</i> )	Talairach coordinates (mm)			<i>T</i> score
				<i>x</i>	<i>y</i>	<i>z</i>	
W1 – L1							
	R	Ventral striatum	400	24	0	–8	8.67
	L	Ventral striatum	88	–26	–2	–12	8.61
	R	Lingual gyrus	10,028*	16	–68	–12	9.47
	L	Lingual gyrus		–16	–86	–2	10.63
	R	Inferior temporal gyrus		40	–78	–12	8.99
	L	Inferior temporal gyrus		–42	–62	–12	9.88
	R	Posterior lobe of cerebellum		–10	–72	–20	9.45
	L	Posterior lobe of cerebellum		–34	–64	–32	8.52
L1 – W1							
		NS					
W4 – L4							
		NS					
L4 – W4							
		NS					

Note. Maximally activated voxels in brain areas where significantly higher activities were found in W1 than in L1 ( $P < 0.05$ , corrected). No significant activation was found in L1 – W1, W4 – L4, and L4 – W4. NS, not significant.

\* This amount is the total number of activated voxels for the bilateral lingual gyrus, inferior temporal gyrus, and posterior lobe of the cerebellum in W4 – L1.

reward or penalty is received, that is, the sequence of each reward/penalty-related event (i.e., four wins in a row, four losses in a row, a win following four losses in a row, and a loss following four wins in a row).

2. Depending on the nature of the unique context (i.e., repetition of win or loss vs. switching from a winning or losing streak to the opposite event), specific brain regions were activated, unrelated to the nature of event (i.e., reward-related vs. penalty-related).

a. The medial aspect of the prefrontal cortex and the anterior cingulate cortex showed significant activations both in a series of reward-related events and in a series of penalty-related events. In these brain regions, there were significant correlations between fMRI signals and the subjects' arousal.

b. The right dorsolateral prefrontal cortex showed significant activation in switching events, such as a reward-related event following a series of penalty-related event and a penalty-related event following a series of reward-related event.

We highlight the finding that the overlapping activation of brain regions was induced in a specific context, whether the event was reward-related or penalty-related. This means that information regarding abstract rewards and penalties is represented as a certain context in the human brain independent of its nature (i.e., reward-related or penalty-related). Then, what is the meaning of this context-dependent brain activity? Why is a specific brain region activated in the unique context, even if the magnitude of each reward or penalty is the same?

Table 3  
Brain activities for all wins versus all losses

	L/R	Region	Number of voxels ( <i>k</i> )	Talairach coordinates (mm)			<i>T</i> score
				<i>x</i>	<i>y</i>	<i>z</i>	
all W — all L							
	R	Ventral striatum	26	26	2	–10	6.88
	L	Ventral striatum	64	–24	4	–12	7.51
	R	Posterior lobe of cerebellum	2426*	38	–76	–28	8.69
	L	Posterior lobe of cerebellum		–28	–78	–20	8.15
	L	Lateral occipital cortex		–22	–92	–6	7.86
all L — all W							
		NS					

Note. Maximally activated voxels in brain areas where significantly higher activities were found in all W than in all L ( $P < 0.05$ , corrected). No significant activation was found in all L — all W. NS, not significant. all W, all wins including W1, W2, W3, W4, and SW. all L, all losses including L1, L2, L3, L4, and SL.

\* This amount is the total number of activated voxels for the bilateral posterior lobe of the cerebellum and the lateral occipital cortex in all W — all L.

Table 4  
Brain activities that correlate with the subjects' arousal

L/R	Region	Number of voxels ( <i>k</i> )	Talairach coordinates (mm)			<i>T</i> score
			<i>x</i>	<i>y</i>	<i>z</i>	
<b>Wm</b>						
L	Medial aspect of superior frontal gyrus	1975 <sup>a</sup>	-8	58	26	10.57
R	Medial aspect of superior frontal gyrus		4	56	24	9.05
L	Medial aspect of superior frontal gyrus		-4	46	10	8.53
L	Medial aspect of superior frontal gyrus	92	-6	20	60	6.90
	Posterior cingulate gyrus	52	0	-24	38	6.51
<b>Lm</b>						
R	Medial aspect of superior frontal gyrus	2528	4	52	24	9.76
R	Middle temporal gyrus	3	58	-14	-22	5.06
L	Middle temporal gyrus	25	-58	-14	-20	5.70
	Vermis	2	6	-62	-6	5.02

Note. Maximally activated voxels in brain areas that significantly correlated with the subjects' arousal ( $P < 0.05$ , corrected). Wm, modulation by the subjects' arousal in the events of win. Lm, modulation by the subjects' arousal in the events of loss.

<sup>a</sup> This amount is the total number of activated voxels for the bilateral medial aspect of superior frontal gyrus in Wm.

### Medial aspect of prefrontal cortex

In our study, the maximum activation during a series of reward-related event and a series of penalty-related event (i.e., four wins in a row and four losses in a row) was localized to the cingulate sulcus, the anatomical border between the anterior cingulate (Brodmann's area 32), and the medial prefrontal cortex (Brodmann's area 9). Studies on cytoarchitectonic characteristics (Vogt et al., 1995) and laminar characteristics (Barbas and Pandya, 1989) of these two areas suggest that a rigid distinction between this portion of the anterior cingulate cortex and the medial prefrontal cortex is misleading. Given that the observed activation extended into both the anterior cingulate and the medial prefrontal cortex, this most likely reflects processes characteristic of these two regions.

Activation of the medial prefrontal cortex observed in our study appears to be involved in the emotionally salient context. Results of other imaging studies are consistent with our data. The medial prefrontal cortex is an area that was

found to be activated in previous PET studies of emotion (George et al., 1995; Lane et al., 1997c). Since the medial prefrontal cortex appears to participate in aspects of emotion unrelated to its type, valence, or method of induction (Lane et al., 1997b; Reiman et al., 1997), it may be involved in the arousal component of emotional response as opposed to valence appraisal (Lang and Greenwald, 1988).

Several studies showed that the anterior cingulate cortex is related to pain (Peyron et al., 1999; Rainville et al., 1997), performance monitoring (Shima and Tanji, 1998), error detection (Carter et al., 1998), conflict monitoring (Bothvinick et al., 1999), and response selection (Turken and Swick, 1999). The activation we observed in our study when subjects experience four wins in a row or four losses in a row, however, is more anterior to the foci reported in these studies. Our interpretation of the data is that the activation of the anterior cingulate cortex is related to emotional processing and subjective experience of emotion (Lane et al., 1997a). Some lesion studies support this idea. Ventromedial prefrontal lesions are associated with the ab-

Table 5  
Brain activities for a win following four losses in a row and a loss following four wins in a row

L/R	Region	Number of voxels ( <i>k</i> )	Talairach coordinates (mm)			<i>T</i> score
			<i>x</i>	<i>y</i>	<i>z</i>	
<b>SW - W1</b>						
R	Middle frontal gyrus	850 <sup>a</sup>	34	22	52	7.52
R	Middle frontal gyrus		26	42	38	5.98
<b>SL - L1</b>						
R	Middle frontal gyrus	317 <sup>b</sup>	36	30	36	5.38
R	Middle frontal gyrus		26	16	54	4.94

Note. Maximally activated voxels in brain areas where significantly higher activities were found in SW than in W1 and in SL than in L1 ( $P < 0.05$ , corrected).

<sup>a</sup> This amount is the total number of activated voxels for the right middle frontal gyrus in SW - W1.

<sup>b</sup> This amount is the total number of activated voxels for the right middle frontal gyrus in SL - L1.



sence of autonomic response to positively as well as negatively charged pictures (Damasio et al., 1990). Ventral prefrontal lesions are related to changes in patients' ability to feel emotions compared with their premorbid state (Homak et al., 1996). This is consistent with Papez' suggestion (Papez, 1937) that this region is "the seat of emotional experience."

A study on single neuronal signal in the anterior cingulate cortex of monkeys demonstrated responses that progressively changed strength with reward expectancy (Shidara and Richmond, 2002). The authors hypothesized that the anterior cingulate cortex activity reflects or perhaps regulates the degree of motivation and/or reward expectancy as the multistage task progresses. A recent study of event-related brain potentials (ERPs) in human participants performing a monetary gambling task also suggests that the anterior cingulate cortex activity is involved in assessing the motivational impact of the outcome events (Gehring and Wilboughby, 2002). Taking these data into account, the activations of the anterior cingulate cortex in our study seem to be sequential-context-dependent, reflecting the strength of subjects' reward expectancy (and/or motivation) or premonition of penalty.

#### *Lateral prefrontal cortex*

On the other hand, the right dorsolateral prefrontal cortex appears to represent a different context in the human brain. Many studies indicate that the right dorsolateral prefrontal cortex is activated during the execution of the Wisconsin Card Sorting Test (WCST; Milner, 1963, 1971; Berman et al., 1995; Nagahama et al., 1996; Meyer-Lindenberg et al., 2002). The WCST is the standard task paradigm for detecting human frontal lobe dysfunction. The WCST has been used to assess the ability to shift a cognitive set from one perceptual attribute of a complex visual stimulus to another (i.e., set-shifting ability). For example, after a prescribed number of successive correct responses to one dimension (e.g., color), the previously irrelevant dimension (i.e., shape) becomes relevant and the subject must shift the sorting criterion to the other one.

In our data, the right dorsolateral prefrontal cortex showed significant activation in the event of switching between reward and penalty, such as "a win following four losses in a row" and "a loss following four wins in a row." Essentially, both in our study and in the WCST, subjects are challenged on switching events. For example, a succession of wins through a sequence of trials has the effect of promoting the development of an attentional and/or emotional bias toward the reward-related event. This bias is then challenged on a loss that requires the subject's attention and/or emotion to be directed away from reward-related event toward the newly penalty-related event and vice versa.

Thus, taking these data into account, the activation of the right dorsolateral prefrontal cortex may depend on the context in which the nature of event is switched from one to

another. In other words, this region may be a critical area in higher-order shifting of attention and/or emotion between different conceptual categories of abstract values. This proposal is consistent with the result in previous studies that the right dorsolateral prefrontal cortex is associated with unpredictability and surprise violations of the learned association (Fletcher et al., 2001; Zeki and Marini, 1998; Fink et al., 1999). This is also consistent with the findings of single-unit studies of the monkey; that is, neurons in the lateral prefrontal cortex are sensitive to the presence or absence of an expected reward in delayed reaction time tasks (Watanabe, 1990, 1992). Thus, the dorsolateral prefrontal cortex may be the key region for monitoring an ongoing outcome and essential for adaptive behavior.

#### *Other related issues*

Our study was based on the previous study performed by Elliott et al. (2000); however, our results are different from theirs. Elliott et al. examined the differential responses to wins occurring as a part of a winning streak and the differential responses to losses occurring as a part of a losing streak. This is conceptually similar to our results of the W4 – W1 and L4 – L1 subtractions, but our results diverged from those of Elliott's. A possible explanation for this divergence is the method of analysis and the design of our experiment (i.e., the sequence of win or loss events), which are different from theirs. Elliott and colleagues focused on the brain activities that were modulated by "how fast the reward level is changing." Moreover, they reported that there were no neural responses associated with a main effect of winning or losing streak. Thus, the regions they detected to be responsive to a winning or losing streak were activated only in contexts in which there was a congruence between the rate of change and the reward level. On the other hand, we took a different approach to the analysis, using a more simple analytical method: we focused on brain activities that were modulated by how each win or loss event was repeated, regardless of the current reward level. Note that, to minimize the effect of this reward level, we designed the win/loss sequence such that the list of scores at which each W4 (i.e., four wins in a row) occurs is the same as that of scores at which each L4 (i.e., four losses in a row) occurs across the trial. This is why Fig. 1 shows point symmetry. Furthermore, we set the winning and losing streaks as four wins in a row and four losses in a row, respectively. This fixed design is more likely to produce subjects' feelings over the winning/losing streaks and enabled us to more efficiently detect the brain activities associated with these streaks. In fact, increased neuronal activities were observed at the anterior cingulate in a single-unit animal study in which the sequence of reward/no-reward events were fixed (Shidara and Richmond, 2002). This is consistent with our results.

Our results are also different from those of another relevant gambling task study carried out by Breiter and col-

leagues 2001). In their study, increased neural activities were observed in the nucleus accumbens during the period of anticipation. In our study, however, no activation was detected in these regions. This seems to be due to difference in our experimental design from theirs. First, there was a variation in the magnitude of each reward/penalty in their studies as opposed to the consistency in the magnitude of each reward/penalty was in our study. Second, the phase of anticipation toward a reward/penalty and the phase of outcome of a reward/penalty were clearly distinguishable in their task design. On the other hand, we cannot clearly distinguish the brain activities associated with the response to the outcome of a reward/penalty from that associated with the anticipation of a reward/penalty in our task design. Because of the consecutive trials, it was not clear when the anticipation/motivation for a reward/penalty of the next trial starts to occur in our task. The possibility cannot be denied that the reaction to the outcome of a reward/penalty and the anticipation of the next reward/penalty occur together during the second part of each trial as shown in Fig. 1. This is a limitation in our study. For the analyses of our data, however, the onset of hemodynamic response function was at the point when subjects found out whether the outcome was a win or a loss in each trial. The model was based on our assumption that the subjects' emotional reactions start at that point.

Previous PET studies found that the human orbitofrontal cortex can be activated by monetary reward (Thut et al., 1997). A recent fMRI study also demonstrated that an increase in the activity of the medial orbitofrontal cortex is related to the subjects' receipt of monetary reward and deactivation following penalty (O'Doherty et al., 2001). The authors obtained the converse pattern of activation in the lateral orbitofrontal cortex (activation following penalties, deactivation following rewards). In our data, however, no significant activation was observed in the orbitofrontal cortex in both winning events and losing events. This may be due to the difference in our experimental design from theirs. That is, we did not aim to detect brain regions related to reward or penalty itself, but those regions that are context-dependent in which reward or penalty is received. In our data, however, the results of comparison between all wins and all losses (shown in Table 3) demonstrate the activations in the bilateral ventral striatum, which is consistent with the findings of previous studies (Apicella et al., 1991; Shidara et al., 1998). Another possible reason is that susceptibility and distortion artifacts in the orbitofrontal cortex may have made it difficult to accurately acquire data for this region.

#### *Clinical implications*

Although this is speculative, because our study is not a clinical study, we would like to consider the possible clinical relevance of our findings in this paragraph. Taking our results and the results of the previous studies into account,

the medial aspect of the prefrontal cortex and the anterior cingulate cortex play a fundamental role in generating and/or regulating appropriate emotional reactions, such as general (i.e., unrelated to reward/penalty) excitement or tension, which are necessary for adjusting oneself to new circumstances. Some clinical studies showed that patients with major depression show a significant decrease in activity of the anterior cingulate cortex (Bench et al., 1993) and hypoperfusion of the anterior cingulate cortex and the medial aspect of the prefrontal cortex (Awata et al., 1998). Patients with posttraumatic stress disorder (PTSD) exhibit a lower anterior cingulate cortex activation in response to disorder-specific stimuli than trauma-exposed normal subjects (Shin et al., 1999). Moreover, patients with major depression or PTSD show abnormality in subjective experience of emotion. Markedly diminished interest or pleasure, complaint of having no feelings, decreased energy or motivation are common symptoms of depression, while anxiety, intrusive memories of the trauma, and increased arousal are major symptoms of PTSD (American Psychiatric Association, 2000). Thus, dysfunction of the anterior cingulate cortex and the prefrontal cortex may lead to failure in regulating mood and in subjective experience of emotion. An emotional or behavioral representation might become inappropriate and result in vulnerability to maladjustment to a new environment. This could be a possible explanation for the observation that the onset of a mood disorder is often induced by both a series of positive life events and a series of negative life events (Holmes and Rahe, 1967; Gelder et al., 2001).

Our data regarding the dorsolateral prefrontal cortex may also have relevance to other mental disorders. Typically, patients with frontal lobe damage show perseverative or "stuck in set" errors in the WCST (Milner, 1963), while schizophrenic patients show a poor performance in the WCST and reduced frontal activation particularly in the right dorsolateral prefrontal cortex in comparison with healthy controls (Riehemann et al., 2001; Wolkin et al., 1992; Volz et al., 1997). This reduction in activation in the right dorsolateral prefrontal cortex is not a pure effect of neuroleptic treatment (Riehemann et al., 2001). These results suggest that the dysfunction of the right dorsolateral prefrontal cortex could be related to the tendency of patients with schizophrenia to show vulnerability to sudden environmental changes (Norman and Malla, 1993).

#### **Conclusion**

In conclusion, we demonstrated that there exist brain activities associated with the event-sequence context in which abstract reward or penalty is received. Our data also suggest that information regarding this context is dissociably represented in the human brain, regardless of the nature of event (i.e., reward-related or penalty-related). The anterior cingulate cortex, the medial aspect of the prefrontal

cortex, and the right dorsolateral prefrontal cortex appear to have an important role in these context-dependent activities. Taking the relevant findings about the function of these regions into account, these context-dependent activities are crucial to adapting oneself to new circumstances and may account for the clinical symptoms of various mental illnesses in which dysfunction of these regions has been reported.

### Acknowledgments

We thank Y. Satoh, O. Imaizumi, and A. Nosaka for support in fMRI data acquisition and K. Satoh, A. Harada, M. Fujikawa, M. Kubota, F. Hashimoto, and T. Satoh for help in data analysis. This study was supported by JST/RISTEX, R&D promotion scheme for regional proposals promoted by TAO, a Grant-Aid for Scientific Research on Priority areas (C)—Advanced Brain Science Project—from MEXT, and the 21st Century Center of Excellence (COE) Program (Ministry of Education, Culture, Sports, Science and Technology) entitled “A Strategic Research and Education Center for an Integrated Approach to Language and Cognition” (Tohoku University).

### References

- American Psychiatric Association 2000. Diagnostic and Statistical Manual of Mental disorders, fourth ed., text revision. American Psychiatric Association, Washington, DC.
- Apicella, P., Ljungberg, T., Scamati, E., Schultz, W., 1991. Responses to reward in monkey dorsolateral and ventral striatum. *Exp. Brain Res.* 85, 491–500.
- Awata, S., Ito, H., Konno, M., Ono, S., Kawashima, R., Fukuda, H., Sato, M., 1998. Regional cerebral blood flow abnormalities in late-life depression: relation to refractoriness and chronification. *Psychiatr. Clin. Neurosci.* 52, 97–105.
- Barbas, H., Pandya, D.N., 1989. Architecture and intrinsic connections of the prefrontal cortex in the rhesus monkey. *J. Comp. Neurol.* 286, 353–375.
- Bechara, A., Damasio, A.R., Damasio, H., Anderson, S.W., 1994. Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition* 50, 7–15.
- Bechara, A., Damasio, H., Tranel, D., Damasio, A.R., 1997. Deciding advantageously before knowing the advantageous strategy. *Science* 275, 1293–1295.
- Bechara, A., Tranel, D., Damasio, H., Damasio, A.R., 1996. Failure to respond autonomically to anticipated future outcomes following damage to the prefrontal cortex. *Cereb. Cortex* 6, 215–225.
- Bench, C.J., Friston, K.J., Brown, R.G., Scott, L.C., Frackowiak, R.S.J., Dolan, R.J., 1993. The anatomy of melancholia—focal abnormalities of cerebral blood flow in major depression. *Psychol. Med.* 22, 607–615.
- Berman, K.F., Ostern, J.L., Randolph, C., Gold, J., Goldberg, T.E., Coppola, R., Carson, R.E., Herscovitch, P., Weinberger, D.R., 1995. Physiological activation of a cortical network during performance of the Wisconsin Card Sorting Test: a position emission tomography study. *Neuropsychologia* 33, 1027–1046.
- Botvinick, M., Nystrom, L.E., Fissell, K., Carter, C.S., Cohen, J.D., 1999. Conflict monitoring versus selection-for-action in anterior cingulate cortex. *Nature* 402, 179–181.
- Breiter, H.C., Aharon, I., Kahneman, D., Dale, A., Shizgal, P., 2001. Functional imaging of neural responses to expectancy and experience of monetary gains and losses. *Neuron* 30, 619–639.
- Carter, C.S., Braver, T.S., Barch, D.M., Botvinick, M.M., Noll, D., Cohen, J.D., 1998. Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science* 280, 747–749.
- Critchley, H.D., Elliott, R., Mathias, C.J., Dolan, R.J., 2000. Neural activity relating to generation and representation of galvanic skin conductance responses: a functional magnetic resonance imaging study. *J. Neurosci.* 20, 3033–3040.
- Damasio, A.R., Tranel, D., Damasio, H., 1990. Individuals with sociopathic behavior caused by frontal damage fail to respond autonomically to social stimuli. *Behav. Brain Res.* 41, 81–94.
- Elliott, R., Friston, K.J., Dolan, R.J., 2000. Dissociable neural responses in human reward systems. *J. Neurosci.* 20, 6159–6165.
- Fink, G.R., Marshall, J.C., Halligan, P.W., Frith, C.D., Driver, J., Frackowiak, R.S.J., Dolan, R.J., 1999. The neural consequences of conflict between intention and the senses. *Brain* 122, 497–512.
- Fletcher, P.C., Anderson, J.M., Shanks, D.R., Honey, R., Carpenter, T.A., Donovan, T., Papadakis, N., Bullmore, E.T., 2001. Responses of human frontal cortex surprising events are predicted by formal associative learning theory. *Nat. Neurosci.* 4, 1043–1048.
- Gehring, W.J., Willoughby, A.R., 2002. The medial frontal cortex and the rapid processing of monetary gains and losses. *Science* 295, 2279–2282.
- Gelder, M., Mayou, R., Cowen, P., 2001. Shorter Oxford Textbook of Psychiatry, 4th ed. Oxford University Press, New York.
- George, M.S., Ketter, T.A., Parekh, P.I., Horwitz, B., Herscovitch, P., Post, R.M., 1995. Brain activity during transient sadness and happiness in healthy women. *Am. J. Psychiatry* 152, 341–351.
- Holmes, T.H., Rahe, R.H., 1967. The social readjustment rating scale. *J. Psychosomatic Res.* 11, 213–218.
- Homak, J., Rolls, E.T., Wade, D., 1996. Face and voice expression identification in patients with emotional and behavioural changes following ventral frontal lobe damage. *Neuropsychologia* 34, 247–261.
- Knutson, B., Adams, C.M., Fong, G.W., Hommer, D., 2001a. Anticipation of increasing monetary reward selectively recruits nucleus accumbens. *J. Neurosci.* 21, 1–5.
- Knutson, B., Fong, G.W., Adams, C.M., Vamer, J.L., Hommer, D., 2001b. Dissociation of reward anticipation and outcome with event-related fMRI. *NeuroReport* 12, 3683–3687.
- Lane, R.D., Fink, G.R., Chau, P.M.L., Dolan, R.J., 1997a. Neural activation during selective attention to subjective emotional responses. *NeuroReport* 8, 3969–3972.
- Lane, R.D., Reiman, E.M., Ahem, G.L., Schwartz, G.E., Davidson, R.J., 1997b. Neuroanatomical correlates of happiness, sadness, and disgust. *Am. J. Psychiatry* 154, 926–933.
- Lane, R.D., Reiman, E.M., Bradley, M.M., Lang, P.J., Ahem, G.L., Davidson, R.J., Schwartz, G.E., 1997c. Neuroanatomical correlates of pleasant and unpleasant emotion. *Neuropsychologia* 35, 1437–1444.
- Lang, P.J., Greenwald, M.K., 1988. The International Affective Picture System Standardization Procedure and Initial Group Results for Affective Judgments, technical report A, Center for Research in Psychophysiology, University of Florida, Gainesville, FL.
- Ljungberg, T., Apicella, P., Schultz, W., 1992. Responses of monkey dopamine neurons during learning of behavioural reactions. *J. Neurophysiol.* 67, 145–163.
- Meyer-Lindenberg, A., Miletich, R.S., Kohn, P.D., Esposito, G., Carson, R.E., Quarantelli, M., Weinberger, D.R., Berman, K.F., 2002. Reduced prefrontal activity predicts exaggerated striatal dopaminergic function in schizophrenia. *Nat. Neurosci.* 5, 267–271.
- Milner, B., 1963. Effects of different brain lesions on card sorting: the role of frontal lobes. *Arch. Neurol.* 9, 90–100.
- Milner, B., 1971. Interhemispheric differences in the localization of psychological processes in man. *Br. Med. Bull.* 27, 272–277.

- Mirenovicz, J., Schultz, W., 1994. Importance of unpredictability for reward responses in primate dopamine neurons. *J. Neurophysiol.* 72, 1024–1027.
- Nagahama, Y., Fukuyama, H., Yamauchi, H., Matsuzaki, S., Konishi, J., Shibasaki, H., Kinura, J., 1996. Cerebral activation during performance of a card sorting test. *Brain* 119, 1667–1675.
- Niki, H., Sakai, M., Kubota, K., 1972. Delayed alternation performance and unit activity of the caudate head and medial orbitofrontal gyrus in the monkey. *Brain Res.* 38, 345–353.
- Norman, R.M., Malla, A.K., 1993. Stressful life events and schizophrenia—a review of the research. *Br. J. Psychiatry* 162, 161–166.
- O'Doherty, J., Kringelbach, M.L., Rolls, E.T., Homak, J., Andrews, C., 2001. Abstract reward and punishment representations in the human orbitofrontal cortex. *Nat. Neurosci.* 4, 95–102.
- Papez, J.W., 1937. A proposed mechanism of emotion. *Arch. Neurol. Psychiatry* 38, 725–734.
- Peyron, R., García-Larrea, L., Grégoire, M.C., Costes, N., Convers, P., Lavenne, F., Mauguière, F., Michel, D., Laurent, B., 1999. Hemodynamic brain response to acute pain in humans: sensory and attentional networks. *Brain* 122, 1765–1779.
- Rainville, P., Duncan, G.H., Price, D.D., Carrier, B., Bushnell, M.C., 1997. Pain affect encoded in human anterior cingulate but not somatosensory cortex. *Science* 277, 968–971.
- Reiman, E.M., Lane, R.D., Ahem, G.L., Schwartz, G.E., Davidson, R.J., Friston, K.J., Yun, L.S., Chen, K., 1997. Neuroanatomical correlates of externally and internally generated human emotion. *Am. J. Psychiatry* 154, 918–925.
- Riehemann, S., Volz, H.P., Stüzer, P., Smesny, S., Gaser, C., Sauer, H., 2001. Hypofrontality in neuroleptic-naïve schizophrenia patients during the Wisconsin Card Sorting Test—a fMRI study. *Eur. Arch. Psychiatr. Clin. Neurosci.* 251, 66–71.
- Rolls, E.T., Homak, J., Wade, D., McGrath, J., 1994. Emotion-related learning in patients with social and emotional changes associated with frontal lobe damage. *J. Neurol. Neurosurg. Psychiatry* 57, 1518–1524.
- Romo, R., Schultz, W., 1990. Dopamine neurons of the monkey midbrain: contingencies of responses to active touch during self-initiated arm movements. *J. Neurophysiol.* 63, 592–606.
- Schultz, W., 1986. Responses of midbrain dopamine neurons to behavioural trigger stimuli in the monkey. *J. Neurophysiol.* 56, 1943–1962.
- Schultz, W., Apicella, P., Scamati, E., Ljungberg, T., 1992. Neuronal activity in monkey ventral striatum related to the expectation of reward. *J. Neurosci.* 12, 4595–4610.
- Schultz, W., Romo, R., 1990. Dopamine neurons of the monkey midbrain: contingencies of responses to stimuli eliciting immediate behavioural reactions. *J. Neurophysiol.* 63, 607–624.
- Shidara, M., Aigner, T.G., Richmond, B.J., 1998. Neuronal signals in the monkey ventral striatum related to progress through a predictable series of trials. *J. Neurosci.* 18, 2613–2625.
- Shidara, M., Richmond, B.J., 2002. Anterior cingulate: single neuronal signals related to degree of reward expectancy. *Science* 296, 1709–1711.
- Shima, K., Tanji, J., 1998. Role for cingulate motor area cells in voluntary movement selection based on reward. *Science* 282, 1335–1338.
- Shin, L.M., McNally, R.J., Kosslyn, S.M., Thompson, W.L., Rauch, S.L., Alpert, N.M., Metzger, L.J., Lasko, N.B., Orr, S.P., Pitman, R.K., 1999. Regional cerebral blood flow during script-driven imagery in childhood sexual abuse-related PTSD: a PET investigation. *Am. J. Psychiatry* 156, 575–584.
- Talairach, J., Tournoux, P., 1988. *Coplanar Stereotaxic Atlas of the Human Brain*. George Thieme Verlag, Stuttgart.
- Thut, G., Schultz, W., Roelcke, U., Nienhusmeier, M., Missimer, J., Maguire, P.R., Leenders, K.L., 1997. Activation of the human brain by monetary reward. *NeuroReport* 8, 1225–1228.
- Thorpe, S.J., Rolls, E.T., Maddison, S., 1983. The orbitofrontal cortex: neuronal activity in the behaving monkey. *Exp. Brain Res.* 49, 93–115.
- Tremblay, L., Schultz, W., 1999. Relative reward preference in primate orbitofrontal cortex. *Nature* 398, 704–708.
- Tremblay, L., Schultz, W., 2000. Reward-related neuronal activity during go-no go task performance in primate orbitofrontal cortex. *J. Neurophysiol.* 83, 1864–1876.
- Turken, A.U., Swick, D., 1999. Response selection in the human anterior cingulate cortex. *Nat. Neurosci.* 2, 920–924.
- Vogt, B.A., Nimchinski, E.A., Vogt, L.J., Hof, P.R., 1995. Human cingulate cortex: surface features, flat maps, and cytoarchitecture. *J. Comp. Neurol.* 359, 490–506.
- Volz, H.P., Gaser, C., Häger, F., Rzanny, R., Mentzel, H.J., Kreitschmann-Andermahr, I., Kaiser, W.A., Sauer, H., 1997. Brain activation during cognitive stimulation with the Wisconsin Card Sorting Test—a functional MRI study on healthy volunteers and schizophrenics. *Psychiatr. Res.* 75, 145–157.
- Watanabe, M., 1990. Prefrontal unit activity during associative learning in the monkey. *Exp. Brain Res.* 80, 296–309.
- Watanabe, M., 1992. Frontal units of the monkey coding the associative significance of visual and auditory stimuli. *Exp. Brain Res.* 89, 233–247.
- Watanabe, M., 1996. Reward expectancy in primate prefrontal neurons. *Nature* 382, 629–632.
- Wolkin, A., Sanfilippo, M., Wolf, A.P., Angrist, B., Brodie, J.D., Rotrosen, J., 1992. Negative symptoms and hypofrontality in chronic schizophrenia. *Arch. Gen. Psychiatry* 49, 959–965.
- Zeki, S., Marini, L., 1998. Three cortical stages of color processing in the human brain. *Brain* 121, 1669–1685.

## Social Isolation Stress Significantly Enhanced the Disruption of Prepulse Inhibition in Mice Repeatedly Treated with Methamphetamine

HONGMEI DAI,<sup>a,b</sup> HIROTOMO OKUDA,<sup>a</sup> KENTARO IWABUCHI,<sup>a</sup>  
EIKO SAKURAI,<sup>a</sup> ZHONG CHEN,<sup>a</sup> MOTOHISA KATO,<sup>a</sup> KAZUIE IINUMA,<sup>b</sup>  
AND KAZUHIKO YANAI<sup>a</sup>

*Departments of Pharmacology<sup>a</sup> and Pediatrics,<sup>b</sup> Tohoku University School of Medicine, Sendai 980-8575, Japan*

**ABSTRACT:** Repeated administration of methamphetamine (METH) causes reverse tolerance or behavioral sensitization in mice. However, the effects of social isolation stress on the METH-caused reverse tolerance have not been studied until now. The aim of this study was to investigate the effects of social isolation stress on METH-caused reverse tolerance by examining the prepulse inhibition of startle response (PPI). PPI was tested in socially isolated and grouped mice after repeated METH injections. Locomotor activity and PPI were also examined just after a four-week isolation rearing period as a control experiment. After completing behavioral experiments, the mice were sacrificed, and the contents of monoamines, including histamine in the brain, were measured. Social isolation stress significantly lowered the locomotion and disrupted PPI. Repeated injections of METH enhanced the effects of social isolation on PPI. The content of dopamine and histamine significantly increased in the cortex, and the turnover rate of dopamine decreased significantly. These findings demonstrate that social isolation stress significantly enhances METH-induced behavioral sensitization and that the altered histaminergic neuron system might play an important role in METH-induced behavioral sensitization in addition to dopaminergic and serotonergic neurotransmission. Our data suggest that social isolation is involved in the development of METH-induced psychosis, schizophrenia, and other related psychiatric disorders.

**KEYWORDS:** social isolation; prepulse inhibition; locomotor activity; methamphetamine; dopamine; histamine; behavioral sensitization

### INTRODUCTION

It is known that repeated administration of psychoactive drugs such as methamphetamine (METH) results in a progressive enhancement of behavioral effects. This phenomenon was termed reverse tolerance or behavioral sensitization. It has been widely used as a model of stimulant-induced psychosis, paranoid schizophrenia, as

Address for correspondence: Kazuhiko Yanai, M.D., Ph.D., Department of Pharmacology, Tohoku University School of Medicine, 2-1 Seiryō-machi, Aoba-ku, Sendai 980-8575, Japan. Voice: +81-22-717-8055; fax: +81-22-717-8060.  
yanai@mail.tains.tohoku.ac.jp

Ann. N.Y. Acad. Sci. 1025: 257–266 (2004). © 2004 New York Academy of Sciences.  
doi: 10.1196/annals.1316.032

well as neuronal plasticity in drug abuse. In addition, social isolation in several mammalian species can result in profound alteration of brain development, social behavior, and information processing. Moreover, there is substantial evidence that early-life events influence brain development and subsequently play an important role in the pathogenesis of certain psychiatric disorders including schizophrenia and depression. However, the effects of early social isolation stress on METH-induced behavioral sensitization have not been studied until now. In this study, we examined whether social isolation stress can affect the prepulse inhibition of the acoustic startle response. PPI is the reduction in startle reflex caused by a pre-stimulus, and it is often used as a measure of sensorimotor gating to investigate the underlying pathophysiology in schizophrenia.<sup>1</sup>

Repeated administration of METH increases the levels of dopamine (DA) release, which is related to behavioral changes. Several reports have demonstrated that isolated rats showed elevated DA concentrations<sup>2</sup> and altered DA turnover in the frontal cortex.<sup>3</sup> The histaminergic (HA) neuron system has an inhibitory role on the behavioral effects of METH.<sup>4,5</sup> However, the effects of the HA neuron system on social isolation stress have not been clear, although one study reported that social stress of isolation considerably diminished the number of brain mast cells without a significant increase of the histamine content in the hypothalamus.<sup>6</sup> In this study, we examined the effects of social isolation on METH-induced behavioral sensitization. We also measured the changes of monoamines and their metabolites in the brain.

## MATERIALS AND METHODS

### *Animals*

All experiments were carried out in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals. Male C57BL/6 mice were purchased from Japan (SLC) immediately after weaning (21 postnatal days) and divided into isolated-housing and multiply-housed groups. The multiply-housed groups contained four mice per cage. All cages were made of plastic, 32 × 22 × 15 cm<sup>3</sup> in size with sawdust flooring. The mice were housed at a constant temperature (22 ± 3°C) and humidity (55 ± 10%) in a room on an automatically controlled light cycle (light on, 0600–1800); they had free access to food and water. All experiments were performed during the light phase between 0800 and 1800. Experimental protocols were approved by the Animal Care Committee, Tohoku University School of Medicine, Japan.

### *Drug*

Methamphetamine (Dainippon Pharmaceutical Co, Japan) was dissolved in saline and was administered intraperitoneally (i.p.) in a volume of 5 mL/kg.

### *Apparatus*

Each mouse was placed in a cage that was unfamiliar to him, and its locomotor activity was measured with a photo-beam system (SUPERMEX®, Muromachi kikai Co., Tokyo, Japan). Startle response testing was conducted in a startle response unit

(Responder-X system, Columbus Instruments, OH). The device was attached to a platform on which the animal was placed in an  $11 \times 6 \times 5$  cm<sup>3</sup> ventilated cubicle. A high-frequency loudspeaker inside the startle chamber produced a continuous broadband background noise of 65 dB and various acoustic stimuli. Vibrations caused by the whole-body startle responses were detected by a piezoelectric accelerometer unit attached to the platform; these signals then were digitized and stored in a computer.

#### *Test Session of Startle Response*

Five different trial types were presented to mice: a 40-ms broadband 120-dB burst (PULSE ONLY trial); three different PREPULSE + PULSE trials in which 20-ms-long 3-dB (68 dB, pp3p120), 6-dB (71 dB, pp6p120), or 12-dB (77dB, pp12p120) stimuli above a 65-dB background preceded the 120-dB pulse by 100 ms (onset to onset), and a NO STIMULUS trial in which only the background noise was presented. All trials were presented in a pseudo-random order, with an average of 15 s separating trials. The test session began with a 5-min acclimation period, followed by four consecutive blocks of test trials (total: 62 trials). Blocks one and four consisted of six consecutive PULSE ONLY trials and were omitted from the overall analysis. Blocks two and three each contained five PULSE ONLY trials, five pp3p120 trials, five pp6p120 trials, five pp12p120 trials, and five NO STIMULUS trials.

#### *Measurements of Histamine and Other Neurotransmitters Contents*

After completion of the behavioral experiment, the mice were immediately decapitated and the brains were quickly divided on ice into three regions: cortex, diencephalons, and cerebellum including brainstem. The samples were then stored at  $-80^{\circ}\text{C}$  until assay. The brain samples were weighed and homogenized in 10 vols 0.2 M perchloric acid containing 5 mM Na<sub>2</sub>-EDTA and 500 nM 5-hydroxy-N<sup>w</sup>-methyltryptamine oxalate (internal standard) in a Polytron homogenizer (Kinematica, Lucern, Switzerland) at a maximum setting of 1 min on an ice bath. Then the homogenate was centrifuged at 12,000 rpm for 20 min at  $4^{\circ}\text{C}$ . After filtration (0.45  $\mu\text{m}$ ), the solution was injected into a high-performance liquid chromatography (HPLC) system. Monoamines and their metabolites were separated using an HPLC system at  $30^{\circ}\text{C}$  on a reverse-phase analytical column (ODS-C18, 4.6 mm i.d.  $\times$  25 cm), and detected by an electrochemical detector (Model Coulochem II, ESA Co., MA). The column was eluted with 0.04 M phosphate—0.04 M citrate buffer (PH 3.0) containing 11.7% methanol and 4.7% acetonitrile, 7.5 mM sodium 1-heptanesulfonate, and 0.08 mM Na<sub>2</sub>-EDTA. The following monoamines and their metabolites were measured: norepinephrine, dopamine (DA), 3,4-dihydroxyphenylacetic acid (DOPAC), homovanillic acid (HVA), 3-methoxytyramine (3-MT), 5-hydroxytryptamine (serotonin, 5-HT), and 5-hydroxyindoleacetic acid (5-HIAA). In addition, we measured histamine content fluorometrically using an HPLC system with *O*-phthalaldehyde after separation on an HPLC column (CM/2SW, 4.6 mm i.d.  $\times$  15 cm, Tosoh, Tokyo; particle size 5  $\mu\text{m}$ ).


#### *Experimental Procedures*

Twenty-four C57BL/6 mice were divided into two groups: socially isolated and multiply housed. Twelve mice were subjected to social isolation stress and 12 mice

were used for the control. Four weeks after isolated rearing, the mice were individually placed in a locomotor-activity testing system for 4 h. Then the startle responses were examined the next day. Twenty-four C57BL/6 mice (twelve in each group) were used for the reverse tolerance experiment of METH. Four weeks after isolated and grouped rearing, the mice were injected with 1 mg/kg methamphetamine every two days. One week after completion of the seventh injection, the startle responses were examined 20 min after the final injection of METH. After behavioral experiments, the mice were quickly killed, and their brain samples were stored until assay of histamine and other monoamines.

#### Data Analyses

Percent PPI (%PPI) for the three prepulse intensities was calculated according to the formula:  $\%PPI = 100 - \{[(\text{startle response for PREPULSE} + \text{PULSE trial}) / (\text{startle response for PULSE ONLY trial})] \times 100\}$ . The large values of PPI indicate that the prepulse potentially inhibited the startle response to the pulse stimuli, while a low or negative percent score of PPI imply that the prepulse stimuli had little inhibitory effect on the startle responses. Data were analyzed by the Mann-Whitney test to evaluate the statistical differences between the two groups. The value of  $P < .05$  was considered significant.



**FIGURE 1.** Effect of social isolation rearing on spontaneous locomotor activity. Data are expressed as the mean  $\pm$  SEM of 12 mice in each group. Statistical significance of difference between two groups was analyzed by the Mann-Whitney test after the calculation of area under the curve (AUC);  $P < .001$ .



## RESULTS AND DISCUSSION

### *Effects of Social Isolation Rearing on Spontaneous Locomotion and PPI*

The locomotor activity among the socially isolated mice was significantly lower than that of the multiply housed mice. In fact, the isolated mice barely moved at all 1.5 h after entering the observation chamber (FIG. 1). Prepulse inhibition was significantly disrupted in the prepulse of 77 dB when compared to the grouped mice. The PPI in 68-dB and 71-dB trials had a tendency to decrease, but it was not significant (FIG. 2a). The socially isolated mice showed a higher magnitude to the startle response than the grouped mice (FIG. 2b). The brain content of monoamines was not significantly different (data not shown), but the histamine content showed a tendency to increase in the brain. The increase was in particular statistically significant in the brain stem (data not shown). The results were essentially consistent with previous studies showing that chronic stress increased the amount of histamine content in the brain.<sup>7</sup>


### *Effect of Social Isolation on METH-Induced Behavioral Sensitization*

We examined METH-induced behavioral sensitization after 4 weeks of social isolation rearing. The locomotion activity was measured 30 min after the METH treatment. The locomotion gradually increased in isolated mice and grouped mice, suggesting that METH-induced behavioral sensitization could be observed in both groups. The isolated mice showed lower locomotion than the grouped mice (data not shown). After the development of behavioral sensitization, PPI was compared between the isolated and grouped mice. Treatment with METH is known to disrupt the PPI responses. The PPI was significantly more disrupted in isolated mice than in the grouped mice (FIG. 3a). Among the isolated mice treated with METH, the disruption of PPI was significantly enhanced in the 71-dB and 77-dB prepulse trials while the startle amplitude was not significantly changed (FIG. 3b). These data demonstrate that social isolation stress effectively enhances the disruption of prepulse inhibition. In accordance with the behavioral data, the content of norepinephrine (NE), dopamine (DA), and serotonin (5-HT) in the cortex of repeatedly METH-treated mice were significantly higher than those in the cortex of the grouped mice. The turnover rate of dopamine decreased significantly in the cortex of isolated mice when compared to the grouped mice (TABLE 1). The turnover rate of 5-HT showed a tendency to decrease in the cortex of isolated mice, but the change was not statistically significant. It is well known that repeated METH treatment increases the rate of dopamine release and results in the reduction of DA transporters; a decrease in transporters has also been found in METH abusers by PET studies.<sup>8</sup> In the present study, the content of NE, DA, and 5-HT were higher in socially isolated mice than the grouped mice, suggesting that early social isolation increases the content of NE, DA, and 5-HT with a decrease of their turnover rates. In parallel, histamine content also increased significantly in the cortex (FIG. 4). These data suggest that isolation stress might generally decrease monoaminergic transmission.


TABLE 1. Contents of monoamines and their metabolites from isolated and grouped mice after MAP-induced behavioral sensitization

Region	Condition	Norepinephrine	DOPAC	Dopamine	HIAA	HVA	3-MT	5-HT	(DOPAC + HVA + 3-MT)/DA	HIAA/5-HT
Cortex	Group	2.1 ± 0.2	1.8 ± 0.7	7.3 ± 1.1	3.5 ± 2.2	3.0 ± 0.6	0.79 ± 0.18	1.0 ± 0.2	0.77 ± 0.13	3.4 ± 1.5
	Isolation	2.4 ± 0.3*	1.7 ± 0.4	9.8 ± 1.7**	3.2 ± 0.4	2.4 ± 0.6	0.85 ± 0.16	1.2 ± 0.2*	0.49 ± 0.08***	2.7 ± 0.4
Diencephalon Group	Group	3.9 ± 0.7	3.2 ± 1.7	4.0 ± 2.0	9.9 ± 2.6	3.7 ± 1.2	0.43 ± 0.22	2.1 ± 0.4	2.0 ± 0.7	4.7 ± 1.2
	Isolation	3.9 ± 1.2	2.6 ± 1.2	3.2 ± 1.6	9.7 ± 4.3	3.3 ± 0.6	0.38 ± 0.20	2.1 ± 0.6	2.2 ± 0.8	4.5 ± 0.8
BS + CB	Group	3.1 ± 0.6	1.0 ± 0.3	0.91 ± 0.16	4.5 ± 1.1	3.7 ± 1.4	0.16 ± 0.09	1.1 ± 0.3	5.4 ± 1.9	4.1 ± 1.0
	Isolation	3.4 ± 0.6	1.1 ± 0.2	1.1 ± 0.3	4.7 ± 1.2	3.9 ± 1.7	0.17 ± 0.11	1.1 ± 0.2	5.0 ± 2.1	4.3 ± 1.2

Values (nmol/g brain tissue) are expressed as mean ± SD from 10 mice. Statistical significance between the two groups was analyzed by Mann-Whitney test. \* $P < .05$ , \*\* $P < .01$ , \*\*\* $P < .001$ . BS + CB: brain stem + cerebellum.



**FIGURE 2.** Effect of social isolation on the prepulse inhibition (A) and startle amplitude (B) of startle responses in different prepulse conditions. The PPI and startle amplitude caused by 120 dB in different prepulse conditions of 68, 71, and 77 dB. Data are expressed as the mean  $\pm$  SEM of 12 mice in each group. Statistical significance of difference between two groups was analyzed by the Mann-Whitney test. \* $P < .05$ .



**FIGURE 3.** The PPI (A) and startle response (B) in METH-induced behavioral sensitization of socially isolated mice. The PPI and startle amplitude caused by 120 dB in different prepulse conditions of 68, 71, and 77 dB. Data are expressed as the mean  $\pm$  SEM of isolated and grouped mice ( $n = 12$ ). Statistical significance of difference between the two groups was analyzed by the Mann-Whitney test. \* $P < .05$ .