

加藤元一郎、青柳 裕、阿内 謙吉、小野寺 晋志：
アルツハイマー病に類似した症状を呈する疾患の脳
機能画像統計解析、臨床放射線、49:1203-1210, 2004

2. 学会発表

Yutaka Kato, Motoichiro Kato, Masuro Shintani: An
earlier component of face perception detected by
seeing-as-face task.

15th Cogress of the International Society for
Brain Electromagnetic Topography, Aprill 11-14,
2004

Hotel MIRACOSTA, Chiba, Japan
Program and Abstracts, p68, 2004

Akiyama T, Kato M, Nakachi R, Saito F, Muramatsu
T, Kashima H: Gaze discrimination and the upper
bank of the right superior temporal sulcus.
Cognitive Neuroscience Society Annual Meeting
2004, April 18-20, 2004, San Francisco, USA,
Cognitive Neuroscience Society Annual Meeting
program 2004, p46, 2004

Nakachi R, Akiyama T, Kato M, Saito F, Yoshino F,
Muramatsu T, Kashima H: Inability to recognize
stimuli within categories in progressive
prosopagnosia.
Cognitive Neuroscience Society Annual Meeting
2004, April 18-20, 2004, San Francisco,
USA, Cognitive Neuroscience Society Annual
Meeting program 2004, p152, 2004

Kazue Igarashi, Yuko Sakai, Motoichiro Kato:
Study of Executive Function and Working Memory
in the Patients with ADHD (Attention Deficit /
Hyperactivity Disorders) .
Second International Conference on Working
Memory, August 17-20, 2004, Kyoto International
Conference Hall, Kyoto, Program and Abstracts,
p60, 2004

Motoichiro Kato: Verbal executive impairment in
dynamic aphasia.
Second International Conference on Working
Memory, Symposion 9,
August 17-20, 2004
Kyoto International Conference Hall, Kyoto,
Program and Abstracts, p14, 2004

藤永 直美、加藤 元一郎、村松 太郎、吉野 文
浩、三村 將、鹿島 晴雄：
生物カテゴリーに特異的な視覚失認例における視覚
的特徴（知識）の想起障害について
第 28 回日本神経心理学会 2004 年 9 月 17・18 日、
広島国際会議場

第 28 回日本神経心理学会総会プログラム予稿集、50

秋山 知子、斎藤 文恵、仲地 良子、村松 太郎、
加藤 元一郎、鹿島 晴雄：
片側扁桃体損傷における表情認知障害の左右差－
「怒り」と「恐怖」「悲しみ」の乖離

第 28 回日本神経心理学会 2004 年 9 月 17・18 日、
広島国際会議場

第 28 回日本神経心理学会総会プログラム予稿集、51

瀬間 久美子、早川 裕子、加藤 元一郎、三村
將：

左手に認められた病的把握現象の回復メカニズム
一症例の経過から

第 28 回日本神経心理学会 2004 年 9 月 17・18 日、
広島国際会議場

第 28 回日本神経心理学会総会プログラム予稿集、
102

船山 道隆、加藤 元一郎、鹿島 晴雄、三村 將：
脳梁損傷による半側空間無視

第 28 回日本神経心理学会 2004 年 9 月 17・18 日、
広島国際会議場

第 28 回日本神経心理学会総会プログラム予稿集、
116

加藤元一郎：失行症状の発現機序と責任病巣、
シンポジウム 「わかりやすい高次脳機能」、
第 45 回日本神経学会総会、
2004 年 5 月 11 日－14 日、東京
プログラム・抄録集、p 67、2004

H. 知的財産権の出願・登録状況
特になし。

Ⅲ. 研究成果の刊行に関する一覧表

研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Takahashi H., Koeda M., Oda K., Matsuda T., Matsushima E., Matsuura M., Asai K., Okubo Y	An fMRI study of differential neural response to affective pictures in schizophrenia	NeuroImage	22	1247-1254	2004
Takahashi H, Yahata N, Koeda M, Matsuda T, Asai K, Okubo Y	Brain activation associated with evaluative processes of guilt and embarrassment: an fMRI study	NeuroImage	23	967-974	2004
Watanabe H, Murata Y, Ohashi I, Oda K, Matsushima E, Okubo Y, Shibuya H	Long-term change in size of cerebral infarction: Predictive value of brain perfusion SPECT using statistical Parametric mapping	Cerebrovasc Dis	18	22-29	2004
Sakamoto A., Ogawa R., Suzuki H., Kimura M., Okubo Y., Fujiya T	Landiolol attenuates acute hemodynamic responses but does not reduce seizure duration during maintenance electroconvulsive therapy	Psychiatry Clin Neurosci	58	630-635	2004
Yasuno F., Suhara T., Okubo Y., Sudo Y., Inoue M., Ichimiya T., Takano A., Nakayama K., Halldin C., Farde L.	Low dopamine D ₂ receptor binding in subregions of the thalamus in schizophrenia	Am J Psychiatry	161	1016-1022	2004
Matsuura M, Adachi N, Oana Y, Okubo Y, Kato M, Nakano T, Takei N	A polydiagnostic and dimensional comparison of epileptic psychoses and schizophrenia spectrum disorders	Schizophr Res	69	189-201	2004
Zhang M.-R., Maeda J., Ogawa M., Noguchi J., Ito T., Yoshida Y., Okauchi T., Obayashi S., Suhara T., Suzuki K.	Development of a new radioligand, N-(5-Fluoro-2-phenoxyphenyl)-N-(2-[¹⁸ F]fluoroethyl-5-methoxybenzyl)acetamide, for PET imaging of peripheral benzodiazepine receptor in primate brain	J Med Chem	47	2228-2235	2004
Sasaki S., Kurosaki F., Haradahira T., Yamamoto F., Maeda J., Okauchi T., Suzuki K., Suhara T., Maeda M.	Synthesis of ¹¹ C-labelled bis(phenylalkyl)amines and their in vitro and in vivo binding properties in rodent and monkey brains	Bilo Pharm Bull	27	531-537	2004
Semba J., Akanuma N., Wakuta M., Tanaka N., Suhara T.	Alterations in the expressions of mRNA for GDNF and its receptors in the ventral midbrain of rats exposed to subchronic phencyclidine	Mol Brain Res	124	88-95	2004

Maeda J., Suhara T. , Zhang M.-R., Okauchi T., Yasuno F., Ikoma Y., Inaji M., Nagai Y., Takano A., Obayashi S., Suzuki K.	Novel peripheral benzodiazepine receptor ligand [¹¹ C]DAA1106 for PET: an imaging tool for glial cells in the brain	Synapse	52	283-291	2004
Takano A., Suhara T. , Maeda J., Ando K., Okauchi T., Obayashi S., Nakayama T., Kapur S.	Relationship between cortical dopamine D ₂ receptor occupancy and suppression of conditioned avoidance response in non-human primate	Psychiatry Clin Neurosci	58	330-332	2004
Yoshizaki T., Inaji M., Kouike H., Shimazaki T., Sawamoto K., Ando K., Date I., Kobayashi K., Suhara T. , Uchiyama Y., Okano H.	Isolation and transplantation of dopaminergic neurons generated from mouse embryonic stem cells	Neurosci Lett	363	33-37	2004
Hatakeyama H., Akita H., Maruyama K., Suhara T. , Harashima H.	Factors governing the in vivo tissue uptake of transferring- coupled polyethylene glycol liposomes in vivo	Int J Pharm	281	25-33	2004
Zhang H., Yoshikawa K., Tamura K., Sagou K., Tian M., Suhara T. , Kandatsu S., Suzuki K., Tanada S., Tsujii H.	Carbon-11-methionine positron emission tomography imaging of chordoma	Skeletal Radiol	33	524-530	2004
Semba J., Wakuta M., Suhara T.	Long-term suppression of methamphetamine-induced c-Fos expression in rat striatum by the injection of c-fos antisense oligodeoxynucleotides absorbed in water-absorbent polymer	Psychiatry Clin Neurosci	58	531-535	2004
Takano A., Suhara T.	The necessary parameters for estimating the time-course of receptor occupancy	Int J Neuropsychopharmacol	8	143-144	2004
Matsuda T, Matsuura M. , Ohkubo T, Ohkubo H, Matsushima E, Inoue K, Taira M, Kojima T	Functional MRI mapping of brain activation during visually guided saccades and antisaccades	Psychiatry Research: Neuroimaging	131	147-155	2004
Ohtsuki T, Tanaka S, Ishiguro H, Noguchi E, Arinami T, Tanabe E, Yara K, Ohkubo T, Takahashi S, Matsuura M , Sakai T, Muto M, Kojima T, Matsushima E, Toru M, Inada T	Failure to find association between PRODH deletion and schizophrenia	Schizophrenia Research	67	111-113	2004
Moriyama Y., Muramatsu T., Kato M. , Kashima H., Mimura M.	Repeated clinical episodes of Wernicke-Korsakoff syndrome	Australian and New Zealand Journal of Psychiatry	38	653	2004

Kato A., Kato M., Ishii H., Ichimiya Y., Suzuki K., Kawasaki H., Yamamoto S., Kumashiro R., Yamamoto K., Kawamura N., Hayashi N., Matsuzaki S., Terano A., Okita K., Watanabe A.	Development of quantitative neuropsychological tests for diagnosis of subclinical hepatic encephalopathy in liver cirrhosis patients and establishment of diagnostic criteria—multicenter collaborative study in Japanese	Hepatology Research	30	71-78	2004
Yahata N, Takahashi H, Okubo Y	Pharmacological modulations on the Human cognitive processes: an fMRI study	J Nippon Med Sch	72 (1)	2-3	2005
Zhang M.-R., Maeda J., Ito T., Ogawa M., Noguchi J., Suhara T. , Halldin C., Suzuki K.	Synthesis and evaluation of N-(5-fluoro-2-phenoxyphenyl)-N-(2-[¹⁸ F]fluoromethoxy-d(2)-5-methoxybenzyl)acetamide: a deuterium-substituted radioligand for peripheral benzodiazepine receptor	Bioorg Med Chem	13	1811-1818	2005
Yasuno F, Suhara T. , Okubo Y, Ichimiya T, Takano A, Inoue M, Sudo Y	Abnormal inter-regional relation of dopamine D2 receptor binding in schizophrenia	Psychiatry Res		in press	
Oda K, Matsushima E, Okubo Y, Ohta K, Murata Y, Koike R, Miyasaka N, Kato M	Abnormal Regional Cerebral Blood Flow in Systemic Lupus Erythematosus Patients with Psychiatric Symptoms	The Journal of Clinical Psychiatry		in press	
Matsuura M. , Adachi N, Muramatsu R, Kato M, Onuma T, Okubo Y. , Oana Y, Hara T	Intellectual Disability and Psychotic Disorders of Adult Epilepsy	Epilepsia		in press	
Aung W., Okauchi T., Sato M., Saito T., Nakagawa H., I shihara H., Ikota N., Suhara T. , Anzai K.	In vivo PET imaging of inducible D2R reporter transgene expression using [¹¹ C]FLB 457 as reporter probe in living rat	Nucl Med Com		in press	
Inaji M., Yoshizaki T., Okauchi T., Maeda J., Nagai Y., Okano H., Nariai T., Ohno K., Ando K., Obayashi S., Suhara T.	<i>In vivo</i> PET measurements with [¹¹ C]PE2I to evaluate fetal mesencephalic transplantations to unilateral 6-OHDA-lesioned rats	Cell Transplantation		in press	
Umeda S., Akine Y., Kato M. , Muramatsu T., Mimura M., Kandatsu S., Tanada S., Obata T., Ikehira H., Suhara T.	Functional network in the prefrontal cortex during episodic memory retrieval	NeuroImage		in press	
Takano A., Suhara T. , Kushumi I., Takahashi Y., Koyama T., Yasuno F., Ichimiya T., Inoue M., Sudo Y.	Time course of dopamine D2 receptor occupancy by clozapine in patients with schizophrenia	Prog. Neuro-Psychopharmacol. & Biol.		in press	

Mimura M., Komatsu S., Kato M., Yoshimasu H., Moriyama Y., Kashima H.	Further evidence for a memory advantage of self-performed tasks in person with alcoholic Korsakoff's syndrome	Journal of the International Neuropsychological Society		in press	
Mimura M., Watanabe R., Kato M., Kashima H.	Selective memory impairment for personally familiar colors following encephalitis	Psychiatry and Clinical Neuroscience		in press	
Fujinaga N., Muramastu T., Ogano M., Kato M.	A 3-year follow-up study of 'orientation agnosia'	Neuropsychologia		in press	
Umeda S., Nagumo Y., Kato M	Dissociative contributions of the medial temporal region and frontal cortex to prospective remembering	Reviews in the Neurosciences		in press	
Akiyama T., Kato M., Muramatsu T, Saito F, Nakachi R., Kashima H.	A deficit in discriminating gaze direction in a case with right superior temporal gyrus lesion	Neuropsychologia		in press	
成重竜一郎, 館野 周, 大久保善朗	非定型抗精神病薬投与中clonazepam減量に伴いParkinson症候群を呈した1症例	精神医学	46 (10)	1113-1115	2004
関根瑞保, 鈴木博子, 竹沢健司, 館野 周, 朝山健太郎, 大久保善朗	救命救急センターに搬送された自殺未遂症例の検討	総合病院精神医学	16 (3)	257-263	2004
伊藤敬雄, 葉田道雄, 木村美保, 黒川 顕, 黒澤 尚, 大久保善朗	高次救命救急センターに入院した自殺未遂患者とその追跡調査－精神科救急対応の現状を踏まえた1考察－	精神医学	46 (4)	389-396	2004
伊藤敬雄, 大久保善朗	アルツハイマー型痴呆患者におけるメラトニン療法	Geriatric Medicine	42 (9)	1201-1206	2004
小坂 淳, 須原哲也, 西條朋行, 高野晶寛, 安野史彦, 生駒洋子, 大久保善朗	PETによる気分障害患者の病態と治療法の作用機序に関する研究	INNERVISION	7	18	2004
大久保善朗, 伊藤逸生, 織田健司	精神疾患の脳形態画像	精神神経学雑誌	106 (7)	900-905	2004
須原哲也	画像から見た薬物治療の評価	精神神経学雑誌	106 (7)	916-920	2004
松浦雅人, 松田哲也, 大久保起延, 大久保博美, 根本安人, 松田玲子, 鹿中紀子, 小島卓也, 福本真衣, 松島英介, 泰羅雅登	統合失調症の眼球運動異常の機能的MRI	精神経誌	106 (7)	906-909	2004
根本安人, 松田哲也, 松浦雅人, 本下真衣, 大久保起延, 大久保博美, 鈴木正泰, 鹿中紀子, 松島英介, 小島卓也	探索眼球運動の神経機構	日大医誌	63 (7)	352-359	2004
松浦雅人	道路交通法改正後のてんかんをもつ人における運転免許, 精神科医からみた問題点と課題	てんかん研究	22 (1)	64-65	2004
松浦雅人	てんかんと抑うつ症状	ともしび	3	4-13	2004

松浦雅人	てんかんのメンタルヘルス, 診断をめぐって	波	9	279-280	2004
小島卓也, 大久保起延, 大久保博美, 鹿中紀子, 根本安人, 鈴木正泰, 松田哲也, 本下真衣, 松島英介, 松浦雅人	統合失調症の基本障害と眼球運動異常	脳精神医学	15 (4)	421-426	2004
加藤元一郎	失行の発現機序と責任病巣	臨床神経学	44	839-841	2004
春原則子, 宇野彰, 金子真人, 加藤元一郎, 吉野文浩	英語学習の困難さを主訴とした中学生・高校生の認知機能	神経心理学	20	264-271	2004
加藤元一郎, 青柳裕, 阿内謙吉, 小野寺晋志	アルツハイマー病に類似した症状を呈する疾患の脳機能画像統計解析	臨床放射線	49	1203-1210	2004
加藤元一郎, 梅田聡, 秋根良英	虚記憶に関する神経心理学的障害と機能的脳画像所見	精神神経学雑誌	106 (7)	910-915	2004
前田貴記, 加藤元一郎	統合失調症の認知機能障害と治療メカニズム - 自己意識の障害について (sense of agencyを中心に) -	Schizophrenia Frontier	5(4)	232-236	2004
鈴木博子, 成重竜一郎, 大久保善朗	Milnacipranが有効であった rapid cyclerの2症例	精神科治療学	20 (2)	203-210	2005
大久保善朗, 浅井邦彦	精神科病床と脱施設化への道	最新精神医学	10 (2)	143-150	2005
大久保善朗, 須原哲也	脳イメージングによる抗精神病薬の薬効評価	Human Science	16 (2)	22-25	2005
大久保善朗, 須原哲也	画像診断からみた薬物療法の評価	カレントセラピー	23 (1)	69-72	2005

書籍

著者氏名	論文タイトル名	書籍全体の 編集者名	書 籍 名	出版社名	出版地	出版 年	ペー ジ
Matsuura M, Fukumoto M, Matsushima M, Matsuda T, Ohkubo T, Ohkubo H, Nemoto Y, Kanaka N, Kojima T, Taira M	Functional MRI study on neural network dysfunction in schizophrenia and epileptic psychosis	Nakagawa M,Hirata K, Koga Y, Nagata K	Frontiers in Human Brain Topography	Elsevier	Amst erda m	2004	311- 314
有竹清夏, 大久保善朗	神経生理学的検査 (脳波・筋電図)	奈良信雄	臨床研修イラス トレイテッド. 3 基本手技	羊土社	東京	2004	159- 162
渡路子, 大久保善朗	不穏とうつ	奈良信雄	臨床研修イラス トレイテッド. 2 救急処置	羊土社	東京	2004	162- 164
松浦雅人	てんかんとうつ	てんかんの精 神症状と行動 研究会	てんかんーその 精神症状と行動	新興医学 出版社	東京	2004	42- 47
松浦雅人	てんかん精神病の 生物学的基礎	山内敏雄	てんかん学の 最前線10	日研化学	東京	2004	
松浦雅人・てんかんの精神症状と行動研究会			てんかんーその 精神症状と行動	新興医学 出版社	東京	2004	
松浦雅人	脳の機能異常, 損傷 および全身性疾患に よる精神障害		脳の病気のすべ てがわかる本	学研	東京	2004	
松浦雅人	解離性障害	井上新平	精神科・神経科 ナースの疾患別 ケアハンドブック	メディカ 出版	大阪	2004	

IV. 研究成果の刊行物・別刷

欧 文

An fMRI study of differential neural response to affective pictures in schizophrenia

Hidehiko Takahashi,^{a,b} Michihiko Koeda,^c Kenji Oda,^a Tetsuya Matsuda,^d Eisuke Matsushima,^d Masato Matsuura,^e Kunihiko Asai,^b and Yoshiro Okubo^{f,*}

^aSection of Psychiatry and Behavioral Science, Graduate School of Medicine, Tokyo Medical and Dental University, Japan

^bAsai Hospital, Japan

^cDepartment of Bioinformatics, Medical Research Institute, Tokyo Medical and Dental University, Japan

^dSection of Liaison Psychiatry and Palliative Medicine, Graduate School of Medicine, Tokyo Medical and Dental University, Japan

^eDepartment of Neuropsychiatry, School of Medicine, Nihon University, Tokyo, Japan

^fDepartment of Neuropsychiatry, Nippon Medical School, Tokyo 113-8603, Japan

Received 20 November 2003; revised 24 January 2004; accepted 8 March 2004

Although emotional dysfunction is considered a fundamental symptom of schizophrenia, studies investigating the neural basis of emotional dysfunction in schizophrenia are few. Using functional magnetic resonance imaging (fMRI) and a task viewing affective pictures, we aimed to examine automatic emotional response and to elucidate the neural basis of impaired emotional processing in schizophrenia. Fifteen healthy volunteers and 15 schizophrenics were studied. During the scans, the subjects were instructed to indicate how each of the presented pictures made them feel. Whole brain activities in response to the affective pictures were measured by fMRI. Controls recruited the neural circuit including amygdaloid–hippocampal region, prefrontal cortex, thalamus, basal ganglia, cerebellum, midbrain, and visual cortex while viewing unpleasant pictures. Despite an equal behavioral result to controls, the patients showed less activation in the components of the circuit (right amygdala, bilateral hippocampal region, medial prefrontal cortex (MPFC), basal ganglia, thalamus, cerebellum, midbrain, and visual cortex). This study demonstrated functional abnormalities in the neural circuit of emotional processing in schizophrenia. In particular, decreased activation in the right amygdala and MPFC appears to be an important finding related to dysfunctional emotional behavior in schizophrenia.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Schizophrenia; Emotion; Functional magnetic resonance imaging; Amygdala; Prefrontal cortex; Affective pictures

Introduction

Emotional dysfunction such as “flattening affect” or “anhedonia” is considered to be one of the key symptoms of schizophrenia

(Andreasen and Flaum, 1991). Recent functional imaging techniques have revealed dysfunction of the neural circuit (interrelationship among cortical region, thalamus, basal ganglia, and cerebellum) in schizophrenia (Andreasen et al., 1999; Schultz and Andreasen, 1999). Most studies focused on cognitive dysfunction (Andreasen et al., 1999), while studies investigating the neural basis of dysfunctional emotional processing in schizophrenia are limited (Gur et al., 2002; Paradiso et al., 2003; Phillips et al., 1999; Schneider et al., 1998; Taylor et al., 2002). Previous neuroimaging studies in normal subjects revealed the neuroanatomical correlates of emotional processing and the crucial role of the amygdala in processing negative emotions. In particular, left-sided activations in the amygdala while processing negative facial expressions have been consistently reported (Calder et al., 2001). Recent functional magnetic resonance imaging (fMRI) studies revealed that schizophrenic patients demonstrated decreased activation in the bilateral amygdala during an emotion induction task by facial expressions (Schneider et al., 1998) or decreased activation in the left amygdala during a discrimination task of emotional facial expressions (Gur et al., 2002). However, discriminating analogous facial expressions is an effortful cognitive process whether subjects categorize the facial expressions by gender or emotion. Cognitive demands for elaborate recognition or detailed rating may modulate the emotional response in the brain (Critchley et al., 2000; Hariri et al., 2000; Keightley et al., 2003; Lange et al., 2003; Taylor et al., 2003). In addition, it should be noted that emotional facial expressions do not necessarily elicit the subjective experience of emotions (Davidson and Irwin, 1999).

Several activation studies used affective pictures to elicit emotion in healthy volunteers (Lane et al., 1997a,b). The task of simply viewing emotionally salient pictures could minimize cognitive demands and would be suitable for examining automatic emotional response. Only a few positron emission tomography (PET) studies have investigated the emotional processing of affective pictures with a task of rating subjective emotional experience (Taylor et al., 2002) or with a task of emotional

* Corresponding author. Department of Neuropsychiatry, Nippon Medical School, 1-1-5, Sendagi, Bunkyo, Tokyo 113-8603, Japan. Fax: +81-3-5814-6280.

E-mail address: okubo-y@nms.ac.jp (Y. Okubo).

Available online on ScienceDirect (www.sciencedirect.com.)

perception (Paradiso et al., 2003) in schizophrenia, and these studies have not established consistent results for a better understanding of dysfunctional emotional processing in schizophrenia. To our knowledge, no fMRI study has examined the neural response across the whole brain to affective pictures in schizophrenia. In the present study, we used fMRI and a task with minimal cognitive demands to identify the neural circuit of automatic emotional processing. We expected the subjects to react to affective pictures without an effortful cognitive process and categorize them roughly according to their subjective emotional experiences. Comparing the neural responses of schizophrenic patients with those of healthy controls, we aimed to elucidate the neural basis of impaired emotional processing in schizophrenia.

Methods

Subjects

Fifteen schizophrenic patients (10 men and 5 women, mean age 29.0 years, SD = 6.9) meeting the DSM-IV criteria for schizophrenia were studied. Diagnoses were made by HT, YO, and the psychiatrists in charge based on a review of their charts and a conventionally semi-structured interview. Exclusion criteria were current or past substance abuse and a history of alcohol-related problems, mood disorder, or organic brain disease. Thirteen patients were recruited from the outpatient unit of Asai Hospital, and two were recruited from the inpatient unit. Eleven of the 15 patients received atypical neuroleptics (mean risperidone equivalent daily dosage = 1.60 mg, SD = 1.31), and the other four received no neuroleptics. The mean illness duration was 4.9 (SD = 4.9) years. Clinical symptoms were assessed by the Brief Psychiatric Rating Scale (BPRS) (Overall and Gorham, 1962). The mean score of BPRS was 17.9 (SD = 5.7). The ratings were reviewed by HT and YO after the patient interview, and disagreements were resolved by consensus; consensus ratings were used in this study. Fifteen normal controls (nine men and six women, mean age 29.1 years, SD = 7.8) were recruited from the surrounding community. The candidates were carefully screened and standardized interviews were conducted by trained psychiatrists (HT and YO). They did not meet criteria for any psychiatric disorder. None of the controls were taking alcohol and medication at the time, nor did they have a history of psychiatric disorder, significant physical illness, head injury, neurological disorder, or alcohol or drug dependence. Patients had a tendency of a lower educational status (patients 13.7 ± 1.5 years, controls 14.9 ± 2.4 years; $P = 0.11$, t test). All of the patients and controls were right-handed, and they all underwent an MRI to rule out cerebral anatomic abnormalities. After the procedures had been fully explained to the subjects, written informed consent was obtained, as approved by the Ethics Committee.

Stimulus and self-rating

Stimulus materials were taken from the International Affective Picture System (IAPS) (Lang et al., 1997). Pictures were divided into three emotional classes (neutral, unpleasant, and pleasant) according to the subjective ratings provided by IAPS. We employed 40 pictures from each class. The mean valence and arousal ratings of 40 pictures were 5.87 (SD = 0.77) and 3.75 (SD = 1.11) for

neutral, 3.07 (SD = 0.69) and 5.75 (SD = 0.83) for unpleasant, and 7.42 (SD = 0.36) and 4.65 (SD = 0.95) for pleasant pictures. Slides of the three emotional classes were matched for content (faces, human figures, animals, objects, and scenery). The pictures were projected via a computer and a telephoto lens onto a screen with a mirror mounted on a head-coil. The experimental design consisted of eight blocks for each of the three conditions (neutral, unpleasant, and pleasant) interleaved with 20-s rest periods. During the rest condition, subjects viewed a crosshair projected at the center of the screen. In each 20-s block, five different pictures of the same emotional class were presented for 3.5 s each, with an interstimulus interval of 0.5 s. During the scans, the subjects were instructed to indicate how each picture made them feel by categorizing their subjective emotions into three emotional classes (neutral, unpleasant, and pleasant) using buttons. Signals from the buttons were transmitted to a computer outside the shielded room via infrared rays to confirm whether expected emotions were evoked in response to individual affective pictures. The rate of the appearance of expected categorizations from the 40 pictures of the same emotional class was calculated for each emotional condition. We compared the percentages of expected categorizations between the patient and control groups.

fMRI acquisition

The images were acquired with a 1.5-T Signa system (General Electric, Milwaukee, WI). Functional images of 240 volumes were acquired with T2*-weighted gradient echo planar imaging sequences sensitive to blood oxygenation level-dependent (BOLD) contrast. Each volume consisted of 40 transaxial contiguous slices with a slice thickness of 3 mm to cover almost the whole brain (flip angle, 90°; TE, 50 ms; TR, 4 s; matrix, 64 × 64; field of view, 24 × 24).

Analysis of functional imaging data

Data analysis was performed with statistical parametric mapping software package (SPM99) (Wellcome Department of Cognitive Neurology, London, UK) that runs with MATLAB (Mathworks, Natick, MA). All volumes were realigned to the first volume of each session to correct for subject motion and were spatially normalized to the standard space defined by the Montreal Neurological Institute (MNI) template. After normalization, all scans had a resolution of $2 \times 2 \times 2$ mm³. Functional images were spatially smoothed with a 3D isotropic Gaussian kernel (full width at half maximum of 8 mm). Low frequency noise was removed by applying a high-pass filter (cut-off period = 240 s) to the fMRI time series at each voxel. A temporal smoothing function was applied to the fMRI time series to enhance the temporal signal-to-noise ratio. These images were scaled to give a grand mean signal of 100 across all voxels in all images to remove global effects. Significant hemodynamic changes for each condition were examined using the general linear model with boxcar functions convoluted with a hemodynamic response function. Statistical parametric maps for each contrast of the t statistic were calculated on a voxel-by-voxel basis. The t values were then transformed to unit normal distribution, resulting in z scores.

We assessed the neutral vs. rest, the unpleasant vs. rest, and the pleasant vs. rest contrasts. To assess the specific condition effect, we used the contrasts by subtracting the neutral condition from the pleasant condition and the unpleasant condition. A random effects

model, which estimates the error variance for each condition across the subjects, was implemented for group analysis. This procedure provides a better generalization to the population from which data are obtained. The contrast images were obtained from single-subject analysis and were entered into the group analysis. A one-

sample *t* test was applied to determine group activation for each effect. Significant clusters of activation were determined using the conjoint expected probability distribution of the height and extent of *z* scores with the height and extent threshold. In addition, we tested for relative differences in the pattern of neural activation by




Fig. 1. Images showing brain regions of significant activation during the neutral and unpleasant conditions relative to the rest condition in 15 controls and 15 schizophrenic patients. The bar shows the range of the *z* score. Within the image, L indicates left and R indicates right. Significant differences were accepted at a height threshold ($z > 4.75$; $P < 0.000001$, uncorrected) and extent threshold (30 voxels). Numbers in the bottom row indicate the *z* coordinates of the Montreal Neurological Institute brain.

Table 1
Brain regions of significant activation during unpleasant condition relative to neutral in 15 controls and 15 schizophrenic patients

	Brain region	Brodmann's area	Coordinates ^a			z score ^b	
			x	y	z		
Controls	Left lingual gyrus	17, 18, 19	-16	-88	-9	5.61	
	Right lingual gyrus	17, 18, 19	8	-82	-1	5.33	
	Left fusiform gyrus	19, 37	-32	-76	-8	5.8	
	Right fusiform gyrus	18, 37	22	-88	-11	4.95	
	Posterior cingulate	31	-2	-53	23	4.39	
	Left hippocampal region	34, 35	-18	-26	-12	5.31	
	Right hippocampal region	27, 28	24	-24	-9	5.55	
	Left amygdala		-24	-3	-18	4.29	
	Right amygdala		28	-1	-18	5.02	
	Left thalamus		-10	0	7	4.71	
	Right thalamus		18	-8	10	4.18	
	Left caudate nucleus		-14	10	11	4.7	
	Medial prefrontal cortex	9	-4	50	25	4.73	
	Right orbitofrontal cortex	47	32	27	-6	4.65	
	Cerebellum		32	-77	-18	6.18	
	Midbrain		-6	-33	-8	5.35	
	Schizophrenia	Left lingual gyrus	17, 18, 19	-18	91	3	4.68
		Right lingual gyrus	17	14	-90	6	4.07
		Left fusiform gyrus	19	-20	-80	-11	4.3
Right fusiform gyrus		19	24	-80	-11	4.33 ^c	
Left amygdala			-22	-6	-18	4.56 ^c	

^a Talairach and Tournoux coordinates in the local point of maximal activation included in the cluster.

^b Significant differences were accepted at a height threshold ($z > 3.89$; $P < 0.00005$, uncorrected) and extent threshold (30 voxels).

^c Right visual cortex and left amygdala in schizophrenia survived the height threshold but not the extent threshold.


subtracting the unpleasant minus neutral (U – N) contrasts of the patients from the U – N contrasts of the controls and vice versa. Between-group comparison was performed with a two-sample *t* test. Using the effect sizes, representing the percent signal change, of the U – N contrasts at the regional maxima uncovered in the between-group comparisons, we analyzed whether the BOLD signal change was correlated with the dosage of neuroleptics and the BPRS score. Coordinates of activation were converted from MNI coordinates to Talairach and Tournoux (1988) coordinates using the mni2tal algorithm (M. Brett, Cambridge, MA). Contrast images were overlaid onto a group mean anatomy image provided by SPM for viewing.

Results

Self-rating

The mean percentages of expected categorizations of the controls for the neutral, unpleasant, and pleasant pictures were 85.3% (SD = 8.3), 88.8% (SD = 11.3), and 59.0% (SD = 22.8), respectively, and those of the schizophrenic patients were 81.3% (SD = 8.3), 92.0% (SD = 9.5), and 55.0% (SD = 34.8), respectively. Two-way repeated-measures analysis of variance of the percentages of expected categorizations showed a significant main effect of condition ($F = 27.6$, $df = 2, 84$, $P < 0.001$), but not a significant main effect of group ($F = 0.26$, $df = 1, 84$, $P = 0.61$) or interaction ($F = 0.44$, $df = 2, 84$, $P = 0.64$). A post hoc test revealed that the percentage of expected category for the pleasant pictures was lower than those for the neutral and unpleasant pictures. That is, the subjects did not categorize the pleasant pictures as we had expected. Most of the remaining pictures not regarded as pleasant were categorized as neutral.

Controls



Schizophrenic patients




Fig. 2. Images showing brain regions of significant activation during unpleasant condition relative to neutral in 15 controls and 15 schizophrenic patients. The bar shows the range of the z score. Within the image, L indicates left and R indicates right. Significant differences were accepted at a height threshold ($z > 3.89$; $P < 0.00005$, uncorrected) and extent threshold (30 voxels). (Right visual cortex and left amygdala in the schizophrenic patients survived the height threshold but not the extent threshold.) Numbers in the bottom row indicate the z coordinates of the Montreal Neurological Institute brain.

fMRI data

Within-group comparisons

Both groups showed similar activation patterns in the neutral vs. rest contrast and in the pleasant vs. rest contrast (Fig. 1, top and bottom). Moreover, activation patterns in the neutral vs. rest contrast and the pleasant vs. rest contrast were also similar. In fact, significant activation in response to pleasant pictures relative to neutral pictures was seen only in the visual cortex (lingual gyrus) across groups (height threshold: $z > 4.75$ and extent threshold > 30 voxels). However, the two groups showed different activation patterns during the unpleasant condition compared to the rest condition (Fig. 1, middle). The controls demonstrated significant activation in response to unpleasant pictures relative to neutral pictures in the bilateral primary and secondary visual cortex, bilateral amygdala, bilateral hippocampal regions, medial prefrontal cortex (MPFC), right orbitofrontal cortex (OFC), bilateral thalamus, left caudate nucleus, cerebellum, and midbrain. Patients demonstrated significant activation in response to unpleasant pictures relative to neutral pictures in the bilateral primary and secondary visual cortex and left amygdala (Table 1 and Fig. 2).

Between-group comparisons

The group comparison of the U – N contrasts showed that schizophrenic patients demonstrated less activation in the right amygdala, bilateral hippocampal regions, MPFC, left visual cortex, left putamen, left caudate nucleus, left posterior thalamus, cerebellum, and midbrain (Table 2 and Fig. 3). No significantly greater activation was identified in schizophrenic patients in the between-group comparison of the U – N contrasts. We did not use between-group analysis for the pleasant minus neutral (P – N) contrast due to its meager activation across groups, possibly resulting from insufficient elicitation of pleasantness.

Correlations with BOLD signal change

There were no correlations between dosage of neuroleptics and signal change in the brain regions where patients showed decreased activation. In addition, no correlations were found between the BPRS score and signal change in these regions (Pearson's correlation analysis, $P > 0.05$).

Table 2
Brain regions with relatively less activation (unpleasant minus neutral) in 15 schizophrenic patients compared with 15 normal controls

Brain region	Brodmann's area	Coordinates ^a			z score ^b
		x	y	z	
Left lingual gyrus	18	-8	-88	-11	3.18
Left hippocampal region	30, 35	-14	-32	-12	3.83
Right hippocampal region	28	26	-24	-9	3.14
Right amygdala		24	-3	-13	3.39
Left thalamus		-22	-25	1	3.08
Left putamen		-20	12	9	2.88
Left caudate nucleus		-16	14	14	2.87
Medial prefrontal cortex	9	-2	54	25	3.21
Cerebellum		-14	-46	-23	3.06
Midbrain		-6	-26	-10	3.15

^a Talairach and Tournoux coordinates in the local point maximal activation included in the cluster.

^b Activation differences were considered significant at height threshold ($z > 2.57$; $P < 0.005$, uncorrected) and extent threshold (30 voxels).




Fig. 3. Images showing brain area of relatively less activation (unpleasant minus neutral) in 15 schizophrenic patients compared with 15 normal controls. The bar shows the range of the z score. Within the image, L indicates left and R indicates right. Activation differences were considered significant at height threshold ($z > 2.57$; $P < 0.005$, uncorrected) and extent threshold (30 voxels). Numbers in the bottom row indicate the z coordinates of the Montreal Neurological Institute brain.

Discussion

In control subjects, we identified the neural circuit of the automatic emotional response to unpleasant pictures in the amygdaloid–hippocampal region, thalamus, OFC, MPFC, basal ganglia, cerebellum, midbrain, and visual cortex. We used evocative pictures with minimal cognitive demands so as to examine the automatic emotional response that requires no elaborate rating or categorization of stimuli for the subjects. Facial expressions do not necessarily elicit strong emotions, and cognitive demands such as discriminating analogous facial expressions might affect brain activations (Critchley et al., 2000; Hariri et al., 2000; Keightley et al., 2003; Lange et al., 2003; Taylor et al., 2003). Passive emotional tasks with minimal cognitive demands might activate the amygdala and other subcortical regions more often than emotional tasks with greater cognitive demands (Phan et al., 2002). Thus, we successfully observed robust activation in widespread cortical and subcortical regions as reported in previous studies (Lane et al., 1997a,b).

The cortical–basal ganglia–thalamic circuit has been implicated in cognitive or emotional processing (Alexander et al., 1986). The circuit involving the components of the cortical–basal ganglia–thalamic circuit along with the amygdala appears to be involved in the control of emotional behavior (Groenewegen and Uylings, 2000; Price et al., 1996), and dysfunction of this circuit is considered to cause mood disorder (Drevets, 2001).

Despite similar categorizations of pictures for the controls, patients demonstrated less activation in the amygdaloid–hippocampal region, MPFC, thalamus, basal ganglia, cerebellum, midbrain, and visual cortex. This finding represented evidence of functional abnormalities in the neural circuit involving the cortical–basal ganglia–thalamic circuit and the amygdala in schizophrenia. Our patients showed decreased cerebellar activation as well. Considering the fact that neuroimaging studies of schizophrenia using a variety of cognitive tasks have demonstrated a disruption in the cortical–cerebellar–thalamic–cortical circuit

(CCTCC) leading to cognitive deficits (Andreasen et al., 1999), the functional abnormalities in the CCTCC along with the limbic area might lead to the emotional dysfunction in schizophrenia. Our results could also be interpreted in this manner, supporting the notion that schizophrenic patients have disruption in the distributed neural circuit, although cortical regions vary depending on the task (Andreasen et al., 1999).

The amygdala and PFC are considered to be key nodes of the neural circuit of emotional processing, the former a main signal generator and the latter a modulator (Davidson, 2002; Drevets, 2000). Previous fMRI studies using facial expressions showed decreased activation in the bilateral amygdala (Schneider et al., 1998) or left amygdala (Gur et al., 2002) in schizophrenia. However, our patients showed significantly less activation in the right amygdala. These inconsistent findings might be due to differences in the emotional tasks. The left amygdala activation has been consistently reported in the processing of emotional facial expressions (Calder et al., 2001), and the right amygdala has been suggested to have a higher affinity with picture processing (Keightley et al., 2003; Markowitsch, 1998). It has also been suggested that cognitive or attention-demanding aspects of the emotional task could attenuate amygdala activation (Critchley et al., 2000; Hariri et al., 2000; Keightley et al., 2003; Taylor et al., 2003). Using affective pictures and minimizing cognitive demands, we demonstrated robust activation in the right amygdala as well as in the left amygdala in controls. This result concerning right amygdala activation could be attributed to group differences. Another interpretation of patients showing less activation in the right amygdala might be possible. Several lines of evidence have suggested the functional laterality of the amygdala, that is, the right amygdala may engage in a rapid automatic processing of ambiguous information, while the left amygdala may participate in conscious evaluation of significant stimuli (Critchley et al., 2000; Markowitsch, 1998; Morris et al., 1998, 1999; Phelps et al., 2001; Wright et al., 2001). Taking this into account, schizophrenic patients might have relatively intact function of conscious processing of significant information, leading to a categorization of pictures similar to that of controls, but impairment of the rapid, automatic processing of salient stimuli. In other words, patients could assign significance to stimuli through conscious processing, but they might have diminished automatic emotional response to external stimuli.

A PET study using IAPS reported that schizophrenic patients showed decreased activation in the right amygdala in response to non-aversive pictures (Taylor et al., 2002). In that study, non-aversive pictures elicited robust activation in the bilateral amygdala and aversive pictures failed to elicit greater activation in the amygdala relative to non-aversive pictures. In other words, their non-aversive pictures were not “neutral” and they might have contained emotionally salient features. In this regard, our result is consistent with this previous finding. By contrast, another recent PET study using IAPS reported decreased activation in the left amygdala in schizophrenia (Paradiso et al., 2003). Unfortunately, the study did not set up a neutral condition. Without such a condition, it remains unclear whether the decreased activation in response to unpleasant pictures stems from impairment in emotional processing of unpleasant pictures or in a more basic cognitive function such as visual perception or object recognition. We ruled out the latter possibility by comparing the activation in response to neutral stimuli across groups. Obviously, more research is needed on the abnormal amygdala function in schizophrenia.

Within the PFC, another key node of the neural circuit of emotional processing, we found decreased activation in the MPFC in patients. The MPFC was commonly activated in studies about emotional response in healthy subjects, and its activation was not specific to particular emotion or induction methods with or without cognitive components (Phan et al., 2002). The MPFC is assumed to play general roles in emotional processing such as attention to emotion, identification or regulation of emotion (Reiman et al., 1997; Teasdale et al., 1999), and guiding motivational behavior by modulating or appraising autonomic emotional responses (Drevets, 2001; Epstein et al., 1999; Phillips et al., 2003). The decreased activation in the MPFC in our patients appears to be an important finding with respect to abnormal motivational behavior in schizophrenia. In contrast, the PET study using IAPS showed hyperactivation in the MPFC in schizophrenia, contrary to the author's expectation (Taylor et al., 2002). These contradictory results therefore emphasize the need for further studies of the activation of the MPFC in schizophrenia.

The present study has several limitations. First, most of the patients were taking neuroleptic medications, possibly affecting neural activation. They were, however, taking atypical neuroleptics, and at relatively low doses. To our knowledge, there has been no previous study on the effect of neuroleptics on the BOLD response of emotional processing. Compared to typical neuroleptics, atypical neuroleptics have shown less influence on BOLD contrast in the motor area or thalamus during a finger-tapping task (Braus et al., 1999; Muller and Klein, 2000). Future studies with neuroleptic-naïve patients, where the effect of neuroleptics can be controlled, will clarify this possible limitation. Second, we could not demonstrate any correlation between signal changes and BPRS scores in patients, possibly due to a lack of dispersion in the psychopathology of the patients, most of them being non-deficit outpatients with mild psychiatric symptoms. Third, the unpleasant pictures contained emotional features ranging from fear to disgust, and we could not differentiate the processing of particular emotions. In the processing of fear, the amygdala plays a central role. In contrast, the basal ganglia rather than the amygdala is considered to be essential in the processing of disgust (Calder et al., 2001; Phan et al., 2002). Thus, activation in the components of the neural circuit, the amygdala and basal ganglia, might reflect both emotional processing. Finally, we have difficulties in measuring emotional behaviors. This point has implications for the interpretation of the discrepancy between normal behavioral result and the abnormal neural activations observed in schizophrenic patients, as was also reported in previous studies (Gur et al., 2002; Paradiso et al., 2003; Schneider et al., 1998; Taylor et al., 2002). Our task could be regarded as an emotion-induction task. However, the finding needs to be interpreted cautiously because, strictly speaking, our task was testing the access to autothetic perception of elicited emotions. It might be possible that the ability of schizophrenic patients to access their emotions (categorization of feeling) was different from that of normal controls. Our behavioral results might not necessarily reflect gut-level elicited emotion that drives emotional behavior. Autonomic data such as skin conductance responses would help to measure gut-level emotional response.

In conclusion, we investigated the automatic emotional response in healthy controls and schizophrenic patients. By using a task with minimal cognitive demands, we identified robust activation across the neural circuit of emotional processing including the amygdaloid–hippocampal region, prefrontal cortex, thalamus, and basal ganglia in response to unpleasant stimuli in the controls.

Schizophrenic patients demonstrated less activation in the components of the circuit. In particular, decreased activation in the right amygdala and MPFC, the key structures in the circuit, could be related, respectively, to diminished automatic emotional response to external stimuli and impairment in regulating emotional responses to guide emotional behavior in schizophrenia.

Acknowledgments

We thank Professor Toru Nishikawa of the Section of Psychiatry and Behavioral Science, Graduate School of Tokyo Medical and Dental University. The staffs of the Section of Biofunctional Informatics, Graduate School of Tokyo Medical and Dental University, and of Asai Hospital, are gratefully acknowledged. This work was supported by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science and Technology (15390348), a research grant for nervous and mental disorders (14B-3), and a Health and Labor Sciences Research Grant for Research on Psychiatric and Neurological Diseases and Mental Health (H15-K0K0R0-003) from the Japanese Ministry of Health, Labor and Welfare.

References

- Alexander, G.E., DeLong, M.R., Strick, P.L., 1986. Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu. Rev. Neurosci.* 9, 357–381.
- Andreasen, N.C., Flaum, M., 1991. Schizophrenia: the characteristic symptoms. *Schizophr. Bull.* 17, 27–49.
- Andreasen, N.C., Nopoulos, P., O'Leary, D.S., Miller, D.D., Wassink, T., Flaum, M., 1999. Defining the phenotype of schizophrenia: cognitive dysmetria and its neural mechanisms. *Biol. Psychiatry* 46, 908–920.
- Braus, D.F., Ende, G., Weber-Fahr, W., Sartorius, A., Krier, A., Hubrich-Ungureanu, P., Ruf, M., Stuck, S., Henn, F.A., 1999. Antipsychotic drug effects on motor activation measured by functional magnetic resonance imaging in schizophrenic patients. *Schizophr. Res.* 39, 19–29.
- Calder, A.J., Lawrence, A.D., Young, A.W., 2001. Neuropsychology of fear and loathing. *Nat. Rev. Neurosci.* 2, 352–363.
- Critchley, H., Daly, E., Phillips, M., Brammer, M., Bullmore, E., Williams, S., Van Amelsvoort, T., Robertson, D., David, A., Murphy, D., 2000. Explicit and implicit neural mechanisms for processing of social information from facial expressions: a functional magnetic resonance imaging study. *Hum. Brain Mapp.* 9, 93–105.
- Davidson, R.J., 2002. Anxiety and affective style: role of prefrontal cortex and amygdala. *Biol. Psychiatry* 51, 68–80.
- Davidson, J.R., Irwin, W., 1999. The functional neuroanatomy of emotion and affective style. *Trends Cogn. Sci.* 3, 11–21.
- Drevets, W.C., 2000. Neuroimaging studies of mood disorders. *Biol. Psychiatry* 48, 813–829.
- Drevets, W.C., 2001. Neuroimaging and neuropathological studies of depression: implication for cognitive-emotional features of mood disorders. *Curr. Opin. Neurobiol.* 11, 240–249.
- Epstein, J., Stern, E., Silbersweig, D., 1999. Mesolimbic activity associated with psychosis in schizophrenia. Symptom-specific PET studies. *Ann. N. Y. Acad. Sci.* 877, 562–574.
- Groenewegen, H.J., Uylings, H.B., 2000. The prefrontal cortex and the integration of sensory, limbic and autonomic information. *Prog. Brain Res.* 126, 3–28.
- Gur, R.E., McGrath, C., Chan, R.M., Schroeder, L., Turner, T., Turetsky, B.L., Kohler, C., Alsop, D., Maldjian, J., Ragland, J.D., Gur, R.C., 2002. An fMRI study of facial emotion processing in patients with schizophrenia. *Am. J. Psychiatry* 159, 1992–1999.
- Hari, A.R., Bookheimer, S.Y., Mazziotta, J.C., 2000. Modulating emotional responses: effects of a neocortical network on the limbic system. *NeuroReport* 11, 43–48.
- Keightley, M.L., Winocur, G., Graham, S.J., Mayberg, H.S., Hevenor, S.J., Grady, C.L., 2003. An fMRI study investigating cognitive modulation of brain regions associated with emotional processing of visual stimuli. *Neuropsychologia* 41, 585–596.
- Lane, R.D., Reiman, E.M., Ahern, G.L., Schwartz, G.E., Davidson, R.J., 1997a. Neuroanatomical correlates of happiness, sadness, and disgust. *Am. J. Psychiatry* 154, 926–933.
- Lane, R.D., Reiman, E.M., Bradley, M.M., Lang, P.J., Ahern, G.L., Davidson, R.J., Schwartz, G.E., 1997b. Neuroanatomical correlates of pleasant and unpleasant emotion. *Neuropsychologia* 35, 1437–1444.
- Lang, P.J., Bradley, M.M., Cuthbert, B.N., 1997. International Affective Picture System (IAPS): Technical Manual and Affective Ratings University of Florida, Center for Research in Psychophysiology, Gainesville.
- Lange, K., Williams, L.M., Young, A.W., Bullmore, E.T., Brammer, M.J., Williams, S.C., Gray, J.A., Phillips, M.L., 2003. Task instructions modulate neural responses to fearful facial expressions. *Biol. Psychiatry* 53, 226–232.
- Markowitsch, H.J., 1998. Differential contribution of right and left amygdala to affective information processing. *Behav. Neurol.* 11, 233–244.
- Morris, J.S., Ohman, A., Dolan, R.J., 1998. Conscious and unconscious emotional learning in the human amygdala. *Nature* 393, 467–470.
- Morris, J.S., Ohman, A., Dolan, R.J., 1999. A subcortical pathway to the right amygdala mediating “unseen” fear. *Proc. Natl. Acad. Sci. U. S. A.* 96, 1680–1685.
- Muller, J.L., Klein, H.E., 2000. Neuroleptic therapy influences basal ganglia activation: a functional magnetic resonance imaging study comparing controls to haloperidol- and olanzapine-treated inpatients. *Psychiatry Clin. Neurosci.* 54, 653–658.
- Overall, J.E., Gorham, D.R., 1962. The brief psychiatric rating scale. *Psychol. Rep.* 10, 799–812.
- Paradiso, S., Andreasen, N.C., Creso-Facorro, B., O'Leary, D.S., Watkins, G.L., Boles Ponto, L.L., Hichwa, R.D., 2003. Emotions in unmedicated patients with schizophrenia during evaluation with positron emission tomography. *Am. J. Psychiatry* 160, 1775–1783.
- Phan, K.L., Wager, T., Taylor, S.F., Liberzon, I., 2002. Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *NeuroImage* 16, 331–348.
- Phelps, E.A., O'Connor, K.J., Gatenby, J.C., Gore, J.C., Grillon, C., Davis, M., 2001. Activation of the left amygdala to a cognitive representation of fear. *Nat. Neurosci.* 4, 437–441.
- Phillips, M.L., Williams, L., Senior, C., Bullmore, E.T., Brammer, M.J., Andrew, C., Williams, S.C., David, A.S., 1999. A differential neural response to threatening and non-threatening negative facial expressions in paranoid and non-paranoid schizophrenics. *Psychiatry Res.* 92, 11–31.
- Phillips, M.L., Drevets, W.C., Rauch, S.L., Lane, R., 2003. Neurobiology of emotion perception I: the neural basis of normal emotion perception. *Biol. Psychiatry* 54, 504–514.
- Price, J.L., Carmichael, S.T., Drevets, W.C., 1996. Networks related to the orbital and medial prefrontal cortex; a substrate for emotional behavior? *Prog. Brain Res.* 107, 523–536.
- Reiman, E.M., Lane, R.D., Ahern, G.L., Schwartz, G.E., Davidson, R.J., Friston, K.J., Yun, L.S., Chen, K., 1997. Neuroanatomical correlates of externally and internally generated human emotion. *Am. J. Psychiatry* 154, 918–925.
- Schneider, F., Weiss, U., Kessler, C., Salloum, J.B., Posse, S., Grodd, W., Muller-Gartner, H.W., 1998. Differential amygdala activation in schizophrenia during sadness. *Schizophr. Res.* 34, 133–142.
- Schultz, S.K., Andreasen, N.C., 1999. Schizophrenia. *Lancet* 353, 1425–1430.
- Talairach, J., Tournoux, P., 1988. *Co-Planar Stereotaxic Atlas of the Human Brain: Three-Dimensional Proportional System* Thieme Medical, New York.
- Taylor, S.F., Liberzon, I., Decker, L.R., Koeppe, R.A., 2002. A func-

- tional anatomic study of emotion in schizophrenia. *Schizophr. Res.* 58, 159–172.
- Taylor, S.F., Phan, K.L., Decker, L.R., Liberzon, I., 2003. Subjective rating of emotionally salient stimuli modulates neural activity. *NeuroImage* 18, 650–659.
- Teasdale, J.D., Howard, R.J., Cox, S.G., Ha, Y., Brammer, M.J., Williams, S.C., Checkley, S.A., 1999. Functional MRI study of the cognitive generation of affect. *Am. J. Psychiatry* 156, 209–215.
- Wright, C.I., Fischer, H., Whalen, P.J., McInerney, S.C., Shin, L.M., Rauch, S.L., 2001. Differential prefrontal cortex and amygdala habituation to repeatedly presented emotional stimuli. *NeuroReport* 12, 379–383.

Brain activation associated with evaluative processes of guilt and embarrassment: an fMRI study

Hidehiko Takahashi,^{a,b} Noriaki Yahata,^c Michihiko Koeda,^d Tetsuya Matsuda,^e
Kunihiko Asai,^b and Yoshiro Okubo^{c,*}

^aSection of Psychiatry and Behavioral Science, Graduate School of Medicine, Tokyo Medical and Dental University, Japan

^bAsai Hospital, Japan

^cDepartment of Neuropsychiatry, Nippon Medical School, Japan

^dDepartment of Bioinformatics, Medical Research Institute, Tokyo Medical and Dental University, Japan

^eBrain Activity Imaging Center, Tamagawa University Research Institute, Japan

Received 21 January 2004; revised 5 June 2004; accepted 19 July 2004
Available online 6 October 2004

We aimed to investigate the neural substrates associated with evaluative process of moral emotions. Using functional magnetic resonance imaging (fMRI), we examined the similarities and differences between evaluative process of guilt and that of embarrassment at the neural basis level. Study of the neural basis of judgments of moral emotions might contribute to a better understanding of the amoral behavior observed in neurological and psychiatric disorders. Nineteen healthy volunteers were studied. The participants read sentences carrying neutral, guilty, or embarrassing contents during the scans. Both guilt and embarrassment conditions commonly activated the medial prefrontal cortex (MPFC), left posterior superior temporal sulcus (STS), and visual cortex. Compared to guilt condition, embarrassment condition produced greater activation in the right temporal cortex (anterior), bilateral hippocampus, and visual cortex. Most of these regions have been implicated in the neural substrate of social cognition or Theory of Mind (ToM). Our results support the idea that both are self-conscious emotions, which are social emotions requiring the ability to represent the mental states of others. At the same time, our functional fMRI data are in favor of the notion that evaluative process of embarrassment might be a more complex process than that of guilt.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Guilt; Embarrassment; Social cognition; Theory of Mind; Medial prefrontal cortex; Superior temporal sulcus

Introduction

Although there have been numerous neuroimaging studies on primary emotions (fear, disgust, happiness, and sadness) that have led to a better understanding of the neuroanatomical correlates of

emotions (Phan et al., 2002), only a few studies on complex social emotions such as guilt, shame, and embarrassment have been reported. These social emotions have been viewed as moral emotions because they occur in response to moral violation and promote moral behavior, interpersonal etiquette, and personal hygiene (Eisenberg, 2000; Haidt, 2003). At the same time, these emotions inhibit transgression of social standards and motivate reparative action such as apology, confession, and atonement.

Impairment of possessing the mental states of these moral emotions could lead to amoral, inappropriate behaviors observed in neurological and psychiatric disorders such as brain injuries (Anderson et al., 1999; Beer et al., 2003), frontotemporal dementia (Miller et al., 2003; Snowden et al., 2002), autism (Capps et al., 1992; Frith, 2001; Hillier and Allinson, 2002), and antisocial personality (Brower and Price, 2001; Moll et al., 2003). Studying the neural substrates of judgments of moral emotions should add to the understanding of the neural basis of amoral behaviors observed in neurological and psychiatric disorders.

From a psychological point of view, guilt, shame, embarrassment, and pride are categorized into the same emotion family, “self-conscious emotions”. “Self-conscious emotions” are emotions founded in social relationship and arise from concerns about others’ opinions of self or the behavior of self (Eisenberg, 2000; Haidt, 2003; Tangney and Dearing, 2002). Negative evaluation of self or the behavior of self is fundamental to guilt, shame, and embarrassment, while positive evaluation of self leads to pride. In other words, one needs the ability to represent the mental states of others (intention/emotion), that is, Theory of Mind (ToM), to recognize self-conscious emotions. The recognition of negative self-conscious emotions involves understanding of the violation of social norms and the negative evaluation of self, both important aspects of ToM. Children with autism demonstrating impaired ToM showed impaired recognition of self-conscious emotions (Heerey et al., 2003). In line with this notion, a recent functional magnetic resonance imaging (fMRI) study demonstrated activation in the medial prefrontal cortex (MPFC), temporal regions, and orbito-

* Corresponding author. Department of Neuropsychiatry, Nippon Medical School, 1-1-5, Sendagi, Bunkyo-ku, Tokyo, 113-8603 Japan. Fax: +81 3 5814 6280.

E-mail address: okubo-y@nms.ac.jp (Y. Okubo).

Available online on ScienceDirect (www.sciencedirect.com.)

frontal cortex (OFC) during the emotional judgments of embarrassment (Berthoz et al., 2002). These areas have been implicated in ToM, social cognition, and moral judgment (Adolphs, 2001; Allison et al., 2000; Frith, 2001; Frith and Frith, 1999; Greene and Haidt, 2002; Greene et al., 2001; Moll et al., 2003; Pinkham et al., 2003). However, a previous positron emission tomography (PET) study using a guilt-related script reported a slightly different activation pattern in anterior paralimbic regions during the experience of guilt (Shin et al., 2000). In the former fMRI study, an emotional judgment task was used. Participants read various kinds of stories depicting embarrassing situations and were instructed to imagine what the story protagonist (participant himself or a third person in the story) would feel. In the latter PET study, an emotion induction method was used. Participants listened to audio-taped personal events involving the most guilt they had actually experienced. They recalled and imagined the event as if they were actually participating in it again. These two studies differed in the emotional tasks and measurement methods, making it difficult to compare the results directly. To our knowledge, no neuroimaging study has as yet investigated the different types of self-conscious emotions and compared the neural activation patterns directly in one session.

Although the distinctions among guilt, shame, and embarrassment are not clear-cut, psychologists have challenged this issue. Although embarrassment has traditionally been considered to be a variant of shame (Lewis, 1993), recent psychological data support the notion that embarrassment is an emotion distinct from other self-conscious emotions (Keltner and Buswell, 1997). Embarrassment has higher affinity to violation of social conventions, while guilt and shame have higher affinity to violation of a moral norm (Eisenberg, 2000; Haidt, 2003; Tangney et al., 1996). In this sense, among the negative self-conscious emotions, distinction between guilt and embarrassment is considered to be relatively clear-cut. Therefore, we focused on these two emotions.

We used block-design fMRI to measure regional activation associated with judgments of guilt and embarrassment during an emotional judgment task presenting short sentences. Emotion processing is composed of evaluative, experiential, and expressive components. We did not intend to induce emotional states because we thought it would be difficult to induce emotional states of guilt or embarrassment by merely having the subjects read short sentences. Moreover, it would be difficult to control the situation so as not to induce emotions other than guilt and embarrassment (e.g., anger, shame, sadness) as reported in the previous induction study (Shin et al., 2000). We aimed to elucidate the similarities and differences between the evaluative process of guilt and that of embarrassment at the neural basis level by measuring neural activation during judgments of both emotions in a session using fMRI.

We hypothesized that both emotional conditions would commonly activate the components of the neural substrates (MPFC, superior temporal sulcus (STS)) that have been implicated in social cognition (Adolphs, 2001; Allison et al., 2000) or ToM (Frith, 2001; Frith and Frith, 1999), and at the same time, would show differences in the extent of activation of the components.

Method

Participants

Nineteen healthy right-handed Japanese subjects (10 men, mean age 30.8 years, SD = 6.2; nine women, mean age 25.1 years, SD =

3.2) were recruited from the surrounding community. Their mean educational achievement level was 16.2 years (SD = 2.1). They did not meet criteria for any psychiatric disorder. None of the controls were taking alcohol or medication at the time, nor did they have a history of psychiatric disorder, significant physical illness, head injury, neurological disorder, or alcohol or drug dependence. All subjects underwent an MRI to rule out cerebral anatomic abnormalities. After complete explanation of the study, written informed consent was obtained from all subjects, and the study was approved by the Ethics Committee.

Materials

Because our experimental design was a block design, we aimed to control readability, the number of words, and luminance across blocks using short sentences. In addition, we expected participants to make emotional judgments repeatedly in a block. For this reason, we used short sentences instead of other forms, for example, stories. Three types of short sentences were provided (neutral, guilt, and embarrassment). Each sentence was written in Japanese and in the first person, past tense. Each sentence was expected to carry guilt, embarrassment, or no prominent emotional content. To validate our expected results, other healthy volunteers (10 men and 10 women, mean age 28.6 years, SD = 3.7) than the subjects participating in this fMRI study were screened. They read each sentence and rated the described situations according to how guilty or embarrassing they seemed using a 6-point analog scale (1 = none, 6 = extremely intense). As we predicted, the mean ratings of guilt and embarrassment for neutral sentences were 1.0 (SD = 0.1) and 1.0 (SD = 0.1), for guilt-related sentences 4.4 (SD = 0.4) and 1.6 (SD = 0.4), and for embarrassing sentences 1.5 (SD = 0.3) and 3.5 (SD = 0.5), respectively. Examples of the sentences are shown in Table 1. The sentences were projected via a computer and a telephoto lens onto a screen mounted on a head coil. The subjects were instructed to read the sentences silently and were told that they would rate the described situations according to how guilty or embarrassing they seemed. After reading each sentence, the subjects were instructed to press a selection button with the right index finger, indicating that they had read and understood it. The experimental design consisted of six blocks for each of the three conditions (neutral, guilt, and embarrassment) interleaved with 20-s rest periods. The order of presentation for the three conditions was fixed in the neutral–guilt–embarrassment sequence (Fig. 1). During the rest condition,

Table 1
Examples of sentences

Neutral	I used a cellular phone in the park.
	I used a computer on the internet.
	I change into pajamas at night.
	I washed my clothes.
Guilt	I had dinner at the restaurant.
	I used a cellular phone in the hospital.
	I sent a computer virus by e-mail.
	I shoplifted a dress from the store.
Embarrassment	I betrayed my friend.
	I left the restaurant without paying.
	I was not dressed properly for the occasion.
	I mistook a stranger for my friend.
	I noticed that the zipper of my pants was open.
	I soiled my underwear.
I did not know the right behavior at the restaurant.	