

# Association of Mitochondrial Complex I Subunit Gene *NDUFV2* at 18p11 with Bipolar Disorder in Japanese and the National Institute of Mental Health Pedigrees

Shinsuke Washizuka, Kazuya Iwamoto, An-a Kazuno, Chihiro Kakiuchi, Kanako Mori, Mizue Kametani, Kazuo Yamada, Hiroshi Kunugi, Osamu Tajima, Tsuyoshi Akiyama, Shinichiro Nanko, Takeo Yoshikawa, and Tadafumi Kato

**Background:** Linkage with 18p11 is one of the replicated findings in molecular genetics of bipolar disorder. Because mitochondrial dysfunction has been suggested in bipolar disorder, *NDUFV2* at 18p11, encoding a subunit of the complex I, reduced nicotinamide adenine dinucleotide (NADH) ubiquinone oxidoreductase, is a candidate gene for this disorder. We previously reported that a polymorphism in the upstream region of *NDUFV2*,  $-602G > A$ , was associated with bipolar disorder in Japanese subjects; however, functional significance of  $-602G > A$  was not known.

**Methods:** We screened the further upstream region of *NDUFV2*. We performed a case-control study in Japanese patients with bipolar disorder and control subjects and a transmission disequilibrium test in 104 parent and proband trios of the National Institute of Mental Health (NIMH) Genetics Initiative pedigrees. We also performed the promoter assay to examine functional consequence of the  $-602G > A$  polymorphism.

**Results:** The  $-602G > A$  polymorphism was found to alter the promoter activity. We found that the other haplotype block surrounding  $-3542G > A$  was associated with bipolar disorder. The association of the haplotypes consisting of  $-602G > A$  and  $-3542G > A$  polymorphisms with bipolar disorder was seen both in Japanese case-control samples and NIMH trios.

**Conclusion:** Together these findings indicate that the polymorphisms in the promoter region of *NDUFV2* are a genetic risk factor for bipolar disorder by affecting promoter activity.

**Key Words:** Bipolar disorder, haplotype, mitochondria, NADH ubiquinone oxidoreductase, promoter assay, transmission disequilibrium test

The etiology of bipolar disorder (BD) is still unknown, but family, twin, and adoption studies strongly suggest the involvement of genetic risk factors (Goochwin and Jamison 1990). Linkage studies have revealed a number of loci to be linked with BD. Of those, several investigators confirmed 18p11 as one susceptibility loci for BD (Benetini et al 1997; Gershon et al 1996; Nothen et al 1999; Stine et al 1995; Turecki et al 1999). Nominally significant linkage of BD with chromosome 18 was also found in a recent extensive meta-analysis (Segurado et al 2003). Thus, 18p is one of the targets of the genetic association study of BD.

We have proposed a mitochondrial dysfunction hypothesis of BD (Kato and Kato 2000) on the basis of the following evidence: altered brain energy metabolism in patients with BD detected by phosphorus-31 magnetic resonance spectroscopy (Kato et al 1993), increased ratio of the mitochondrial DNA (mtDNA) deletion in the brains of patients with BD (Kato et al 1997), association with mtDNA polymorphisms causing amino acid substitutions in the subunits of complex I (reduced nicotinamide adenine dinucleotide [NADH] ubiquinone oxidoreductase; Kato et al 2001).

From the Laboratories for Molecular Dynamics of Mental Disorders (SW, KI, AK, CK, KM, MK, TK) and Molecular Psychiatry (KY, TY), Brain Science Institute, RIKEN, Wako, Saitama; Department of Mental Disorder Research (HK), National Institute of Neuroscience; Kyorin University School of Health Sciences (OT); Department of Neuropsychiatry (TA), NTT East Kanto Medical Center; and Department of Psychiatry and Genome Research Center (SN), Teikyo University School of Medicine, Tokyo, Japan. Address reprint requests to Tadafumi Kato, M.D., Ph.D., Laboratory for Molecular Dynamics of Mental Disorders, Brain Science Institute, RIKEN, Hirosawa 2-1, Wako, Saitama, 351-0198, Japan; E-mail: kato@brain.riken.go.jp.

Received March 9, 2004; revised June 22, 2004; accepted July 5, 2004.

0006-3223/04/\$30.00

doi:10.1016/j.biopsych.2004.07.004

Complex I catalyzes the transfer of electrons from NADH to ubiquinone and the largest and most complicated enzyme in the mitochondrial electron transport chain, consisting of at least 43 subunits. Whereas seven subunits of complex I are coded in the mtDNA, the others are coded in the nuclear genome (Smeitink et al 2001). Of those, *NDUFV2* is located at 18p11 (de Coe et al 1995; Hattori et al 1995) and is a candidate gene for BD. Recently, Nakatani et al (2004) examined the gene expression patterns in the frontal cortex and hippocampus in animal models of depression and reported that *NDUFV2* was one of two genes altered in both regions. Moreover, Karry et al (2004) reported that protein levels of 24kDa subunit of complex I encoded by *NDUFV2* were altered in the autopsied brains of BD patients. These findings suggested a possible role of *NDUFV2* in mood disorders.

We previously screened mutations and polymorphisms in all exons and the 1-kb upstream region of *NDUFV2* in BD patients and reported that a polymorphism,  $-602G > A$ , in the upstream region was significantly associated with BD in Japanese (Washizuka et al 2003). The mRNA expression of *NDUFV2* was also significantly decreased in the lymphoblastoid cells of patients with bipolar I disorder.

In this study, we further screened the 4kb-upstream region of the *NDUFV2* and examined the association with BD in a Japanese case-control samples. Furthermore, we performed a promoter assay to examine the functional significance of the  $-602G > A$  polymorphism, which determines the major haplotypes associated with BD. We then examined whether a similar association was found in the National Institute of Mental Health (NIMH) Initiative Genetics Bipolar Pedigrees by the haplotype transmission disequilibrium test (TDT).

## Methods and Materials

### Japanese Case-Control Samples

The subjects with BD were 189 unrelated patients (117 women and 72 men, 136 with bipolar I disorder (BDI) and 53

with bipolar II disorder (BDII; 49.8 ± 13.8 years) who were followed at the hospitals or clinics participating in this study. Their age at onset was 34.9 ± 13.4 years. Consensus diagnosis by at least two senior psychiatrists according to the DSM-IV criteria was made for each patient using a nonstructured interview and by reviewing medical records. The 222 unrelated control subjects (117 women and 105 men, 30.2 ± 8.3 years old) were recruited from hospital staff and students. Control subjects were not assessed for psychiatric symptoms by any structured interview method, but they showed good social functioning and reported themselves to be in good health. All the subjects were Japanese with characteristics as described previously (Washizuka et al 2003). The objective of this study was clearly explained, and written informed consent was obtained from all subjects.

Genomic DNA was extracted from leukocytes using standard methods. There was no evidence for the presence of population substructure in either control subjects or BD using the method of Prichard (2000; Kakiuchi et al 2003). The ethics committees of the Brain Science Institute and participating institutes approved this study.

**NIMH Genetics Initiative Pedigrees**

For TDT, 105 trio samples (94 trios with BDI probands and 11 trios with BDII probands) were obtained from NIMH Genetics Initiative Bipolar Pedigrees. Each trio was obtained from a larger NIMH family independent of each other. Of those, results of genotyping were inconsistent with the parent-child relationship in one pedigree with BDI. Thus, this pedigree was omitted from the analysis.

**Mutation Screening of the *NDUFV2* Gene by Sequencing**

Polymorphisms of the upstream region of the *NDUFV2* (GenBank accession number NT\_010859) were screened in 20 randomly selected Japanese subjects (9 BDI, 3 BDII, and 8 control subjects). For the scanning of the 5'-upstream region, the following primer sets were used: 5'-TATAGGTCATGAACTCAAAAAGACG and 5'-GCCACACTGTTACCTTCC. These primers amplified a 3983bp product. Polymerase chain reaction (PCR) was performed in a 25-µL volume containing 20 ng of genomic DNA, .2 µmol/L of each primer, 100 µmol/L of each dNTP, 12.5 µL of 2 × GC buffer I (Takara, Shiga, Japan), and 1.25 units of LA-Taq DNA polymerase (Takara). After an initial denaturation at 95°C for 2 min, 27 cycles consisting of 30 sec at 94°C, 30 sec at 62°C, and 4 min at 72°C were performed. An extension at 72°C for 5 min followed. Sequencing of the PCR products was conducted using the BigDye terminator sequencing kit (Applied Biosystems, Foster City, California) and an ABI 3700 DNA sequencer (Applied Biosystems). For this analysis, 15 sequence primers were used.

**Genotyping**

Five single nucleotide polymorphisms (SNPs) detected in the screening analysis, -3542G> A, -3245T> C, -3041T> G, -2694A> G, and -1020G> T, were genotyped in Japanese samples and NIMH bipolar pedigrees. For genotyping of the former four polymorphisms, genomic DNA was amplified by using the upstream primer 5'-AAACTAGCCCTTCCATTCTCCTT and the downstream primer 5'-CCTTCTGTCTCATTTGGCT-TACA. These primers amplified a 1547bp product. We performed PCR in a 15-µL volume containing 15 ng of genomic DNA, .1 µmol/L of each primer, 25 µmol/L of each dNTP, 1.5 µL of 10 × Ex-Taq buffer (Takara), and .72 units of Ex-Taq DNA polymerase (Takara). After an initial denaturation at 95°C for 2 min, 35 cycles

Table 1. Pairwise Linkage Disequilibrium Between Polymorphisms in the *NDUFV2*

	-3542G>A	-3245T>C	-3041T>G	-2694A>G	-1020G>T	-796C>G	-795T>G	-602G>A	-233T>C	86C>T
-3542G>A	.32									$r^2 = .59$
-3245T>C	.32	$r^2 = .97$								$r^2 = .59$
-3041T>G	.28	$D' = 1.00$	$r^2 = .17$							$r^2 = .21$
-2694A>G	.38	$D' = 1.00$	$r^2 = .16$	$r^2 = 1.00$						$r^2 = .60$
-1020G>T	.16	$D' = 1.00$	$D' = 1.00$	$D' = 1.00$	$r^2 = .11$					$r^2 < .01$
-796C>G	.41	$D' = .96$	$D' = .94$	$D' = 1.00$	$D' = 1.00$	$r^2 = .33$				$r^2 = .16$
-795T>G	.41	$D' = .96$	$D' = .94$	$D' = .97$	$D' = 1.00$	$r^2 = .29$	$r^2 = .33$			$r^2 = .16$
-602G>A	.32	$D' = .81$	$D' = .77$	$D' = .98$	$D' = .92$	$r^2 = .23$	$r^2 = .29$	$r^2 = .91$		$r^2 = .43$
-233T>C	.28	$D' = .84$	$D' = .65$	$D' = 1.00$	$D' = .29$	$r^2 = .23$	$r^2 = .29$	$r^2 = .63$	$r^2 = .20$	$r^2 = .13$
86C>T	.39	$D' = .87$	$D' = 1.00$	$D' = .87$	$D' < .01$	$r^2 = .29$	$r^2 = 1.00$	$r^2 = .11$	$r^2 = .08$	$D' = 1.00$

**Table 2.** Genotypic and Allele Distributions of the Additional *NDUF2* Gene Polymorphisms and -602G>A in Japanese Controls and Bipolar Patients

Polymorphisms		Subject Counts (%)				p Value <sup>b</sup>
		Controls	BP Total	BPI	BPII	
<b>-3542G&gt;A</b>						
Genotype	G/G	15 (.07)	25 (.13)	13 (.09)	12 (.23)	.02
	G/A	102 (.46)	80 (.42)	63 (.46)	17 (.32)	
	A/A	104 (.47)	85 (.45)	61 (.45)	24 (.45)	
P Value			.09	.62	.003	
Allele	G	132 (.30)	130 (.34)	89 (.32)	41 (.39)	.21
	A	310 (.70)	250 (.66)	185 (.68)	65 (.61)	
	P Value		.20	.50	.08	
<b>-3245T&gt;C</b>						
Genotype	T/T	15 (.07)	24 (.13)	12 (.09)	12 (.23)	.01
	T/C	101 (.47)	80 (.42)	64 (.47)	16 (.30)	
	C/C	97 (.46)	84 (.45)	59 (.44)	25 (.47)	
P Value			.14	.80	.002	
Allele	T	131 (.31)	128 (.34)	88 (.33)	40 (.38)	.38
	C	295 (.69)	248 (.66)	182 (.67)	66 (.62)	
	P Value		.32	.61	.20	
<b>-3041T&gt;G</b>						
Genotype	T/T	106 (.49)	98 (.52)	73 (.54)	25 (.47)	.009
	T/G	88 (.40)	86 (.46)	60 (.44)	26 (.49)	
	G/G	23 (.11)	4 (.02)	2 (.02)	2 (.04)	
P Value			.001	.002	.23	
Allele	T	300 (.69)	282 (.75)	206 (.76)	76 (.72)	.12
	G	134 (.31)	94 (.25)	64 (.24)	30 (.28)	
	P Value		.07	.04	.63	
<b>-2694A&gt;G</b>						
Genotype	A/A	16 (.07)	24 (.13)	12 (.09)	12 (.23)	.03
	A/G	99 (.46)	79 (.42)	62 (.46)	17 (.32)	
	G/G	100 (.47)	84 (.45)	60 (.45)	24 (.45)	
P Value			.19	.86	.006	
Allele	A	131 (.31)	128 (.34)	88 (.33)	40 (.38)	.38
	G	295 (.69)	248 (.66)	182 (.67)	66 (.62)	
	P Value		.32	.61	.20	
<b>-1020G&gt;T</b>						
Genotype	G/G	160 (.72)	131 (.69)	93 (.66)	38 (.72)	.77
	G/T	58 (.26)	52 (.28)	39 (.29)	13 (.25)	
	T/T	4 (.02)	6 (.03)	4 (.03)	2 (.03)	
P Value			.64	.62	.60	
Allele	G	378 (.85)	314 (.83)	225 (.83)	89 (.84)	.66
	T	66 (.15)	64 (.17)	47 (.17)	17 (.16)	
	P Value		.44	.39	.76	
<b>-602G&gt;A<sup>a</sup></b>						
Genotype	G/G	17 (.08)	27 (.14)	15 (.11)	12 (.23)	.02
	G/A	106 (.48)	77 (.41)	60 (.44)	17 (.32)	
	A/A	99 (.44)	85 (.45)	61 (.45)	24 (.45)	
P Value			.07	.51	.003	
Allele	G	140 (.32)	131 (.35)	90 (.33)	41 (.39)	.37
	A	304 (.68)	247 (.65)	182 (.67)	65 (.61)	
	P Value		.34	.66	.15	

BP, bipolar disorder


<sup>a</sup>Data of Washizuka et al (2003).<sup>b</sup>Differences in genotype distributions or allele frequencies among patients with BPI, BPII, and controls.

consisting of 30 sec at 94°C, 30 sec at 61°C, and 2 min at 72°C were performed. An extension at 72°C for 3 min followed.

For genotyping of the -1020G>T polymorphism, the primer sets were used as follows: 5'-ACCAAGGCATTGGTATCTATTCT and 5'-ATGTTTGGTTATCTCTGGAAA. We performed PCR in a 25- $\mu$ L volume containing 25 ng of genomic DNA, .1  $\mu$ mol/L of each primer, 25  $\mu$ mol/L of each dNTP, 2.5  $\mu$ L of 10  $\times$  Ex-Taq buffer, and 1.2 units of Ex-Taq DNA polymerase. After an initial

denaturation at 95°C for 2 min, 35 cycles consisting of 20 sec at 94°C, 30 sec at 60°C, and 30 sec at 72°C were performed. An extension at 72°C for 3 min followed.

In addition, our previously reported polymorphisms of the *NDUF2* gene (-796C>G, -795T>G, -602G>A, -233T>C, and 86C>T) were also genotyped in NIMH samples. The primer set and the PCR condition that were used in genotyping of these SNPs are shown in our previous article (Washizuka et al 2003).



**Figure 1.** Haplotype associations in Japanese case control samples: (a) three-marker haplotype analysis; (b) two-marker haplotype analysis. *P* value indicates the global *P* analyzed using COCAPHASE program. Gray squares indicate statistically significant global *p* value.

Sequencing of the PCR products was conducted using the BigDye terminator sequencing kit (Applied Biosystems) and an ABI 3700 DNA sequencer (Applied Biosystems).

#### Statistical Procedures


Deviations from Hardy-Weinberg equilibrium (HWE) were evaluated by use of the Arlequin program (<http://anthropologie.unige.ch/arlequin/methods.html>). Genotype distributions and allele frequencies between patients and control subjects were computed using Fisher's Exact Probability Test, which was applied using SPSS software (SPSS, Tokyo, Japan). For other analyses, UNPHASED programs (COCAPHASE and TDTPHASE; <http://www.rfcgr.mrc.ac.uk/~fdudbrid/software/unphased/>) were used. The normalized LD coefficient *D'* and the squared correlation coefficient *r*<sup>2</sup> were calculated using COCAPHASE program. For TDT of NIMH trio samples, the McNemar Test was used. For the computation of haplotype frequencies, evaluation of haplotypic distributions, and TDT analysis of the multimeric haplotypes, the COCAPHASE and TDTPHASE programs were used. To evaluate the data appropriately, we reanalyzed the significant result using the permutation test implemented in COCAPHASE and TDTPHASE. Sequences were searched for potential transcription factor binding sites using the Match program (<http://www.gene-regulation.de/>).

#### Promoter Assay

A 1106-bp fragment (-1111 to -6) of the upstream from the initiation codon of the *NDUFV2* gene was amplified by PCR and cloned into the *MluI/BglII* site of pGL3-Basic vector (Promega, Madison, Wisconsin). Two kinds of reporter plasmids, having either -602G or -602A were prepared. A 586-bp fragment (-591 to -6) lacking the -602G>A site was also amplified and cloned into the same vector. HeLa-S3 and HEK293 cells cultured in a 96-well plate were transfected using Superfect (Qiagen, Valencia, California) with .5 mg of the reporter plasmid, .05 mg of a reference plasmid (pRL-TK), and the pGL3-Basic vector carrying no insert. After 36 hours incubation, luciferase activities were measured with the aid of Dual-Glo luciferase assay system (Promega). Four independent experiments were performed for each condition, and the mean and SEM values were presented.

#### Results

We previously reported that -602G>A, among four polymorphisms (-796C>G, -795T>G, -602G>A, -233T>C) in



**Figure 2.** Promoter assay. Promoter activity of three kinds of reporter plasmids, having -602G or -602A and lacking the -602G>A site, were examined by the luciferase assay. Two kinds of cells, HeLa-S3 and HEK293, were used. Four independent experiments were performed for each condition, and the mean and SEM values were presented. The transcription activity of the -602G construct was significantly higher than that of the -602A in both HeLa-S3 and HEK293. The short construct lacking in the -602G>A site presented nearly equal activity of the -602A construct and had significantly smaller activity compared with -602G constructs. \*\**p* < .01. \**p* < .05. +*p* = .05.

the upstream region of *NDUFV2*, showed the significant association with BD (Washizuka et al 2003). In this study, four novel (-3542G>A, -3245T>C, -3041T>G, -2694A>G) and one known (-1020G>T) polymorphisms were detected by sequencing the upstream region extending up to 3963 bp of the transcription initiation site in the *NDUFV2*. We then genotyped these polymorphisms in Japanese patients with BD (*n* = 189) and control subjects (*n* = 222). The genotype frequencies of these five polymorphisms were in HWE in control subjects, although -3041T>C polymorphism was not in BDI patients, and -3245T>G was not in BDII patients. There was no significant difference of genotype frequencies of each SNP between male and female subjects. Detected polymorphisms in this study and in our previously reported polymorphisms (-796C>G, -795T>G, -602G>A, -233T>C, and 86C>T) were in strong linkage disequilibrium with each other (Table 1).

The genotype and allele distributions of the polymorphisms in Japanese population are shown in Table 2. The data for -602G>A polymorphism were cited from our previous paper (Washizuka et al 2003). Statistically significant differences in genotype distribution were observed between patients with BD and control subjects for -3041T>G (*p* = .001). The -3542G>A, -3245T>G, and -2694A>G polymorphisms showed significant genotypic association with BDII (*p* = .003, *p* = .002, and *p* = .006, respectively). These SNPs tended to be associated with BD or BDII even after Bonferroni correction. There was also a nominally significant difference in allelic distribution of -3041T>G polymorphism between patients with BDI and control subjects (*p* = .04).

Haplotype analysis consisting of all 10 polymorphisms revealed a statistically significant association in Japanese samples (global *p* < .0001). To explore which part of the *NDUFV2* gene

**Table 3.** Estimated Haplotype Frequencies of *NDUFV2* in Japanese Analyzed by Using the COCAPHASE Program

Haplotype <sup>a</sup>	Case	Frequency	Control	Frequency	$\chi^2$	P Value	Common
A-A	244	.64	270	.61	.34	.55	+
G-A	3	.008	31	.07	20.35	<.0001	+
A-G	4	.01	37	.08	23.81	<.0001	+
G-G	127	.33	99	.22	10.79	.001	+
Global P						<.0001 (<.0001) <sup>b</sup>	

<sup>a</sup>Haplotypes of -3542G>A and -602G>A.

<sup>b</sup>The global P value in parentheses shows the global significance by permutation test.

contributes most to this overall association, we employed the sliding window approach in which each set of two or three consecutive polymorphisms were tested for association with BD (two- or three-marker haplotype analysis). This analysis showed evidence of association with BD in two limited regions around -3542G>A and -602G>A (most significant haplotype  $p < .0001$ , global  $p < .0001$ , and most significant haplotype  $p = .008$ , global  $p = .001$ , for the three-marker analysis, and global  $p < .0001$  and global  $p = .002$ , respectively, for the two-marker analysis; Figure 1).

Because these two SNPs were located at the putative promoter region, we supposed that these polymorphisms might alter the transcription activity. At the beginning, we prepared a 3983bp fragment containing those two polymorphisms and tried to ligate this fragment into the pGL3-basic vector; this was not successful, however. Then we examined whether the -602G>A polymorphism had functional significance. Based on our previous analysis indicating that the two major haplotypes, C-T-A-T and C-T-G-T (consisting of -796C>G, -795T>G, -602G>A, and -233T>C polymorphisms of *NDUFV2*) were associated with BD, constructs of these two haplotypes were analyzed. Promoter activity was examined in two cell lines, HeLa-S3 and HEK293. The transcription activity of the -602G construct was significantly higher than that of the -602A both in HeLa-S3 and HEK293 ( $p = .03$  for HeLa-S3, and  $p = .05$  for HEK293). The short construct lacking in the -602G>A site presented nearly equal activity of the -602A construct but had significantly smaller activity compared with the -602G construct ( $p = .0009$  and  $p = .005$ , respectively; Figure 2).

Because we could not experimentally examine the functional significance of the -3542G>A, we examined whether this site affects the putative binding sites of transcription factors using the Match program. The -3542G>A was predicted to be within the putative binding site for HSF (heat shock transcription factor). HSF1 is known to affect the expression of several other nuclear encoded mitochondrial complex I subunit genes (e.g., *NDUFB8*,

*NDUFA10*, *NDUFAB1*, and *NDUFS1*). Recently, the binding sequence of HSF1 was well characterized (TTCT/C][G/A]GAANN-TC[T/C]; the bases similar to this site of *NDUFV2* promoter was italicized; Trinklein et al 2004). When the -3542 site is G, the core sequence of putative binding site for HSF1, GAA, is lost, and probability of binding was predicted to be decreased.

Thus, the frequency of haplotypes consisting of -3542G>A and -602G>A polymorphisms was also estimated. Distributions of haplotype frequencies differed significantly between patients with BD and control subjects (global  $p < .0001$ ) (Table 3). Among the haplotypes, the G-G haplotype was significantly more frequently seen in BD ( $p = .001$ ), whereas G-A and A-G haplotypes were significantly less common in patients with BD compared with control subjects ( $p < .0001$ ). The results were basically similar when younger control subjects were excluded to match ages of the subjects.

We then performed a TDT in NIMH Genetics Initiative Bipolar Pedigrees. The distributions of genotypes of all 10 detected polymorphisms of the probands, fathers, and mothers were in HWE. We could not detect any allele that was significantly overtransmitted from patients to affected offspring in the NIMH trio samples of BD (Table 4).

We then examined the transmission of haplotypes consisting of -3542G>A and -602G>A polymorphisms from patients to affected offspring by using TDTPHASE program. We found significant association of the *NDUFV2* haplotypes with BD (global  $p < .0001$ ). Two haplotypes (G-A and A-G) tended to be undertransmitted in parents-proband trios of NIMH samples (nominal  $p = .04$  and  $p = .01$ , respectively; Table 5).

## Discussion

We identified four novel polymorphisms (-3542G>A, -3245T>G, -3041T>C, and -2694A>G) associated with BD in this study. Haplotype analysis revealed that two haplotype blocks surrounding the -3542G>A and -602G>A polymor-

**Table 4.** Transmission Disequilibrium Test in National Institute of Mental Health Initiative Bipolar Pedigrees

Polymorphism	Allele	Tr	Not Tr	Ratio	$\chi^2$	P	Number of Trios
-3542G>A	G	20	27	.74	1.04	.30	85
-3245T>C	T	20	30	.67	2.00	.15	91
-3041T>G	T	39	44	.89	.30	.58	90
-2694A>G	A	19	26	.73	1.08	.29	81
-1020G>T	G	10	16	.63	1.38	.23	95
-796C>G	C	42	48	.88	.40	.52	94
-602G>A	G	31	24	1.29	.89	.34	98
-233T>C	T	43	50	.86	.52	.46	99
86C>T	C	34	25	1.36	1.37	.24	99

Tr, transmitted.

**Table 5.** Transmission of Haplotypes in National Institute of Mental Health Initiative Bipolar Pedigrees Analyzed by using TDTPHASE

Haplotype <sup>a</sup>	Tr	Frequency	Not Tr	Frequency	$\chi^2$	P Value	Common
A-A	146	.84	137	.80	1.61	.20	+
G-A	<.01	<.01	3	.02	4.18	.04	+
A-G	<.01	<.01	4	.02	5.59	.01	+
G-G	26	.15	28	.16	.08	.76	+
Global P						.01 (.11) <sup>b</sup>	

<sup>a</sup>Haplotypes of -3542G>A and -602G>A.

<sup>b</sup>The global P value in parentheses shows the global significance by permutation test.

phisms were associated with BD. The haplotype of these two SNPs were significantly associated with BD in Japanese subjects. In NIMH trios, no individual SNP was associated, and the overtransmission of the risk haplotype in Japanese, G-G, was not observed. Although the observed trend of undertransmission of two haplotypes, G-A and A-G, might be due to the small number of trios, it is noteworthy that the trends of undertransmission of these two haplotypes seen in the NIMH bipolar trio samples were in the same direction to the significant decrease of these haplotypes in Japanese BD subjects.

Although the mechanism by which -602A>G changed the promoter activity is unknown, it would be of interest to note that the -602G polymorphism loses the putative binding site of a transcription factor, p300 (CCACTC). The finding that the -602G haplotype is more common in BD, although yielding a significantly higher promoter activity, is apparently inconsistent with our mitochondrial dysfunction hypothesis in BD; however, the direction of change of promoter activity in the luciferase assay cannot be directly compared with that in vivo. First, promoter activity can be affected by neighboring sequences because several transcription factors form a complex. When only a part of the promoter sequence is subcloned into the luciferase vector, as in this study, the promoter activity does not directly represent the activity in vivo. Second, regulation of gene expression is complex. Although we reported that mRNA expression of *NDUFV2* was decreased in the lymphoblastoid cells, Karry et al (2004) reported that the protein encoded by *NDUFV2* was up-regulated in autopsied BD brains. Thus, it cannot be concluded what kind of mitochondrial dysfunction is caused by polymorphisms of *NDUFV2* promoter. Even though the direction of the change may not represent the promoter activity in vivo, the results of promoter assay indicate that this region has some functional activity only when the -602 position is G.

We could not determine whether -3542G>A affects promoter activity, and this would be a worthwhile topic for future study. Because it has been reported that lithium enhances HSF1 activity (Carmichael et al 2002), it would be particularly interesting to examine the effects of HSF1.

It is most important to test whether the association of *NDUFV2* with BD is replicated using independent BD case-control or trio samples. In addition, 18p11 is a common linkage locus for BD and schizophrenia (Berrettini 2000; Lewis et al 2003), and mRNA and protein expression of *NDUFV2* is also altered in schizophrenia (Karry et al 2004). It would thus be interesting to examine the association between *NDUFV2* and schizophrenia.

In conclusion, the haplotypes consisting of -3542G>A and -602G>A polymorphisms in the upstream region of *NDUFV2* were associated with BD commonly in two ethnicities. Together with altered promoter activity, these findings indicate the role of *NDUFV2* as a genetic risk factor of bipolar disorder.

*Data and biomaterials of the National Institute of Mental Health (NIMH) pedigrees were collected in four projects that participated in the NIMH Bipolar Disorder Genetics Initiative. From 1991 to 1998, the principal investigators and co-investigators were as follows: Indiana University, Indianapolis, Indiana, U01 MH46282, John Nurnberger, M.D., Ph.D., Marvin Miller, M.D., and Elizabeth Bowman, M.D.; Washington University, St. Louis, MO, U01 MH46280, Theodore Reich, M.D., Allison Goate, Ph.D., and John Rice, Ph.D.; Johns Hopkins University, Baltimore, Maryland U01 MH46274, J. Raymond DePaulo, Jr., M.D., Sylvia Simpson, M.D., MPH, and Colin Stine, Ph.D.; NIMH Intramural Research Program, Clinical Neurogenetics Branch, Bethesda, Maryland, Elliot Gershon, M.D., Diane Kazuba, B.A., and Elizabeth Maxwell, M.S.W.*

- Berrettini WH (2000): Susceptibility loci for bipolar disorder: Overlap with inherited vulnerability to schizophrenia. *Biol Psychiatry* 47:245–251.
- Berrettini WH, Ferraro TN, Goldin LR, Detera-Wadleigh SD, Choi H, Muniec D, et al (1997): A linkage study of bipolar illness. *Arch Gen Psychiatry* 54:27–35.
- Carmichael J, Sugars KL, Bao YP, Rubinsztein DC (2002): Glycogen synthase kinase-3beta inhibitors prevent cellular polyglutamine toxicity caused by the Huntington's disease mutation. *J Biol Chem* 277:33791–33798.
- de Coo R, Buddiger P, Smeets H, Geurts van Kessel A, Morgan-Hughes J, et al (1995): Molecular cloning and characterization of the active human mitochondrial NADH: Ubiquinone oxidoreductase 24-kDa gene (*NDUFV2*) and its pseudogene. *Genomics* 26:461–466.
- Gershon ES, Bander JA, Detera-Wadleigh SD, Ferraro TN, Berrettini WH (1996): Maternal inheritance and chromosome 18 allele sharing in unilineal bipolar illness pedigrees. *Am J Med Genet* 67:202–207.
- Goodwin FK, Jamison KR (1990): *Manic-Depressive Spectrum*. New York: Oxford University Press.
- Hattori N, Suzuki H, Wang Y, Minoshima S, Shimizu N, Yoshino H, et al (1995): Structural organization and chromosomal localization of the human nuclear gene (*NDUFV2*) for the 24-kDa iron-sulfur subunit of complex I in mitochondrial respiratory chain. *Biochem Biophys Res Commun* 216:771–777.
- Kakiuchi C, Iwamoto K, Ishiwata M, Bundo M, Kasahara T, Kusumi I, et al (2003): Impaired feedback regulation of XBP1 as a genetic risk factor for bipolar disorder. *Nat Genet* 35:171–175.
- Karry R, Klein E, Ben Shachar D, (2004): Mitochondrial complex I subunits expression is altered in schizophrenia: A postmortem study. *Biol Psychiatry* 55:676–684.
- Kato T, Kato N (2000): Mitochondrial dysfunction in bipolar disorder. *Bipolar Disord* 2:180–190.
- Kato T, Kunugi H, Nanko S, Kato N (2001): Mitochondrial DNA polymorphisms in bipolar disorder. *J Affect Disord* 62:151–164.
- Kato T, Stine OC, McMahon FJ, Crowe RR (1997): Increased levels of a mitochondrial DNA deletion in the brain of patients with bipolar disorder. *Biol Psychiatry* 42:871–875.
- Kato T, Takahashi S, Shioiri T, Inubushi T (1993): Alterations in brain phosphorus metabolism in bipolar disorder detected by in vivo 31P and 7Li magnetic resonance spectroscopy. *J Affect Disord* 27:53–59.
- Lewis CM, Levinson DF, Wise LH, DeLisi LE, Straub RE, Hovatta I, et al (2003): Genome scan meta-analysis of schizophrenia and bipolar disorder, part II: Schizophrenia. *Am J Hum Genet* 73:34–48.

- Nakatani N, Aburatani H, Nishimura K, Semba J, Yoshikawa T (2004): Comprehensive expression analysis of a rat depression model. *Pharmacogenomics J* 4:114–126.
- Nothen MM, Cichon S, Rohleder H, Hemmer S, Franzek E, Fritze J, et al (1999): Evaluation of linkage of bipolar affective disorder to chromosome 18 in a sample of 57 German families. *Mol Psychiatry* 4:76–84.
- Pritchard JK, Stephens M, Donnelly P (2000): Inference of population structure using multilocus genotype data. *Genetics* 155:945–959.
- Segurado R, Detera-Wadleigh SD, Levinson DF, Lewis CM, Gill M, Nurnberger JI, et al (2003): Genome scan meta-analysis of schizophrenia and bipolar disorder, part III: Bipolar disorder. *Am J Hum Genet* 73:49–62.
- Smeitink J, Sengers R, Trijbels F, van den Heuvel L (2001): Human NADH: ubiquinone oxidoreductase. *J Bioenerg Biomembr* 33:259–266.
- Stine OC, Xu J, Koskela R, McMahon FJ, Gschwend M, Friddle C, et al (1995): Evidence for linkage of bipolar disorder to chromosome 18 with a parent-of-origin effect. *Am J Hum Genet* 57:1384–1394.
- Trinklein ND, Murray JI, Hartman SJ, Botstein D, Myers RM (2004): The role of heat shock transcription factor 1 in the genome-wide regulation of the mammalian heat shock response. *Mol Biol Cell* 15:1254–1261.
- Turecki G, Grof P, Cavazzoni P, Duffy A, Grof E, Martin R, et al (1999): Lithium responsive bipolar disorder, unilinearity and chromosome 18: A linkage study. *Am J Med Genet* 88:411–415.
- Washizuka S, Kakiuchi C, Mori K, Kunugi H, Tajima O, Akiyama T, et al (2003): Association of mitochondrial complex I subunit gene *NDUFV2* at 18p11 with bipolar disorder. *Am J Med Genet* 120B:72–78.

# A Family-Based Association Study and Gene Expression Analyses of Netrin-G1 and -G2 Genes in Schizophrenia

Mika Aoki-Suzuki, Kazuo Yamada, Joanne Meerabux, Yoshimi Iwayama-Shigeno, Hisako Ohba, Kazuya Iwamoto, Hitomi Takao, Tomoko Toyota, Yumiko Suto, Noriaki Nakatani, Brian Dean, Sachiko Nishimura, Kenjiro Seki, Tadafumi Kato, Shigeyoshi Itoharu, Toru Nishikawa, and Takeo Yoshikawa

**Background:** The netrin-G1 (NTNG1) and -G2 (NTNG2) genes, recently cloned from mouse, play a role in the formation and/or maintenance of glutamatergic neural circuitry. Accumulating evidence strongly suggests that disturbances of neuronal development and the N-methyl-D-aspartate receptor-mediated signaling system might represent a potential pathophysiology in schizophrenia. We therefore set out to examine the genetic contribution of human NTNG1 and NTNG2 to schizophrenia.

**Methods:** Twenty-one single nucleotide polymorphisms (SNPs) from NTNG1 and 10 SNPs from NTNG2 were analyzed in 124 schizophrenic pedigrees. All genotypes were determined with the TaqMan assay. The expression levels of NTNG1 and NTNG2 were examined in the frontal (Brodmann's Area [BA]11 and BA46) and temporal (BA22) cortices from schizophrenic and control postmortem brains. The isoform-specific expression of NTNG1 splice variants was assessed in these samples.

**Results:** Specific haplotypes encompassing alternatively spliced exons of NTNG1 were associated with schizophrenia, and concordantly, messenger ribonucleic acid isoform expression was significantly different between schizophrenic and control brains. An association between NTNG2 and schizophrenia was also observed with SNPs and haplotypes that clustered in the 5' region of the gene.

**Conclusions:** The NTNG1 and NTNG2 genes might be relevant to the pathophysiology of schizophrenia.

**Key Words:** Neurodevelopment, laminin-1, laminin-2, postmortem brain, real-time quantitative polymerase chain reaction, alternative splicing

Schizophrenia is a severe brain disorder that usually produces a lifetime of disability, with emotional and cognitive distress for the affected individuals (Lewis and Lieberman 2000). Although schizophrenia has a worldwide prevalence of approximately 1%, the specific factors that underlie its predisposition remain elusive. A number of studies have focused on identifying genetic and environmental components that separately, or in combination, contribute to the manifestation of the disease (Lewis and Levitt 2002).

A strongly supported theory is that schizophrenia evolves from a fixed brain lesion occurring early in neurodevelopment that responds abnormally to later maturational stimuli (Chua and Murray 1996; Marengo and Weinberger 2000; Weinberger 1987). Developmental neurobiology has made great strides in detailing the molecular machinery that leads to the assembly of brain systems (Jessell 2000; Tessier-Lavigne and Goodman 1996). Several gene families that are responsible for encoding cues for

guiding axonal growth cones and cell migration have been identified. These genes include members of the immunoglobulin superfamily, ephrins, semaphorins, slits, and netrins (Chisholm and Tessier-Lavigne 1999; Tessier-Lavigne and Goodman 1996). For the cellular and signaling bases for schizophrenia pathophysiology, disturbances in N-methyl-D-aspartate (NMDA) and dopamine neurotransmission have also been strongly implicated.

The netrins are a family of diffusible axon guidance molecules related to laminin that are conserved from *Caenorhabditis elegans* to vertebrates (Ishii et al 1992; Serafini et al 1994). A netrin-related molecule, netrin-G1 (Ntng1) (also called laminin-1), has been identified in mice and shown to be distinct from the classical netrins in several aspects (Nakashiba et al 2000; Yin et al 2002). Unlike classical netrins, Ntng1 is predominantly tethered to the membrane through a carboxyl-terminal glycosyl phosphatidyl-inositol anchor. It generates several splice variants, none of which bind deleted in colorectal carcinoma (DCC) or uncoordinated 5 (UNC5), the best characterized netrin receptors. Nakashiba et al (2002) and Yin et al (2002) have recently identified a close paralogue, netrin-G2 (Ntng2) (also called laminin-2). Both *Ntng1* and *Ntng2* are elaborated within the vertebrate lineage. Comparative analysis of mouse *Ntng1* and *Ntng2* revealed complementary expression in the brain, implying nonredundant roles (Nakashiba et al 2002; Yin et al 2002). Our ongoing studies with mutant mice devoid of these proteins suggest that *Ntng1* and *Ntng2* are necessary in correct NMDA receptor functioning and that mutant mice show behavioral phenotypes related to schizophrenia (Nishimura et al 2004).

To define the possible roles of *NTNG1* and *NTNG2* molecules in schizophrenia, we mapped the genomic layout of human *NTNG1* and *NTNG2*, confirmed their chromosomal localization, and detected multiple splice variants of *NTNG1* transcripts. On the basis of these findings, we performed a

From the Laboratories for Molecular Psychiatry (MAS, KY, JM, YIS, HO, HT, TT, NN, TY), Molecular Dynamics of Mental Disorders (KI, TK), and Behavioral Genetics (SN, KS, SI), RIKEN Brain Science Institute, Wako; Section of Psychiatry and Behavioral Sciences (MAS, TN), Tokyo Medical and Dental University Graduate School, Tokyo; Department of Pediatric Cardiology (YS), Tokyo Women's Medical University, Tokyo, Japan; and The Rebecca L. Cooper Research Laboratories (BD), Mental Health Research Institute of Victoria, Parkville, Victoria, Australia.

Address reprint requests to Takeo Yoshikawa, M.D., Ph.D., RIKEN Brain Science Institute, Laboratory for Molecular Psychiatry, 2-1 Hirosawa, Wako-city, Saitama 351-0198, Japan; E-mail: takeo@brain.riken.go.jp.

Received April 21, 2004; revised November 4, 2004; accepted November 11, 2004.



genetic and expression study of these two closely related genes in schizophrenia.

## Methods and Materials

### Subjects

The schizophrenia pedigrees were all derived from a geographic area located in central Japan. The probands, both in- and outpatients, were followed up by hospital doctors for at least 6 months. The samples consisted of 124 families with 376 members, of whom 163 were affected. This included 1) 80 independent and complete trios (schizophrenic offspring and their parents); 2) 15 probands with one parent; 3) 13 probands with affected siblings; and 4) 30 probands with discordant siblings (Yamada et al 2004). There was some overlap in the samples from 1), 3), and 4). Consensual diagnosis was made according to DSM-IV criteria by at least two experienced psychiatrists, on the basis of direct interviews, available medical records, and information from hospital staff and relatives. None of the patients had additional Axis I disorders as defined by DSM-IV, and none of the present family members suffered from neurodegenerative disorders, including Parkinson's and Alzheimer's diseases.

The present study was approved by the ethics committee of RIKEN Brain Science Institute. All patients and family members gave informed and written consent to participate in the study.

### Determination of the Exon/Intron Structures of NTNG1 and NTNG2

The complete genomic structure of *NTNG1* was not available from the databases and was determined as described (Meerabux et al, unpublished data): briefly, mouse netrin-G1a complementary deoxyribonucleic acid (cDNA) sequence NM\_030699 and G1d sequence AB038664, as well as human *NTNG1* clones BC030220 and AB023193, were aligned with human genomic bacterial artificial chromosome (BAC) clones RP11-270C12, RP11-396N10, and RP11-436H6 (GenBank identification [ID] numbers AC114491, AL590427, and AL513187, respectively; <http://www.ncbi.nlm.nih.gov/Entrez/index.html>) using the National Center for Biotechnology Information (NCBI) BLAST 2 sequences algorithm (<http://www.ncbi.nlm.nih.gov/blast/bl2seq/bl2.html>). For *NTNG2* genomic organization, we consulted the Ensembl Human Genome Browser ([http://www.ensembl.org/Homo\\_sapiens/](http://www.ensembl.org/Homo_sapiens/)), which provided annotation on the longest cDNA clone available in the databases (Vega Gene ID: OTTHUMG00000020835).

### Fluorescent In Situ Hybridization (FISH)

For fluorescent in situ hybridization (FISH) analysis, *NTNG1* cDNA probe was prepared by polymerase chain reaction (PCR) amplification of Marathon-Ready Human adult brain cDNA (Clontech, Palo Alto, California) with a forward primer (5'-AGGGGGCGACTTGCAGGAGGC, 3' end at nucleotide (nt) -570: A of the ATG initiation codon is counted as +1) and a reverse primer (5'-GCAGTGCCTTTGGGGATGGGG, 3' end at nt 1032 in BC030220). The procedure was repeated for *NTNG2* (forward primer: 5'-CTGAGAGGTTCTGCTCCCATG, 3' end at nt 3; reverse primer: GGGTAGCAGCCCTGGCGCCAG, 3' end at nt 1313 in AB058760). The PCR products were separated on an agarose gel, and the DNA was excised, purified, and sequenced to check the fidelity of the probes. The purified DNA samples were fluorescently labeled under standard conditions and used as probes against normal metaphase chromosomes.

### Detection of Ntng1 Transcript Isoforms

On the basis of information regarding the mouse *Ntng1* splice variation, which is generated by alternative splicing of downstream exons (Nakashiba et al 2000), we designed a PCR assay to detect human splice variants from Marathon-Ready Human adult brain cDNA (Clontech), using a forward primer designed to exon 4 and a reverse primer in exon 10 (Meerabux et al, unpublished data). The PCR products were cloned into pCR2.1 (Invitrogen, Carlsbad, California) and sequenced with the DYEnamic ET terminator cycle sequencing kit (Amersham Pharmacia Biotech, Piscataway, New Jersey). The exonic composition of each fragment was determined by aligning each sequence result against the known *NTNG1* exon sequences with SEQUENCHER software (Gene Codes, Ann Arbor, Michigan).

We also performed semiquantitative analysis of *NTNG1* messenger ribonucleic acid (mRNA) splice variants by separating the PCR fragments with an Agilent 2100 bioanalyzer with the DNA1000 LabChip kit (Agilent Technologies, Palo Alto, California). The bioanalyzer software automatically calculates migration time (sec), area, size (bp), concentration (ng/ $\mu$ L), and molarity (nmol/L) of each separated band and displays the results in real time. The identities of the detected peaks were determined by comparison with the cloned isoforms run in a separate well on the same chip.

### Mutation Screening

Genomic DNA was isolated from blood samples according to standard methods. The complete coding region and exon/intron boundaries of *NTNG1* and *NTNG2* were screened for polymorphisms by direct sequencing of PCR products, from 40 unrelated schizophrenia samples. Primers used for amplification are listed in Appendix 1 (available online). Polymerase chain reaction was performed with an initial denaturation at 94°C for 1 min, followed by 35 cycles at 94°C for 15 sec, 50°–70°C (optimized for each primer pair) for 15 sec, 72°C for 45 sec, and a final extension at 72°C for 2 min, with TaKaRa Taq polymerase (Takara Bio, Shiga, Japan). Detailed information on amplification conditions is available upon request. Direct sequencing of PCR products was performed with the BigDye Terminator Cycle Sequencing FS Ready Reaction kit (Applied Biosystems, Foster City, California) and the ABI PRISM 3700 Genetic Analyzer (Applied Biosystems). Polymorphisms were detected with SEQUENCHER. The Japanese Single Nucleotide Polymorphisms (<http://snp.ims.u-tokyo.ac.jp/index.html>), The SNP Consortium (<http://snp.cshl.org/news/>), and the NCBI (<http://www.ncbi.nlm.nih.gov/>) databases were searched for additional single nucleotide polymorphisms (SNPs).

### SNP Genotyping

Single nucleotide polymorphisms were typed in all 124 families with the TaqMan system (Applied Biosystems). Polymerase chain reactions were performed in an ABI 9700 thermocycler, and fluorescence was determined with an ABI 7900 sequence detector single point measurement and SDS v2.2 software (Applied Biosystems). Each marker was checked for allele-inheritance inconsistency within a pedigree with PEDCHECK software (O'Connell and Weeks 1998), and conflicts or flagged alleles were resolved by resequencing.

### Statistical Analyses

The initial panel of 124 families was analyzed with the pedigree disequilibrium test (PDT) program, v3.12 (<http://www.chg.duke.edu/software/pdt.html>) (Martin et al 2000, 2001). The statistically more conservative follow-up study was carried out

**Table 1.** Demographic and Treatment Data for the Australian Schizophrenic and Control Brains (BA11 and BA22)

Subject ID	Age (y)	Gender	PMI (h)	pH	Cause of Death	FRAD	FAPDD
<b>Schizophrenic Patients</b>							
1	79	F	26	6.27	Hypothermia	Fluphenazine decanoate	89
2	57	M	24	6.06	Coronary artery atheroma	Fluphenazine decanoate	45
3	65	M	41	6.57	Ischemic heart disease	Fluphenazine decanoate	45
4	47	F	50	6.31	Pneumonia	Resperidone	1200
5	65	F	50	6.35	Rupture of abdominal aortic aneurysm	Fluphenazine decanoate + haloperidol	339
6	69	M	48	6.44	Suicide: Carbon monoxide poisoning	Haloperidol	350
Mean ± SD	63.67 ± 10.86		39.83 ± 11.97	6.33 ± .17			
<b>Control Subjects</b>							
1	77	F	17	6.32	Hypertension heart disease		
2	57	M	27	6.43	Ischemic heart disease		
3	68	M	41	6.06	Aortic stenosis		
4	39	F	65	6.38	Mitral valve prolapse		
5	62	F	40	6.45	Ischemic heart disease		
6	68	M	69	6.59	Coronary artery atheroma		
Mean ± SD	61.83 ± 13.04		43.17 ± 20.52	6.37 ± .18			

BA, Brodmann's Area; ID, identification number; PMI, postmortem interval; FRAD, final recorded antipsychotic drug; FAPDD, final recorded antipsychotic drug dose (mg chlorpromazine equivalents/day); M, male; F, female.

with the complete 80-trio set and the extended transmission disequilibrium test (ETDT) algorithm, v2.2 (Sham and Curtis 1995). Empirical significance levels of the ETDT results were simulated from 10,000 Monte Carlo permutations with the MCETDT program, v1.3 (<http://www.mds.qmw.ac.uk/statgen/dcurtis/software.html>) (Zhao et al 1999). The TRANSMIT program, v2.5.4 (Clayton 1999; Clayton and Jones 1999) was used for haplotype-based transmission disequilibrium testing. Uncorrected *p* values for individual haplotypes (haplotypic *p*) and corrected *p* values for multi-allele testing (global *p*) were calculated with the TRANSMIT program.

Genomic linkage disequilibrium (LD) patterns retained in the Japanese population were determined by pairwise LD examination of markers within *NTNG1* and *NTNG2*, in 186 unrelated individuals from our schizophrenic pedigree panel. The standardized disequilibrium coefficient (*D'*) (Lewontin 1988) and the squared correlation coefficient (*r*<sup>2</sup>) were calculated with COCAPHASE software (<http://www.hgmp.mrc.ac.uk/~fdudbrid/software/unphased/>) (Dudbridge 2003).

#### Collection of Human Central Nervous System Tissues

Two sources of postmortem brain samples were used in this study. In one sampling, Brodmann's Area (BA)11 (lateral orbitofrontal cortex) and BA22 (superior temporal cortex) were collected by the Victorian Institute of Forensic Medicine, affiliated with the State Coroner's Office in Australia. At autopsy, brains were removed, and 1-cm coronal slices from the left hemispheres were rapidly frozen to  $-80^{\circ}\text{C}$ . The postmortem diagnoses were confirmed according to DSM-IV criteria by a psychiatrist and a senior psychologist using the Diagnostic Instrument for Brain Studies (Keks et al 1999). Ethical approval for the study was given by the North-Western Healthcare Network Human Ethics Committee. The gender distribution, mean age, mean postmortem interval, and pH for the tissues were not significantly different between schizophrenia (*n* = 6) and control (*n* = 6) groups (Table 1). The control subjects had no known history of psychiatric illness.

The second set of samples consisted of RNA extracted from BA46 (dorsolateral prefrontal cortex [DLPFC]), obtained from the

Stanley Foundation brain collection. Detailed information on the subjects and the extraction method are described on the web site, [http://www.stanleyresearch.org/programs/brain\\_collection.asp](http://www.stanleyresearch.org/programs/brain_collection.asp). Both schizophrenia and control groups consisted of 27 subjects. Diagnoses were made according to DSM-IV. A summary of the demographic information on these subjects is shown in Table 2. There were no significant demographic differences between schizophrenic and control brains (Torrey et al 2000). This study was performed unblinded.

#### RNA Preparation and cDNA Synthesis

Total RNA was extracted from the brain sample with an acid guanidium thiocyanate/phenol chloroform extraction method (ISOGEN; NIPPON Gene, Toyama, Japan). In the Australian samples, single-stranded cDNA was synthesized with SuperScript III RT (Invitrogen, Carlsbad, California) and oligo(dT) and random hexamers. For the Stanley Foundation brain collection samples, the extracted RNA was purified by RNAeasy column (Qiagen, Hilden, Germany). Single-stranded cDNA was synthesized by SuperScript II RT (Invitrogen) and oligo(dT) primers (Invitrogen) and converted to double-stranded cDNA by T4 DNA polymerase (Invitrogen).

#### Real-Time Quantitative PCR

The mRNA levels were determined by real-time quantitative PCR, with TaqMan universal PCR mastermix, transcript-specific minor groove binding (MGB) probes (Assays-on-Demand, Applied Biosystems), and an ABI 7900 sequence detection system, according to the manufacturer's instructions. The MGB probe for total *NTNG1* (detects all isoforms) was derived from exons 3 and 4 and that for *NTNG2* from exons 1 and 2. The probe specific to each *NTNG1* isoform is described in the corresponding sections.

The  $\beta$ 2-microglobulin gene was chosen as a control from the TaqMan Human Endogenous Control Plate (Applied Biosystems) after evaluation of 11 available internal control probes. The PCR assay was performed simultaneously with test and standard samples and no template controls in the same plate. A standard curve plotting the cycle of threshold values against input quantity (log scale) was constructed for both the  $\beta$ 2-microglobulin gene

**Table 2.** Demographic and Treatment Data for the Stanley Schizophrenic and Control Brains (BA46)

Sample ID	Age (y)	Gender	PMI (h)	pH	Cause of Death	Lifetime Antipsychotics <sup>a</sup>	Sample ID	Age (y)	Sex	PMI (h)	pH	Cause of Death
Schizophrenic Patients (n = 27)							Control Subjects (n = 27)					
S1	45	F	52	8.51	Sucide: jumped	20,000	C1	49	M	48	8.5	Cardiac disease
S2	40	M	34	6.18	Pneumonia	75,000	C2	53	M	9	6.4	Cardiac disease
S3	51	M	43	6.63	Cardiac disease	130,000	C3	37	M	13	6.5	Cardiac disease
S4	19	M	28	6.73	Overdose	2,500	C4	51	M	31	6.7	Cardiac disease
S5	53	F	13	8.49	Cardiac disease	15,000	C5	53	M	28	6	Cardiac disease
S6	37	M	30	6.8	Cardiac disease	20,000	C6	38	F	33	6	Cardiac disease
S7	52	M	10	8.1	Cardiac disease	100,000	C7	38	F	26	6.7	Cardiac disease
S8	24	M	15	8.2	Sucide: overdose	12,000	C8	60	M	47	6.6	Cardiac disease
S9	39	M	80	6.6	Motor vehicle accident	120,000	C9	35	M	62	6.7	Myocarditis
S10	33	M	29	6.5	Cardiac disease	20,000	C10	34	M	22	6.48	Cardiac disease
S11	50	M	9	6.2	Cardiac disease	34,000	C11	45	M	29	6.94	Cardiac disease
S12	43	M	18	6.3	Cirrhosis	90,000	C12	34	F	24	6.67	Cardiac disease
S13	32	F	38	6.6	Sucide: jumped	10,000	C13	42	M	37	6.91	Cardiac disease
S14	35	M	47	6.4	Cardiac disease	200,000	C14	44	F	10	6.2	Cardiac disease
S15	44	M	32	6.67	Cardiac disease	20,000	C15	57	M	26	6.4	Cancer
S16	47	M	13	6.3	Acute pancreatitis	300,000	C16	45	M	18	8.81	Cardiac disease
S17	45	M	35	6.66	Cardiac disease	50	C17	49	M	23	6.93	Cardiac disease
S18	36	F	27	8.49	Sucide: hanging	600	C18	49	F	45	6.72	Cardiac disease
S19	53	M	38	8.17	Cardiac disease	120,000	C19	33	F	29	6.52	Asthma
S20	54	F	42	6.65	Pneumonia	400,000	C20	48	M	31	6.60	Cardiac disease
S21	44	F	26	8.58	Pulmonary thrombosis	50,000	C21	50	M	49	6.75	Cardiac disease
S22	39	M	28	6.6	Sucide: hanging	48,000	C22	32	M	13	8.57	Cardiac disease
S23	38	M	35	6.68	Overdose	15,000	C23	47	M	11	8.8	Cardiac disease
S24	43	M	65	6.67	Sucide: hanging	70,000	C24	48	M	31	6.67	Cardiac disease
S25	42	M	19	6.48	Cardiac disease	18,000	C25	48	M	24	6.91	Cardiac disease
S26	46	M	30	6.72	Pneumonia	200,000	C26	39	F	58	8.48	Cardiac disease
S27	59	F	38	6.93	Cardiac disease	30,000	C27	47	M	38	8.57	Cardiac disease
Mean ±	42.3 ±		32.1 ±	8.53 ±		78,524 ±		44.8 ±		29.7 ±	8.81 ±	
SD	9.1		16.3	.23		97,532		7.6		13.3	.28	

BA, Brodmann's Area; PMI, postmortem interval; F, female; M, male.

<sup>a</sup>Lifetime neuroleptic dose in fluphenazine milligram equivalent dose.

and the target molecules (*NTNG1* and *NTNG2*) for each PCR assay. All real-time quantitative PCR data was captured with SDS v2.2 (Applied Biosystems). The ratio of the relative concentration of the target molecule to  $\beta 2$ -microglobulin gene (target molecule/ $\beta 2$ -microglobulin gene) was calculated. We used the Mann-Whitney *U* test (two-tailed) to detect significant changes in gene expression levels for each gene.

## Results

### Genomic Organizations and Chromosomal Localizations of *NTNG1* and *NTNG2*

The FISH analysis showed that human *NTNG1* and *NTNG2* map to chromosomes 1p13.3 and 9q34, respectively (Figure 1), in keeping with their database assignment, although the *NTNG2* probe showed weak signal at 12q24.3 in 20% of the cells. Genomic analysis demonstrated that the *NTNG1* spans 341 kilobases (kb) and is composed of 10 exons (Figure 2), whereas human *NTNG2* is encoded by eight exons and spans an interval of 82 kb (Figure 3). Both genes are located in schizophrenia linkage regions (Lewis et al 2003; Kaufman et al 1998).


### Isoform Variation of *NTNG1* Transcripts

Detection of human *NTNG1* mRNA is generally limited to brain and kidney by Northern analysis (Lin et al 2003). An initial assay of *NTNG1* splice products in human adult brain detected at least nine alternatively spliced transcripts, involving exons 5, 6, 7, 8, and 9 (Figure 4). The relative abundance of each isoform in


postmortem brain samples is shown in Figure 5A. The main isoforms in the brain were G1a, G1c, and G1d (Figure 5B).

### SNP Identification and Genetic Analyses

Our mutation screening and database search detected a total of 21 SNPs in *NTNG1* (2 novel SNPs in our mutation screening) (Figure




**Figure 1.** Fluorescence in situ hybridization of netrin-G1 (A) and netrin-G2 (B) sequences to normal metaphase human chromosomes. (A) The arrow denotes the positive signal at 1p13. (B) The thin arrows show the signal at 9q34, and the thick arrow shows weak signal at 12q24.




**Figure 2.** Genomic structure and location of polymorphic sites for human *NTNG1*. Exons are denoted by boxes, with untranslated regions in white and translated regions in black. The sizes of exons (bp) and introns (kb) are also shown. The rs number of each single nucleotide polymorphism (SNP) is the National Center for Biotechnology Information SNP cluster identification number from the dbSNP database (<http://www.ncbi.nlm.nih.gov/SNP/>).

2) and 10 SNPs in *NTNG2* (6 SNPs in our screening) (Figure 3). SNPs -400G>C (G1-02) and IVS5-43A>G (G1-15) in *NTNG1* and IVS4-15G>A (G2-09) in *NTNG2* were novel. No TaqMan typing probe could be designed for G2-09 in *NTNG2*. All other SNPs were unambiguously typed, with the exceptions of IVS4+2434A>G (G1-12) and Stop+1406C>T (G1-20) in *NTNG1*. The detailed information on these SNPs including heterozygosity is shown in Table 3. For subsequent genetic analyses, we chose 12 SNPs in *NTNG1* and 8 SNPs in *NTNG2*, on the basis of their relative abundance, with minor allele frequencies greater than .04 (Table 3).

Two different pairwise LD statistics, *D'* (normalized *D*) and *r*<sup>2</sup> (squared correlation coefficient) values, both of which measure LD values between 0 and 1, were computed between markers within the two genes, in 186 unrelated individuals from the present pedigree panel. Abecasis et al (2001) suggested a *D'* value greater than .33 as the minimum useful amount of LD, whereas Nakajima et al (2002) designated *r*<sup>2</sup> > .1 as the criterion for useful LD. In *NTNG1*, LD blocks with moderate strength exist in the intervals of G1-01 through to G1-03 and G1-11 through to G1-18 (Table 4). In *NTNG2*, intermediate LD measured by *D'*,



**Figure 3.** Genomic structure and locations of polymorphic sites for human *NTNG2*. Exons are denoted by boxes, with untranslated regions in white and translated regions in black. The sizes of exons (bp) and introns (kb) are also shown. The rs number of each single nucleotide polymorphism (SNP) is the National Center for Biotechnology Information SNP cluster identification number.



**Figure 4.** Splicing patterns of *NTNG1*. (A) All exons of *NTNG1*, with untranslated regions denoted in white and translated regions in color. (B) The nine different messenger ribonucleic acid isoforms.

was observed among the markers in the 5' region of the gene (G2-01~G2-06) (Table 5).

Table 6 shows the results of family-based association tests for the entire pedigree panel. The PDT program computes two statistical measures, PDT-sum and PDT-ave. Briefly, PDT-sum gives more weight to larger families, whereas PDT-ave places equal weight on all families. The suitability of both statistical methods depends on family structure and genetic models (Martin et al 2001). In *NTNG1*, the SNP G1-19 showed significant association with schizophrenia (nominal  $p = .02$ ) by PDT-sum. In *NTNG2*, the SNP G2-02 displayed significant association with schizophrenia with both PDT-sum ( $p = .008$ ) and PDT-ave ( $p = .014$ ). The SNPs G2-03 ( $p = .022$ ) and G2-06 ( $p = .045$ ) detected nominally significant associations by PDT-sum. After corrections for multiple SNP testing, the empirical  $p$  values for SNP transmission of *NTNG1* were .703 (PDT-ave) and .416 (PDT-sum), and those for *NTNG2* were .144 (PDT-ave) and .060 (PDT-sum). To

test the SNP associations in a more stringent manner, we performed ETD analysis on the 80 complete triad families. This detected a significant association between the G2-02 of *NTNG2* and schizophrenia ( $p = .045$ ) (Table 6).


The transmission of individual haplotypes (haplotypic  $p$ ) was assessed with the TRANSMIT program (Table 6). Significant distortions of haplotype transmission were observed for multiple haplotype blocks in *NTNG1* and *NTNG2* based on haplotypic  $p$  values, but the global  $p$  values were significant only with the (G1-14)~(G1-17)~(G1-18) haplotype of *NTNG1* ( $p = .049$ ) and with the G2-02~G2-03 haplotype of *NTNG2* ( $p = .049$ ). After Bonferroni correction for multiple testing, these transmission deviations were not significant. Further inspection of individual haplotypes revealed that the G-C-A haplotype (G1-14~G1-17~G1-18) of *NTNG1* was significantly less frequently transmitted ( $p = .002$ ) and the A-T haplotype (G2-02~G2-03) of *NTNG2* was excessively transmitted ( $p = .016$ ) to patients (Table 7), suggesting a protective role for the former and a risk factor for the latter.

#### Expression Analyses of *NTNG1* and *NTNG2* in Postmortem Brains

We quantified *NTNG1* and *NTNG2* mRNA expression levels in the two frontal regions BA11 (lateral orbitofrontal cortex) and BA46 (DLPFC) and one temporal region BA22 (superior temporal cortex) from schizophrenic and control postmortem brains, using MGB reactions. We analyzed six BA11 and six BA22 tissues each from schizophrenic patients and control subjects and 27 BA46 samples each from patients and control subjects. For *NTNG1*, transcript expressions were not significantly different between schizophrenic patients and control subjects in the three brain regions (Figure 6); nor were there significant differences in *NTNG2* expression levels between schizophrenic patients and normal subjects in the brain regions examined (Figure 6).

#### Expression Analysis of *NTNG1* Isoforms

The *NTNG1* expression assay designed to detect the common exons 3 and 4 showed no significant changes between schizophrenic and control brains (Figure 6); however, because the haplotype spanning the alternatively spliced exons in *NTNG1* and composed of genetic variants (G1-14, G1-17, and G-18)



**Figure 5.** Demographic examples of semiquantitative analysis of *NTNG1* isoforms by the Agilent 2100 bioanalyzer using the DNA1000 LabChip kit. (A) Each analysis was represented by one lane in the gel-like image. The brain samples were Brodmann's Area 46 from the Stanley Foundation brain collection. Each sample number corresponds to the "Sample ID" shown in Table 2. The 15-bp and 1500-bp size markers were included in all samples for calibration of size and total amount. (B) The analysis of the S25 sample is presented as an electropherogram.

**Table 3.** Allele Frequencies for SNP Markers Located Within *NTNG1* and *NTNG2*

Marker No.	Polymorphism <sup>a</sup>	Genome Location by UCSC (bp) <sup>b</sup>	Distance from Neighboring SNP (bp)	Minor Allele Frequency <sup>c</sup>	Heterozygosity <sup>d</sup>
<i>NTNG1</i>					
Chromosome 1p13.3					
G1-01	IVS1+2656T/C	107398460		.339	.448
G1-02	-400G/C	107402858	4398	.095	.171
G1-03	IVS2+1706T/C	107405209	2351	.289	.411
G1-04	IVS2+29544G/A	107433047	27838	.209	.330
G1-05	IVS2+81357A/C	107484861	51814	.048	.091
G1-06	IVS2-66691A/T	107512251	27390	0	0
G1-07	IVS2-48075A/G	107530867	18616	0	0
G1-08	IVS2-24065G/T	107554881	24014	.462	.497
G1-09	IVS2-6059G/A	107572887	18006	0	0
G1-10	IVS3-9907G/T	107639911	67024	0	0
G1-11	IVS3-179C/T	107649639	9728	.238	.363
G1-12	IVS4+2434A/G	107652424	2785	ND	
G1-13	IVS4-2418T/C	107659928	7504	.238	.363
G1-14	IVS5+2275T/G	107664647	4719	.238	.363
G1-15	IVS5-43A/G	107873201	8554	.035	.068
G1-16	IVS6+1905G/A	107675214	2013	.016	.032
G1-17	IVS8-1467C/T	107689862	14648	.347	.453
G1-18	IVS9+10464A/G	107701927	12085	.323	.437
G1-19	IVS9-4386C/T	107730889	28962	.194	.312
G1-20	Stop+1406C/T	107736910	6021	ND	
G1-21	Stop+6299C/A	107741803	4893	0	0
<i>NTNG2</i>					
Chromosome 9q34					
G2-01	IVS1-2180T/C	132069110		.054	.102
G2-02	IVS1-2128T/A	132069162	52	.173	.286
G2-03	IVS-1700G/T	132069590	428	.284	.407
G2-04	IVS2+149G/A	132072134	2544	.071	.132
G2-05	IVS2-13866G/C	132089041	16907	.116	.205
G2-06	738C/T	132103431	14390	.390	.476
G2-07	IVS3+13340T/G	132116890	13459	0	0
G2-08	876C/T	132131808	14918	.465	.498
G2-09	IVS4-15G>A	132135498	3690	ND	
G2-10	Stop+1085T/G	132148138	12640	.446	.494

SNP, single nucleotide polymorphism; ND, genotypes were not unambiguously determined.

<sup>a</sup>Second allele is a minor allele.

<sup>b</sup>Data on May, 2004; UCSC: University of California, Santa Cruz.

<sup>c</sup>Based on the genotype data from 186 unrelated subjects.

showed a preferential transmission to schizophrenic probands (global  $p = .049$ ) (Table 6), we examined the composition of mRNA isoforms in BA46 of postmortem brains in a semiquantitative manner, using the Agilent bioanalyzer system (Figure 5A).

Because semiquantitative analysis suggested differential expressions of the three main isoforms, G1a, G1c, and G1d, between schizophrenic and control brains (data not shown), we measured these transcripts in a more stringent manner, by design-

**Table 4.** Pairwise Marker-to-Marker LD Statistics of *NTNG1*

SNP Marker	G1-01	G1-02	G1-03	G1-04	G1-05	G1-08	G1-11	G1-13	G1-14	G1-17	G1-18	G1-19
G1-01		.939 <sup>a</sup>	.887 <sup>a</sup>	.386 <sup>a</sup>	.180	.046	.052	.052	.052	.039	.200	.084
G1-02	.189 <sup>a</sup>		1.000 <sup>a</sup>	.069	.873 <sup>a</sup>	.671 <sup>a</sup>	.036	.036	.036	.214	.114	.013
G1-03	.628 <sup>a</sup>	.260 <sup>a</sup>		.150	.193	.231	.055	.055	.055	.104	.074	.153
G1-04	.020	.002	.002		.067	.670 <sup>a</sup>	.006	.006	.008	.084	.020	.084
G1-05	.003	.004	.005	.001		1.000 <sup>a</sup>	.144	.144	.144	.429 <sup>a</sup>	.013	.217
G1-08	.001	.041	.019	.101	.042		.319 <sup>a</sup>	.319 <sup>a</sup>	.319 <sup>a</sup>	.057	.238	.083
G1-11	.000	.000	.000	.000	.003	.027		1.000 <sup>a</sup>	1.000 <sup>a</sup>	1.000 <sup>a</sup>	.901 <sup>a</sup>	.143
G1-13	.000	.000	.000	.000	.003	.027	1.000 <sup>a</sup>		1.000 <sup>a</sup>	1.000 <sup>a</sup>	.901 <sup>a</sup>	.143
G1-14	.000	.000	.000	.000	.003	.027	1.000 <sup>a</sup>	1.000 <sup>a</sup>		1.000 <sup>a</sup>	.901 <sup>a</sup>	.143
G1-17	.001	.002	.008	.001	.005	.002	.166 <sup>a</sup>	.166 <sup>a</sup>	.166 <sup>a</sup>		.960 <sup>a</sup>	.253
G1-18	.010	.001	.001	.000	.000	.031	.120 <sup>a</sup>	.120 <sup>a</sup>	.120 <sup>a</sup>	.233 <sup>a</sup>		.148
G1-19	.001	.000	.002	.006	.010	.002	.002	.002	.002	.008	.011	

For each pair of markers, the standardized  $D'$  is shown above the diagonal, and  $r^2$  is shown below the diagonal. LD, linkage disequilibrium; SNP, single nucleotide polymorphism.

<sup>a</sup> $D'$  values of  $>.3$  and  $r^2$  values of  $>.1$ .

**Table 5.** Pairwise Marker-to-Marker LD Statistics of *NTNG2*

SNP Marker	G2-01	G2-02	G2-03	G2-04	G2-05	G2-06	G2-08	G2-09
G2-01		1.000 <sup>a</sup>	1.000 <sup>a</sup>	.025	1.000 <sup>a</sup>	1.000 <sup>a</sup>	.661 <sup>a</sup>	.092
G2-02	.012		1.000 <sup>a</sup>	1.000 <sup>a</sup>	.049	.794 <sup>a</sup>	.119	.154
G2-03	.146 <sup>a</sup>	.506 <sup>a</sup>		.787 <sup>a</sup>	.448 <sup>a</sup>	.443 <sup>a</sup>	.116	.164
G2-04	.000	.017	.121 <sup>a</sup>		.046	.416 <sup>a</sup>	.280	.418 <sup>a</sup>
G2-05	.008	.002	.010	.000		.643 <sup>a</sup>	.080	.215
G2-06	.036	.199 <sup>a</sup>	.122 <sup>a</sup>	.020	.085		.136	.004
G2-08	.029	.003	.006	.005	.001	.014		.176
G2-10	.001	.004	.008	.011	.005	.000	.022	

For each pair of markers, the standardized  $D'$  is shown above the diagonal, and  $r^2$  is shown below the diagonal. LD, linkage disequilibrium; SNP, single nucleotide polymorphism.

<sup>a</sup> $D'$  values of  $>.3$  and  $r^2$  values of  $>.1$ .

**Table 6.** Transmission Analyses of Each SNP and Haplotype for *NTNG1* and *NTNG2* by PDT and ETDT

Genes and Polymorphisms	PDT		ETDT	TRANSMIT				
	SUM PDT	AVE PDT	Allele	Specific Haplotype $p$ (2 SNPs) <sup>a</sup> Global $p^c$		Specific haplotype $p$ (3 SNPs) <sup>b</sup> Global $p^c$		
<i>NTNG1</i>								
G1-01	.337	.168	.172	.162				
G1-02	.199	.156	.222	.371	.099		.177	
G1-03	.284	.247	.117		.216	.039 <sup>d</sup>	.492	.061
G1-04	1.000	.509	.504			.177	.181	.026 <sup>d</sup>
G1-05	.695	1.000	.667	.124			.402	.303
G1-08	.758	.992	.392	.256	.335		.207	.095
G1-11	.833	.626	.891		.795	.809	.563	.517
G1-13	.753	.542	1.000			.809	.809	.361
G1-14	.753	.542	1.000	.045 <sup>d</sup>			.809	.809
G1-17	.120	.172	.199	.112	.068		.002 <sup>a</sup>	.045 <sup>d</sup>
G1-18	.855	.647	.692		.289	.154	.049 <sup>d</sup>	.112
G1-19	.020 <sup>d</sup>	.088	.095			.284		
<i>NTNG2</i>								
G2-01	.655	.571	.761	.046 <sup>d</sup>				
G2-02	.008 <sup>d</sup>	.014 <sup>d</sup>	.045 <sup>d</sup>	.119	.018 <sup>d</sup>		.016 <sup>d</sup>	
G2-03	.022 <sup>d</sup>	.061	.264		.049 <sup>d</sup>	.062	.098	.016 <sup>d</sup>
G2-04	.748	.571	1.000	.056		.090		.057
G2-05	.121	.118	.076	.143	.019 <sup>d</sup>			.043 <sup>d</sup>
G2-06	.045 <sup>d</sup>	.055	.199		.088	.099	.035 <sup>d</sup>	.098
G2-08	.421	.749	.535	.235		.373	.362	.041 <sup>d</sup>
G2-10	.571	.724	.802	.664			.707	.068

SNP, single nucleotide polymorphism; PDT, pedigree disequilibrium test; ETDT, extended transmission DT.

<sup>a</sup>For two-marker analysis, a sliding window of two markers was tested, with one-marker overlap.


<sup>b</sup>For three-marker analysis, a sliding window of three markers was tested, with two-marker overlap.

<sup>c</sup>The global  $p$  values represent the overall significance when the observed versus expected frequencies of all of the haplotypes are considered together.

<sup>d</sup> $p < .05$ .

**Table 7.** Transmission of Individual *NTNG1* and *NTNG2* Haplotypes that Showed Global Significance in Transmission Analyses

Gene and Haplotype	Observed Transmissions	Expected Transmissions	Var (O-E)	$\chi^2$	p Value (1 df)
<i>NTNG1</i> (G1-14)-(G1-17)-(G1-18)					
G-C-A	7.0	15.3	7.0	9.741	.002 <sup>a</sup>
T-C-A	36.0	35.0	13.0	.077	.781
G-T-A	63.0	56.7	18.3	2.145	.143
G-C-G	51.0	50.5	16.7	.016	.900
T-C-G	1.0	1.0	.5	.000	.989
G-T-G	2.0	1.5	.8	.325	.569
<i>NTNG2</i> (G2-02)-(G2-03)					
T-G	107.0	111.9	15.9	1.518	.218
A-G	0	.5	.3	1.028	.311
T-T	18.0	20.6	6.8	.981	.322
A-T	35.0	27.0	11.0	5.817	.016 <sup>a</sup>

<sup>a</sup> $p < .05$ .**Figure 6.** Messenger ribonucleic acid levels for *NTNG1* and *NTNG2* in the indicated brain regions (Brodmann's Area [BA]11, BA22, and BA46) of control and schizophrenic subjects. Each data point represents the relative expression level against that of the internal control ( $\beta$ 2-microglobulin) for each sample. Horizontal bars indicate the mean value. BA11 and BA22 samples were from the Victorian Institute of Forensic Medicine in Australia ( $n = 6$  for each group) and BA 46 were obtained from the Stanley Foundation brain collection ( $n = 27$  for each group). The  $p$  values were calculated by Mann-Whitney  $U$  test (two-tailed).

ing isoform-specific PCR primers and TaqMan MGB probes (Figure 7). This quantitative analysis showed that the expression levels of G1c ( $p = .013$ ) and G1d ( $p = .015$ ) were significantly decreased in schizophrenic compared with control brains (Figure 7).


## Discussion

In contrast to the classical netrin genes, netrin-G1 and -G2 are specific for vertebrates. In situ hybridization studies in mice show that *Ntng1* and *Ntng2* expression overlaps in limited brain areas (Nakashiba et al 2000, 2002). Mouse *Ntng1* is expressed with greatest abundance in the dorsal thalamic nuclei and the inferior colliculus, followed by perforant pathways of the entorhinal cortex and the olfactory pathway in the piriform cortex. Strikingly, mouse *Ntng2* expression occurs widely in the cerebral cortex and in the habenular nucleus and superior colliculus but not in regions that show *Ntng1* positivity. Northern analysis of human tissue also indicates that in the brain, *Ntng1* is expressed maximally in thalamic regions (Lin et al 2003). This spatial dissociation is suggestive of a specific role for each gene in these areas, and the strong expression in the thalamus and cortical areas hint that the role of netrin-Gs could include maintaining neuronal plasticity associated with sensory and/or cognitive functioning (Nakashiba et al 2002).

Abnormalities in measures of attention and cognition have been observed in patients with schizophrenia. These deficits might be linked to impairments in sensorimotor gating, whereby intrusive, overwhelming stimuli are improperly filtered (Braff and Geyer 1990). Interestingly, *Ntng1* knockout mice exhibit a reduced level of prepulse inhibition (Nishimura et al 2004), a behavioral paradigm that examines sensory filtering function (Geyer et al 2001). Furthermore, in the same study of mice lacking *Ntng1*, animals displayed an augmented behavioral sensitization to repeated administration of a noncompetitive NMDA receptor antagonist, dizocilpine (MK-801), and mice devoid of *Ntng1* or *Ntng2* showed reduced NMDA receptor-mediated postsynaptic responses in electrophysiologic analysis of brain slices (Nishimura et al 2004 and unpublished data). These data demonstrate that the proteins Ntng1 and Ntng2 are imperative for NMDA receptor function, lending further support to the disturbance of NMDA neurotransmission hypothesis for schizophrenia.

Our association analysis of the *NTNG1* gene detected a nominal association of the most 3' variant G1-19 with schizophrenia. More importantly, the haplotype block constructed





**Figure 7.** Isoform-specific expression analyses of *NTNG1*. (A) The design of polymerase chain reaction primers and TaqMan minor groove binding probes to analyze the three main isoforms G1a, G1c, and G1d. (B) Results of quantitative analysis of *NTNG1* isoforms. Each data point represents the relative expression level of a splice variant against that of the internal control ( $\beta 2$ -microglobulin) for each brain sample. The data for failed amplifications are omitted. Horizontal bars indicate the mean value. The brain samples were Brodmann's Area 46 from the Stanley Foundation brain collection. The  $p$  values were calculated by the Mann-Whitney  $U$  test.

by G1-14~G1-17~G1-18, located in the 3' region of the gene showed a globally significant excess of transmission to the disease. These results suggest that a potential disease-causing variant(s) might reside in the 3' region of the gene. The mouse genomic structure of *Ntn1* is very similar to that of the human orthologue, as is the generation of multiple mRNA isoforms by alternative splicing (Nakashiba et al 2000). In humans, splicing involves the exons 5, 6, 7, 8, and 9. These exons are within the same LD block and overlap with the associated haplotype block. Therefore it is tempting to speculate that aberrant transcript processing might underpin this genetic association. In support of this speculation, splice variant netrin-G1c and -G1d distributions were significantly different between the DLPFC (BA46) of schizophrenic patients and control subjects. Although netrin-G1c and -G1d bind equally well to the NTNG1 ligand (NGL-1), the first molecule known to interact extracellularly with NTNG1 (Lin et al 2003; Nishimura et al, unpublished data), these isoforms might have distinct affinities for other unknown interacting molecules.

Both individual SNP and haplotype transmissions suggest a possible association between *NTNG2* and disease. The association signals were clustered within the same LD block located in the 5' region of the gene. These results imply that the risk variant(s) might influence the regulation of gene expression. Our quantification assay, however, did not detect significantly different expression of *NTNG2* mRNA in BA46 of schizophrenic and control brains. A larger number of brain samples need to be examined before firm conclusions regarding differential gene expression between groups (schizophrenic vs. control) and regarding haplotype-dependent expressional levels of both

*NTNG1* and *NTNG2* in each group can be drawn. Having stated this, it is practically and technically difficult to determine the exact haplotypes of genetically independent brain samples in the case of *NTNG1* (the SNPs G1-14, G1-17, and G1-18 of *NTNG1* span approximately 40 kb in the genome).

The anatomic significance of the frontal cortex in the pathogenesis of schizophrenia has long been recognized (Wong 2003). The DLPFC circuit involves the DLPFC in the cortex, the dorso-lateral caudate nucleus of the striatum, the lateral dorsal medial nucleus of the globus pallidus, the posterolateral nucleus of the substantia nigra, and the ventral anterior, medial dorsalis, pars parvocellularis subnuclei of the dorsal medial nucleus of the thalamus. Defects in any of these structures could produce alterations in intrinsic and extrinsic functional connectivity that might be related to the symptomology of schizophrenia (Bunney et al 2000). In this context, the present findings of a potential disturbance of at least *NTNG1* gene regulation at the transcriptional level might suggest a molecular contribution by netrin-G gene(s) to the disrupted higher-order brain functions in schizophrenia.

In conclusion, our data suggest a possible involvement of human *NTNG1* and *NTNG2* in the vulnerability to schizophrenia; however, to draw a robust conclusion regarding the disease-promoting role of these vertebrate-specific genes, further replication studies are warranted in independent samples. In addition, it would be interesting to include the gene for transmembrane protein NGL-1, which has recently been identified as a specific binding partner of NTNG1 (Lin et al 2003), in future genetic studies to determine the role of the netrin-G system in schizophrenia.

This work was supported by RIKEN BSI Funds, the RIKEN President Fund, Research on Brain Science Funds from the Ministry of Health Labor and Welfare, and CREST funds from the Japan Science and Technology Agency, Japan.

This article is dedicated to Dr. Masayuki Fukasawa (1973–2004).

- Abecasis GR, Noguchi E, Heinzmann A, Traherne JA, Bhattacharyya S, Leaves NI, et al (2001): Extent and distribution of linkage disequilibrium in three genomic regions. *Am J Hum Genet* 68:191–197.
- Braff DL, Geyer MA (1990): Sensorimotor gating and schizophrenia. Human and animal model studies. *Arch Gen Psychiatry* 47:181–188.
- Bunney WE, Bunney BG (2000): Evidence for a compromised dorsolateral prefrontal cortical parallel circuit in schizophrenia. *Brain Res Rev* 31:138–146.
- Chisholm A, Tessier-Lavigne M (1999): Conservation and divergence of axon guidance mechanisms. *Curr Opin Neurobiol* 9:603–615.
- Chua SE, Murray RM (1996): The neurodevelopmental theory of schizophrenia: Evidence concerning structure and neuropsychology. *Ann Med* 28:547–555.
- Clayton D (1999): A generalization of the transmission/disequilibrium test for uncertain-haplotype transmission. *Am J Hum Genet* 65:1170–1177.
- Clayton D, Jones H (1999): Transmission/disequilibrium tests for extended marker haplotypes. *Am J Hum Genet* 65:1161–1169.
- Dudbridge F (2003): Pedigree disequilibrium tests for multilocus haplotypes. *Genet Epidemiol* 25:115–121.
- Geyer MA, Krebs-Thomson K, Braff DL, Swerdlow NR (2001): Pharmacological studies of prepulse inhibition models of sensorimotor gating deficits in schizophrenia: A decade in review. *Psychopharmacology* 156:117–154.
- Ishii N, Wadsworth WG, Stern BD, Culotti JG, Hedgecock EM (1992): UNC-6, a laminin-related protein, guides cell and pioneer axon migrations in *C. elegans*. *Neuron* 9:873–881.
- Jessell TM (2000): Neuronal specification in the spinal cord: Inductive signals and transcriptional codes. *Nat Rev Genet* 1:20–29.
- Kaufman CA, Suarez B, Malaspina D, Pepple J, Svrakic D, Markel PD (1998): NIMH genetic initiative millennium schizophrenia consortium: Linkage analysis of African-American pedigrees. *Am J Med Genet* 81:282–289.
- Keks NA, Hill C, Opeskin K, Copolov DL, Dean B: Psychiatric diagnosis after death: The problems of accurate diagnosis from the case history review and relative interviews. In: Dean B, Kleinman JE, Hyde TM, editors. *Using CNS Tissue in Psychiatric Research*. Amsterdam: Harwood Academic Press, 19–37.
- Lewontin RC (1988): On measures of gametic disequilibrium. *Genetics* 120:849–852.
- Lewis CM, Levinson DF, Wise LH, DeLisi LE, Straub RE, Hovatta I, et al (2003): Genome scan meta-analysis of schizophrenia and bipolar disorder, part II: Schizophrenia. *Am J Hum Genet* 73:34–48.
- Lewis DA, Levitt P (2002): Schizophrenia as a disorder of neurodevelopment. *Ann Rev Neurosci* 25:409–432.
- Lewis DA, Lieberman JA (2000): Catching up on schizophrenia: Natural history and neurobiology. *Neuron* 28:325–334.
- Lin JC, Ho WH, Gurney A, Rosenthal A (2003): The netrin-G1 ligand NGL-1 promotes the outgrowth of thalamocortical axons. *Nat Neurosci* 6:1270–1276.
- Marengo S, Weinberger DR (2000): The neurodevelopmental hypothesis of schizophrenia: Following a trail of evidence from cradle to grave. *Dev Psychopathol* 12:501–527.
- Martin ER, Monks SA, Warren LL, Kaplan NL (2000): A test for linkage and association in general pedigrees: The pedigree disequilibrium test. *Am J Hum Genet* 67:146–154.
- Martin ER, Bass MP, Kaplan NL (2001): Correcting for a potential bias in the pedigree disequilibrium test. *Am J Hum Genet* 68:1065–1067.
- Nakajima T, Jorde LB, Ishigami T, Umemura S, Emi M, Lalouel JM, et al (2002): Nucleotide diversity and haplotype structure of the human angiotensinogen gene in two populations. *Am J Hum Genet* 70:108–123.
- Nakashiba T, Ikeda T, Nishimura S, Tashiro K, Honjo T, Culotti JG, et al (2000): Netrin-G1: A novel glycosyl phosphatidylinositol-linked mammalian netrin that is functionally divergent from classical netrins. *J Neurosci* 20:6540–6550.
- Nakashiba T, Nishimura S, Ikeda T, Itoharu S (2002): Complementary expression and neurite outgrowth activity of netrin-G subfamily members. *Mech Dev* 111:47–60.
- Nishimura S, Seki K, Watanabe A, Niimi K, Yoshikawa T, Nakashiba T, et al (2004): Axonal netrin-G1 regulates NMDA-receptor dependent synaptic functions in selected neural circuits. *Abstr Soc Neurosci* 723:19.
- O'Connell JR, Weeks DE (1998): PedCheck: A program for identification of genotype incompatibilities in linkage analysis. *Am J Hum Genet* 63:259–266.
- Serafini T, Kennedy TE, Galko MJ, Mirzayan C, Jessell TM, Tessier-Lavigne M (1994): The netrins define a family of axon outgrowth-promoting proteins homologous to *C. elegans* UNC-6. *Cell* 78:409–424.
- Sham PC, Curtis D (1995): An extended transmission/disequilibrium test (TDT) for multi-allele marker loci. *Ann Hum Genet* 59:323–336.
- Tessier-Lavigne M, Goodman CS (1996): The molecular biology of axon guidance. *Science* 274:1123–1133.
- Torrey EF, Webster M, Knable M, Johnston N, Yolken RH (2000): The Stanley Foundation brain collection and Neuropathology Consortium. *Schizophrenia Res* 44:151–155.
- Weinberger DR (1987): Implications of normal brain development for the pathogenesis of schizophrenia. *Arch Gen Psychiatry* 44:660–669.
- Wong AH, Van Tol HH (2003): Schizophrenia: From phenomenology to neurobiology. *Neurosci Biobehav Rev* 27:269–306.
- Yamada K, Iwayama-Shigeno Y, Yoshida Y, Toyota T, Itokawa M, Hattori E, et al (2004): Family-based association study of schizophrenia with 444 markers and analysis of a new susceptibility locus mapped to 11q13.3. *Am J Med Genet* 127B:11–19.
- Yin Y, Miner JH, Sanes JR (2002): Laminins: Laminin- and netrin-related genes expressed in distinct neuronal subsets. *Mol Cell Neurosci* 19:344–358.
- Zhao JH, Sham PC, Curtis D (1999): A program for the Monte Carlo evaluation of significance of the extended transmission/disequilibrium test. *Am J Hum Genet* 64:1484–1485.

**Appendix 1. PCR Primers Used to Search for Nucleotide Variants in *NTNG1* and *NTNG2***

Region	Exon Length (bp)	Primers	Product Size (bp)	3' End of Primer
<i>NTNG1</i>				
Exon 2	771	(F) 5'-TGCTTTATATTGCATCAGACCTC-3'	480	-603 (intron 1)
		(R) 5'-GACCTCAAAGCAGATCCCAAAA-3'		-167 (exon 2)
Exon 3	641	(F) 5'-AGTATGTTAGGCTTCCACCAA-3'	568	-218 (exon 2)
		(R) 5'-GTCTTCACACTCACCACATC-3'		+309 (intron 2)
Exon 4	173	(F) 5'-TAGGGCAAATAAAAATGA-3'	584	+175568 (intron 2)
		(R) 5'-AAAAACGCGAACCTGTC-3'		+176118 (exon 3)
Exon 5	27	(F) 5'-ACGAACATGGCAGCCCTATCAG-3'	529	+175966 (exon 3)
		(R) 5'-AATGCCTTCAGAACCTACT-3'		+178445 (intron 3)
Exon 6	66	(F) 5'-GGCCTGCAAATCTATCTCTACTA-3'	511	+246439 (intron 3)
		(R) 5'-GATGACTGAATTTTACTGAT-3'		+246858 (intron 4)
Exon 7	60	(F) 5'-TGCACCTGTATTTGTGTGTGTC-3'	283	+258962 (intron 4)
		(R) 5'-CCTATTACATCAGAAATGGACACT-3'		+259198 (intron 5)
Exon 8	168	(F) 5'-AATTGCTTGCTCTTGT-3'	347	+269832 (intron 5)
		(R) 5'-TTTCAAAGACATAGCATTTCAT-3'		+270142 (intron 6)
Exon 9	135	(F) 5'-CTTAATTTAGGGCTACTTTTCA-3'	254	+272404 (intron 6)
		(R) 5'-TACACTTCACAGATATCCAGATT-3'		+272813 (intron 7)
Exon 10	1404	(F) 5'-ATGCCATTCCACCGTCTTT-3'	406	+282063 (intron 7)
		(R) 5'-AGGATATTTTCTACATTGAG-3'		+282431 (intron 8)
Exon 11	135	(F) 5'-TCATTAATGGACATCTTT-3'	352	+287968 (intron 8)
		(R) 5'-GGATCTTTTCTGCTCTGA-3'		+288282 (intron 9)
Exon 12	1404	(F) 5'-GGCTGAAAACATGATGTACCAGATG-	453	+331947 (intron 9)
		(R) 5'-AGGCCTTCTTAGTTGTACTCTGTC-		+332351 (exon 10)
<i>NTNG2</i>				
Exon 2	698	(F) 5'-GTTTGCAAAGCTTCAGTGCTCG-3'	432	-133 (intron 1)
		(R) 5'-CAAGGATCTCCTCCAGACCTCCT-3'		+257 (intron 2)
Exon 3	644	(F) 5'-ACAGAGCAGGTTTCTCGGTTCG-3'	494	+31048 (intron 2)
		(R) 5'-GAGTACTCCTCGGTGACAGCA-3'		+31499 (exon 3)
Exon 4	173	(F) 5'-CTGGCAGCCCTACCAAGTTCTA-3'	469	+31421 (exon 3)
		(R) 5'-ACCAGGTAACATCCAGGTATC-3'		+31848 (intron 3)
Exon 5	24	(F) 5'-GGAATCAAGGAGTTTCTGGCCT-3'	418	+59870 (intron 3)
		(R) 5'-ATCCACAGCTCCTGGGATTGA-3'		+60248 (intron 4)
Exon 6	168	(F) 5'-CACCCTGTCCTCCACAC-3'	320	+63577 (intron 4)
		(R) 5'-GGCCCAAGTCCCAGAGG-3'		+63864 (intron 5)
Exon 7	135	(F) 5'-CCTGGGGTGAGTCCTTCC-3'	273	+72205 (intron 5)
		(R) 5'-TGCCGTGTCCGTGCCCTCTC-3'		+72435 (intron 8)
Exon 8	2659	(F) 5'-AACCGCTGCAGCTACATTGA-3'	247	+72316 (exon 6)
		(R) 5'-ACAGTCTCCCCAGGTGATTCT-3'		+72523 (intron 6)
Exon 9	135	(F) 5'-AGTCTTCTCTCCAGGCCTGGCCA-3'	444	+73900 (intron 6)
		(R) 5'-TCTTGCTAGGCCCAAGTCCACC-3'		+74301 (intron 7)
Exon 10	2659	(F) 5'-AGCCTCTACATCCCCGGCCCA-3'	489	+75766 (intron 7)
		(R) 5'-GCCCTGCTGAGTAGCACCTGGGA-3'		+757167 (exon 8)

Nucleotide positions are counted from A of the start codons on the genomic stretches of *NTNG1* (GenBank accession No. NM\_014917) and *NTNG2* (NM\_032536). F, forward; R, reverse.



## Extended analyses support the association of a functional (GT)<sub>n</sub> polymorphism in the *GRIN2A* promoter with Japanese schizophrenia

Yoshimi Iwayama-Shigeno<sup>a</sup>, Kazuo Yamada<sup>a</sup>, Masanari Itokawa<sup>a,b</sup>, Tomoko Toyota<sup>a</sup>, Joanne M.A. Meerabux<sup>a</sup>, Yoshio Minabe<sup>c</sup>, Norio Mori<sup>c</sup>, Toshio Inada<sup>d</sup>, Takeo Yoshikawa<sup>a,\*</sup>

<sup>a</sup> Laboratory for Molecular Psychiatry, RIKEN Brain Science Institute, Saitama, Japan

<sup>b</sup> Department of Schizophrenia Research, Tokyo Institute of Psychiatry, Tokyo, Japan

<sup>c</sup> Department of Psychiatry and Neurology, Hamamatsu University School of Medicine, Hamamatsu, Japan

<sup>d</sup> Department of Psychiatry and Psychobiology, Nagoya University Graduate School of Medicine, Nagoya, Japan

Received 5 November 2004; received in revised form 6 December 2004; accepted 7 December 2004

### Abstract

Dysfunction of the *N*-methyl-*D*-aspartate (NMDA) type glutamate receptor has been proposed as a mechanism in the etiology of schizophrenia. Recently, we identified a variable (GT)<sub>n</sub> repeat in the promoter region of the NMDA NR2A subunit gene (*GRIN2A*), and showed its association with schizophrenia in a case-control study, together with a correlation between the length of the repeat and severity of chronic outcome. In this study, we extended our analyses, by increasing the number of case-control samples to a total of 672 schizophrenics and 686 controls, and excluded potential sample stratification effects. We confirmed the significant allelic association between the repeat polymorphism and disease ( $P=0.011$ ), and as in the previous study, we observed an over-representation of longer alleles in schizophrenia. These results suggest a probable genetic effect for the *GRIN2A* promoter (GT)<sub>n</sub> variation on the predisposition to schizophrenia in Japanese cohorts.

© 2004 Elsevier Ireland Ltd. All rights reserved.

**Keywords:** NMDA receptor; NR2A subunit; Chromosome 16; Bipolar disorder; Linkage

Schizophrenia is a chronic and disabling brain disease of unknown etiology. Multiple genes with weak to moderate effects are thought to play a role in pathogenesis. A number of candidate genes for the disease have been proposed, of which, genes for the *N*-methyl-*D*-aspartate (NMDA) receptor are of special interest [19,22]. The NMDA receptor contains a non-competitive binding site for psychotomimetics, such as phenylcyclidine, MK-801 and ketamine. These antagonists trigger psychotic behavior, including both schizophrenia-like positive (delusions and hallucinations) and negative (affective blunting and specific cognitive defects) symptoms in normal subjects [10]. Conversely, the NMDA receptor co-agonists, such as glycine, *D*-serine and *D*-cycloserine, are reported to improve the negative symptoms of schizophrenia to varying degrees [6,11].

At least seven genes encoding NMDA receptor subunits have been identified in humans, NR1 (*GRIN1*), NR2A to D (*GRIN2A* to *D*) and NR3A and B (*GRIN3A* and *B*) [2]. Functionally distinct NMDA receptor subtypes are formed by heteromeric assembly of NR1 and NR2 subunits [16]. Expression of NR2A occurs predominately in the neocortices [13] and begins around puberty [20], corresponding to the period of schizophrenia onset. Mice lacking the  $\epsilon 1$  subunit gene, orthologous to human *GRIN2A*, exhibited phenotypes resembling both the positive and negative symptoms of schizophrenia [12]. In a previous study, we identified a variable (GT)<sub>n</sub> repeat in the 5'-regulatory region of the *GRIN2A*, and showed that the repeat sequence repressed transcriptional activity in a length-dependent manner, such that the longer the repeat, the greater the repression of promoter activity [8]. Case-control genetic analysis detected marginally significant association of the repeat polymorphism with schizophrenia in Japanese cohorts ( $P=0.05$ ), and showed that longer alle-

\* Corresponding author. Tel.: +81 48 467 5968; fax: +81 48 467 7462.  
E-mail address: [takeo@brain.riken.jp](mailto:takeo@brain.riken.jp) (T. Yoshikawa).