

which were apparently lower than those in the present study (Table 3). In general, PCB contamination in general population in Japan is in the moderate to higher levels among developed nations and somewhat higher than that in developing world. Our compilation and data analysis on worldwide human exposure in adipose tissues and liver also demonstrated this feature of contamination (Minh *et al.* 2000a, 2001).

The local and worldwide contamination pattern of OC insecticides in human blood was somewhat different from that of PCBs (Table 4). DDT residue levels in Miyako, Saku and Tottori were comparable to those reported in human blood samples from Sapporo, northern Japan (Takasuga *et al.* 2002). However, the levels were less than those in rural areas in Akita prefecture previously examined by Hanaoka *et al.* (2002). Sasaki *et al.* (1991) also reported elevated concentrations of DDTs in blood of residents from Okinawa in a survey conducted during 1986-88 (Table 4). Mean concentrations of DDTs in Okinawa subjects were 5.5 ng/ml (range 1.5 – 19 ng/ml), which was 4 times greater than that measured in this study for people in Miyako, a district in Okinawa prefecture (Table 4). Although the locations of samples are different between two studies, this can be considered as a hint of significant decline in human exposure to DDTs from Okinawa prefecture. Similar pattern was observed for HCHs and HCB, showing lower levels in the present study than those reported previously in other locations in Japan.

On a global scale, DDT contamination in human blood from Japan was less pronounced when compared to developing countries such as China, India, Egypt, Mexico and Brazil, where heavy usage of DDT for various agricultural and public health purpose have been common during the past four decades. Similar feature of contamination was also observed in adipose tissues and human milk, showing much higher DDT concentrations in populations from developing countries (Minh TB *et al.*, 2000a; Minh NH *et al.* 2004).

Interestingly, CHL concentrations in human from Miyako, Saku and Tottori, Japan were among the highest values reported for the countries examined. Chlordane was banned

relatively late (in 1986) in Japan as compared to other persistent OCs and therefore, high contamination of CHLs in Japanese humans still continues to be of considerable concern. Our previous time trend monitoring studies using archived adipose tissues revealed that CHL residues exhibited increasing trend until 1985 (Loganathan *et al.* 1993). Data from Hirai and Tomokuni (1991) and our previous investigation (Minh *et al.* 2001) showed higher concentrations of CHLs in Japanese adipose tissues examined in 1999 than those collected in 1989. The steady trend of CHL contamination in Japanese human samples during the last four decades reflect the remarkable increase in the usage of CHLs as termite control agent before it has been officially banned in 1986. Statistical data from the Ministry of the Environment, Japan showed a significant increase in the amount of CHL imported to Japan, and the import reached the maximum amount of approximately > 2000 tons in 1985 (Ministry of the Environment, Japan 2002). Therefore, although CHL was banned in Japan in 1986, elevated contamination of these compounds still remain in Japanese human samples.

Major route of human exposure to persistent OC compounds is via dietary intake (mainly fatty food, fish, meat and dairy product). Therefore, it is worthy to compare the residue concentrations of OCs in human tissues in relation to the dietary intake. A compilation of OC concentrations in human blood samples and dietary intake via foodstuffs of different countries is given in Figure 2. Data for dietary intake were cited from an extensive review article done by Kannan *et al.* (1997). Since dietary intakes can be substantially different among locations and time of investigations as a result of effective restriction and prohibition of OCs usage, data on blood concentrations and intakes reported at similar periods of time were cited for comparison as far as possible. In general, intakes of PCBs, DDTs and HCHs more or less reflect the contamination status in human blood samples (Figure 2). Some inconsistent results observed in intake and blood levels of OCs in Finland and United States (for PCBs, HCHs and DDTs), which could be due to the different foods and seafood consumption, or kinds of food used for estimating the dietary intakes. It can be emphasized that CHL compounds have lower dietary

intakes via foodstuffs. CHL compounds were used in the United States and Japan as termite control agent, and technical chlordane was sprayed directly to the households to eradicate termite. In the United States, it was estimated that over 80 million people lived in homes treated with CHLs and these compounds were detected in the indoor air of homes even 15 years after treatment (ATSDR 1994). In Japan, a previous occupational investigation revealed immediate increase in CHL residues in blood and sebum of residents of the treated houses and pest control operators (Wariishi and Nishimura 1989). Direct exposure via indoor air and house walls (inhalation or skin contact) is a major route of intake of CHLs into human body, in addition to the dietary intake via foods. These facts explain the differences in CHL residue concentrations in human blood and dietary intake estimated via foodstuffs as described in Figure 2.

Accumulation pattern and time trend of chlordane compounds in Japanese human blood samples

In this study, *p,p'*-DDE and β -HCH were predominant chemicals among DDT compounds and HCH isomers, respectively. In all the plasma samples analyzed, they account for > 90 % of the total DDT and HCH concentrations. This result is common in human samples and is similar to the pattern observed in adipose tissues from Tokyo, Japan, which we previously examined (Minh *et al.* 2001). Therefore, in this section, we examined the accumulation pattern of chlordane compounds in comparison with data from previous investigations to understand the time trend of contamination in Japanese human blood. Sasaki *et al.* (1991) in a 1986-88 survey reported CHL residues in human blood from Okinawa prefecture; and for this study, data are available for only three compounds, oxychlordane, *trans*- and *cis*-nonachlor. Wariishi *et al.* (1986) reported CHL residues in human blood from Tokushima, a small city located at the southern part of Japan (Shikoku island), and compared CHL exposure in subjects having no special exposure to CHLs, and those who were exposed to CHLs following termite control

treatment in their houses. Subsequent occupational study done by the same authors (Wariishi and Nishiyama 1989) provided an extensive investigation on the changes in CHL blood levels in pest control operators from Tokushima, who were occupationally exposed to elevated CHL concentrations. Residents from Okinawa, who were not occupationally exposed, are considered as controls in this investigation. Total residue levels of CHLs and proportions of the related compounds (oxychlordane, *trans*-chlordane, *cis*-chlordane, *trans*-nonachlor and *cis*-nonachlor) to total CHL concentrations reported in these investigations and in the present study were compared to understand the geographical differences in accumulation pattern, the possible implications of occupational exposure on the changes of accumulation pattern, and the time trend of CHLs in Okinawa residents during the last decade (Figure 3). Since lipid-normalized concentrations were not reported in the previous studies (Wariishi *et al.* 1986; Wariishi and Nishiyama 1989; Sasaki *et al.* 1991), concentrations on a wet weight basis were used for comparison. In general, oxychlordane and *trans*-nonachlor were predominant compounds. In Miyako, Okinawa, proportions of these chemicals were higher than those in Saku and Tottori (Figure 3). Similar pattern was noticed in Tokushima in an investigation during the period of 5 years after the CHL treatment for termite control (Figure 3) (Wariishi *et al.* 1986). Occupational analysis on subjects from Tokushima revealed substantial differences in residues concentrations and pattern (proportions to total CHL concentrations) in pest control operators (workers who are occupationally exposed to elevated CHL levels) and in those not occupationally exposed (controls). Proportions of oxychlordane in workers were much lower than those in controls (Wariishi and Nishiyama 1989). Oxychlordane is the terminal metabolic product of other chlordane compounds such as *trans*-chlordane and *cis*-chlordane (Wariishi *et al.* 1986). Immediate changes in the levels of *trans*-nonachlor and *trans*-chlordane in human blood were observed after termite control treatment and *trans*-nonachlor residues remained relatively constant even after 5 months from occupational exposure (Wariishi and Nishiyama

1989). This phenomenon can explain the higher proportions of *trans*-nonachlor and oxychlordane in high exposure groups.

Data on human plasma in Miyako analyzed in this study were compared to those in residents from Okinawa reported by Wariishi and Nishiyama (1989) and Sasaki *et al.* (1991) to discern the time trend of CHLs during the last decade (Figure 3). Residue concentrations of total CHLs declined substantially; about 2 times lower than in 1986-88 survey (Sasaki *et al.* 1991) and 10 times lower than in 1980-85 investigation (Wariishi and Nishiyama 1989). Although the backgrounds of samples and analytical methods of each study are different, results of this paper suggested that the CHL residue concentrations in human blood from Okinawa, Japan exhibited a declining trend during the last decade. Proportions of *trans*-chlordane and *cis*-chlordane, the two major components of technical chlordane, in 1999 survey (present study) were also apparently lower than those in the previous investigation (Wariishi and Nishiyama 1989). Loganathan *et al.* (1993) reported a relatively rapid increasing trend in CHL concentrations in human adipose tissues from Japan during 1955-1985, which is parallel with the remarkable increase in the amount of CHLs imported for agricultural usage and termite control. The amount of technical CHL imported to Japan for agricultural usage increased about 3 times during 1963-68. Subsequently, from 1979 a remarkable increase in import amount of CHL for termite control was noticed, reached a maximum level of > 2,000 tons in 1985 before it was officially banned in 1986 (Ministry of the Environment, Japan 2002). Thus, monitoring studies using human samples as bio-indicators clearly reflect the status of CHL usage and the effect of the ban of CHLs on the accumulation pattern in human blood.

PCB isomer pattern

Isomer-specific analysis of PCB revealed the presence of 60 congeners. CB-118, 138, 153 and 180 were predominant congeners, and this result is common for human samples. In females, we found that relative concentrations of higher chlorinated congeners such as hepta- and

octachlorobiphenyls were lower than those in males (although no statistical differences were noticed), suggesting a possibility of preferential elimination of higher chlorinated biphenyls in females (Figure 4). In contrast, proportions of lower chlorinated biphenyl congeners (tetra- and pentachlorobiphenyls) were higher in females than those in males. This male – female difference in PCB patterns was also observed in Dall's porpoises collected from off Pacific coasts of Japan (Subramanian *et al.* 1988), suggesting the similarity in accumulation kinetics of PCBs in higher trophic animals.

Age and sex dependent accumulation

To understand the accumulation kinetics of persistent OC compounds in healthy persons and diseased persons, we compared the age and gender differences in OC residue levels in the subjects examined in the present study and in autopsied adipose tissues of dead subjects who lived in Tokyo and nearby areas, which were analyzed in our previous study employing the same analytical methods (Minh *et al.* 2001) (Table 5 and 6, Figure 5 and 6). Although human tissues were different among two studies, it has been reported previously that blood residues indeed reflect adipose tissue concentrations, because significant correlations of OC concentrations in blood and adipose tissues were observed (review by Pelletier *et al.* 2003). In general, concentrations of OC compounds tended to increase with age in both males and females, which are similar to those found in human adipose tissue previously reported (Minh *et al.* 2001) (Figure 5 and 6). This result indicates long term accumulation of persistent and highly bio-accumulative OC compounds in humans despite phasing out of the usage of these chemicals in Japan more than 30 years ago. Similar age-dependent accumulation was also observed in human samples from other countries such as Netherlands (Greve and van Zoonen 1990), Spain (Ferrer *et al.* 1992), Italy (Focardi *et al.* 1986), Belgium (Voorspoels *et al.*, 2002) and Korea (Kang *et al.* 1997).

A Spearman rank correlation analysis revealed that PCB, DDT and TCPMe concentrations in blood from Miyako and in adipose tissues from Tokyo correlated significantly with age (Table 6). Interestingly, TCPMe showed the highest correlation with age in all the locations examined (Table 6), which further indicate high bio-accumulation potential of this compound in humans. This is quite consistent to the result observed for adipose tissue (Minh *et al.* 2001). Good positive correlation between concentrations and age suggest that body growth seems to be an important factor influencing the TCPMe burdens in humans. In marine mammals, it has been also suggested that differences in metabolic capacity among different species play a crucial role in the accumulation kinetics of TCPMe (Minh *et al.* 2000b).

Apparent gender differences were observed in samples from three locations, with higher concentrations in males than in female. A Mann-Whitney *U* test showed significant gender differences for PCBs in Miyako and Saku, for CHLs and TCPMe in Saku and Tottori (Table 5, Figure 5 and 6). In the previous study on sex-dependent accumulation in autopsied adipose tissues of patients, no apparent gender differences were noticed (Minh *et al.* 2001). Based on these different results, we suggest that accumulation kinetics of persistent OC compounds may be different among healthy and diseased persons. In diseased subjects, nutrient deficient conditions were observed with extremely low lipid content in adipose tissue, enhancing concentrations of OCs in the remaining lipids (Minh *et al.* 2001). Mobilization of stored contaminants from adipose tissue takes place during starvation or lactation and as a result of diseases (Thomas and Colborn 1992). It has been reported that increases in plasma OC concentrations significantly correlated with decreased in fat mass, suggesting that OCs are released into the blood during lipid mobilization (Pelletier *et al.* 2003). Bioaccumulation kinetics (age- and sex-dependent accumulation) of persistent OCs in human body is therefore strongly influenced by the lipid deficiency and weight loss, which is commonly observed in diseased persons.

Conclusions and Recommendations

To our knowledge, this is the first report on the specific accumulation of persistent OCs including TCPMe in human blood from Japan. Our result demonstrated elevated accumulation of PCBs and CHLs in Japanese humans, despite the ban on these chemicals for more than two decades in Japan. Given that limited data on OC residues in Japanese human blood, the result of this study can be a considerable contribution to the database of human exposure to persistent OCs in Japan. On the basis of the results of this study, we suggest that a systematic national survey on persistent organic pollutants (particularly PCBs, DDTs and CHLs) should be conducted by the Ministry of Health, Labor and Welfare, Japan.

Acknowledgements

This study was supported by the Health Sciences Research Grants of Ministry of Health, Labor and Welfare, Japan entitled "Research on Environmental Health - Human Exposure by Endocrine Disruptors in Japan". Financial support was also provided by a Grant-in-Aid from the Scientific Research Programs (Project Nos. 12308030 and 12055101), and the 21st Centre of Excellence Program (COE) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- Anderson HA, Falk C, Hanrahan L, Olson J, Burse VW, Needham L, Paschal D, Patterson D, Hill RHJ, *et al.* (1998) Profile of Great Lakes critical pollutants: a sentinel analysis of human blood and urine. *Environ Health Perspect* 106: 279-289
- ATSDR (1994) Toxicological Profile for Chlordane. Atlanta, GA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry. Available at: <http://www.atsdr.cdc.gov/toxprofiles/tp31-p.pdf>. Accessed: 2004-07-05
- ATSDR (2000) Toxicological Profile for Polychlorinated Biphenyls (PCBs). Atlanta, GA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic

Substances and Disease Registry. Available at:

<http://www.atsdr.cdc.gov/toxprofiles/tp17.html>. Accessed: 2004-07-07

ATSDR (2002) Toxicological Profile for DDT, DDE and DDD. Atlanta, GA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry. Available at: <http://www.atsdr.cdc.gov/toxprofiles/tp35.html>. Accessed: 2004-07-05

ATSDR (2003) Toxicological Profile for Hexachlorocyclohexanes (HCHs). Atlanta, GA. U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry. Available at: <http://www.atsdr.cdc.gov/toxprofiles/tp43.html>. Accessed: 2004-07-05

Colborn T, Voom Saal FS, Soto AM (1993) Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspec* 101:378-384.

De Voto E, Fiore BJ, Millikan R, Anderson HA, Sheldon L, Sonzogni WC, Longnecker MP (1997) Correlations among human blood levels of specific PCB congeners and implications for epidemiologic studies. *Am J Ind Med* 32: 606-613.

Dua VK, Kumari R, Sharma VP, Subbarao SK (2001) Organochlorine residues in human blood from Nainital (U.P.), India. *Bull Environ Contam Toxicol* 67: 42-45.

FAOSTAT Data (2004) Fish, Seafood: Food Balance Sheet (Year 2000). FAO Statistical Databases. Rome, Italy. Food and Agriculture Organization of the United Nations (FAO). Available at: <http://apps.fao.org/faostat/servlet/XteServlet3?OutputLanguage=english&Areas=ALL&Items=2960&Years=2000&Domain=FBS&ItemTypes=FBS&language=EN&Codes=ShowCodes>. Accessed: 2004-07-07

Ferrer A, Bona MA, Castellano M, To-Figueras J, Brunet M (1992) Organochlorine residues in human adipose tissue of the population of Zaragora (Spain). *Bull Environ Contam Toxicol* 48:561-566

- Focardi S, Fossi C, Leonzio C, Romei R. (1986) PCB congeners, hexachlorobenzene, and organochlorine insecticides in human fat in Italy. *Bull Environ Contam Toxicol* 36:644-650
- Gammon MD, Wolff MS, Neugut AI, Eng SM, Teitelbaum SL, Britton JA, Terry MB, Levin B, Stellman SD, Kabat GC, Hatch M, Senie R, Berkowitz G, Bradlow HL, Garbowski G, Maffeo C, Montalvan P, Kemeny M, Citron M, Schnabel F, Schuss A, Hajdu S, Vinceguerra V, Niguidula N, Ireland K, Santella RM (2002) Environmental toxins and breast cancer on Long Island. II. Organochlorine compound levels in blood. *Cancer Epidemiol Biomark Prev* 11: 686-697.
- Glynn AW, Wolk A, Aune M, Atuma S, Zettermark S, Mehle-Schmidt M, Darnerud PO, Becker W, Vessby B, Adami H (2000). Serum concentrations of organochlorines in men: a search for markers of exposure. *Sci Total Environ* 263:197-208.
- Glynn AW, Granath F, Aune M, Atuma S, Darnerud PO, Bjerselius R, Vainio H, Weiderpass E (2003) Organochlorines in Swedish women: determinants of serum concentrations. *Environ Health Perspect* 111: 349-355.
- Greve PA, van Zoonen P (1990) Organochlorine pesticides and PCBs in tissues from Dutch citizens (1968-1986). *Inter J Environ Anal Chem* 38: 265-277.
- Hanaoka T, Takahashi Y, Kobayashi M, Sasaki S, Usuda M, Okubo S, Hayashi M, Tsugane S. (2002) Residuals of beta-hexachlorocyclohexane, dichlorodiphenyltrichloroethane, and hexachlorobenzene in serum, and relations with consumption of dietary components in rural residents in Japan. *Sci Total Environ* 286:119-127.
- Hirai Y, Tomokuni K (1991) Levels of chlordane, oxychlordane, and nonachlor in human adipose tissues. *Bull Environ Contam Toxicol* 47:173-176.
- Hopper K, She J (2003) Lessons from the polybrominated diphenyl ethers (PBDEs): precautionary principle, primary prevention, and the value of community-based body-burden monitoring using breast milk. *Environ Health Perspect* 111: 109-114.

- Kang Y, Matsuda M, Kawano M, Wakimoto T, Min B (1997) Organochlorine pesticides, polychlorinated biphenyls, polychlorinated dibenzo-*p*-dioxins and dibenzofurans in human adipose tissue from western Kyungnam, Korea. *Chemosphere* 35:2107-2117.
- Kannan K, Tanabe S, Giesy JP, Tatsukawa R (1997) Organochlorine pesticides and polychlorinated biphenyls in foodstuffs from Asian and Oceanic countries. *Rev Environ Contam Toxicol* 152: 1-55.
- Kitamura K, Kikuchi Y, Watanabe S, Waechter G, Sakurai H, Takada T (2000) Health effects of chronic exposure to polychlorinated dibenzo-*p*-dioxins (PCDD), Dibenzofurans (PCDF) and coplanar PCB (Co-PCB) of municipal waste incinerator worker. *J Epidemiol* 10: 262-270.
- Koopman-Essenboom C, Huisman M, Weisglas-Kuperus N, Van der Paauw C, Tuinstra L, Boersma E, *et al.* (1994) PCB and dioxin levels in plasma and human milk of 418 Dutch women and their infants. Predictive value of PCB congener levels in maternal plasma for *fet al* and infant's exposure to PCBs and dioxins. *Chemosphere* 28: 1721-1732.
- Korrick SA, Chen C, Damokosh AI, Ni J, Liu X, Cho S, Altshul L, Ryan L, Xu X (2001) Association of DDT with spontaneous abortion: a case control study. *Ann Epidemiol* 11:491-496.
- Kunisue T, Someya M, Watanabe M, Toyota T, Kuroda Y, Nagayama J, Tanabe S (2004) Contamination by persistent organic pollutants (POPs) in human breast milk collected from primiparas in Ehime Prefecture [in Japanese]. *Environ Sci* 17:37-48.
- Laden F, Hankinson SE, Wolff MS, Colditz GA, Willett WC, Speizer FE, Hunter DJ (2001) Plasma organochlorine levels and the risk of breast cancer: an extended follow-up in the nurses health study. *Int J Cancer* 91: 568-574.
- Lascombe I, Beffa D, Ruegg U, Tarradellas J, Wahli W (2000) Estrogenic activity assessment of environmental chemicals using in vitro assays: identification of two new estrogenic compounds. *Environ Health Perspect* 108:621-629.

- Li YF (1999) Global technical hexachlorocyclohexane usage and its contamination consequences in the environment: from 1948 to 1997. *Sci Total Environ* 232:121-158.
- Loffler G, Bavel BV (2000) Potential pathways and exposure to explain the human body burden of organochlorine compounds: a multivariate statistical analysis of human monitoring in Wurzburg, Germany. *Chemosphere* 40: 1075-1082.
- Loganathan BG, Tanabe S, Hidaka Y, Kawano M, Hidaka H, Tatsukawa R (1993) Temporal trends of persistent organochlorine residues in human adipose tissue from Japan, 1928-1985. *Environ Pollut* 81:31-39.
- Ministry of the Environment, Government of Japan (2002) The Environmental Monitoring Report on the Persistent Organic Pollutants (POPs) in Japan. Tokyo, Japan: Ministry of the Environment, Government of Japan.
- Minh NH, Someya M, Minh TB, Kunisue T, Iwata H, Watanabe M, Tanabe S, Viet PH, Tuyen BC (2004) *Environ Pollut* 129:431-441.
- Minh TB, Watanabe M, Tanabe S, Yamada T, Hata J, Watanabe S (2000a) Occurrence of *tris*(4-chlorophenyl)methane, *tris*(4-chlorophenyl)methanol and other persistent organochlorines in Japanese human adipose tissue. *Environ Health Perspect* 108:599-603.
- Minh TB, Watanabe M, Tanabe S, Miyazaki N, Jefferson TA, Prudente M, Subramanian A, Karuppiyah S (2000b) Widespread contamination by *tris*(4-chlorophenyl)methane and *tris*(4-chlorophenyl)methanol in cetaceans from the North Pacific and Asian coastal waters. *Environ Pollut* 110:459-468.
- Minh TB, Watanabe M, Tanabe S, Yamada T, Hata J, Watanabe S (2001) Specific accumulation and elimination kinetics of *tris*(4-chlorophenyl)methane, *tris*(4-chlorophenyl)methanol and other persistent organochlorines in humans from Japan. *Environ Health Perspect* 109: 927-935.

- Miyako, Okinawa. Wikipedia Encyclopedia 2004. FL, USA. Wikimedia Foundation Inc.
Available at: http://en.wikipedia.org/wiki/Miyako_district%2C_Okinawa. Accessed:
2004-07-07.
- Monirith I, Nakata H, Watanabe S, Takahashi S, Tanabe S, Tana TS (2000) Organochlorine contamination in fish and mussels from Cambodia and other Asian countries. *Wat Sci Technol* 42: 241-252.
- Monirith I, Ueno D, Takahashi S, Nakata H, Sudaryanto A, Subramanian A, Karuppiyah S, Ismail A, Muchtar A, Zheng J, Richardson BJ, Prudente M, Hue ND, Tana TS, Tkalin AV, Tanabe S (2003) Asia-Pacific mussel watch: monitoring contamination of persistent organochlorine compounds in coastal waters of Asian countries. *Mar Pollut Bull* 46: 281-300.
- Moysich KB, Ambrosone CB, Mendole P, Kostyniak PL, Greizerstein HB, Vena JE, Menezes RJ, Swede H, Shields PG, Freudenheim JL (2002) *Am J Ins Med* 41: 102-110.
- North East Asian Regional (NEAR) Member, Tottori Prefecture (2002) Tottori, Japan: The Association of North East Asia Regional Government (NEAR). Available at: <http://www.near.khb.ru/eng/members/japan/tottori.html>. Accessed: 2004-07-06.
- Pauwels A, Covaci A, Weyler J, Delbeke L, Dhont M, De Sutter P, D'Hooghe T, Schepens PJC (2000) Comparison of persistent organic pollutant residues in serum and adipose tissue in female population in Belgium, 1996-1998. *Arch Environ Contam Toxicol* 39: 265-270.
- Pelletier C, Imbeault P, Tremblay A (2003) Energy balance and pollution by organochlorines and polychlorinated biphenyls. *Obesity Rev* 4: 17-24.
- Prudente MS, Tanabe S, Watanabe M, Subramanian A, Miyazaki N, Suarez P, Tatsukawa R (1997) Organochlorine contamination in some odontoceti species from the North Pacific and Indian Ocean. *Mar Environ Res* 44:415-427.

- Rahman MS, Montanarella L, Johansson B, Larson BR (1993) Trace levels of tris(4chlorophenyl)methanol and methane in human milk. *Chemosphere* 27:1487-1497.
- Rathore M, Bhatnagar P, Mathur D, Saxena GN (2002) Burden of organochlorine pesticides in blood and its effect on thyroid hormones in women. *Sci Total Environ* 295: 207-215.
- Sarcinelli PN, Pereira ACS, Mesquita SA, Oliveira-Silva J, Meyer A, Menezes MAC, Alves SR, Mattos RCOC, Moreira JC, Wolff M (2003) Dietary and reproductive determinants of plasma organochlorine levels in pregnant women in Rio de Janeiro. *Environ Res* 91: 143-150.
- Sasaki K, Ishizaka T, Suzuki T, Takeda M, Uchiyama M (1991) Accumulation levels of organochlorine pesticides in human adipose tissue and blood. *Bull Environ Contam Toxicol* 46: 662-669.
- Soliman AS, Wang X, DiGiovanni J, Eissa S, Morad M, Vulimiri S, Mahgoub KG, Johnston DA, Do KA, Seifeldin IA, Boffetta P, Bondy ML (2003) Serum organochlorine levels and history of lactation in Egypt. *Environ Res* 92: 110-117.
- Subramanian A, Tanabe S, Tatsukawa R (1988) Use of organochlorines as chemical tracers in determining some reproductive parameters in Dalli-type Dall's porpoise *Phocoenoides dalli*. *Mar Environ Res* 25:161-174.
- Takasuga T, Tsuji H, Nagayama J (2002) Gender specific dynamics of PCDD/DFs, PCBs, PBDEs and organochlorines in blood of Japanese families over two-year study period. *Organohalogen Comp* 58: 297-300.
- Thomas KR, Colborn T (1992) Organochlorine endocrine disruptors in human tissue. In: Colborn T, Clement C (eds) *Chemically Induced Alterations in Sexual and Functional Development: the Wildlife/Human Connection*. Princeton, Princeton Scientific Publishing Co., Inc., NJ, USA, p 365-394.

- Van Oostdam J, Gilman A, Dewailly E, Usher P, Wheatley B, Kuhnlein H, Neve S, Walker J, Tracy B, Feeley M, Jerome V, Kwavnick B (1999) Human health implications of environmental contaminants in Arctic Canada: a review. *Sci Total Environ* 230: 1-82.
- Van Oostdam J, Tremblay N. (2002) Biological monitoring: human tissue levels of environmental contaminants. In: AMAP Assessment 2002: Human Health in the Arctic. Arctic Monitoring and Assessment Programme (AMAP). Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, p 31-51.
- Voorspoels S, Covaci A, Maervoet J, Schepens P (2002) Relationship between age and levels of organochlorine contaminants in human serum of a Belgian population. *Bull Environ Contam Toxicol* 69: 22-29.
- Walizewski SM, Aguire AA, Infanzon RM, Silva CS, Siliceo J (2001) Organochlorine pesticide levels in maternal adipose tissue, maternal blood serum, umbilical blood serum and milk from inhabitants of Veracruz, Mexico. *Arch Environ Contam Toxicol* 40:432-438.
- Walker JB, Seddon L, McMullen E, Houseman J, Tofflemire K, Corriveau A, Weber J, Mills C, Smith S, Van Oostdam J (2003) Organochlorine levels in maternal and umbilical cord blood plasma in Arctic Canada. *Sci Total Environ* 302, 27-52.
- Wariishi M, Suzuki Y, Nishiyama K (1986) Chlordane residues in normal human blood. *Bull Environ Contam Toxicol* 36: 635-643.
- Wariishi M, Nishiyama K (1989) Observations on the progress of chlordane contamination in humans by blood and sebum analysis. *Arch Environ Contam Toxicol* 18: 501-507.
- Website Nagano Prefecture 2001. Nagano, Japan. Nagano Prefectural Government. Available at: <http://www.pref.nagano.jp/english/indexe.htm#e03>. Accessed: 2004-07-07
- Zhao YF, Wu YN, Wang XQ, Gao JQ, Chen JS (2003) Study of dietary pesticide residues in Chinese residents [in Chinese, Abstract]. *Zhonghua Liu Xing Binh Xue Za Zhi* 24: 661-664.

Table 1. Concentrations (ng/g lipid wt) of persistent organochlorines in human blood from Miyako, Saku and Tottori, Japan.

Sample Code	Sex	Age	PCBs	DDTs	HCHs	CHLs	HCB	TCFMs
Miyako (n = 23)								
210	M	27	160	100	16	27	6.0	< 0.1
243	M	28	180	88	13	28	4.6	< 0.1
203	M	30	230	85	16	55	8.5	2.0
247	M	38	440	190	33	60	8.2	2.6
230	M	41	620	1400	95	150	20	8.1
250	M	43	240	56	20	70	7.5	5.7
204	M	46	240	620	72	70	8.3	< 0.1
206	M	46	430	500	85	150	12	< 0.1
219	M	49	250	110	7.6	24	4.8	4.7
225	M	51	340	230	39	59	8.7	4.7
205	M	55	380	990	20	92	6.6	< 0.1
229	M	57	340	350	120	77	5.8	5.6
221	M	58	360	280	24	43	5.2	5.9
216	F	24	200	230	23	25	7.1	< 0.1
208	F	25	120	76	22	150	8.9	0.7
246	F	25	180	80	17	34	6.5	1.6
245	F	30	170	120	40	53	8.1	2.1
226	F	31	160	73	36	58	7.8	1.8
234	F	35	320	150	65	74	11	5.9
254	F	38	290	240	28	29	11	5.1
244	F	42	290	400	24	77	7.2	5.4
222	F	49	170	350	32	99	6.1	3.5
217	F	52	360	300	130	110	12	6.7
Mean ^a		40	280	310	43	70	8.3	3.2
Median		41	250	238	28	60	7.8	3.1
Geometric mean		38	268	218	32	60	7.9	1.4
Range		24 - 58	120 - 620	56 - 1400	7.6 - 130	24 - 150	4.6 - 20	< 0.1 - 8.1
Saku (n = 32)								
335	M	30	240	64	35	13	5.1	0.82
301	M	34	580	130	15	22	15	1.8
329	M	39	390	79	1.7	23	6.2	0.72
332	M	40	350	300	27	31	14	2.5
345	M	40	350	390	22	65	7.9	2.3
362	M	46	720	360	37	47	4.6	3.4
316	M	51	730	900	68	68	32	3.9
323	M	51	780	180	0.81	29	6.7	2.3
328	M	52	440	270	23	36	11	2.1
343	M	52	590	440	32	34	18	3.7
339	M	53	480	460	37	55	24	3.7
360	M	55	530	150	100	44	20	2.4
348	M	57	210	35	32	16	3.3	2.1
327	M	59	450	370	41	25	9.1	2.6
308	M	59	990	2700	64	79	27	6.2
306	F	28	260	230	44	17	14	0.62
356	F	32	160	140	17	4.8	4.1	< 0.1
349	F	35	440	840	49	26	12	1.2
359	F	37	220	75	28	20	8.3	1.1
312	F	43	270	210	54	21	14	1.2
366	F	48	560	1300	53	42	13	2.0
338	F	51	270	230	35	19	14	1.4
310	F	52	310	130	36	31	15	0.54
321	F	53	250	320	35	23	15	3.4
342	F	53	290	130	22	19	17	1.9
354	F	54	240	180	33	21	13	0.64
358	F	54	140	310	22	21	11	0.56
309	F	54	260	290	24	26	19	1.9
367	F	56	320	64	22	47	3.3	1.5
313	F	57	310	200	39	34	5.8	1.3
303	F	58	450	190	64	35	8.2	1.5
317	F	59	320	32	30	26	< 1.0	4.5
Mean		48	400	370	36	32	12	2.1
Median		51	340	228	34	26	13	1.9
Geometric mean		47	360	220	28	28	10	1.6
Range		28 - 59	140 - 990	32 - 2700	0.81 - 100	4.8 - 79	< 1.0 - 32	< 0.1 - 6.2
Tottori (n = 22)								
470	M	30	200	34	9.8	24	5.1	0.42
416	M	37	250	96	24	32	9.5	1.2
420	M	45	650	210	37	49	16	3.8
407	M	47	560	250	95	46	12	1.8
431	M	48	350	200	26	33	11	2.1
405	M	52	420	85	26	38	14	2.6
408	M	56	300	200	38	34	10	2.9
469	M	56	520	420	42	68	8.9	3.4
450	M	57	550	190	69	24	11	4.9
429	M	59	780	440	46	67	13	4.2
471	F	30	220	50	17	12	5.9	1.4
454	F	35	480	380	64	30	12	1.8
418	F	39	230	130	41	21	11	2.2
440	F	41	240	89	9.2	14	7.6	0.9
446	F	43	270	70	34	44	6.4	4.8
426	F	45	490	490	32	28	10	2.8
438	F	48	200	140	24	16	8.2	2.1
457	F	49	290	210	32	27	9.5	1.9
451	F	49	400	75	34	17	9.1	1.7
449	F	50	510	350	44	28	13	2.0
458	F	55	400	230	42	19	9.5	0.9
436	F	59	210	190	28	18	8.1	2.7
Mean		47	398	210	37	31	10	2.4
Median		48	388	200	34	28	9.8	2.1
Geometric mean		46	360	160	32	28	9.7	2.1
Range		30 - 59	200 - 780	34 - 490	9.2 - 95	12 - 68	5.1 - 16	0.42 - 4.9

^aConcentration values below the detection limit were assigned as equal to detection limit for calculation of mean, median and geometric mean.

Table 2. Production and usage of persistent organochlorine compounds in Japan

Organochlorine compound	Approximate amount of production/usage (tons)	Year of starting usage/production	Year of ban	Source of information/Reference
PCBs	58,787/54,001	early 1950s	1972	Ministry of the Environment, Japan (2002)
DDTs	45,000	1945	1971	Ministry of the Environment, Japan (2002)
HCHs	400,000	1950	1972	Li (1999)
CHLs	> 2,000	1955	1986	Ministry of the Environment, Japan (2002)
HCB	4,000 (peak annual demand)	NA ^a	1979	Ministry of the Environment, Japan (2002)

^aNA: no data available.

Table 3. Comparison of the concentrations of polychlorinated biphenyl congeners (PCBs) in human blood from different locations in the world.*

Location	Year	Sex	Age (year)	n	Tissue	Unit	CB-118	CB-138	CB-153	CB-180	Total PCBs	Reference	Remark
Miyako, Okinawa, Japan	1999	M&F ^b	24 - 58	23	plasma	ng/g lipid	12 (4.3 - 39) ^c	33 (21 - 140)	50 (21 - 140)	43 (13 - 99)	280 (120 - 620)	Present study	Total PCBs: sum of 60 congeners
						ng/ml	0.048 (0.017 - 0.16)	0.13 (0.084 - 0.56)	0.2 (0.084 - 0.56)	0.17 (0.052 - 0.40)	1.1 (0.48 - 2.5)		
Saku, Nagano, Japan	1999	M&F	28 - 59	32	plasma	ng/g lipid	23 (4.3 - 65)	53 (11 - 150)	77 (17 - 210)	55 (11 - 150)	400 (140 - 990)	Present study	Total PCBs: sum of 60 congeners
						ng/ml	0.09 (0.017 - 0.25)	0.2 (0.043 - 0.59)	0.3 (0.067 - 0.84)	0.21 (0.043 - 0.59)	1.6 (0.55 - 3.9)		
Tottori, Tottori, Japan	1999	M&F	30 - 59	22	plasma	ng/g lipid	20 (9.1 - 50)	50 (20 - 100)	80 (37 - 180)	66 (33 - 150)	390 (200 - 780)	Present study	Total PCBs: sum of 60 congeners
						ng/ml	0.08 (0.036 - 0.2)	0.2 (0.08 - 0.4)	0.32 (0.15 - 0.72)	0.26 (0.13 - 0.6)	1.6 (0.8 - 3.1)		
Sapporo, Japan	NA	M&F	37-48	18	whole blood	ng/g lipid	NA ^d	NA	NA	NA	270 (62 - 970)	Takasuga et al. (2002)	NA
Anqing, China	1998	F	25 ± 1.2 ^e	15	serum	ng/ml	NA	NA	NA	NA	0.2 ± 0.1	Korrick et al. (2001)	Total PCBs: sum of 67 congeners
Great Lakes Region, US	1993	M&F	36 - 76	30	serum	ng/ml	0.4 (0.02 - 1.1)	0.8 (0.1 - 2.2)	1.1 (0.3 - 2.9)	0.4 (0.09 - 1.0)	5.2 (1.2 - 15)	Anderson et al. (1998)	Total PCBs: sum of 30 congeners
New York, US	1991	F	45 - 85	192	serum	ng/ml	NA	NA	NA	NA	4.1 ± 2.2	Moysich et al. (2002)	Total PCBs: sum of 69 congeners
Long Island, NY, US	1996-97	F	> 20	1556	serum	ng/g lipid	56 ± 2.0	85 ± 2.5	150 ± 1.7	79 ± 1.8	370 ± 1.7	Grammon et al. (2002)	Total PCBs: sum of 4 congeners
US, 11 states	1989-90	F	30-55	370	plasma	ng/g lipid	68	97	110	75	540	Laden et al. (2001)	Total PCBs: sum of 16 congeners
North Carolina, US	1993-1995	F	28-74	68	plasma	ng/ml	0.34	0.44	0.49	0.34	1.4	DeVoto et al. (1997)	Total PCBs: sum of 26 congeners
Wisconsin, US	1991	M&F	20-71	100	serum	ng/ml	0.62	1.4	1.7	1	6.9	DeVoto et al. (1997)	Total PCBs: sum of 26 congeners
Canadian Arctic	1994-99	F	15 - 45	385	plasma	ng/g lipid	8.6 (<2.2 - 71)	27 (<2.2 - 350)	48 (2.2 - 890)	23 (2.2 - 450)	140 (12 - 2300)	Walker et al. (2003)	Total PCBs: sum of 14 congeners
Greenland Arctic	1997	M	19-60	15	plasma	ng/ml	1.0 (0.18 - 4.3)	6.0 (1.0 - 19)	14 (2.4 - 44)	9.6 (1.4 - 31)	41 (6.6 - 130)	Van Oostdam & Tremblay (2003)	Total PCBs: sum of 14 congeners
Alaskan Arctic	2001	M&F	18-91	166	serum	ng/ml	NA	NA	NA	NA	7.7 (nd ^f - 54)	Van Oostdam & Tremblay (2003)	Total PCBs: sum of 36 congeners
Russian Arctic	1999	F	NA	30	plasma	ng/ml	0.29 (0.1 - 1.0)	0.42 (0.12 - 1.5)	0.39 (0.11 - 1.7)	0.13 (0.04 - 1.0)	1.7 (0.6 - 6.8)	Van Oostdam & Tremblay (2003)	Total PCBs: sum of 13 congeners
Norwegian Arctic	1999	F	NA	50	plasma	ng/ml	0.11 (0.01 - 0.67)	0.79 (0.24 - 2.3)	0.47 (0.17 - 1.2)	0.4 (0.13 - 0.84)	2.3 (0.8 - 6.2)	Van Oostdam & Tremblay (2003)	Total PCBs: sum of 11 congeners
Finland	1996-98	F	NA	13	plasma	ng/ml	0.07 (0.04 - 0.11)	0.3 (0.15 - 0.4)	0.43 (0.21 - 0.61)	0.24 (0.15 - 0.33)	1.5 (0.8 - 2.0)	Van Oostdam & Tremblay (2003)	Total PCBs: sum of 14 congeners
Sweden, central region	NA	M	40 - 74	120	serum	ng/g lipid	42 (4.3 - 140)	140 (3.1 - 340)	300 (23 - 630)	220 (71 - 480)	750 (150 - 1500)	Glynn et al. (2000)	Total PCBs: sum of 10 congeners
Sweden	1996-97	F	50 - 74	205	serum	ng/g lipid	43 (5.0 - 180)	100 (18 - 270)	220 (60 - 610)	150 (56 - 400)	550 (150 - 1600)	Glynn et al. (2003)	Total PCBs: sum of 7 congeners
Netherlands	1990-92	F	NA	418	plasma	ng/ml	0.16	0.60	0.91	0.54	2.2	Koopman-Esseboom et al. (1994)	Total PCBs: sum of 4 congeners
Flanders, Belgium	1996-98	F	24-42	96	serum	ng/g lipid	27 ± 16	71 ± 33	91 ± 39	70 ± 31	260	Pauwels et al. (2000)	Total PCBs: sum of 4 congeners
Belgium	2001	M&F	20-54	132	serum	ng/g lipid	13	64	99	60	240	Voorspoels et al. (2002)	Total PCBs: sum of 4 congeners
Wurzburg, Germany	1992-95	M&F	NA	309	whole blood	ng/ml	NA	0.39 ± 0.22	0.49 ± 0.29	0.24 ± 0.16	1.2	Löffler et al. (2000)	Total PCBs: sum of 4 congeners

*Cited values were rounded to 2 significant digits for comparison. ^{NA}, Male, F, Female. ^cRange concentration. ^dNA, no data available. ^eMean ± standard deviation (SD). ^fnd, not detected; detection limit was not reported.

Table 4. Comparison of organochlorine insecticide residue levels in human blood from various locations in the world.^a

Location	Year	Sex	Age (year)	n	Tissue	Unit	DDTs	HCHs	CHLs	HCB	Ref. ^b	Remark
Miyako, Okinawa, Japan	1999	M&F	24-58	23	plasma	ng/g lipid	310 (56-1400) ^c	43 (7.6-130)	70 (24-150)	8.2 (4.6-12)	(1)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: α -, β - and γ -HCH CHL: trans- and cis-nonachlor, trans- and cis-chlordane, and oxychlorane
Sakan, Nagano, Japan	1999	M&F	28-59	32	plasma	ng/ml	1.3 (0.22-5.6)	0.17 (0.03-5.2)	0.28 (0.096-0.6)	0.033 (0.018-0.048)	(1)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: α -, β - and γ -HCH CHL: trans- and cis-nonachlor, trans- and cis-chlordane, and oxychlorane
Tottori, Tottori, Japan	1999	M&F	30-59	22	plasma	ng/ml	370 (32-2700)	36 (0.81-100)	32 (4.8-79)	12 (<1.0-32)	(1)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: α -, β - and γ -HCH CHL: trans- and cis-nonachlor, trans- and cis-chlordane, and oxychlorane
Sapporo, Japan	NA	M&F	37-48	18	whole blood	ng/g lipid	1.4 (0.12-11)	0.14 (0.0032-0.39)	0.12 (0.019-0.31)	0.047 (<0.004-0.12)	(1)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: α -, β - and γ -HCH CHL: trans- and cis-nonachlor, trans- and cis-chlordane, and oxychlorane
Akita, Japan	1999	M&F	41-69	41	serum	ng/ml	210 (34-490)	37 (9.2-95)	3.0 (1.2-6.8)	10 (5.1-16)	(1)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: α -, β - and γ -HCH CHL: trans- and cis-nonachlor, trans- and cis-chlordane, and oxychlorane
Akita, Japan	1986-88	M&F	54-78	12	whole blood	ng/ml	0.84 (0.14-2.0)	0.15 (0.037-0.38)	0.12 (0.05-0.27)	0.04 (0.02-0.064)	(2)	NA
Okawa, Japan	1986-88	M&F	19-67	11	whole blood	ng/ml	350 (51-1600)	140 (14-710)	43 (7.6-150)	18 (2.7-120)	(3)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: β -HCH only
Anqing, China	1998	F	25 \pm 1.2 ^f	15	serum	ng/ml	3000 (430-15000)	260 (24-710)	NA ^g	110 (9.5-330)	(3)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: β -HCH only
Japur city, India	1997-98	F	11-50	100	whole blood	ng/ml	6.3 (0.9-31)	0.54 (0.05-1.5)	NA	0.24 (0.02-0.7)	(4)	DDT: p,p'-DDE only; HCH: β -HCH only
Nainital, India	2000	M&F	18-50	39	whole blood	ng/ml	4.3 (1.9-8.2)	2.2 (0.8-5.5)	0.30 (0.15-0.96)	NA	(4)	CHL: trans- and cis-nonachlor, heptachlor and oxychlorane
Egypt, rural area	1997-98	F	18-44	60	serum	ng/ml	5.5 (1.5-19)	1.8 (0.7-4.2)	0.88 (0.37-2.8)	0.3 \pm 0.1	(5)	DDT: p,p'-DDE, p,p'-DDD, p,p'-DDT, o,p'-DDE and o,p'-DDT; CHL: oxychlorane only
Egypt, urban area	1997-98	F	18-44	60	serum	ng/ml	13 \pm 6	NA	0.01 \pm 0.01	0.3 \pm 0.1	(5)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; CHL: oxychlorane only
Rio de Janeiro, Brazil	1997	F	16-49	64	plasma	ng/ml	6900 \pm 550	3900 \pm 350	1400 \pm 150	NA	(6)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: α -, β - and γ -HCH; CHL: heptachlor only
Great Lakes Region, US	1993	M&F	36-76	30	serum	ng/ml	4600 (1000-13000)	1300 (250-7500)	NA	NA	(7)	DDT: p,p'-DDE, p,p'-DDD, p,p'-DDT and o,p'-DDT; HCH: α -, β - and γ -HCH
New York, US	1991	F	45-85	192	serum	ng/ml	18 (0.3-140)	2.6 (0.3-18)	NA	NA	(8)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: β -HCH only
Canadian Arctic	1994-99	F	15-45	385	plasma	ng/g lipid	9.9 (0.7-62)	1.3 (0.3-9.5)	NA	NA	(8)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: β -HCH only
Greenland Arctic	1997	M	19-60	15	plasma	ng/ml	16 \pm 31	1.6 \pm 2.0	NA	1.0 \pm 0.8	(9)	DDT: p,p'-DDE, p,p'-DDD, p,p'-DDT, and o,p'-DDT; HCH: α -, β - and γ -HCH
Alaskan Arctic	2001	M&F	18-91	166	serum	ng/ml	3.0 (0.52-18)	1.0 (0.22-5.1)	0.18 (0.015-0.75)	0.084 (0.008-0.0585)	(10)	DDT: p,p'-DDE, p,p'-DDD and p,p'-DDT; HCH: α -, β - and γ -HCH; CHL: heptachlor only
Russian Arctic	1995-96	F	NA	18	plasma	ng/ml	5.6 (0.68-25)	0.05 (0.04-1.2)	1.1 (0.16-3.6)	0.1 (0.02-0.2)	(11)	DDT: p,p'-DDE, p,p'-DDT and o,p'-DDT; HCH: β -HCH only; CHL: oxychlorane, heptachlor epoxide, trans- and cis-nonachlor
Norwegian Arctic	1999	F	NA	50	plasma	ng/ml	11 \pm 10	NA	NA	0.42 \pm 0.19	(12)	DDT: p,p'-DDE only
Finland	1996-98	F	NA	13	plasma	ng/ml	200 (13-4100)	25 (<4.3-4200)	66 (8.6-1200)	38 (2.2-490)	(13)	DDT: p,p'-DDE and p,p'-DDT; HCH: β -HCH only; CHL: α -, γ -chlordane, trans-, cis-nonachlor, heptachlor epoxide, and oxychlorane
Sweden	1996-97	F	50-74	205	serum	ng/g lipid	11 (1.6-33)	0.56 (0.17-1.8)	6.9 (0.81-65)	1.5 (0.4-3.9)	(14)	DDT: p,p'-DDE and p,p'-DDT; HCH: β -HCH only; CHL: oxychlorane and trans-nonachlor
Sweden, central region	NA	M	40-74	120	serum	ng/g lipid	14 (nd ^h -160)	0.35 (nd-2.6)	2.4 (nd-18)	NA	(14)	DDT: p,p'-DDE and p,p'-DDT; HCH: β -HCH only; CHL: oxychlorane, trans-nonachlor and heptachlor epoxide
Whurzburg, Germany	1992-95	M&F	NA	309	whole blood	ng/ml	1.2 (0.43-2.1)	0.35 (0.02-0.76)	0.09 (0.04-0.15)	0.2 (0.02-0.71)	(14)	DDT: p,p'-DDE and p,p'-DDT; HCH: β -HCH only; CHL: trans-nonachlor only
Flanders, Belgium	1996-98	F	24-42	101	serum	ng/g lipid	0.95 (0.20-5.1)	0.05 (0.02-0.36)	0.14 (0.06-0.43)	NA	(14)	DDT: p,p'-DDE only; HCH: β -HCH only; CHL: oxychlorane and trans-nonachlor
							0.6 (0.21-0.83)	0.07 (0.02-0.09)	0.08 (0.04-0.12)	0.19 (0.12-0.31)	(14)	DDT: p,p'-DDE and p,p'-DDT; HCH: β -HCH only; CHL: oxychlorane and trans-nonachlor
							490 (32-2500)	51 (7-744)	36 (9-120)	65 (15-350)	(15)	DDT: p,p'-DDE only; HCH: β -HCH only; CHL: oxychlorane and trans-nonachlor
							840 (25-4100)	52 (12-210)	44 (12-130)	83 (23-1500)	(15)	DDT: p,p'-DDE, p,p'-DDD, p,p'-DDT, o,p'-DDE and o,p'-DDT; HCH: α -, β - and γ -HCH; CHL: trans-nonachlor and oxychlorane
							1.5 \pm 1.3	0.27 \pm 0.16	NA	1.5 \pm 1.5	(16)	DDT: p,p'-DDE, p,p'-DDD, p,p'-DDT, o,p'-DDE, o,p'-DDD and o,p'-DDT; HCH: β - and γ -HCH
							1300 \pm 1500	NA	NA	NA	(17)	DDT: p,p'-DDE only

^aCited values were rounded to 2 significant digits for comparison. ^bReference: (1) present study; (2) Takayama et al. (2002); (3) Hamada et al. (2002); (4) Suzuki et al. (1991); (5) Kerrick et al. (2002); (6) Rathore et al. (2002); (7) Dua et al. (2002); (8) Sohrman et al. (2003); (9) Waliszewski et al. (2001); (10) Sarcinelli et al. (2003); (11) Anderson et al. (1998); (12) Moysich et al. (2002); (13) Walker et al. (2003); (14) Van Oostdam and Tremblay (2003); (15) Glynos et al. (2003); (16) Löffler et al. (2000); (17) Pauwels et al. (2000); ^cM, male; F, female. ^dRange concentration. ^eNA, no data available. ^fMean \pm standard deviation (SD). ^gnd, not detected; detection limit was not reported.

Table 5. Gender differences in organochlorine accumulation in human blood from Miyako, Saku and Toitori and in human adipose tissues from Tokyo and nearby areas, Japan*.

Location	Age	n	Concentration (ng/g lipid wt) [median, mean \pm SD (range)]b						
			PCBs	DDTs	HCHs	CHLs	HCB	TCrMs	
Miyako									
Male	44 \pm 11 (27 - 58)	13	340, 320 \pm 130 (160 - 620)*	230, 380 \pm 410 (56 - 1400)	24, 43 \pm 37 (7.6 - 120)	60, 70 \pm 41 (24 - 150)	7.5, 8.2 \pm 4.1 (4.6 - 20)	2.6, 3.1 \pm 2.8 (<0.1 - 8.1)	
Female	35 \pm 10 (24 - 52)	10	190, 230 \pm 81 (120 - 360)*	190, 200 \pm 120 (73 - 400)	30, 42 \pm 34 (17 - 130)	66, 71 \pm 40 (25 - 150)	8.0, 8.6 \pm 2.1 (6.1 - 12)	2.8, 3.3 \pm 2.3 (<0.1 - 6.7)	
Saku									
Male	48 \pm 9.2 (30 - 59)	15	480, 520 \pm 210 (210 - 990)**	300, 460 \pm 660 (35 - 2700)	32, 36 \pm 26 (0.81 - 100)	34, 39 \pm 20 (13 - 79)*	11, 14 \pm 8.9 (3.3 - 32)	2.4, 2.7 \pm 1.4 (0.72 - 6.2)**	
Female	48 \pm 9.7 (25 - 59)	17	270, 300 \pm 100 (140 - 560)**	200, 290 \pm 320 (32 - 1300)	35, 36 \pm 13 (17 - 64)	23, 25 \pm 10 (4.8 - 47)*	13, 11 \pm 5.1 (1.0 - 19)	1.3, 1.5 \pm 1.1 (<0.1 - 4.5)**	
Toitori									
Male	49 \pm 9.4 (30-59)	10	470, 460 \pm 190 (200 - 780)	200, 210 \pm 130 (34 - 440)	38, 41 \pm 25 (9.8 - 95)	36, 42 \pm 16 (24 - 68)**	11, 11 \pm 3.0 (5.1 - 16)	2.9, 2.7 \pm 1.4 (0.42 - 4.9)	
Female	45 \pm 8.2 (30 - 59)	12	280, 330 \pm 120 (200 - 510)	170, 200 \pm 140 (50 - 490)	33, 33 \pm 14 (9.2 - 64)	20, 23 \pm 8.9 (12 - 44)**	9.3, 9.2 \pm 2.0 (5.9 - 13)	2.0, 2.1 \pm 1.0 (0.9 - 4.8)	
Tokyo and nearby areas*									
Male	57 \pm 19 (17 - 81)	12	1700, 2000 \pm 1600 (230 - 6600)	1800, 1800 \pm 1200 (200 - 4800)	230, 340 \pm 270 (47 - 900)*	250, 410 \pm 550 (90 - 2100)	36, 44 \pm 25 (21 - 110)*	13, 17 \pm 11 (2.7-44)	
Female	70 \pm 14 (48 - 87)	8	2400, 2200 \pm 880 (400 - 3200)	1800, 2900 \pm 2800 (160 - 8100)	850, 1200 \pm 1000 (130 - 3200)*	340, 400 \pm 240 (42 - 730)	86, 85 \pm 41 (17 - 160)*	18, 21 \pm 11 (5.7-39)	

*Mann Whitney's U Test was used for evaluating gender differences in OC concentrations: *p < 0.05, **p < 0.005. bMedian, mean \pm standard deviation (SD), range in parentheses.

*Data for adipose tissues in diseased persons from Tokyo and nearby areas were cited from our previous study (Minh et al. 2001)