

Table 4 Adjusted odds ratios (OR) and 95% confidence interval (95% CI) of smoking status according to genotype of genes selected by stepwise logistic analysis. Allele A represents major allele; allele a represents minor allele

Gene symbol	SNP (rs number)	Smoking status			OR ^a (95% CI)	
		Never smokers	Past smokers	Current smokers	Current versus nonsmokers	Current versus never smokers
Genotype (Aa + aa) %						
OGG1	rs1052133	89.1 (49/55)	67.6 (48/71)	66.2 (141/213)	0.6 (0.4–1.0)	0.2 (0.1–0.6)
SLC6A4	rs717742	6.1 (3/49)	6.5 (4/62)	18.9 (36/191)	3.5 (1.5–8.2)	3.3 (1.0–11.2)
CYP17A1	rs743572	70.9 (39/55)	81.7 (58/71)	66.0 (140/212)	0.6 (0.4–1.0)	0.8 (0.4–1.6)
Genotype aa %						
EPHX1	rs2292566	12.7 (7/55)	14.1 (10/71)	5.6 (12/213)	0.4 (0.2–0.8)	0.5 (0.2–1.2)
ESR1	rs1913474	20.4 (11/54)	11.9 (8/67)	26.0 (53/204)	1.9 (1.1–3.4)	1.4 (0.7–2.9)

^aOR was adjusted for alcohol drinking status (never, past, and current drinking)

allele of rs743572 SNP in CYP17A1, among current smokers compared with past smokers. The findings may suggest involvement of the ESR1 and/or the CYP17A1 SNP in smoking, as well as in difficulty in quitting, among homozygous minor allele carriers of the ESR1 SNP and/or major allele carriers of the CYP17A1 SNP. An American study indicated that the ESR1 gene might play a role in anxiety (Comings et al. 1999). Anxiety, perhaps, drives individuals to use tobacco for comfort.

The present population-based study using a random Japanese sample screened genes related to habitual smoking and drinking. The findings provided basic but essential information on the association between candidate genes and smoking or drinking behavior for future case-only study, which could determine the interaction between genetics and behavior. The findings also confirmed that ALDH2 gene polymorphism is associated with habitual alcohol consumption, suggesting for the first time that polymorphism in the OGG1, SLC6A4, EPHX1, ESR1, CYP17A1, and NUDT1 genes may influence tobacco and/or alcohol use among Japanese men. Allelic association is a powerful method for detecting genetic influence on complex traits such as cigarette smoking and drinking; yet it cannot formally identify the functional polymorphisms responsible for the phenotype because the association analysis is based on the linkage disequilibrium between the marker SNPs and the causative genetic variation. Further studies are needed to confirm our findings, as well as to expose the underlying molecular mechanism.

Acknowledgements This study was supported mainly by the Program for Promotion of Fundamental Studies in Health Sciences of the Organization for Pharmaceutical Safety and Research, and the Grant-in-Aid for Cancer Research, and Risk Analysis Research on Food and Pharmaceuticals from the Japanese Ministry of Health, Labour and Welfare.

References

- Carmelli D, Swan GE, Robinette D, Fabsitz R (1992) Genetic influence on smoking—a study of male twins. *N Engl J Med* 327:829–833
- Comings DE, Muhleman D, Johnson P, MacMurray JP (1999) Potential role of the estrogen receptor gene (ESR1) in anxiety. *Mol Psychiatry* 4:374–377
- Eriksson CJ (2001) The role of acetaldehyde in the actions of alcohol (update 2000). *Alcohol Clin Exp Res* 25:15S–32S
- Goedde HW, Agarwal DP, Fritze G, Meier-Tackmann D, Singh S, Beckmann G, Bhatia K, Chen LZ, Fang B, Lisler R et al (1992) Distribution of ADH2 and ALDH2 genotypes in different populations. *Hum Genet* 88:344–346
- Goode EL, Ulrich CM, Potter JD (2002) Polymorphisms in DNA repair genes and associations with cancer risk. *Cancer Epidemiol Biomarkers Prev* 11:1513–1530
- Heath AC, Meyer J, Jardine R, Martin NG (1991) The inheritance of alcohol consumption patterns in a general population twin sample: II. Determinants of consumption frequency and quantity consumed. *J Stud Alcohol* 52:425–433
- Hopfer CJ, Stallings MC, Hewitt JK (2001) Common genetic and environmental vulnerability for alcohol and tobacco use in a volunteer sample of older female twins. *J Stud Alcohol* 62:717–723
- Hussain SP, Hofseth LJ, Harris CC (2001) Tumor suppressor genes: at the crossroads of molecular carcinogenesis, molecular epidemiology and human risk assessment. *Lung Cancer* 34 (Suppl 2):S7–S15
- Ishikawa H, Ohtsuki T, Ishiguro H, Yamakawa-Kobayashi K, Endo K, Lin YL, Yanagi H, Tsuchiya S, Kawata K, Hamaguchi H, Arinami T (1999) Association between serotonin transporter gene polymorphism and smoking among Japanese males. *Cancer Epidemiol Biomarkers Prev* 8:831–833
- Le Marchand L, Donlon T, Lum-Jones A, Seifried A, Wilkens LR (2002) Association of the hOGG1 Ser326Cys polymorphism with lung cancer risk. *Cancer Epidemiol Biomarkers Prev* 11:409–412
- Lerman C, Shields PG, Audrain J, Main D, Cobb B, Boyd NR, Caporaso N (1998) The role of the serotonin transporter gene in cigarette smoking. *Cancer Epidemiol Biomarkers Prev* 7:253–255
- McGue M, Iacono WG, Legrand LN, Elkins I (2001) Origins and consequences of age at first drink. II. Familial risk and heritability. *Alcohol Clin Exp Res* 25:1166–1173
- Munaka M, Kohshi K, Kawamoto T, Takasawa S, Nagata N, Itoh H, Oda S, Katoh T (2003) Genetic polymorphisms of tobacco- and alcohol-related metabolizing enzymes and the risk of hepatocellular carcinoma. *J Cancer Res Clin Oncol* 129:355–360
- Muramatsu T, Wang ZC, Fang YR, Hu KB, Yan H, Yamada K, Higuchi S, Harada S, Kono H (1995) Alcohol and aldehyde dehydrogenase genotypes and drinking behavior of Chinese living in Shanghai. *Hum Genet* 96:151–154
- Pianezza ML, Sellers EM, Tyndale RF (1998) Nicotine metabolism defect reduces smoking. *Nature* 393:750
- Ross P, Hall L, Smirnov I, Haff L (1998) High level multiplex genotyping by MALDI-TOF mass spectrometry. *Nat Biotechnol* 16:1347–1351

- Sabol SZ, Nelson ML, Fisher C, Gunzerath L, Brody CL, Hu S, Sirota LA, Marcus SE, Greenberg BD, Lucas FRt, Benjamin J, Murphy DL, Hamer DH (1999) A genetic association for cigarette smoking behavior. *Health Psychol* 18:7-13
- Schlesselman JJ (1982) *Case-control studies: design, conduct, analysis*. Oxford University Press, New York
- Shields PG, Lerman C, Audrain J, Bowman ED, Main D, Boyd NR, Caporaso NE (1998) Dopamine D4 receptors and the risk of cigarette smoking in African-Americans and Caucasians. *Cancer Epidemiol Biomarkers Prev* 7:453-458
- Sugimura H, Wakai K, Genka K, Nagura K, Igarashi H, Nagayama K, Ohkawa A, Baba S, Morris BJ, Tsugane S, Ohno Y, Gao C, Li Z, Takezaki T, Tajima K, Iwamasa T (1998) Association of Ile462Val (Exon 7) polymorphism of cytochrome P450 IA1 with lung cancer in the Asian population: further evidence from a case-control study in Okinawa. *Cancer Epidemiol Biomarkers Prev* 7:413-417
- Tsugane S, Gey F, Ichinowatari Y, Miyajima Y, Ishibashi T, Matsuishima S, Hirota Y, Inami T, Yamaguchi M, Karita K, Kabuto M, Takeshima Y, Todoriki H, Tsuda M, Akabane M, Furuichi Y, Hamada G, Watanabe S (1992a) Cross-sectional epidemiologic study for assessing cancer risks at the population level I. Study design and participation rate. *J Epidemiol* 2:75-81
- Tsugane S, Gey F, Ichinowatari Y, Miyajima Y, Ishibashi T, Matsuishima S, Hirota Y, Inami T, Yamaguchi M, Karita K, Kabuto M, Takeshima Y, Todoriki H, Tsuda M, Akabane M, Furuichi Y, Hamada G, Watanabe S (1992b) Cross-sectional epidemiologic study for assessing cancer risks at the population level II. Baseline data and correlation analysis. *J Epidemiol* 2:83-89
- Wu X, Hudmon KS, Detry MA, Chamberlain RM, Spitz MR (2000) D2 dopamine receptor gene polymorphisms among African-Americans and Mexican-Americans: a lung cancer case-control study. *Cancer Epidemiol Biomarkers Prev* 9:1021-1026
- Yokoyama A, Omori T (2001) Alcohol and oropharyngolaryngeal and digestive tract cancer. *Nihon Arukoru Yakubutsu Igakkai Zasshi* 36:551-566
- Yoshida K, Hamajima N, Kozaki K, Saito H, Maeno K, Sugiura T, Ookuma K, Takahashi T (2001) Association between the dopamine D2 receptor A2/A2 genotype and smoking behavior in the Japanese. *Cancer Epidemiol Biomarkers Prev* 10:403-405
- Yoshimura K, Hanaoka T, Ohnami S, Kohno T, Liu Y, Yoshida T, Sakamoto H, Tsugane S (2003) Allele frequencies of single nucleotide polymorphisms (SNPs) in 40 candidate genes for gene-environment studies on cancer: data from population-based Japanese random samples. *J Hum Genet* 48:654-658

Association between endometriosis and genetic polymorphisms of the estradiol-synthesizing enzyme genes *HSD17B1* and *CYP19*

M.Tsuchiya^{1,2}, H.Nakao¹, T.Kato^{1,6}, H.Sasaki³, M.Hiroshima³, T.Tanaka³, T.Matsunaga⁴, T.Hanaoka⁵, S.Tsugane⁵ and T.Ikenoue²

Departments of ¹Public Health and ²Obstetrics and Gynecology, Miyazaki Medical College, University of Miyazaki, 5200 Kihara, Kiyotake, Miyazaki, ³Department of Obstetrics and Gynecology, The Jikei University School of Medicine, 3-25-8 Nishi-shinbashi, Minato-ku, Tokyo, ⁴Department of Biotechnology, Tokyo University of Agriculture and Technology, 2-24-16 Naka-cho, Koganei, Tokyo and ⁵Epidemiology and Prevention Division, Research Center for Cancer Prevention and Screening, National Cancer Center, 5-1-1 Tsukiji, Chuo-ku, Tokyo, Japan

⁶To whom correspondence should be addressed. E-mail: katoht@med.miyazaki-u.ac.jp

BACKGROUND: Endometriosis, an estrogen-dependent disease, is believed to be influenced by multiple genetic and environmental factors. Here, we evaluated whether the risk and severity of endometriosis are associated with polymorphisms in estradiol-synthesizing enzyme genes: the Ser312Gly polymorphism in 17-beta-hydroxysteroid dehydrogenase type 1 (*HSD17B1*) and the Arg264Cys polymorphism in cytochrome P450, subfamily XIX (*CYP19*). **METHODS:** All participants underwent diagnostic laparoscopy, and the stage of endometriosis was determined according to the Revised American Fertility Society classification. Of the 138 women enrolled, 59 had no endometriosis, 21 had stage I, 10 had stage II, 23 had stage III and 25 had stage IV. SNPs were discriminated by allele-specific oligonucleotide hybridization. **RESULTS:** Individuals having at least one A-allele (A/G or A/A genotype) of *HSD17B1* showed a significantly increased risk of endometriosis (A/G genotype: adjusted OR, 3.06; 95% CI 1.21–7.74; A/A genotype: adjusted OR, 3.02; 95% CI 1.08–8.43). There was a significant trend associating A/G + A/A genotypes with severity of endometriosis (*P* for trend <0.01). No statistically significant association was found for the *CYP19* polymorphism. **CONCLUSIONS:** Evidence for association between the Ser312Gly polymorphism in *HSD17B1* and endometriosis was found in a Japanese population. The A-allele of *HSD17B1* appears to confer higher risk for endometriosis.

Key words: CYP19/endometriosis/estrogen synthesis/genetic polymorphism/HSD17B1

Introduction

Endometriosis, one of the most common causes of female infertility and chronic pelvic pain, is defined as the presence of endometrial tissue outside the uterus. Three predominant theories have been proposed for the etiology of this disease: Mullerian remnants, metaplasia and direct implantation of endometrial cells (El-Mahgoub and Yaseen, 1980; Murphy *et al.*, 1986; Fujii *et al.*, 1991). Although the exact prevalence is still not known, endometriosis affects up to 5–10% of women of reproductive age (Wheeler *et al.*, 1989). The prevalence of endometriosis is as high as 20–50% in infertile women (Strathy *et al.*, 1982; Rawson *et al.*, 1991).

Endometriosis is regarded as a complex trait, in which genetic and environmental factors contribute to the disease phenotype (Kennedy *et al.*, 1998). A variety of factors affect the development of endometriosis, including hormonal status and genetic factors. For example, women with shorter intervals between menstruation and longer duration of menses are

at higher risk for endometriosis (Vercellini *et al.*, 1997). The risk of endometriosis is seven times higher if a first-degree relative has been affected by endometriosis (Simpson *et al.*, 1980). However, the interaction between genetic susceptibility and environmental factors is not yet adequately understood.

The development of endometriosis is estrogen-dependent. Endometrial implants contain estrogen and progesterone receptors (Lessey *et al.*, 1989) and respond to ovarian hormonal changes, causing local bleeding, inflammation and formation of adhesions. The three main estrogens are estradiol, estrone and estriol. Estradiol, the most active form, is produced either from estrone via 17- β -hydroxysteroid dehydrogenase type 1 (*HSD17B1*) or from testosterone via cytochrome P450, subfamily XIX (*CYP19*, aromatase) (Mitrunen and Hirvonen, 2003).

Current evidence indicates that polymorphisms in genes of drug-metabolizing enzymes can affect phenotypic metabolic

variations. The *HSD17B1* gene is located in chromosome 17q12 and has a polymorphism consisting of an A to G substitution in exon 6, resulting in an amino acid change of Ser312Gly (Puranen *et al.*, 1994). The *CYP19* gene, located in chromosome 15q21, has a polymorphism consisting of C to T substitution in exon 7, resulting in an amino acid change of Arg264Cys (Toda *et al.*, 1990).

Genetic polymorphisms involved in estrogen synthesis and metabolism may play an important role in the variation of endometriosis among individuals by altering local estrogen production or circulating levels of estrogen. Here, we evaluate whether the Ser312Gly polymorphism in *HSD17B1* and Arg264Cys polymorphism in *CYP19* are associated with the risk and severity of endometriosis. A case-control study was conducted on these two polymorphisms in patients with different stages of endometriosis and controls.

Materials and methods

The protocol for the study was approved by the Institutional Review Board of University of Miyazaki, The Jikei University School of Medicine and National Cancer Center. All subjects gave their written informed consent before the laparoscopic examination.

Study participants

This study was a part of a case-control study of endometriosis. During the years 1999–2000, 139 women were recruited at the Department of Obstetrics and Gynecology, The Jikei University School of Medicine Hospital. Participants were patients between the ages of 20 and 45 who had complained of infertility and attended the hospital. The study protocol excluded all women from the study who had ever given birth or lactated. Of the 139 women recruited, only one was excluded from subsequent analysis because a DNA sample was not available.

All participants underwent diagnostic laparoscopy, and the stage of endometriosis was determined according to the Revised American Fertility Society classification (r-AFS) (American Fertility Society, 1985). Of the 138 women enrolled, 59 women had no endometriosis, 21 had stage I, 10 had stage II, 23 had stage III and 25 had stage IV.

Cases and controls were similar in several confounding factors. Risk factors for endometriosis include age, shorter menstrual cycles and longer duration of menstrual flow (Vessey *et al.*, 1993; Eskenazi and Warner, 1997). The mean age of the cases was 32.4 years and 33.1 in the controls ($P = 0.35$). No significant difference was observed in the duration of menstruation. However, there was a significant trend towards cases having shorter menstrual cycles compared to the controls (28.8 days for cases and 30.7 for controls, $P = 0.03$).

Genotyping

Blood samples were obtained before the laparoscopic examination. Genomic DNA samples were extracted from peripheral white blood cells by using a DNA Extractor WB Kit (Wako, Osaka, Japan).

A 67 bp fragment in *HSD17B1*, including an SNP site located at 27 bases from the 5' end, was amplified using sense (5'-CTGGGGC-AGAGGACGAGG) and biotin-labeled antisense (5'-GCGGCCGG-AGGATCG) primers. A 56 bp fragment of *CYP19*, including an SNP site located at 31 bases from the 5' end, was amplified using biotin-labeled sense (5'-GCCATAGAAGTTCTGATAGCAG) and antisense (5'-AGTTTCTCTTCTGTGGAAATCCT) primers. PCR

amplifications were performed using a TPC-200 thermal cycler (MI Research Inc., Watertown, MA) in a total reaction volume of 25 μ l containing 20 ng of DNA sample, 0.6 U AmpliTaq DNA polymerase (Applied Biosystems), 0.25 mM dNTPs, 0.2 μ M primers and PCR buffer [1 \times GC buffer II (Takara Bio Inc., Otsu, Japan) for *HSD17B1* and 1 \times GeneAmp PCR buffer (Applied Biosystems) for *CYP19*]. The amplification protocol comprised initial denaturation at 95 $^{\circ}$ C for 5 min, then 35 cycles of denaturation at 95 $^{\circ}$ C for 15 s and annealing at 55 $^{\circ}$ C for 30 s.

SNP discriminations were conducted in a manner similar to that described previously (Maruyama *et al.*, 2004) with details as follows, based on allele-specific oligonucleotide hybridization using bio-nano magnetite particles (Takeyama *et al.*, 2000; Matsunaga *et al.*, 2001). Cy3- and Cy5-labeled detection probes were designed for each SNP as follows: Cy3-labeled *HSD17B1* A-allele detection probe (5'-CCGGGCGCAGTGGCGTG), Cy5-labeled *HSD17B1* G-allele detection probe (5'-CCGGGCGCGGTGCGGTG), Cy3-labeled *CYP19* T-allele detection probe (5'-AATCCTGCATCTT-TTTT) and Cy5-labeled *CYP19* C-allele detection probe (5'-AAAT-CCTGCGTCTTTTTT). All the following experiments were performed using the semi-automated SNP detection processor (Tanaka *et al.*, 2003).

Biotinylated PCR product (12.5 μ l) and streptavidin-immobilized bio-nano magnetic particles (25 μ g/12.5 μ l), which were prepared according to the method described by Yoshino *et al.* (2002), were mixed and incubated for 15 min at room temperature for capturing the PCR products on the magnetic particles. PCR products were denatured into single strands by alkali treatment (10 mM NaOH). After neutralization of the mixture by neutralization buffer (100 mM Tris-HCl pH 7.5, 3 mM EDTA, 0.1% BSA), 25 μ l of hybridization buffer (1 M tetramethylammonium chloride, 37.5 mM Tris-HCl pH 7.5 and 3 mM EDTA) containing 12.5 pmoles of Cy3-labeled and Cy5-labeled detection probes was added to the PCR products captured on magnetic particles.

The mixture was rapidly heated up to 70 $^{\circ}$ C, and then cooled slowly to 25 $^{\circ}$ C over 10 min to allow hybridization of the detection probes and biotinylated PCR products. The optimum temperature for dissociating single-base mismatched probes was determined by analysis of dissociation curves. The mixture was heated to the optimum temperature: 56 $^{\circ}$ C for *HSD17B1* or 54 $^{\circ}$ C for *CYP19*, and the detection probes dissociated were removed at this temperature using an automated SNP detection processor (Maruyama *et al.*, 2004). Finally, the mixture was heated to 80 $^{\circ}$ C to liberate the hybridized detection probe into the supernatant.

The fluorescence intensity of the liberated detection probe was measured at Ex: 540 nm, Em: 570 nm for Cy3 and Ex: 645 nm, Em: 675 nm for Cy5 by a microplate reader (BMG Labtech, Offenburg, Germany), respectively. The samples were classified into three distinct categories according to the signal ratio of Cy5/Cy3: (i) those with values >2 , representing samples with the homozygous G/G genotype in *HSD17B1* or the homozygous C/C genotype in *CYP19*; (ii) values <0.5 , representing samples with the homozygous A/A genotype in *HSD17B1* or the homozygous T/T genotype in *CYP19*; (iii) intermediate values, representing samples with the heterozygous genotype.

Statistical analysis

To estimate the risk of endometriosis, crude odds ratios (ORs) and 95% confidence intervals (95% CIs) were calculated for subgroups with the different genotypes with respect to the Ser312Gly polymorphism in *HSD17B1* and the Arg264Cys polymorphism in *CYP19*. ORs were adjusted for three endometriosis risk factors: age, menstrual cycle and duration of menstruation, using multiple logistic

regression analysis by SPSS for Windows software (version 11.0.1J, SPSS Japan, Tokyo, Japan).

The Wilcoxon rank-sum test for trend was also used to estimate the associations between the two polymorphisms and stage of endometriosis (Goodman *et al.*, 1954). All statistical tests were based on two-tailed probability. $P < 0.05$ was considered statistically significant.

Results

Of the 138 DNA samples, 10 could not be reliably genotyped and were excluded from subsequent statistical analysis (six subjects for HSD17B1, four subjects for CYP19). The observed genotype distributions were in Hardy-Weinberg equilibrium in the control subjects.

Test for association with risk of endometriosis

Table I shows the genotypic and allelic distributions of HSD17B1 and CYP19 polymorphisms. Individuals with at least one HSD17B1 A-allele (A/G or A/A genotype) showed a significantly increased risk of endometriosis (A/G genotype: adjusted OR, 3.06; 95%CI 1.21-7.74; A/A genotype: adjusted OR, 3.02; 95%CI 1.08-8.43). A similar result was obtained in comparison of the combined A/A + A/G genotypes with the G/G genotype (adjusted OR, 3.05; 95%CI 1.30-7.14). No significant association was observed between the Arg264Cys polymorphism in CYP19 and risk of endometriosis (C/T genotype: adjusted OR, 1.35; 95%CI 0.62-2.95; T/T genotype: adjusted OR, 0.41; 95%CI 0.09-1.85).

Test for association with severity of endometriosis

To evaluate whether the Ser312Gly polymorphism in HSD17B1 and the Arg264Cys polymorphism in CYP19 are associated with the severity of endometriosis, participants were categorized into three groups according to r-AFS classification: controls, stage I-II and stage III-IV (Table II). There was a significant trend between the combined A/G + A/A genotypes and stage of endometriosis (P for trend < 0.01). No statistically significant association was found between the Arg264Cys polymorphism in CYP19 and stage of endometriosis (data not shown).

Discussion

In the present study, we evaluated whether two polymorphisms in estradiol-synthesizing enzyme genes are associated with the risk and severity of endometriosis. The results of this study suggested that the Ser312Gly polymorphism in HSD17B1 is associated with both risk and severity of endometriosis, while no associations were found for the Arg264Cys polymorphism in CYP19.

In this study, we applied strict clinical criteria for the definition of cases and controls. One third of women with endometriosis are asymptomatic, and laparoscopy or laparotomy is indispensable for the diagnosis of endometriosis. All participants in the present study underwent diagnostic laparoscopy and were diagnosed according to the r-AFS classification. In addition, bias from confounding variables was minimized by adjusting ORs for the risk and stage of

Table I. Genotypic and allelic distribution of HSD17B1 and CYP19 polymorphisms

Polymorphisms	Cases n (%)		Controls n (%)		Crude OR (95%CI)	Adjusted OR ^a (95%CI)
	Genotype	Allele	Genotype	Allele		
HSD17B1 Ser312Gly						
G/G	13 (17.3)	G allele: 66 (44.0)	22 (38.6)	G allele: 65 (57.0)	1	1
A/G	40 (53.3)		21 (36.8)		3.22** (1.36-7.66)	3.06* (1.21-7.74)
A/A	22 (29.3)	A allele: 84 (56.0)	14 (24.6)	A allele: 49 (43.0)	2.66* (1.02-6.94)	3.02* (1.08-8.43)
A/G, A/A	62 (82.7)		35 (61.4)		3.00** (1.35-6.68)	3.05** (1.30-7.14)
CYP19 Arg264Cys						
C/C	35 (46.0)	C allele: 107 (70.4)	29 (50.0)	C allele: 81 (69.8)	1	1
C/T	37 (48.7)		23 (39.7)		1.33 (0.65-2.73)	1.35 (0.62-2.95)
T/T	4 (5.3)	T allele: 45 (29.6)	6 (10.3)	T allele: 35 (30.2)	0.56 (0.14-2.15)	0.41 (0.09-1.85)
C/T, T/T	41 (54.0)		29 (50.0)		1.17 (0.59-2.32)	1.13 (0.54-2.36)

^aORs adjusted for age and menstrual characteristics.

* $P < 0.05$.

** $P < 0.01$.

Table II. Severity of endometriosis associated with HSD17B1 polymorphism

Endometriosis	G/G genotype	A/G, A/A genotype	Crude OR (95%CI)	Adjusted OR ^a (95%CI)
Controls				
n (%)	22 (39.7)	35 (60.3)	1	1
Stage ?-?				
n (%)	7 (23.3)	23 (76.7)	2.07 (0.76-5.62)	2.07 (0.69-6.20)
Stage ?-?				
n (%)	6 (13.3)	39 (86.7)	4.08** (1.49-11.22)	3.99** (1.41-11.26)

^aORs adjusted for age and menstrual characteristics.

P for trend < 0.01 .

** $P < 0.01$.

endometriosis for variables known to affect endometriosis: age, menstrual cycle and duration of menstrual flow. Detailed questionnaires were designed to determine patients' menstrual cycle and duration of menstruation. The questionnaires were administered by a trained interviewer before the laparoscopic examination to minimize recall bias.

Retrograde menstruation is thought to be one of the main causes of endometriosis. However, retrograde menstruation is a common phenomenon (Kruitwagen *et al.*, 1991), and not all the women of reproductive age are affected by endometriosis. In short, women with endometriosis can be considered to have defects in the regulation of endometrial proliferation: (i) a tendency to proliferate endometrial tissue, and (ii) impaired clearance of abnormal endometrial tissue (Vinatier *et al.*, 2001).

Development of endometriosis is estrogen-dependent, and several features of this disease can be explained on the basis of overproduction of estrogen. Current therapy consists of hormone treatments aimed at lowering circulating estrogen. Genetic polymorphisms in the estrogen-synthesis or estrogen-metabolizing enzymes may cause inter-individual variation of levels and activity of circulating estrogen.

Estradiol, the most physiologically active form of estrogen, stimulates the proliferation of the endometrium during the ovulatory phase of the menstrual cycle. The *HSD17B1* genotypes found to be associated with endometriosis may confer increased activity or expression of HSD17B1 enzyme, causing increased exposure to estradiol. Our results suggest that possessing at least one A-allele of the Ser312Gly polymorphism in *HSD17B1* increases the risk of endometriosis. Furthermore, AG + AA genotypes showed a significantly increased risk for severe endometriosis (P for trend < 0.01). These findings support the idea that the Ser312Gly polymorphism in *HSD17B1* is associated with the risk and progression of endometriosis, especially in terms of the inter-individual variation of estradiol synthesis.

Breast and endometrial cancer, like endometriosis, are considered to be estrogen-dependent diseases. The results of this study are consistent with those of a previous study that examined the Ser312Gly polymorphism in *HSD17B1* and breast cancer risk. When the *HSD17B1* A-allele and *CYP17* A2-allele were considered as the high-risk alleles, the risk of advanced breast cancer among women carrying four high-risk alleles (*HSD17B1* AA and *CYP17* A2A2) was 2.21 compared with that of women who carried none (Feigelson *et al.*, 2001).

The *CYP19* gene, encoding aromatase, plays a crucial role in estradiol synthesis. However, we could not find any association between the Arg264Cys polymorphism in *CYP19* and endometriosis. A possible explanation for this negative result might be a lack of a functional effect for this polymorphism. A previous study reported that aromatase activity was not affected by the Arg264Cys polymorphism (Watanabe *et al.*, 1997).

Three major limitations must be considered when evaluating the results of this study. First is the relatively small number of subjects. Although we found statistically significant differences, the 95% confidence intervals were relatively

wide, reflecting the small number of cases. The sample size was sufficient to detect odds ratios of three or larger with 80% power at the 5% level of significance. The lack of a significant association with the *CYP19* Arg264Cys polymorphism means only that we failed to detect a difference. It remains unclear whether this polymorphism affects endometriosis.

Secondly, the distribution of *HSD17B1* alleles in the control group deviated considerably from the expected Hardy-Weinberg equilibrium, a difference that was not significant ($P = 0.06$), but was on the edge of being so. The discrepancy may result from the small number of subjects or from the characteristics of the control group. The setting of this study is a hospital, and the control group is women complaining of infertility. Because of the requirement for a surgical diagnosis, the selection of a control group in case-control studies of endometriosis has been particularly difficult (Zondervan *et al.*, 2002), and it is difficult to exclude a selection bias completely. Development of non-invasive methods for diagnosis or a prospective randomized trial will minimize any sampling bias.

Lastly, endogenous estrogen levels were not measured in this study. *HSD17B1* is not expressed in normal endometrium or endometrial hyperplasia (Utsunomiya *et al.*, 2001). One possible explanation of the apparent influence of *HSD17B1* is that the *HSD17B1* A-allele increases the level of circulating estradiol. Although an *in vitro* study failed to demonstrate any change in HSD17B1 catalytic activity produced by the Ser312Gly polymorphism (Puranen *et al.*, 1994), a recent molecular epidemiological study showed that *HSD17B1* A/A genotype was associated with higher estradiol levels in lean women (Setiawan *et al.*, 2004). The functional effects of the Ser312Gly polymorphism in *HSD17B1* and the Arg264Cys polymorphism in *CYP19* have not yet been clearly established. Further information on functional changes and more epidemiologic studies will help clarify the association between these polymorphisms and endometriosis.

Many alleles have been reported to vary in frequency among different ethnic or geographic populations. In the present study, allelic frequencies of *HSD17B1* Ser312Gly polymorphism in control individuals were similar to previously reported distributions in Chinese and American populations: A 0.43 and G 0.57 in Japanese and Chinese populations, and A 0.51 and G 0.49 in an American population (Wu *et al.*, 2003; Setiawan *et al.*, 2004). On the other hand, large variations were found between Japanese and Caucasians in the *CYP19* Arg264Cys polymorphism: C 0.70 and T 0.30 in a Japanese population, versus C 0.96 and T 0.04 in a Caucasian population (Hefler *et al.*, 2004).

In summary, we demonstrated that the Ser312Gly polymorphism in *HSD17B1* was associated with the risk and severity of endometriosis in a Japanese population. The A-allele of the *HSD17B1* gene is considered to be a high-risk allele for endometriosis. However, no association was found between the Arg264Cys polymorphism in *CYP19* and endometriosis. The results of this study are not conclusive and further investigation is warranted. Endometriosis is a complex trait. The many factors contributing to the disease

phenotype make its pathophysiology very difficult to understand. Progress in the genetics of the disease, including understanding of genetic polymorphisms, will facilitate research on endometriosis.

Acknowledgements

We thank Drs Haruko Takeyama and Kouhei Maruyama, Department of Biotechnology, Tokyo University of Agriculture and Technology, for genetic analysis. This work was supported by a Grant-in-Aid for Risk Analysis Research on Food and Pharmaceuticals and Cancer Research from the Ministry of Health, Labour and Welfare of Japan.

References

- American Fertility Society (1985) Revised American Fertility Society classification of endometriosis. *Fertil Steril* 43,351–352.
- El-Mahgoub S and Yaseen S (1980) A positive proof for the theory of coelomic metaplasia. *Am J Obstet Gynecol* 137,137–140.
- Eskenazi B and Warner ML (1997) Epidemiology of endometriosis. *Obstet Gynecol Clin North Am* 24,235–258.
- Feigelson HS, McKean-Cowdin R, Coetzee GA, Stram DO, Kolonel LN and Henderson BE (2001) Building a multigenic model of breast cancer susceptibility: CYP17 and HSD17B1 are two important candidates. *Cancer Res* 61,785–789.
- Fujii S (1991) Secondary mullerian system and endometriosis. *Am J Obstet Gynecol* 165,219–225.
- Goodman LA and Kruskal WH (1954) Measures of association for cross classifications. *J Am Stat Assoc* 49,732–764.
- Hefler LA, Tempfer CB, Grimm C et al. (2004) Estrogen-metabolizing gene polymorphisms in the assessment of breast carcinoma risk and fibroadenoma risk in Caucasian women. *Cancer* 101,264–269.
- Kennedy S (1998) The genetics of endometriosis. *J Reprod Med* 43, 263–268.
- Kruitwagen RF, Poels LG, Willemsen WN, de Ronde IJ, Jap PH and Rolland R (1991) Endometrial epithelial cells in peritoneal fluid during the early follicular phase. *Fertil Steril* 55,297–303.
- Lessey BA, Metzger DA, Haney AF and McCarty KS, Jr (1989) Immunohistochemical analysis of estrogen and progesterone receptors in endometriosis: comparison with normal endometrium during the menstrual cycle and the effect of medical therapy. *Fertil Steril* 51,409–415.
- Maruyama K, Takeyama H, Nemoto E, Tanaka T, Yoda T and Matsunaga T (2004) Single nucleotide polymorphism detection in aldehyde dehydrogenase 2 (ALDH2) gene using bacterial magnetic particles based on dissociation curve analysis. *Biotechnol Bioeng* in press.
- Matsunaga T, Nakayama H, Okochi M and Takeyama H (2001) Fluorescent detection of cyanobacterial DNA using bacterial magnetic particles on a MAG-microarray. *Biotechnol Bioeng* 73,400–405.
- Mitrunen K and Hirvonen A (2003) Molecular epidemiology of sporadic breast cancer. The role of polymorphic genes involved in oestrogen biosynthesis and metabolism. *Mutat Res* 544,9–41.
- Murphy AA, Green WR, Bobbie D, dela Cruz ZC and Rock JA (1986) Unsuspected endometriosis documented by scanning electron microscopy in visually normal peritoneum. *Fertil Steril* 46,522–524.
- Puranen TJ, Poutanen MH, Peltoketo HE, Vihko PT and Vihko RK (1994) Site-directed mutagenesis of the putative active site of human 17 beta-hydroxysteroid dehydrogenase type 1. *Biochem J* 304,289–293.
- Rawson JM (1991) Prevalence of endometriosis in asymptomatic women. *J Reprod Med* 36,513–515.
- Setiawan VW, Hankinson SE, Colditz GA, Hunter DJ and De Vivo I (2004) HSD17B1 gene polymorphisms and risk of endometrial and breast cancer. *Cancer Epidemiol Biomarkers Prev* 13,213–219.
- Simpson JL, Elias S, Malinak LR and Buttram VC Jr (1980) Heritable aspects of endometriosis I. Genetic studies. *Am J Obstet Gynecol* 137, 327–331.
- Strathy JH, Molgaard CA, Coulam CB and Melton LJ 3rd (1982) Endometriosis and infertility: a laparoscopic study of endometriosis among fertile and infertile women. *Fertil Steril* 38,667–672.
- Takeyama H, Tsuzuki H, Chow S, Nakayama H and Matsunaga T (2000) Discrimination between Atlantic and Pacific subspecies of northern bluefin tuna (*Thunnus thynnus*) by magnetic-capture hybridization using bacterial magnetic particles. *Mar Biotechnol* 2,309–313.
- Tanaka T, Maruyama K, Yoda K, Nemoto E, Udagawa Y, Nakayama H, Takeyama H and Matsunaga T (2003) Development and evaluation of an automated workstation for single nucleotide polymorphism discrimination using bacterial magnetic particles. *Biosens Bioelectron* 19,325–330.
- Toda K, Terashima M, Kawamoto T et al. (1990) Structural and functional characterization of human aromatase P-450 gene. *Eur J Biochem* 193, 559–565.
- Utsunomiya H, Suzuki T, Kaneko C et al. (2001) The analyses of 17beta-hydroxysteroid dehydrogenase isozymes in human endometrial hyperplasia and carcinoma. *J Clin Endocrinol Metab* 86,3436–3443.
- Vercellini P, De Giorgi O, Aimi G, Panazza S, Uglietti A and Crosignani PG (1997) Menstrual characteristics in women with and without endometriosis. *Obstet Gynecol* 90,264–268.
- Vessey MP, Villard-Mackintosh L and Painter R (1993) Epidemiology of endometriosis in women attending family planning clinics. *BMJ* 306, 182–184.
- Vinatier D, Orazi G, Cosson M and Dufour P (2001) Theories of endometriosis. *Eur J Obstet Gynecol Reprod Biol* 96,21–34.
- Watanabe J, Harada N, Suemasu K, Higashi Y, Gotoh O and Kawajiri K (1997) Arginine-cysteine polymorphism at codon 264 of the human CYP19 gene does not affect aromatase activity. *Pharmacogenetics* 7, 419–424.
- Wheeler JM (1989) Epidemiology of endometriosis-associated infertility. *J Reprod Med* 34,41–46.
- Wu AH, Seow A, Arakawa K, Van Den Berg D, Lee HP and Yu MC (2003) HSD17B1 and CYP17 polymorphisms and breast cancer risk among Chinese women in Singapore. *Int J Cancer* 104,450–457.
- Yoshino T, Tanaka T, Takeyama H and Matsunaga T (2003) Single nucleotide polymorphism genotyping of aldehyde dehydrogenase 2 gene using a single bacterial magnetic particles. *Biosens Bioelectron* 18,661–666.
- Zondervan KT, Cardon LR and Kennedy SH (2002) What makes a good case-control study? Design issues for complex traits such as endometriosis. *Hum Reprod* 17,1415–1423.

Submitted on November 17, 2004; accepted on December 8, 2004

Running Title: AhRR polymorphism and endometriosis

Analysis of the AhR, ARNT and AhRR gene polymorphisms: Genetic contribution to susceptibility to and severity of endometriosis

REVISED: Masaki Tsuchiya, M.D.,^{a,b} Takahiko Katoh, M.D, Ph.D.,^a Hiroshi Motoyama, M.D.,^c Hiroshi Sasaki, M.D, Ph.D.,^c Shoichiro Tsugane, M.D, Ph.D.^d and Tsuyomu Ikenoue, M.D, Ph.D.^b

Departments of ^aPublic Health, and ^bObstetrics and Gynecology, Miyazaki Medical College, University of Miyazaki, 5200 Kihara, Kiyotake, Miyazaki, Japan

^cDepartment of Obstetrics and Gynecology, The Jikei University School of Medicine, 3-25-8 Nishi-shinbashi, Minato-ku, Tokyo, Japan

^dEpidemiology and Prevention Division, Research Center for Cancer Prevention and Screening, National Cancer Center, 5-1-1 Tsukiji, Chuo-ku, Tokyo, Japan

Corresponding author: Takahiko Katoh, M.D, Ph.D., Department of Public Health, Miyazaki Medical College, University of Miyazaki, 5200 Kihara, Kiyotake, Miyazaki, Japan

Fax: +81-985-85-6258; E-mail: katoht@med.miyazaki-u.ac.jp

CAPSULE

A polymorphism in the aryl hydrocarbon receptor repressor (*AhRR*) gene was found to be associated with risk and severity of endometriosis. Polymorphisms in *AhR* and *ARNT* showed no such association.

ABSTRACT

Objective: To explore whether polymorphisms in AhR, ARNT and AhRR contribute to the susceptibility to and severity of endometriosis.

Design: Case-control study.

Setting: Hospital.

Patients: 138 women with or without endometriosis diagnosed endoscopically. Women who had ever given birth or lactated were not eligible. All women were staged into two groups: cases, with r-AFS stage I - IV endometriosis, and controls without endometriosis.

Main Outcome Measures: AhR, ARNT and AhRR polymorphisms were genotyped using real-time PCR analysis. Odds ratios and 95% confidence intervals were calculated for AhR, ARNT and AhRR genotypes to evaluate the risk of endometriosis. Associations between these polymorphisms and stage of endometriosis were also examined.

Results: The C/G + G/G genotypes at codon 185 of AhRR showed a significant association with risk of endometriosis (adjusted odds ratio, 2.53; 95% confidence interval, 1.16-5.55). Furthermore, there was a significant trend associating the C/G + G/G genotypes with the clinical stage of endometriosis (P for trend: 0.02). No

statistically significant association was observed between AhR codon 554 or ARNT codon 189 polymorphisms and endometriosis.

Conclusion: AhRR codon 185 polymorphism was associated with susceptibility to and severity of endometriosis in Japanese women.

Key Words: Endometriosis, genetic polymorphisms, AhR, ARNT, AhRR

INTRODUCTION

Endometriosis is a gynecologic condition that occurs in approximately 10% of women in the general population (1) and in 40% in infertile women (2). The most common symptoms associated with pelvic endometriosis are dysmenorrhea, chronic pelvic pain and infertility.

Endometriosis is regarded as a complex trait, in which genetic and environmental factors contribute to the disease phenotype (3). Genetic understanding of endometriosis has recently begun to progress rapidly, particularly through analysis of genetic polymorphisms. Genetic polymorphisms associated with endometriosis include drug metabolizing enzymes, growth factors and hormone receptor genes (4-7).

The aryl hydrocarbon receptor (AhR) is a ligand-dependent transcription factor that regulates cell differentiation and the induction of the phase I and II drug metabolizing enzymes (8-9). The AhR signaling pathway regulates induction of CYP1A1 and CYP1B1, representative phase I drug metabolizing enzymes (10-11). These isoforms catalyze the conversion of 17β -estradiol to 2- or 4-hydroxyestradiol. Alterations in the AhR signaling pathway could affect the risk of endometriosis through altered expression of CYP1A1 and CYP1B1 or increased proliferation of endometrial cells.

The most well-known AhR ligands are polycyclic aromatic hydrocarbons, including

2,3,7,8-tetrachlorodibenzo-p-dioxin (12). Recently, Ohtake et al. reported functional cross-talk between dioxin-activated AhR and estrogen receptors (13). Exposure to dioxins has been suggested as a risk factor for endometriosis (14). However, several studies reached different conclusions, and the issue remains controversial (15-16).

AhR function is regulated by the AhR nuclear translocator (ARNT) and the AhR repressor (AhRR). Ligand-bound AhR translocates to the nucleus, where it heterodimerizes with ARNT. The AhR-ARNT heterodimer binds to xenobiotic response element sequences and facilitates activation of target genes (17). AhRR competes with AhR in heterodimer formation, thus down-regulating the genes regulated by AhR (18). Both AhR and ARNT are expressed in the female reproductive tract, and changes in their expression have been reported in specific pathological conditions (19).

Polymorphic sites have been identified in the coding regions of the AhR, ARNT and AhRR genes, including *AhR* codon 554 in exon 10 (AGA to AAA, Arg to Lys), *ARNT* codon 189 in exon 7 (GTG to GTC, silent mutation) and *AhRR* codon 185 in exon 5 (CCC to GCC, Pro to Ala) (20-22).

Altered AhR-mediated signaling caused by polymorphisms in AhR, ARNT and/or AhRR may account for individual differences in susceptibility to endometriosis. However, one previous study found no association between *AhR*, *ARNT* and *AhRR*

polymorphisms and endometriosis (23). In the present study, we explored whether these polymorphisms contribute to the susceptibility to and severity of endometriosis. A case-control study was conducted in patients with different stages of endometriosis and controls.

MATERIALS AND METHODS

This study was approved by the Institutional Review Board of the University of Miyazaki, The Jikei University School of Medicine and National Cancer Center. All subjects gave their written informed consent before the laparoscopic examination.

Participants

During 1999-2000, 139 women were recruited at the Department of Obstetrics and Gynecology, Jikei University School of Medicine Hospital. Participants were patients between the ages of 20-45 who had complained of infertility and attended the hospital. The mean ages of cases and controls were similar (32 years for cases; 33 years for controls). No women ever had prior empiric therapy with either progestins or GnRH analogues before laparoscopic examination. Women who had ever given birth or lactated were not eligible for this study. One woman was excluded because a DNA sample was not available, leaving 138 women for the subsequent analysis.

All women underwent diagnostic laparoscopy as part of the infertility work-up. Women were classified into two groups according to the revised American Fertility Society (r-AFS) classification (24): endometriosis = r-AFS stage I - IV; controls = without endometriosis. For the most part, diagnosis was made by a single, trained

gynecologist. The diagnosis of endometriosis was established by visual criteria during laparoscopic examination, and histological confirmation was not always obtained. Of the 138 women enrolled, 59 had no endometriosis, 21 had stage I endometriosis, 10 had stage II, 23 had stage III and 25 had stage IV disease.

Genotyping

Blood samples were obtained before the laparoscopic examination. Genomic DNA was extracted from peripheral leucocytes using a DNA Extractor WB Kit (Wako, Osaka, Japan). Genotyping was performed blinded to case-control status, minimizing measurement bias.

AhR, ARNT and AhRR polymorphisms were genotyped by real-time PCR analysis on an ABI Prism 7000 Sequence Detection System (Applied Biosystems, Japan Ltd) using fluorescently labeled probes (25). All the primers and probes were designed appropriately with Primer Express 2.0 software (Applied Biosystems, Japan Ltd). Two differentially-labeled TaqMan probes and forward and reverse primers were prepared for each reaction. Primers and probes are listed in Table 1.

Reactions were performed with 200 nM of each probe, 900 nM each of forward primer and reverse primer, 1X TaqMan Universal PCR Master Mix (Applied

Biosystems), and 20 ng DNA. PCR cycling conditions consisted of one 2-minute cycle at 50°C, one 10-minute cycle at 95°C, followed by 40 cycles of 95°C for 15 seconds and 60°C for 1 minute. HPLC water was used as a negative PCR control in each amplification.

Statistical Analysis

Crude odds ratios (ORs) and 95% confidence intervals (95% CIs) were calculated for *AhR*, *ARNT* and *AhRR* genotypes to evaluate association with endometriosis. Since endogenous estrogen exposure was thought to be correlated with the risk of endometriosis, ORs were adjusted for risk factors that might affect endogenous estrogen exposure: age (<35, ≥35 years), menstrual cycle (<26, 26-30, ≥31 days) and duration of menstruation (<6, 6-8, ≥9 days) using multiple logistic regression analysis by SPSS for Windows software (version 11.0.1J, SPSS JAPAN, Tokyo, Japan) (26-27). The Wilcoxon rank-sum test for trend was also used to examine the associations between *AhR*, *ARNT* and *AhRR* polymorphisms and stage of endometriosis (28). All statistical tests were based on two-tailed probability. P values <0.05 were considered statistically significant.

RESULTS

The genotype and allele frequencies of each polymorphism are shown in Table 2. The distributions of genotypes among controls were in Hardy-Weinberg equilibrium.

The C/G + G/G genotypes at codon 185 of *AhRR* showed a significant association with risk of endometriosis compared to the C/C genotype (adjusted OR: 2.53; 95% CI, 1.16-5.55). No statistically significant association was observed between the *AhR* codon 554 and *ARNT* codon 189 polymorphisms and risk of endometriosis (Table 3).

Furthermore, we evaluated whether these polymorphisms were associated with the stage of endometriosis. There was a significant trend between C/G + G/G genotypes and the stage of endometriosis (P for trend, 0.02; Table 4). The *AhR* codon 554 and *ARNT* codon 189 polymorphisms were not associated with a higher risk of advanced clinical stage (data not shown).

DISCUSSION

In the present study, we evaluated whether polymorphisms at *AhR* codon 554, *ARNT* codon 189 and *AhRR* codon 185 contribute to the susceptibility to and severity of endometriosis. Our results indicate that the *AhRR* codon 185 polymorphism is associated with both susceptibility and severity. The genotypic distribution of this polymorphism was significantly different in endometriosis and controls. The risk of endometriosis was approximately 2.5 times higher with the *AhRR* C/G + G/G genotype. In addition, the *AhRR* C/G + G/G genotype was more frequently observed in patients with advanced stage endometriosis. No such associations were found with the *AhR* codon 554 and *ARNT* codon 189 polymorphisms. We should not, however, definitively conclude that no association exists between *AhR* codon 554 or *ARNT* codon 189 polymorphisms and susceptibility to and severity of endometriosis, because sample size was too small to allow detection of subtle differences.

In a previous study, Watanabe et al. failed to find a relation between the *AhRR* codon 185 polymorphism and endometriosis (23). There are several reasons that might account for the discrepancy. First and most important is the definition of endometriosis. Because endometriosis is sometimes asymptomatic, it can be diagnosed only by laparoscopy or laparotomy. In the present study, all participants underwent laparoscopy, and