

B-2-3. プロトコール

検査前に15分から20分の座位安静をとらせた後3分間安静時記録し、1分間負荷なしでエルゴメーターを漕いでもらうウォーミングアップの後、50W、3分間の運動負荷を行い、終了後8分間記録した。静脈採血は、負荷前、負荷直後および負荷8分後に行った。

C. 研究結果

C-1. 正常例での記録

血圧、心拍数、皮膚温、中心体温、負荷強度を示す(図 C-1)。負荷中、皮膚温は低下し、その後上昇を認めた。一方中心温は運動後徐々に上昇した。




図 C-1. 正常例での記録

C-2. 心拍数、血圧の変化

心拍数は正常群では不可直後より上昇し、負荷後速やかに減少したが、心不全群では負荷に対する反応が鈍く、負荷後の減少も緩やかであった。平均血圧は、正常群ではやや上昇した後速やかに減少したが、心不全群では心拍と同様、負荷に対する反応が鈍く、負荷後の減少も緩やかであった(図 C-2)。これらの反応は、今まで報告されたものと同様である

と考えられた。




図 C-2. 心拍数、血圧の変化

C-3. 体温の変化

正常群と、心不全群の運動による皮膚温と中心温の変化を示す。運動前の皮膚温は心不全群で低く、正常では皮膚温は運動により速やかに低下し、その後上昇するが、心不全では徐々に低下するのみであった。中心温は正常群では運動後徐々に上昇するのに対し、心不全群では徐々に低下した(図 C-3)。




図 C-3. 体温の変化

C-4. 酸素消費量

心不全群ではややピークの立ち上がりが悪い傾向がみられるが、max VO2 に両群での有意差は認めなかった(図 C-4)。




図 C-4. 酸素消費量

C-5. 血漿ノルエピネフリン値、BNP値

血漿ノルエピネフリン値は負荷前、負荷中および負荷後すべての時点で、心不全群で有意に高値であり、負荷による増加率も心不全群で有意に大であった。負荷前安静時のBNP値も心不全群で有意に高値であった(Table C-4)。

Table C-4

	Normal	CHF
PNE (ng/ml)		
Rest	222 ±69	370 ±190 *
During	317 ±90	547 ±288 *
After	249 ±58	383 ±173 *
Increment	91 ±104	182 ±168 *
BNP (pg/ml)	17 ±9	192 ±300 *

* P<0.05

D.考察

メカニズムとして考えられるアルゴリズムを図に示す(図 D)。正常例では、軽運動負荷により、骨格筋の血管抵抗の低下がおき、それによる平均血圧の低下にともなう圧受容器、交感神経系の緊張による皮膚の血管収縮にて皮膚温の低下がもたらされる。一方、運動による、骨格筋内熱産生および酸素消費量増加により深部体温は上昇する。よって運動により産生されたエネルギーは体温上昇にうまく反映される。一方、心不全患者では、安静時より交感神経活動が亢進しており、皮膚血管は収縮していると考えられる。運動負荷時にはさらに交感神経が緊張するが、安静時よりの皮膚血管収縮により、正常人に見られる収縮反応が傷害される。よって、皮膚からの熱放散が進むため、深部体温は正常人より低下すると考えられる。これらの反応により、同様の酸素消費による産生エネルギーは、心不全

例ではうまく蓄積されることが考えられる。




図 D. メカニズム

E.結論

軽度運動負荷時の体温変化を評価することにより、心不全の血流・神経調節の異常を評価することが可能であった。

F.健康危険情報

なし

G.研究発表

G-1.論文

1. Yamasaki F, Ushida T, Yokoyama T, Ando M, Yamashita K, Sato T. Artificial baroreflex: Clinical application of a bionic baroreflex system. *Circulation* 113: 634-639, 2006
2. Ando M, Katare RG, Kakinuma Y, Zhang D, Yamasaki F, Muramoto K, Sato T. Efferent vagal nerve stimulation protects heart against ischemia-induced arrhythmias by preserving connexin43 protein. *Circulation* 112: 164-170, 2005
3. Kakinuma Y, Ando M, Kuwabara M, Katare RG, Okudela K, Kobayashi M, Sato T. Acetylcholine from vagal

stimulation protects cardiomyocytes against ischemia and hypoxia involving additive nonhypoxic induction of HIF-1 α . FEBS Lett 579: 2111-2118, 2005

G-3.学会発表

1. 安藤元紀, カタレ G ラジエシユ, 柿沼由彦, 佐藤隆幸. 神経インターフェイス技法による心筋細胞死および致死性不整脈防止療法. 第 20 回生体・生理工学シンポジウム. 東京 (2005.09.05-07)
2. Katare RG, Ando M, Kakinuma Y, Yamasaki F, Sato T. Vagal nerve stimulation activates TNF-alpha and protects heart against acute ischemic injury. International Workshop on Gene Expression and Apoptosis-Associated Signal Transduction in Cancer Cells. Kochi. Japan (2005.08.20)
3. Katare RG, Ando M, Kakinuma Y, Yamasaki F, Sato T. Vagal nerve stimulation differentially regulates TNF receptors and protect the heart against acute ischemic injury. Circulation 112(SuppleII): 195-196, 2005. 78th Annual Scientific Sessions of American Heart Association. Dallas, USA (2005.11.13-16)
4. Kakinuma Y, Ando M, Katare RG, Yamasaki F, Sato T. Acetylcholine regulates a HIF-1 α -mediated gene, involved in cardiac energy metabolism suppression and cardioprotection against hypoxia. Circulation 112(SuppleII): 284, 2005. 78th Annual Scientific Sessions of American Heart Association. Dallas, USA (2005.11.13-16)
5. Ando M, Katare RG, Kakinuma Y, Li M, Zheng C, Yamasaki H, Sato T. Donepezil, acetylcholinesterase inhibitor, promotes angiogenesis over infarct area in rats after chronic coronary occlusion. Circulation 112(SuppleII): 288, 2005. 78th Annual Scientific Sessions of American Heart Association. Dallas, USA (2005.11.13-16)
6. 佐藤隆幸, 山崎文靖, 安藤元紀. 神経インターフェイス技法による血圧制御と心不全治療. 第 43 回日本人工臓器学会 34: S-40, 2005. 東京 (2005.11.30-12.02)
7. Katare RG, Ando M, Kakinuma Y, Kuwabara M, Yamasaki F, Sasaguri S, Sato T. Acetylcholine inhibits mitochondrial permeability transition pore and protects myocardium against acute ischemia-reperfusion injury. Circ J 2006;70 Suppl I:155. 第 70 回日本循環器学会 総会・学術集会. 名古屋 (2006.03.24-26)
8. Katare RG, Ando M, Kakinuma Y, Kuwabara M, Yamasaki F, Sato T. Vagal nerve stimulation differentially regulates TNF receptors and protects the heart against acute ischemic injury. Circ J 2006;70 Suppl I:169. 第 70 回日本循環器学会 総会・学術集会. 名古屋 (2006.03.24-26)
9. Katare RG, Ando M, Kakinuma Y, Kuwabara M, Yamasaki F, Sato T.

Acetylcholine inhibits opening of mitochondrial permeability transition pore(PTP) and enhances the functional recovery after long time hypothermic heart preservation. Circ J 2006;70 Suppl I:185. 第 70 回日本循環器学会総会・学術集会. 名古屋 (2006.03.24-26)

10. Katare RG, Ando M, Kakinuma Y, Kuwabara M, Yamasaki F, Sasaguri S, Sato T. Gap junctional communication between cardiomyocyte and fibroblast improves electrical conduction in 3D-engineered remodeled heart. Circ J 2006;70 Suppl I:476. 第 70 回日本循環器学会総会・学術集会. 名古屋 (2006.03.24-26)
11. Kakinuma Y, Ando M, Katare GR, Kuwabara M, Yamasaki F, Sato T. Activation of the non-neuronal cholinergic system by donepezil, an acetylcholinesterase inhibitor, involves angiogenesis through VEGF production. Circ J 2006;70 Suppl I:512. 第 70 回日本循環器学会総会・学術集会. 名古屋 (2006.03.24-26)

研究成果の刊行に関する一覧表レイアウト

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Ando M, Katare RG, Kakinuma Y, Zhang D, Yamasaki F, Muramoto K, Sato T.	Efferent vagal nerve stimulation protects heart against ischemia-induced arrhythmias by preserving connexin43 protein.	Circulation	112	164-170	2005
Yamasaki F, Furuno T, Sato K, Zhang D, Nishinaga M, Sato T, Doi Y, Sugiura T.	Association between arterial stiffness and platelet activation.	J Hum Hypertens	19	527-533	2005
Kakinuma Y, Ando M, Kuwabara M, Katare RG, Okudela K, Kobayashi M, Sato T.	Acetylcholine from vagal stimulation protects cardiomyocytes against ischemia and hypoxia involving additive nonhypoxic induction of HIF-1 α .	FEBS Lett	579	2111-2118	2005
Kudo Y, Yamasaki F, Kataoka H, Doi Y, Sugiura T.	Effect of serum albumin on QRS wave amplitude in patients free of heart disease.	Am J Cardiol	95	789-791	2005
Kawada T, Yamamoto K, Kamiya A, Ariumi H, Michikami D, Shishido T, Sunagawa K, Sugimachi M.	Dynamic characteristics of carotid sinus pressure-nerve activity transduction in rabbits.	Jpn J Physiol	55	157-163	2005
Kawada T, Yamazaki T, Akiyama T, Shishido T, Mori H, Sugimachi M.	Myocardial interstitial choline and glutamate levels during acute myocardial ischaemia and local ouabain administration.	Acta Physiol Scand	184	187-193	2005
Yamasaki F, Ushida T, Yokoyama T, Ando M, Yamashita K, Sato T	Artificial baroreflex: Clinical application of a bionic baroreflex system.	Circulation	113	634-639	2006
Uemura K, Kamiya A, Hidaka I, Kawada T, Shimizu S, Shishido T, Yoshizawa M, Sugimachi M, Sunagawa K.	Automated drug delivery system to control systemic arterial pressure, cardiac output, and left heart filling pressure in acute decompensated heart failure.	J Appl Physiol	100	1278-1286	2006

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
杉本健樹, 小林道也, 岡林雄大, 甫喜本兼弘, 西森秀明, 山崎文靖, 森本利明, 高橋保, 笹栗志朗, 荒木京二郎	Cisplatinの心嚢内注入が奏効した乳癌心嚢転移の1例.	癌と化学療法	32	1311-1313	2005
谷 俊一, 坪屋英志, 池本竜則	末梢神経におけるRapidly reversible block	末梢神経	16(2)	21-27	2005
谷 俊一	整形外科医のための電気生理学的検査入門	臨床雑誌 整形外科	56(2)	201-208	2005
谷 俊一	立体角近似による近接電場電位の解析: 伝導ブロックの波形変化	臨床神経生理学	33(4)	253-257	2005
牛田享宏, 谷 俊一	整形外科慢性疼痛疾患に対する当院集学的治療の試み	中部日本整形外科学会雑誌	48(5)	861-862	2005
川崎元敬, 谷口慎一郎, 牛田享宏, 谷 俊一	術中脊髄誘発電位検査で上位頸髄部において伝導ブロックを示した3例	西日本脊椎研究会誌	31(2)	73-76	2005
川田倫子, 谷口慎一郎, 石田健司, 牛田享宏, 岸本裕樹, 池本竜則, 谷 俊一	腰仙部神経根障害に対する後方手術時に記録した馬尾神経活動電位の検討	脊髄機能診断学	26(1)	109-112	2005
池本竜則, 谷口慎一郎, 牛田享宏, 谷 俊一	多椎間圧迫性頸髄症に対する前方除圧固定術と椎弓形成術の術後成績	中国・四国整形外科学会雑誌	17(2)	265-269	2005

研究成果の刊行物・別刷

Efferent Vagal Nerve Stimulation Protects Heart Against Ischemia-Induced Arrhythmias by Preserving Connexin43 Protein

Motonori Ando, PhD; Rajesh G. Katare, MD; Yoshihiko Kakinuma, MD; Dongmei Zhang, MD; Fumiyasu Yamasaki, MD; Kazuyo Muramoto, PhD; Takayuki Sato, MD

Background—Myocardial ischemia (MI) leads to derangements in cellular electrical stability and the generation of lethal arrhythmias. Vagal nerve stimulation has been postulated to contribute to the antifibrillatory effect. Here, we suggest a novel mechanism for the antiarrhythmogenic properties of vagal stimulation during acute MI.

Methods and Results—Under anesthesia, Wistar rats underwent 30 minutes of left coronary artery (LCA) ligation with vagal stimulation (MI-VS group, n=11) and with sham stimulation (MI-SS group, n=12). Eight of the 12 rats in the MI-SS group had ventricular tachyarrhythmia (VT) during 30-minute LCA ligation; on the other hand, VT occurred in only 1 of the 11 rats in the MI-VS group (67% versus 9%, respectively). Atropine administration abolished the antiarrhythmogenic effect of vagal stimulation. Immunoblotting revealed that the MI-SS group showed a marked reduction in the amount of phosphorylated connexin43 (Cx43), whereas the MI-VS group showed only a slight reduction compared with the sham operation and sham stimulation group ($37 \pm 20\%$ versus $79 \pm 18\%$). Immunohistochemistry confirmed that the MI-induced loss of Cx43 from intercellular junctions was prevented by vagal stimulation. In addition, studies with rat primary-cultured cardiomyocytes demonstrated that acetylcholine effectively prevented the hypoxia-induced loss of phosphorylated Cx43 and ameliorated the loss of cell-to-cell communication as determined by Lucifer Yellow dye transfer assay, which supports the *in vivo* results.

Conclusions—Vagal nerve stimulation exerts antiarrhythmogenic effects accompanied by prevention of the loss of phosphorylated Cx43 during acute MI and thus plays a critical role in improving ischemia-induced electrical instability. (*Circulation*. 2005;112:164-170.)

Key Words: arrhythmia ■ connexins ■ electrical stimulation ■ gap junctions ■ vagus nerve

Acute myocardial ischemia results in a dramatic reduction of tissue pH, an increase in interstitial potassium levels, an increase in intracellular calcium concentration, and neurohumoral changes, all of which contribute to the development of electrical instability that leads to life-threatening cardiac arrhythmias.^{1,2} In particular, cell-to-cell electrical uncoupling of ventricular myocytes plays an important role in arrhythmogenesis during acute and chronic ischemic heart disease.³⁻⁶

Gap-junction channels, composed of highly homologous proteins known as connexins in vertebrate species, have been implicated in the electrical coupling of excitable tissues, such as cardiac muscles. There are essentially 2 subtypes of connexins (Cx), Cx40 and Cx43, in the adult heart muscle. Cx43 is predominantly expressed in ventricular tissue, whereas Cx40 is mainly found in atrial tissue and in the conduction system.⁷ The protein content of ventricular Cx43 is remarkably reduced in ischemia^{8,9} and heart failure.^{10,11}

Gene-targeting studies demonstrate that reduced expression of Cx43 increases the incidence of ventricular tachyarrhythmias¹² and causes a significant reduction in conduction velocity in mice during acute myocardial ischemia.¹³ These results suggest that the dysfunction of Cx43 in cardiomyocytes could be one of the components of the substrate that promotes lethal ventricular tachyarrhythmias.

With regard to life-threatening arrhythmias in acute ischemia, the effect of vagal nerve stimulation (VS) has been reported to prevent ventricular fibrillation in dogs.¹⁴ Recently, VS therapy markedly improved long-term survival in an animal model of chronic heart failure after myocardial infarction,¹⁵ and earlier studies have demonstrated that ventricular arrhythmia is one of the major causes of death in chronic heart failure conditions.¹⁶ However, the mechanisms of VS on myocardial infarction remain unknown. With these in mind, we hypothesized that VS would exert an antiarrhythmogenic effect even in acute myocardial ischemia by target-

Received June 21, 2004; de novo received November 28, 2004; revision received March 25, 2005; accepted March 30, 2005.

From the Department of Cardiovascular Control (M.A., R.G.K., Y.K., D.Z., T.S.), Kochi Medical School, Nankoku, Japan; Department of Clinical Laboratory (F.Y.), Kochi Medical School, Nankoku, Japan; and Department of Integrative Physiology (K.M.), Kochi Medical School, Nankoku, Japan.

Correspondence to Takayuki Sato, Department of Cardiovascular Control, Kochi Medical School, Nankoku, Kochi 783-8505, Japan. E-mail tacsato-kochimed@umin.ac.jp

© 2005 American Heart Association, Inc.

Circulation is available at <http://www.circulationaha.org>

DOI: 10.1161/CIRCULATIONAHA.104.525493

ing the gap junctions. Therefore, in the present study, using both an *in vivo* acute ischemia model in rats and *in vitro* primary cultured cardiomyocytes from neonatal rats, we examined the effects of VS on arrhythmogenesis during acute myocardial ischemia.

Methods

The care and use of animals were in strict accordance with the guiding principles of the Physiological Society of Japan.

In Vivo Arrhythmia Study

Male Wistar rats (SLC, Japan) weighing 270 to 300 g were assigned to 6 groups receiving the following treatments: sham-operated rats treated with sham stimulation (SO-SS, $n=6$), sham-operated rats treated with vagal stimulation (SO-VS, $n=5$), myocardial ischemia rats with sham stimulation (MI-SS, $n=12$), myocardial ischemia rats with vagal stimulation (MI-VS, $n=11$), myocardial ischemia rats with both vagal stimulation and atropine administration (MI-VS-At, $n=6$), and myocardial ischemia rats preconditioned by vagal stimulation (pcVS-MI, $n=9$).

Acute Ischemia Model

After induction of anesthesia, the rat was ventilated artificially with a volume-controlled rodent respirator (model 683, Harvard Apparatus) at 80 strokes per minute. Anesthesia was maintained through the use of 1.2% halothane during surgical procedures and 0.6% halothane during data recording. Left ventricular MI was induced by 30 minutes of left coronary artery (LCA) ligation. In sham-operated rats, we loosely tied a suture around the LCA without arterial occlusion. For measurement of arterial pressure, a polyethylene tubing (PE-10, Becton Dickinson) that was filled with saline and connected to a fluid-filled transducer (DX-300, Viggo-Spectramed) was cannulated into the right femoral artery. Experimental solutions were infused through another polyethylene tubing, which was cannulated into the right femoral vein. Atropine sulfate (1 mg/kg) (Sigma) was administered as vagal efferent muscarinic blockade. For the prevention of dehydration during experiments,¹⁷ physiological saline was continuously infused at a rate of $5 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ with a syringe pump (CFV-3200; Nihon Kohden).

Vagal Nerve Stimulation

The right vagal nerve was identified, isolated, and cut in the neck region. Only the distal end of the vagal nerve was placed on a pair of platinum wires and was used for stimulation to exclude the effects of vagal afferent. The electrode was connected to an isolated constant voltage stimulator (SS-202J and SEN-7203, Nihon Kohden). The vagal nerve was stimulated with electrical rectangular pulses of 0.1-ms duration at 10 Hz during LCA ligation. The electrical voltage of pulses was optimized in each rat to obtain a 10% reduction in heart rate before LCA ligation. The actual electrical voltage was in the range of 2 to 6 V. VS was started at 1 minute before LCA ligation and continued for 30 minutes after LCA ligation. A run of first spontaneous ventricular tachyarrhythmia (VT) was defined as 10 or more beats with a cycle length <100 ms.

To exclude vagally induced bradycardiac effects during MI and to investigate whether or not a heart preconditioned by VS (pcVS) was insusceptible to ischemia-induced VT, we also examined the effect of 10-minute VS only before LCA ligation. After a 5-minute stabilization period for the recovery of heart rate, the rat was subjected to 30-minute LCA ligation.

Risk Area Assessment

To confirm the ischemic area induced by LCA ligation, 2 mL of 2% Evans blue dye was injected via the femoral vein at the end of 30 minutes of coronary artery ligation. The area at risk was determined by negative staining with Evans blue. Area measurements were determined with Image-Pro version 4.0 (Media Cybernetics).

Excised Heart Study

We conducted this protocol separately from the *in vivo* arrhythmia study because the experimental preparation of Evans blue dye injection for risk area assessment did not allow us to perform immunoblotting and immunohistochemical assay of excised hearts. For each group, we prepared and analyzed 5 hearts.

Protein Preparation and Immunoblotting

Pulverized frozen left ventricle samples were suspended in 40 vol of ice-cold 10% trichloroacetic acid and homogenized with a tissue homogenizer (2 bursts, 30 seconds each).¹⁸ Homogenates were centrifuged at $10\,000g$ for 10 minutes. Supernatants were discarded, and the remaining pellets were resuspended in sampling solution (9 mol/L urea, 0.065 mol/L dithiothreitol, 2% Triton X-100) and sonicated. After addition of 0.37 mol/L lithium dodecylsulfate, the protein-containing solution was neutralized by 1 mol/L Tris solution, and sonicated again. For immunoblot analysis, proteins separated by SDS-PAGE with 15% polyacrylamide gels were transferred electrophoretically to polyvinylidene difluoride sheets (Millipore). After 1 hour of blocking in the 4% skimmed milk solution, the membrane was incubated overnight with anti-Cx43 antibody (71-0700, Zymed) diluted 1:1000. Blots were then incubated for 1 hour with horseradish peroxidase-conjugated goat anti-rabbit IgG, developed with an ECL chemiluminescence reagent (Amersham), and exposed to a medical x-ray film. Equal protein content in all the samples was confirmed by Coomassie brilliant blue staining.

Immunohistochemistry and Confocal Microscopy

Transmural blocks of left ventricular myocardium from selected hearts were immersed in a fixative containing 4% paraformaldehyde and 0.1 mol/L phosphate buffer (pH 7.4), embedded in paraffin, and sectioned at a thickness of $4 \mu\text{m}$. Sections were deparaffinized, placed in citrate buffer, and boiled in a microwave oven for 10 minutes to enhance specific immunostaining. The sections were then incubated overnight with anti-Cx43 antibody diluted 1:100 and then incubated for 2 hours in Alexa546-conjugated goat anti-rabbit IgG (Molecular Probes) diluted 1:100. Fluorescence of Alexa546 was observed with a confocal laser scanning microscope system (FV300, Olympus). Reconstructed projection images were obtained from serial optical sections recorded at an interval of $0.5 \mu\text{m}$.

Primary Culture Study

We used neonatal rat primary cultured cardiomyocytes to investigate the direct action of acetylcholine (ACh), a neurotransmitter released by VS, on phosphorylated Cx43. Hearts were excised from 1- to 2-day-old neonatal Wistar rats and digested with 0.03% type IV collagenase (Sigma) and 0.03% trypsin (Invitrogen). After digestion, cardiomyocytes were washed repeatedly in serum containing culture medium and were preplated for 90 minutes in the noncoated culture dish in the presence of DMEM medium (Sigma) with 10% FBS to remove the contaminating noncardiomyocytes. Next, cardiomyocytes were plated and cultured in gelatin-coated culture dishes in DMEM medium with 10% FBS. New medium was replaced every day, and all experiments were performed after 4 days of culture.

Hypoxia and Immunoblotting

For hypoxia experiments, cardiomyocytes in serum-deficient DMEM were transferred into an airtight incubator and maintained at a humidified hypoxic atmosphere of $<2\%$ O_2 with nitrogen and 5% CO_2 for 30 minutes. Cells were treated either with ACh (0.5 mmol/L) alone or with both ACh and atropine (0.1 mmol/L) 10 minutes before the cells were subjected to hypoxia. After 30 minutes, the treated cells were sampled for immunoblotting.

Dye Transfer Assay

To assess the effect of ACh on cell-to-cell communication, we monitored dye transfer in primary cultured cardiomyocytes with Lucifer Yellow CH (LY; lithium salt, Molecular Probes). Cells in the culture medium were viewed on a Zeiss microscope equipped with fluorescence illumination and FITC filters. The emitted fluorescent light was projected onto a CCD camera system (C4742-98-24ER,




Figure 1. Example of run of VT induced by LCA ligation in rat treated with sham stimulation. In this case, ventricular fibrillation was initiated after ventricular tachycardia. SAP indicates systemic arterial pressure.

Hamamatsu Photonics). LY was injected electrophoretically via a conventional intracellular microelectrode made from borosilicate glass capillaries. The outer diameter of microelectrodes was 0.15 to 0.25 μm , and the tip resistance of these electrodes filled with 3% LY was 0.6 to 1.4 $\text{G}\Omega$. An inward square-pulse current (20 to 40 nA) of 0.5-second duration was injected at a frequency of 1 Hz for 2 to 3 minutes using the circuitry of an amplifier (707, WPI) and a voltage pulse generator (SEN7203, Nihon Kohden). For induction of the hypoxic condition, cardiomyocytes were treated for 5 hours with 0.2 mmol/L CoCl_2 ,¹⁹ because microinjection methods are not suitable for gas-tight chamber experiments with N_2 substitution. Cells were treated either with ACh (0.5 mmol/L) alone or with both ACh and atropine (0.1 mmol/L) before chemical hypoxia. Dye injection was performed at room temperature (24°C to 26°C).

Statistical Analysis

Time of onset of the first run of VT was analyzed by the Kaplan-Meier method, and comparisons were made with the Mantel-Haenszel log-rank test. A nonparametric test for comparison of treatments with a control was performed by a Mann-Whitney *U* test with Bonferroni adjustment. Differences were considered significant at $P < 0.05$. Values are expressed as mean \pm SD.

Results

In Vivo Arrhythmia Study

Effects of VS on Acute Ischemia-Induced VT

Figure 1 illustrates an example of the VT observed after LCA ligation with sham stimulation in the MI-SS group. As demonstrated in this case, VT followed by ventricular fibrillation was often detected after LCA ligation without VS. Figure 2 shows the time of onset and the incidence of the first spontaneous VT after LCA ligation. Eight of 12 rats in the MI-SS group experienced VT; on the other hand, only 1 of 11 in the MI-VS group developed VT during the 30-minute LCA ligation (67% versus 9%, $P = 0.005$). VS achieved an 87% reduction in the relative incidence ratio of VT. In all cases, the first runs of VT were observed within 15 minutes after LCA ligation. Atropine administration abolished the anti-VT effects of VS; 4 of 6 rats developed VT (Figure 2). No VT was detected in either the SO-SS or SO-VS groups.

To exclude vagally induced bradycardiac effects on the incidence of VT, a group of 9 rats were subjected to preconditioning by VS (pcVS-MI group; see Methods). There was no significant difference in heart rate between the MI-SS and pcVS-MI groups immediately before and after LCA ligation (Table). Nevertheless, only 2 of 9 rats in the pcVS-MI group developed VT (22%, $P = 0.04$ versus the MI-SS group).




Figure 2. Incidence and onset of first VT after LCA ligation. Myocardial ischemia rats treated with sham stimulation (MI-SS, \bullet , $n = 12$) and with vagal stimulation (MI-VS, \blacksquare , $n = 11$). VS significantly ($P = 0.005$ vs MI-SS group) decreased incidence of ischemia-induced VT. In contrast, atropine administration blocked antiarrhythmic effect by VS during acute MI (MI-VS-Atr, \blacktriangle , $n = 6$). Preconditioning by VS significantly ($P = 0.04$ vs MI-SS group) decreased incidence of ischemia-induced VT (pcVS-MI, \circ , $n = 9$).

Risk Area Assessment and Hemodynamic Parameters

There was no significant difference in risk area between the MI-SS and MI-VS groups ($55 \pm 6\%$ versus $59 \pm 3\%$). The differences in heart rate between groups with or without VS reached ≈ 50 bpm, and after 30-minute ischemia, only the MI-VS group had a significantly lower arterial pressure than the SO-SS group (Table).

Excised Heart Study

Effects of VS on Cx43 Expression

The polyclonal anti-Cx43 antibody in the present study showed closely spaced bands migrating between 43 and 46 kDa and a faint band migrating at 41 kDa. Previous reports²⁰ demonstrated that the higher- and lower-molecular-weight bands represent phosphorylated and nonphosphorylated isoforms of Cx43, respectively. Figure 3A shows a representative immunoblot prepared with anti-Cx43 antibody. The phosphorylated isoform of Cx43 (bands with 43 kDa) in the MI-VS group was preserved compared with that in the MI-SS group. Quantitative densitometric analysis revealed the level of the phosphorylated isoform of Cx43 in the MI-VS group was maintained at $79 \pm 18\%$ of that in the SO-SS group; on the other hand, the level of phosphorylated Cx43 in the MI-SS group was reduced significantly to $37 \pm 20\%$ of that in the SO-SS group (Figure 3B). The phosphorylated Cx43 in the SO-VS group was increased significantly to $120 \pm 8\%$ of that in the SO-SS group. Atropine inhibited the preserving effects of VS on Cx43 in the MI-VS-Atr group ($47 \pm 12\%$). In addition, protein analysis also confirmed that the phosphorylated Cx43 in the pcVS-MI group was at almost the same level as in the SO-SS group ($97 \pm 20\%$).

Effects of VS on Cx43 Localization

To investigate the distribution of Cx43 during acute ischemia with or without VS, we performed confocal image analysis of left ventricular tissues stained with anti-Cx43 antibody. As shown in Figure 4A, localization of immunoreactive signals

Heart Rate and Mean Arterial Pressure Immediately Before and After 30-Minute LCA Ligation

Group	n	Heart Rate, bpm		Mean Arterial Pressure, mm Hg	
		Before	After 30-Minute Ligation	Before	After 30-Minute Ligation
SO-SS	6	404±24	414±24	118±14	116±13
SO-VS	5	362±10*	348±15*	106±9	96±6
MI-SS	12	395±22	365±43	117±7	102±10
MI-VS	11	354±12*	353±31*	102±13	90±6*
MI-VS-Atr	6	409±20	362±48	103±8	94±15
pcVS-MI	9	391±33	372±36	109±11	103±10

VS was started at 1 minute before LCA ligation and continued for 30 minutes after LCA ligation. pcVS was performed during 10 minutes without LCA ligation, and after a 5-minute stabilization period for recovery of heart rate, rats were subjected to a 30-minute LCA ligation. Values are mean±SD immediately before and after 30-minute LCA ligation.

* $P < 0.05$ vs SO-SS group.

in the SO-SS group was restricted to intercellular junctions, consistent with the gap junctions and intercalated disks. In contrast, the Cx43 signal was reduced dramatically in the MI-SS group (Figure 4B); however, the Cx43 signal in the MI-VS group was almost comparable to the level in the SO-SS group (Figure 4C). These observations indicate that the loss of phosphorylated Cx43 during acute ischemia was prevented by VS.

Primary Culture Study

Effects of Hypoxia and ACh on Cx43 Expression

To examine the muscarinic effect on hypoxia-induced loss of phosphorylated Cx43, immunoblot analysis was performed in

primary cultured cardiomyocytes. Under normoxic conditions, addition of ACh for 30 minutes significantly increased the phosphorylated isoform of Cx43 compared with the control group (Figure 5). Although hypoxia for 30 minutes decreased phosphorylated Cx43, ACh treatment prevented this hypoxia-induced loss of Cx43. Atropine inhibited the preserving effects of ACh. Thus, these results indicate that ACh preserved phosphorylated Cx43 through a muscarinic receptor during 30-minute hypoxia.

Effects of Hypoxia and ACh on Intercellular Coupling and Beating Rate

To assess whether the induction of phosphorylated Cx43 by ACh results in a functional effect on cell-to-cell communication, LY dye transfer analysis was performed in primary cultured cardiomyocytes. As shown in Figure 6A, under normoxic conditions, LY injected into a cardiomyocyte diffused into other cardiomyocytes around the injected cell. In contrast, LY injected into a cardiomyocyte under chemical hypoxic conditions was confined to the injected cell, and no dye coupling with other cardiomyocytes was observed (Figure 6B); on the other hand, ACh administration preserved cell-to-cell communication (Figure 6C). Atropine treatment inhibited the preventive effect of ACh on hypoxia-induced uncoupling (Figure 6D). Under normoxic conditions, ACh treatment remarkably induced dye coupling between cardiomyocytes (Figure 6E).

Under normoxic conditions, the rate of spontaneous beating was 62 ± 13 bpm ($n = 5$ experiments), and ACh did not slow the beating rate. Chemical hypoxia with CoCl_2 stopped the beating; however, even under hypoxic conditions, ACh preserved spontaneous beating. In the presence of atropine, ACh failed to prevent the hypoxia-induced cessation of beating. The results of Cx43 immunoblotting under chemical hypoxic conditions were identical to those under hypoxia with $< 2\%$ O_2 (data not shown).

Discussion

In the present study, we investigated whether VS could attenuate acute MI-induced arrhythmogenic properties via modulation of a principal cardiac gap-junction protein, Cx43. Our results provide novel evidence that VS effectively




Figure 3. A, Representative immunoblots of homogenate of left ventricles from SO-SS, SO-VS, MI-SS, MI-VS, MI-VS-Atr, and pcVS-MI rats, probed with polyclonal Cx43 antibody. Arrows indicate position of phosphorylated isoform of Cx43 (43 kDa) and nonphosphorylated isoform of Cx43 (41 kDa) bands, respectively. B, Quantitative densitometric analysis of phosphorylated isoform of Cx43 in SO-SS, SO-VS, MI-SS, MI-VS, MI-VS-Atr, and pcMI-VS groups. Data are expressed as mean±SD. a.u. indicates arbitrary units. $n = 5$ for each group. * $P < 0.05$.




Figure 4. Representative confocal images of rat left ventricles from SO-SS rats (A), MI-SS rats (B), and MI-VS rats (C). Positive immunoreactive signals were concentrated in discrete spots at sites of intercellular apposition (red).

inhibits loss of the phosphorylated isoform of Cx43 during acute MI. Although the precise mechanism by which VS modulates the dephosphorylation of Cx43 remains unknown, it is most likely that VS exerts its antiarrhythmic effects on ischemic ventricular myocytes through the preserved function of Cx43.

VS and Antiarrhythmic Properties

VS has already been reported to prevent ventricular fibrillation in dogs.¹⁴ In the present study, we hypothesized that VS exerts its antiarrhythmic properties by maintaining electrical coupling with ventricular cardiomyocytes as a result of prevention of Cx43 dephosphorylation induced by acute MI. However, because VS simultaneously evokes a bradycardiac effect, the question remains whether the heart rate deceleration caused by VS is a primary mechanism for antiarrhythmic properties during MI. In a preliminary study, we confirmed




Figure 5. Representative immunoblots of homogenates from primary cultured rat cardiomyocytes probed with polyclonal anti-Cx43 antibody. Hypoxic treatment for 30 minutes decreased phosphorylated Cx43. ACh treatment prevented hypoxia-induced loss of phosphorylated Cx43. Atr inhibited ACh effect. Under normoxic conditions, ACh significantly increased phosphorylated isoform of Cx43 compared with control group.

that short-term exposure of cultured cardiomyocytes to ACh only before hypoxia prevented the hypoxia-induced loss of phosphorylated Cx43 (unpublished observations, 2004). Therefore, it is conceivable that ACh had a cardioprotective effect independent of the heart rate–slowing mechanism during hypoxia or ischemia. To further clarify such a preconditioning effect in *in vivo* experiments, we examined whether hearts preconditioned by VS were unsusceptible to ischemia-induced VT and whether pcVS prevented the ischemia-induced loss of Cx43. As expected, we confirmed that pcVS exerted its antiarrhythmic effects and sustained the level of phosphorylated Cx43 during ischemia. These results suggest that VS or ACh had a cardioprotective effect independent of the heart rate–slowing mechanism.

It is well recognized that Cx43, which is the principal component of ventricular gap-junction proteins, contributes to intercellular communication and electrical coupling. Beardslee et al²¹ showed that Cx43 underwent marked dephosphorylation during the process of electrical uncoupling induced by ischemia. Genetically engineered Cx43-deficient (Cx43^{+/-} or Cx43^{-/-}) mice have been reported to be markedly susceptible to ischemia-induced VT.^{12,13,22} In the present study, VS drastically reduced the incidence of VT and prevented the loss of phosphorylated Cx43 during acute MI. Therefore, functional preservation of Cx43 by VS would play an important role in antiarrhythmic properties during acute MI.

The result that ACh administration ameliorated the hypoxia-induced loss of dye coupling in cardiomyocytes is consistent with that of ACh-induced upregulation of phosphorylated Cx43. Under normoxic conditions, ACh did not slow down the spontaneous beating rate of cardiomyocytes. Hypoxia stopped the beating and diminished the phosphorylated isoform of Cx43; however, even under hypoxic conditions, ACh preserved the spontaneous beating and the phosphorylated isoform of Cx43. Therefore, it is conceivable that ACh has a cardioprotective effect independent of the beating rate.

Upregulation of Cx43 has been reported to accelerate spontaneous beating in cultured cardiomyocytes.²³ Moreover,




Figure 6. LY dye transfer assay in primary cultured rat cardiomyocytes. Each panel consists of fluorescent micrograph and transmission light micrograph. A, Control cultured cardiomyocytes. B, Cardiomyocytes treated with CoCl_2 . C, Cardiomyocytes treated with ACh and CoCl_2 . D, Cardiomyocytes treated with Atr, ACh, and CoCl_2 . E, Cardiomyocytes treated with ACh alone. Asterisks indicate dye-injected cells; white arrows, coupled neighbors.

cultured cardiomyocytes from genetically engineered Cx43-deficient ($\text{Cx43}^{-/-}$) mice demonstrated slow spontaneous beating rates and were poorly synchronized with each other compared with wild-type cultured cardiomyocytes.²⁴ From these findings and the present results, we speculate that ACh exerts its antiarrhythmic properties on ischemic or hypoxic hearts by preserving Cx43 and that such a beneficial effect is independent of its bradycardiac effect.

The spatial distribution of Cx43 can influence electrical stability of the heart. A more recent study by Poelzing and Rosenbaum¹¹ has shown that the transmural derangement of Cx43 expression can potentially be an arrhythmogenic substrate in the canine model of pacing-induced heart failure. Although we did not evaluate the transmural heterogeneity of Cx43 expression in the present study, such an analysis would be needed to clarify precise mechanisms for the antiarrhythmic effects of VS.

How Does VS Modulate Phosphorylated Cx43?

Three potential mechanisms might be involved in the linkage between VS and the sustained phosphorylated-protein level

of Cx43 during acute MI. First, VS may activate several protein kinases and induce phosphorylation of Cx43 through muscarinic receptors.²⁵ Liu et al²⁶ demonstrated that ACh prevented ischemic injury in cultured cardiomyocytes by activating protein kinase C, and the protective effect was mediated through a nitric oxide-dependent pathway. Second, VS can block the degradation pathway of Cx43 during acute MI. It has been reported that Cx43 is a short-lived protein with a half-life of only 1 to 3 hours in adult hearts²⁰ and that both lysosomal and proteasomal degradation play distinct roles in the life cycle of Cx43.²⁷ Our observations in the present study suggest that VS may prevent the ischemia-induced loss of Cx43 as a consequence of inhibition of its degradation pathway. Third, VS is also postulated to suppress excessive inflammation. Recently, Tracey and colleagues^{28,29} identified a novel molecular link between the vagus nerve system and an antiinflammatory response to disease. They suggest that VS exerts an antiinflammatory effect via the nicotinic ACh receptor $\alpha 7$ -subunit expressed in macrophages. The release of tumor necrosis factor- α from macrophages is inhibited by nicotinic stimulation. In contrast, the present results indicate that VS exerts its antiarrhythmic effects via muscarinic cholinergic receptors.

Study Limitations

In the present study, we did not measure the myocardial interstitial level of ACh at the ischemic region during VS. Therefore, it is unclear whether the cardiac vagal efferent fiber innervating the ischemic region can release its neurotransmitter in response to electrical stimulation. An *in vivo* microdialysis technique has enabled monitoring of the local concentration of neurotransmitters such as catecholamines, amino acids, and ACh. In the cat heart, Kawada et al showed that VS increased the myocardial interstitial ACh level even in the ischemic region (e-mail, February 12, 2004). Although their previous study³⁰ showed that acute MI induced the nerve-firing independent release of ACh from the vagal terminal, the electrical stimulation of the vagal efferent during acute MI produced the significant additive release of ACh to the myocardial interstitial space. Such an additional release of ACh in response to electrical stimulation during acute MI would play an important role in Cx43 preservation in the ischemic region.

Conclusion and Future Aspects

The present study demonstrates that VS exerts antiarrhythmic effects during acute MI accompanied by prevention of the loss of phosphorylated Cx43. The preserved function of Cx43 may improve electrical instability during acute MI. In view of the present results, we can provide an alternative therapeutic strategy with a neural interface approach. We have already developed the sympathetic interface approach for the treatment of central baroreflex failure in rats.³¹ To establish the therapeutic strategy shown here, further studies are required.

Acknowledgments

This study was supported by Health and Labor Sciences Research Grant (H15-PHYSI-001) for Advanced Medical Technology from

the Ministry of Health, Labor and Welfare of Japan. We thank Ken-ichi Yagyu for his technical assistance.

References

- Zipes DP, Wellens HJ. Sudden cardiac death. *Circulation*. 1998;98:2334–2351.
- Mehta D, Curwin J, Gomes JA, Fuster V. Sudden death in coronary artery disease: acute ischemia versus myocardial substrate. *Circulation*. 1997;96:3215–3223.
- Smith WT IV, Fleet WF, Johnson TA, Engle CL, Cascio WE. The Ib phase of ventricular arrhythmias in ischemic in situ porcine heart is related to changes in cell-to-cell electrical coupling. *Circulation*. 1995;92:3051–3060.
- Cinca J, Blanch P, Carreno A, Mont L, Garcia-Burillo A, Soler-Soler J. Acute ischemic ventricular arrhythmias in pigs with healed myocardial infarction: comparative effects of ischemia at a distance and ischemia at the infarct zone. *Circulation*. 1997;96:653–658.
- Saffitz JE, Schuessler RB, Yamada KA. Mechanisms of remodeling of gap junction distributions and the development of anatomic substrates of arrhythmias. *Cardiovasc Res*. 1999;42:309–317.
- Severs NJ. Gap junction remodeling in heart failure. *J Card Fail*. 2002;8:S293–S299.
- Dhein S, Polontchouk L, Salameh A, Haefliger JA. Pharmacological modulation and differential regulation of the cardiac gap junction proteins connexin 43 and connexin 40. *Biol Cell*. 2002;94:409–422.
- Peters NS, Green CR, Poole-Wilson PA, Severs NJ. Reduced content of connexin43 gap junctions in ventricular myocardium from hypertrophied and ischemic human hearts. *Circulation*. 1993;88:864–875.
- Kaprielian RR, Gunning M, Dupont E, Sheppard MN, Rothery SM, Underwood R, Pennell DJ, Fox K, Pepper J, Poole-Wilson PA, Severs NJ. Downregulation of immunodetectable connexin43 and decreased gap junction size in the pathogenesis of chronic hibernation in the human left ventricle. *Circulation*. 1998;97:651–660.
- Spragg DD, Leclercq C, Loghmani M, Faris OP, Tunin RS, DiSilvestre D, McVeigh ER, Tomaselli GF, Kass DA. Regional alterations in protein expression in the dyssynchronous failing heart. *Circulation*. 2003;108:929–932.
- Poelzing S, Rosenbaum DS. Altered connexin43 expression produces arrhythmia substrate in heart failure. *Am J Physiol*. 2004;287:H1762–H1770.
- Lerner DL, Yamada KA, Schuessler RB, Saffitz JE. Accelerated onset and increased incidence of ventricular arrhythmias induced by ischemia in Cx43-deficient mice. *Circulation*. 2000;101:547–552.
- Gutstein DE, Morley GE, Tamaddon H, Vaidya D, Schneider MD, Chen J, Chien KR, Stuhlmann H, Fishman GI. Conduction slowing and sudden arrhythmic death in mice with cardiac-restricted inactivation of connexin43. *Circ Res*. 2001;88:333–339.
- Vanoli E, De Ferrari GM, Stramba-Badiale M, Hull SS Jr, Foreman RD, Schwartz PJ. Vagal stimulation and prevention of sudden death in conscious dogs with a healed myocardial infarction. *Circ Res*. 1991;68:1471–1481.
- Li M, Zheng C, Sato T, Kawada T, Sugimachi M, Sunagawa K. Vagal nerve stimulation markedly improves long-term survival after chronic heart failure in rats. *Circulation*. 2004;109:120–124.
- Engelstein ED. Prevention and management of chronic heart failure with electrical therapy. *Am J Cardiol*. 2003;91:62F–73F.
- DiBona GF, Sawin LL. Reflex regulation of renal nerve activity in cardiac failure. *Am J Physiol*. 1994;266:R27–R39.
- Kikuchi H, Fujinawa T, Kuribayashi F, Nakanishi A, Imajoh-Ohmi S, Goto M, Kanegasaki S. Induction of essential components of the superoxide generating system in human monoblastic leukemia U937 cells. *J Biochem*. 1994;116:742–746.
- Chun YS, Hyun JY, Kwak YG, Kim IS, Kim CH, Choi E, Kim MS, Park JW. Hypoxic activation of the atrial natriuretic peptide gene promoter through direct and indirect actions of hypoxia-inducible factor-1. *Biochem J*. 2003;370:149–157.
- Beardslee MA, Laing JG, Beyer EC, Saffitz JE. Rapid turnover of connexin43 in the adult rat heart. *Circ Res*. 1998;83:629–635.
- Beardslee MA, Lerner DL, Tadros PN, Laing JG, Beyer EC, Yamada KA, Kleber AG, Schuessler RB, Saffitz JE. Dephosphorylation and intracellular redistribution of ventricular connexin43 during electrical uncoupling induced by ischemia. *Circ Res*. 2000;87:656–662.
- Yao JA, Gutstein DE, Liu F, Fishman GI, Wit AL. Cell coupling between ventricular myocyte pairs from connexin43-deficient murine hearts. *Circ Res*. 2003;93:736–743.
- Ai Z, Fischer A, Spray DC, Brown AM, Fishman GI. Wnt-1 regulation of connexin43 in cardiac myocytes. *J Clin Invest*. 2000;105:161–171.
- Vink MJ, Suadcani SO, Vieira DM, Urban-Maldonado M, Gao Y, Fishman GI, Spray DC. Alterations of intercellular communication in neonatal cardiac myocytes from connexin43 null mice. *Cardiovasc Res*. 2004;62:397–406.
- van Koppen CJ, Kaiser B. Regulation of muscarinic acetylcholine receptor signaling. *Pharmacol Ther*. 2003;98:197–220.
- Liu H, McPherson BC, Zhu X, Da Costa ML, Jeevanandam V, Yao Z. Role of nitric oxide and protein kinase C in ACh-induced cardioprotection. *Am J Physiol*. 2001;281:H191–H197.
- Qin H, Shao Q, Igdoura SA, Alaoui-Jamali MA, Laird DW. Lysosomal and proteasomal degradation play distinct roles in the life cycle of Cx43 in gap junctional intercellular communication-deficient and -competent breast tumor cells. *J Biol Chem*. 2003;278:30005–30014.
- Tracey KJ. The inflammatory reflex. *Nature*. 2002;420:853–859.
- Wang H, Yu M, Ochani M, Amella CA, Tanovic M, Susarla S, Li JH, Wang H, Yang H, Ulloa L, Al-Abed Y, Czura CJ, Tracey KJ. Nicotinic acetylcholine receptor alpha7 subunit is an essential regulator of inflammation. *Nature*. 2003;421:384–388.
- Kawada T, Yamazaki T, Akiyama T, Sato T, Shishido T, Inagaki M, Takaki H, Sugimachi M, Sunagawa K. Differential acetylcholine release mechanisms in the ischemic and non-ischemic myocardium. *J Mol Cell Cardiol*. 2000;32:405–414.
- Sato T, Kawada T, Sugimachi M, Sunagawa K. Bionic technology revitalizes native baroreflex function in rats with baroreflex failure. *Circulation*. 2002;106:730–734.

CLINICAL PERSPECTIVE

Increased cardiac vagal tone reduces ventricular arrhythmias during acute myocardial ischemia and has been linked to a lower risk of sudden arrhythmic death. Although the benefit of bradycardia associated with vagal tone is well appreciated, this may not be the only benefit. Cardiomyocytes are electrically coupled to one another through gap junctions. This coupling is critical to maintenance of cardiac electrical stability. Uncoupling occurs during ischemia and promotes heterogeneity of repolarization and slowing of conduction, with proarrhythmic effects. In the present study, short-term vagal stimulation (VS) was applied before or during acute ischemia in rats. VS protected against ventricular arrhythmias. Furthermore, VS preserved a phosphorylated form of connexin 43 (Cx43), a major subtype of gap-junction proteins in ventricles. In vitro studies of rat primary-cultured cardiomyocytes showed that ACh, a vagal efferent neurotransmitter, effectively prevented hypoxia-induced loss of phosphorylated Cx43 proteins and maintained cell-to-cell communication. Antiarrhythmic properties of VS and ACh were mediated via muscarinic receptors but were independent of heart rate deceleration. That cellular coupling can be improved through neural stimulation may lead to novel therapeutic strategies for preventing ventricular fibrillation during ischemia.

ORIGINAL ARTICLE

Association between arterial stiffness and platelet activation

F Yamasaki¹, T Furuno², K Sato², D Zhang³, M Nishinaga², T Sato³, Y Doi² and T Sugiura¹
¹Department of Clinical Laboratory, Kochi Medical School, Nankoku, Kochi, Japan; ²Department of Medicine and Geriatrics, Kochi Medical School, Nankoku, Kochi, Japan; ³Department of Cardiovascular Control, Kochi Medical School, Nankoku, Kochi, Japan

Increased arterial stiffness is strongly associated with atherosclerosis, while platelet activation is an important trigger of thrombotic events in patients with atherosclerosis. However, little is known about the effect of arterial stiffness on platelet activation. We therefore investigated the association between arterial stiffness and platelet activation in 38 normal volunteers (20 men and 18 women) aged 23–77 years (mean = 49 ± 15 years). Arterial stiffness was assessed by measuring brachial–ankle pulse wave velocity (ba-PWV) and heart–brachial PWV (hb-PWV). Flow cytometric analyses were performed to evaluate platelet activation by measuring surface expression of P-selectin and platelet–neutrophil complexes (PNC) before and after activation by ADP. We also calculated the difference between basal and stimulated states of P-selectin and PNC to assess platelet activation reserve. PWVs were significantly

correlated with age and BP ($r=0.60–0.81$). For platelet activation and activation reserve, correlations with age were less strong but remained significant ($r=0.36–0.61$), with the exception of P-selectin (not significant, NS), and correlations with SBP were similar ($r=0.35–0.53$). A significant correlation was found between PWVs and platelet activation ($r=0.43–0.74$). Multiple regression analysis demonstrated significant correlations between platelet activation and reserve and PWVs (coefficient = 2.17–6.59), when both age and BP were adjusted for simultaneously. In conclusion, platelet activation was associated with arterial stiffness, suggesting that arterial stiffness may play an important role in thrombotic events.

Journal of Human Hypertension (2005) 19, 527–533.

doi:10.1038/sj.jhh.1001861

Published online 7 April 2005

Keywords: arterial stiffness; pulse wave velocity; P-selectin; platelet–neutrophil complexes

Introduction

Platelet activation and aggregation are important triggers of thrombotic events in patients with atherosclerosis. In such patients, platelets are activated at the site of atheroma¹ due to increased shear stress in the narrowed vessels.^{2,3} Increased platelet activation is observed in patients with coronary risk factors and cardiovascular events.^{4–12}

Increased arterial stiffness, measured with pulse wave velocity (PWV), has been shown to be associated with atherosclerosis and risk factors of atherosclerotic cardiovascular disease,^{13–21} and is an independent predictor of cardiovascular events,^{22,23} Therefore, although platelets are likely to be activated in patients with atherosclerotic disease who exhibit increased arterial stiffness, little is known

about the relation of arterial stiffness itself to platelet activation.

Recently, platelet activation has been widely evaluated by measuring soluble P-selectin; a platelet surface molecule also termed CD62P.^{4,6–8,11} Although the measurement of soluble P-selectin is simple and useful, it is an indirect method of evaluating platelet activation. On the other hand, platelet activation can be detected directly by measuring surface antigen CD62P using flow cytometry.^{2,3,5,9,10,12} Furthermore, detection of platelet–neutrophil complexes (PNC), which are formed as a result of interaction with CD62P provides an additional means to detect platelet activation.²⁴

The purpose of this study was to investigate the association between arterial stiffness and platelet activation by measuring PWV, P-selectin, and PNC in subjects without atherosclerotic disease.

Correspondence: Dr F Yamasaki, Department of Clinical Laboratory, Kochi Medical School, Nankoku, Kochi 783-8505, Japan.
E-mail: yamasakf@kochi-ms.ac.jp

This research was supported in part by a grant from the President Research Fund of Kochi Medical School Hospital and the Japan Arteriosclerosis Prevention Fund.

Received 15 March 2004; revised 5 January 2005; accepted 9 February 2005; published online 7 April 2005

Materials and methods

Subjects

We studied 38 healthy nonsmoking volunteers (20 men and 18 women), aged 23–77 years

(mean = 49 ± 15 years) with no evidence of heart disease on physical examination, standard 12-lead electrocardiography, chest radiography, echocardiography, or blood chemistry analysis. Subjects had no self-reported past history or current evidence of cardiovascular disease, hypertension, hypercholesterolaemia, diabetes mellitus or renal disease. Basic characteristics of subjects are shown in Table 1. None of the subjects had frequent ectopic beats or atrial fibrillation and none had taken any medication for at least 10 days. Informed consent was obtained before performing the study and the study protocol was approved by the Local Ethics Committee of Kochi Medical School.

Evaluation of arterial stiffness

Arterial stiffness was evaluated by PWV, measured using volume-plethysmographic apparatus (Colin, Komaki, Japan).^{18–21} Data were acquired with subjects lying supine in a quiet and temperature-controlled room at 11 AM, at least 3 h after breakfast. Surface electrodes were attached to both wrists for ECG measurement, a microphone was positioned at the left sternal edge to detect heart sounds, and cuffs incorporating plethysmographic and oscillometric sensors were fastened around both the brachial regions and ankles to measure pulse wave forms and blood pressure. Brachial–ankle PWV (ba-PWV) and heart–brachial PWV (hb-PWV) were measured as follows. The time interval between the wave foot of the brachial waveform and that of the ankle waveform was defined as the time interval between the brachial region and ankle, while the time interval between the heart and the right brachial

artery was defined as the time interval between the second heart sound and the right brachial waveform. The distance between these sampling points was calculated automatically according to the height of the subject. PWVs were calculated by dividing each distance by the respective time interval. Right brachial blood pressure (systolic and diastolic) and pulse rate were concurrently measured.

Measurement of platelet activation

Sample preparation and measurement of platelet P-selectin (CD62P) and PNC levels were performed according to the method described by Peters *et al*.²⁴ To minimize platelet activation during blood collection, blood was drawn via a 21 G butterfly needle without the use of a tourniquet. After discarding the first 2 ml of blood, a further 2 ml was collected and immediately added to 200 μ l of sodium citrate (3.13%). All antibodies were sourced as follows: Fluorescein isothiocyanate (FITC) labelled IgG1 anti-CD62P from Dainippon Pharmaceutical, Osaka, Japan, phycoerythrin (PE) labelled IgG2a anti-CD42b and FITC labelled IgG1 anti-CD11b from Beckman Coulter, Fullerton, CA, USA. As negative controls, FITC-labelled IgG1 (Beckman Coulter, Fullerton, CA, USA) and double-stained (FITC/PE) IgG1 and IgG2a (Dako, High Wycombe, Bucks, UK) irrelevant antibodies were included.

Sample preparation for the measurement of platelet CD62P level: In all, 5 μ l of blood was added to a round-bottomed polystyrene tube containing 50 μ l of platelet buffer (10 mmol/l HEPES, 145 mmol/l NaCl, 5 mmol/l KCl, 1 mmol/l MgSO₄, pH 7.4), and 5 μ l of anti-CD62P or control IgG1 antibody. Following gentle suspension, samples were incubated in the dark at room temperature for 20 min without stirring. Then 250 μ l of fixative was added and the tubes were incubated for an additional 10 min. The samples were then diluted with 500 μ l of buffer and analysed. Flow cytometric analysis was performed within 1 h of fixation.

Sample preparation for the measurement of PNC level: In all, 50 μ l of blood was added to a round-bottomed polystyrene tube containing 5 μ l of anti-CD42b, and 5 μ l of anti-CD11b or isotype control antibodies. Following gentle mixing, samples were incubated in the dark at room temperature for 10 min without stirring. Then 500 μ l of fixative was added and the tubes were incubated for additional 10 min. Flow cytometric analysis was performed within 1 h of preparation.

Flow cytometric analysis

Blood samples were analysed in a COULTER EPICS XL Profile Flow Cytometer, Miami, FL, USA, using either single or double fluorochromes. The peak emission intensity of FITC fluorescence was

Table 1 Clinical characteristics of subjects

Parameters	All subjects (n = 38)
Age (years)	49 \pm 15
Gender, male/female	20/18
Systolic blood pressure (mmHg)	125 \pm 16
Diastolic blood pressure (mmHg)	77 \pm 10
Pulse rate (bpm)	66 \pm 10
Blood sugar (mg/dl)	98.5 \pm 18.5
Total cholesterol (mg/dl)	192.6 \pm 20.7
Blood urea nitrogen (mg/dl)	14.0 \pm 18.5
Creatinine (mg/dl)	0.69 \pm 0.15
PNC (%)	9.5 \pm 4.9
PNC(ADP) (%)	20.2 \pm 9.9
Δ -PNC	10.7 \pm 6.9
P-selectin (%)	13.1 \pm 1.7
P-selectin(ADP) (%)	36.6 \pm 9.2
Δ -P-selectin	23.6 \pm 9.1
hb-PWV (m/s)	5.3 \pm 0.9
ba-PWV (m/s)	13.8 \pm 3.0

Values are expressed as mean \pm s.d.

PNC = platelet neutrophil complexes; ADP = adenosine diphosphate; Δ -PNC = PNC (ADP) – PNC; Δ -P-selectin = P-selectin (ADP) – P-selectin; hb-PWV = heart–brachial pulse wave velocity; ba-PWV = brachial–ankle pulse wave velocity.

detected at 515 nm and that of phycoerythrin fluorescence at 580 nm.

Measurement of platelet CD62P level: After forward and side scatter measurements were made with gain setting in logarithmic mode, platelet-sized events were counted. CD62P-positive platelets were defined as those with a fluorescence intensity exceeding that of 98% of the platelets staining with control antibody.

Measurement of PNC level: After forward and side scatter measurements were made with gain setting in linear mode, neutrophil-sized events were selected. Results were defined as positive when the fluorescence intensity exceeded that of 98% of the isotype-matched (IgG1 and IgG2a) control antibodies staining. Events positive for both CD11b and CD42b were considered to represent PNCs and were expressed as percentages of events with positive CD11b staining.

Evaluation of platelet activation reserve: We evaluated platelet activation reserve, that is, the ability of the platelets to be activated, in a separate experiment. Platelets were activated with 5 μ l of adenosine diphosphate (ADP). We also calculated the difference between basal and stimulated states of P-selectin expression (Δ -P-selectin) and PNC level (Δ -PNC) to determine activation reserve.

Statistical analysis

Data are presented as mean \pm s.d. Univariate linear correlation analysis and multiple regression analysis were used for statistical evaluation. The variables significantly associated with platelet activation on univariate analysis were included in a multiple regression analysis in order to adjust PWV for each variable. Gender differences were evaluated with ANOVA. *P*-values <0.05 were considered to represent statistical significance.

Results

Both ba-PWV and hb-PWV exhibited significant positive correlations with age, systolic, and diastolic blood pressure ($r=0.60$ – 0.81 , $P<0.05$ or <0.01), and pulse rate ($r=0.44$, $P<0.05$, $r=0.65$, <0.01 , respectively) (Table 2). For platelet activation and activation reserve, correlations with age were less strong but remained significant ($r=0.36$ – 0.61 , $P<0.05$ or <0.01) with the exception of Δ -P-selectin (not significant, NS), and correlations with systolic and diastolic blood pressure were similar ($r=0.35$ – 0.53 , $P<0.05$ or <0.01) with the exception of P-selectin (NS) (Table 3). However, platelet activation and activation reserve exhibited no significant correlation with pulse rate, blood glucose, total cholesterol, blood urea nitrogen or creatinine. No significant gender-related differences were observed in any of these correlations (Tables 2 and 3).

Table 2 Correlation between PWV and clinical indices

	hb-PWV	ba-PWV
Age	0.74**	0.80**
Systolic blood pressure	0.61**	0.81**
Diastolic blood pressure	0.60**	0.74**
Pulse rate	0.44*	0.65**
Blood sugar	-0.05	-0.17
Total cholesterol	-0.03	-0.30
Blood urea nitrogen	-0.32	0.32
Creatinine	0.04	-0.14
Gender		
Male	5.5 \pm 1.0	14.1 \pm 3.0
Female	5.2 \pm 0.8	13.6 \pm 3.1

PNC = platelet neutrophil complexes; ADP = adenosine diphosphate; Δ -PNC = PNC (ADP)-PNC; Δ -P-selectin = P-selectin (ADP)-P-selectin; hb-PWV = heart-brachial pulse wave velocity; ba-PWV = brachial-ankle pulse wave velocity.

For parameters from age to creatinine, values are correlation coefficients.

* $P<0.05$.

** $P<0.01$.

For gender, values are mean \pm s.d., with differences evaluated with ANOVA.

PWVs exhibited significant positive correlations ($r=0.43$ – 0.74 , $P<0.05$ or <0.01) to all indices of platelet activation and reserve (Table 4, Figure 1). When age or blood pressures were adjusted for on multivariate analysis, some indices of platelet activation and reserve were significantly related to PWVs ($r=0.34$ – 0.67 , $P<0.05$ or <0.01). When both age and blood pressures were simultaneously adjusted for, significant correlations remained between platelet activation and reserve and PWVs ($r=2.17$ – 6.59 , $P<0.05$ or <0.01) (Table 4). In other words, although the relationship between PWVs and the indices of platelet activation was strongly affected by age and blood pressure, a significant association remained when these factors were adjusted for.

Discussion

The main finding of this study was that platelet activation and activation reserve were associated with arterial stiffness when analyses were adjusted for age and blood pressure. This suggests that increased arterial stiffness might play an important role in thrombotic events.

Patients with hypertension, cerebrovascular disease, coronary heart disease, diabetes mellitus, and renal failure are recognized to have less arterial compliance than normal subjects.^{13–15,17–19} Increased PWV has also been reported to be an independent predictor of cardiovascular events in patients with hypertension or renal failure, and in elderly subjects.^{22,23} The association between increased arterial stiffness and high incidence of cardiovascular events may be explained by the existence of atherosclerosis. Hirai *et al*²⁵ have demonstrated strong associations between abdominal aortic and

Table 3 Correlation between platelet activation and clinical indices

	PNC	PNC (ADP)	Δ-PNC	P-selectin	P-selectin (ADP)	Δ-P-selectin
Age	0.51**	0.61**	0.52**	0.36*	0.38*	0.32
Systolic blood pressure	0.41*	0.53**	0.48**	0.41*	0.43*	0.35*
Diastolic blood pressure	0.43*	0.49**	0.40*	0.25	0.40*	0.36*
Pulse rate	0.28	0.25	0.16	0.04	0.15	0.15
Blood sugar	0.09	-0.18	-0.31	-0.17	0.13	0.16
Total cholesterol	-0.14	-0.07	0.001	-0.10	-0.13	-0.11
Blood urea nitrogen	-0.01	0.12	0.18	-0.05	0.05	0.06
Creatinine	0.05	-0.13	-0.22	0.04	-0.17	-0.18
Gender						
Male	10.3±5.9	19.7±8.7	9.4±6.9	13.1±1.8	35.5±9.3	22.4±9.0
Female	8.8±3.8	20.7±11.4	11.9±6.8	13.0±1.7	37.7±9.2	24.7±9.3

PNC=platelet neutrophil complexes; ADP=adenosine diphosphate; Δ-PNC=PNC (ADP)-PNC; Δ-P-selectin=P-selectin (ADP)-P-selectin; hb-PWV=heart-brachial pulse wave velocity; ba-PWV=brachial-ankle pulse wave velocity.

For parameters from age to creatinine, values are correlation coefficients.

*P<0.05.

**P<0.01.

For gender, values are mean ± s.d., with differences evaluated with ANOVA.

Table 4 Relation between platelet activations and PWV

	PNC	PNC (ADP)	Δ-PNC	P-selectin	P-selectin (ADP)	Δ-P-selectin
<i>Not adjusted</i>						
hb-PWV	0.62**	0.74**	0.63**	0.45**	0.57**	0.50**
ba-PWV	0.59**	0.71**	0.61**	0.47**	0.51**	0.43*
<i>Adjusted for age</i>						
hb-PWV	2.86**	6.95**	4.09*	0.75	6.55**	5.80*
ba-PWV	0.79	2.01**	1.22*	0.28	1.75*	1.47
<i>Adjusted for systolic blood pressure</i>						
hb-PWV	3.20**	7.23**	4.04**	0.59	5.09*	4.50*
ba-PWV	1.21**	2.64**	1.44*	0.23	1.48	1.25
<i>Adjusted for diastolic blood pressure</i>						
hb-PWV	3.08**	7.67**	4.59**	0.87*	5.32**	4.46*
ba-PWV	0.97**	2.50**	1.54**	0.34*	1.45*	1.10
<i>Adjusted for age and systolic blood pressure</i>						
hb-PWV	2.80*	6.43**	3.63*	0.58	5.93*	5.35*
ba-PWV	1.08	2.32*	1.24	0.24	1.72	1.48
<i>Adjusted for age and diastolic blood pressure</i>						
hb-PWV	2.63*	6.59**	3.97**	0.78	6.06*	5.28*
ba-PWV	0.76	2.17*	1.40	0.40	1.66	1.26

PNC=platelet neutrophil complexes; ADP=adenosine diphosphate; Δ-PNC=PNC (ADP)-PNC; Δ-P-selectin=P-selectin (ADP)-P-selectin; hb-PWV=heart-brachial pulse wave velocity; ba-PWV=brachial-ankle pulse wave velocity.

'Not adjusted' — values are correlation coefficients between PWVs and indices of platelet activation before adjustment.

*P<0.05.

**P<0.01.

Other values are regression coefficients between PWVs and indices of platelet activation adjusted for age and/or blood pressures as indicated.

*P<0.05.

**P<0.01.

carotid arterial stiffness and the degree of coronary artery disease. Popele *et al*²⁶ recently reported that aortic stiffness as measured by PWV is strongly associated with common carotid intima-media thickness, carotid arterial plaques, and the presence of peripheral arterial disease. Moreover, some population-based studies have demonstrated higher blood pressure, increased age, and male gender to be

associated with increased PWV.^{16,20,21} Pulse pressure may also relate to arterial stiffness and cardiovascular events, with higher pulse pressure reflecting elevated systolic pressure and reduced diastolic pressure due to increased arterial stiffness. In the present study, significant relationships were observed between PWVs and age, blood pressure, and pulse rate, in accordance with previous studies.




Figure 1 Correlation between ba-PWV and PNC (upper two panels). PNC=platelet neutrophil complexes; ADP=adenosine diphosphate; Δ -PNC=PNC (ADP)-PNC; Δ -P-selectin=P-selectin (ADP)-P-selectin; hb-PWV=heart-brachial pulse wave velocity; ba-PWV=brachial-ankle pulse wave velocity.

P-selectin is a component of α -granules that is expressed on the platelet surface membrane and released into the plasma upon platelet activation. Although the bulk of circulating soluble P-selectin appears to be platelet derived,²⁷ the substance is also found in the Weibel-Palade bodies of endothelial cells.²⁸ Direct measurement of platelet membrane P-selectin is therefore a more sensitive method of assessing platelet activation. In the present study, we evaluated platelet activation by measuring membrane activation markers using flow cytometry with activation-dependent monoclonal antibodies. PNC levels were also measured using the same method. P-selectin levels in our normal subjects aged 49 ± 15 years were $13.1 \pm 1.7\%$; this was higher than that in quoted by other studies, possibly due to the differences in monoclonal antibodies or in sample manipulation.

P-selectin expressed on activated platelets causes formation of PNC. Moreover, platelets and platelet-derived P-selectin play an important role in thrombus growth at the site of atherosclerosis.² *In vivo* and *in vitro* studies have shown that shear stress and exposure to atherogenic stimuli, such as oxidation by low-density lipoprotein or cigarette smoking, induce rapid P-selectin-dependent aggregation and

accumulation of leukocytes and platelets.^{4,5,11} Activated platelets accumulating in thrombi at the site of ruptured atherosclerotic plaques will express CD62P. In clinical studies, P-selectin has been shown to be a marker of platelet activation related to adverse cardiovascular events such as hypertension, coronary artery disease, cerebrovascular disease, and peripheral arterial disease,^{6,7,10-12} and also to be a predictor of cardiovascular events.^{8,12} PNC, forming as a result of the interaction of platelet P-selectin and neutrophils also promotes platelet activation.²⁴ This is the first study to demonstrate that P-selectin and PNC were significantly correlated with arterial stiffness evaluated by PWV in normal subjects. In an analysis of four randomized trials, Hebert *et al*²⁹ showed that aspirin therapy was beneficial in the primary prevention of vascular disease. Higher levels of other membrane markers such as von Willebrand factor receptor are observed in activated platelets, which are affected by aspirin or ticlopidine.³⁰ Therefore, our results indicate that, in the normal population, antiplatelet agents may play a role in preventing cardiovascular events through factors other than P-selectin.

Although the exact mechanism accounting for the relationship between platelet activation and arterial stiffness is unknown, it is possible to make

the following speculations. When arterial stiffness is raised, shear stress might play an important role in platelet activation. Using cone-plate viscometry,³ Goto *et al* showed that platelet activation (measured by P-selectin surface expression, von Willebrand factor-mediated platelet aggregation and translocation of GP Iba) was induced by high shear rate of 10 800 s⁻¹. Higher arterial stiffness increases blood flow velocity and produces a steep systolic pressure waveform,³¹ and it is possible that the resulting increased shear stress could promote platelet activation. Another possible mechanism is that endothelial dysfunction may interact with arterial stiffness and platelet hyperactivity. Kobayashi *et al*³² showed significant correlation between endothelial dysfunction measured by flow-mediated dilatation and ba-PWV. Platelets are also activated by endothelial dysfunction. On the other hand, activated platelets themselves may cause arterial stiffness via vascular smooth muscle cell growth factors and extracellular matrix modulator released from platelets, that is, PDGF.³³ However, this response also occurs at the site of endothelial injury. Further study is therefore required to clarify whether arterial stiffness causes platelet activation or alternatively whether platelet activation might result in arterial stiffness.

Limitations

Despite the small sample size, it is possible that the broad age range (23–77 years) of our subjects caused outliers in PWV and platelet activation. However, significant correlations were found when age and blood pressure were adjusted for, suggesting that the influence of age did not entirely explain the correlation between PWV and platelet activation. In the present study, ba-PWV was 14.1 ± 3.0 m/s in men and 13.6 ± 3.1 m/s in women; values higher than those reported by Yamashina *et al*.²⁰ Furthermore, it is not known whether such a relationship between arterial stiffness and platelet activation is found in patients with conditions such as hypertension, diabetes mellitus, coronary heart disease, and stroke. Further studies should be therefore performed in such patients, using larger sample sizes.

Acknowledgements

We would like to thank Tadashi Ueta for technical assistance and Misa Nakagawa for her assistance throughout the study.

References

- 1 Tenaglia AN *et al*. Levels of expression of P-selectin, E-selectin, and intercellular adhesion molecule-1 in coronary atherectomy specimens from patients with stable and unstable angina pectoris. *Am J Cardiol* 1997; **79**: 742–747.

- 2 Hagberg IA, Roald HE, Lyberg T. Platelet activation in flowing blood passing growing arterial thrombi. *Arterioscler Thromb Vasc Biol* 1997; **17**: 1331–1336.
- 3 Goto S *et al*. Effects of ticlopidine on von Willebrand factor-mediated shear-induced platelet activation and aggregation. *Platelets* 2001; **12**: 406–414.
- 4 Davi G *et al*. Increased levels of soluble P-selectin in hypercholesterolemic patients. *Circulation* 1998; **97**: 953–957.
- 5 Pernerstorfer T *et al*. Low-dose aspirin does not lower *in vivo* platelet activation in healthy smokers. *Br J Haematol* 1998; **102**: 1229–1231.
- 6 Spencer CG *et al*. Von Willebrand factor, soluble P-selectin, and target organ damage in hypertension: a substudy of the Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT). *Hypertension* 2002; **40**: 61–66.
- 7 Blann AD, Dobrotova M, Kubisz P, McCollum CN. von Willebrand factor, soluble P-selectin, tissue plasminogen activator and plasminogen activator inhibitor in atherosclerosis. *Thromb Haemost* 1995; **74**: 626–630.
- 8 Blann AD, Faragher EB, McCollum CN. Increased soluble P-selectin following myocardial infarction: a new marker for the progression of atherosclerosis. *Blood Coagul Fibrinolysis* 1997; **8**: 383–390.
- 9 Grau AJ *et al*. Increased fraction of circulating activated platelets in acute and previous cerebrovascular ischemia. *Thromb Haemost* 1998; **80**: 298–301.
- 10 Serebruany VL *et al*. Uniform platelet activation exists before coronary stent implantation despite aspirin therapy. *Am Heart J* 2001; **142**: 611–616.
- 11 Parker III C, Vita JA, Freedman JE. Soluble adhesion molecules and unstable coronary artery disease. *Atherosclerosis* 2001; **156**: 417–424.
- 12 Lip GY *et al*. Sequential alterations in haemorrhology, endothelial dysfunction, platelet activation and thrombogenesis in relation to prognosis following acute stroke: The West Birmingham Stroke Project. *Blood Coagul Fibrinolysis* 2002; **13**: 339–347.
- 13 De Cesaris R, Ranieri G, Filitti V, Andriani A. Large artery compliance in essential hypertension. Effects of calcium antagonism and beta-blocking. *Am J Hypertens* 1992; **5**: 624–628.
- 14 Asmar R *et al*. Assessment of arterial distensibility by automatic pulse wave velocity measurement. Validation and clinical application studies. *Hypertension* 1995; **26**: 485–490.
- 15 Lehmann ED, Riley WA, Clarkson P, Gosling RG. Non-invasive assessment of cardiovascular disease in diabetes mellitus. *Lancet* 1997; **350**Suppl 1: S114–S119.
- 16 Asmar J *et al*. Arterial stiffness and cardiovascular risk factors in a population-based study. *J Hypertens* 2001; **19**: 381–387.
- 17 Mourad JJ *et al*. Creatinine clearance, pulse wave velocity, carotid compliance and essential hypertension. *Kidney Int* 2001; **59**: 1334–1341.
- 18 Yamashina A *et al*. Validity, reproducibility, and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. *Hypertens Res* 2002; **25**: 359–364.
- 19 Munakata M, Ito N, Nunokawa T, Yoshinaga K. Utility of automated brachial ankle pulse wave velocity measurements in hypertensive patients. *Am J Hypertens* 2003; **16**: 653–657.
- 20 Yamashina A *et al*. Nomogram of the relation of brachial-ankle pulse wave velocity with blood pressure. *Hypertens Res* 2003; **26**: 801–806.