

Table 2 Comparison among various nutritional markers.

	BMI (m/kg ²), n = 235			Albumin (g/dl), n = 179			Total cholesterol (mg/dl), n = 177			MNA score, n = 235		
	<18.5	≥18.5	P-value*	<3.5	≥3.5	P-value	<150	≥150	P-value	<17	≥17	P-value
Men/women	14/29	53/139	0.549	7/14	41/117	0.473	6/8	41/122	0.150	18/39	49/129	0.514
	Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)	
Age (years)	79.9 (7.5)	78.3 (7.5)	0.188	82.1 (7.9)	77.9 (7.3)	0.017	77.6 (8.5)	78.4 (7.4)	0.700	80.2 (8.1)	78.1 (7.3)	0.065
BMI (kg/m ²)	16.4 (1.6)	23.2 (3.3)	<0.0001	19.1 (4.8)	22.1 (3.9)	0.002	19.5 (3.9)	22.0 (4.1)	0.035	19.1 (3.6)	22.8 (3.8)	<0.0001
MAC (cm)	20.3 (2.3)	25.4 (3.0)	<0.0001	22.1 (4.7)	24.7 (3.3)	0.002	22.6 (3.9)	24.5 (3.6)	0.053	21.6 (3.4)	25.2 (3.2)	<0.0001
TSF (mm)	8.18 (6.6)	14.3 (8.4)	<0.0001	7.8 (6.3)	13.8 (8.8)	0.003	7.8 (6.4)	13.5 (8.8)	0.018	8.1 (6.1)	14.4 (8.4)	<0.0001
CC (cm)	28.6 (4.7)	31.5 (3.6)	<0.0001	27.1 (4.4)	31.4 (3.9)	<0.0001	26.0 (3.7)	31.3 (4.0)	<0.0001	28.9 (4.5)	31.6 (3.5)	<0.0001
Albumin (g/dl)	3.7 (0.9)	4.1 (0.4)	<0.0001	3.1 (0.3)	4.2 (0.3)	<0.0001	3.1 (0.5)	4.1 (0.4)	<0.0001	3.6 (0.6)	4.2 (0.3)	<0.0001
Total cholesterol (mg/dl)	187 (44)	204 (38)	0.026	152 (32)	207 (36)	<0.0001	129 (19)	206 (35)	<0.0001	175 (42)	208 (36)	<0.0001
MNA score	15.2 (4.8)	21.3 (3.7)	<0.0001	12.0 (5.1)	21.1 (3.7)	<0.0001	12.1 (4.9)	20.8 (4.3)	<0.0001	13.0 (3.4)	21.9 (2.7)	<0.0001
Lymphocyte (count/mm ³)	1620 (577)	1789 (680)	0.244	1890 (692)	1754 (666)	0.513	2016 (742)	1749 (664)	0.273	1829 (630)	1748 (675)	0.557

*Student's, *t* test was conducted except for the gender difference (χ^2 -test) and MNA score (Mann-Whitney *U*-test).

except for TSF ($P = 0.001$), but not with TLC ($P = 0.524$).

Discussion

Although the TLC is one of the most commonly used markers for assessing nutritional status, so far little evidence exists as to whether TLC reflects the nutritional status of the elderly. In the present study, we concluded that TLC is not a suitable marker of malnutrition in the elderly. This conclusion was based on the observation that no correlation was detected between TLC and other well-known nutritional parameters including anthropometric measurements, biochemical markers, and MNA score, a comprehensive nutritional screen tool for the elderly. In addition, MNA score was correlated with all of the nutritional markers used in the present study except for TLC. This result is consistent with the previous observation of Goodwin JS that no significant correlation was observed between lymphocyte count and blood levels of specific nutrients including serum albumin in the independently living healthy elderly.¹⁰

It has been shown that the serum albumin and total cholesterol levels, both of which are commonly used as nutritional markers, are sometime discordant with clinical assessments of malnutrition, largely because these biomarkers are influenced by factors such as inflammatory activity, hemoconcentration, and various diseases such as liver cirrhosis and nephritic syndrome. However, in the present study, these biochemical markers for malnutrition were well correlated with anthropometric measurements as well as with MNA score.

There is no general agreement of the effect of aging on TLC. Divergent data have been reported concerning age-related changes in total lymphocyte number.^{5,11,12} This may be due to the heterogeneity of the aging immune system. The present study suggested that TLC was correlated with aging in subjects between 65 and 90 years old, indicating that TLC appears to be reflective of age rather than of nutritional status. Our results that TLC is not a suitable marker of malnutrition in the elderly does not indicate that malnutrition is not a risk factor for the impairment of immune function. In fact, nutritional status has long been recognized as a major factor in age-related immune impairment, and a number of studies have already demonstrated that malnutrition is associated with decreased lymphocyte proliferation, reduced cytokine release, and lower antibody response to vaccines.^{12,13} In addition, an important modification

in T lymphocyte subsets is known to occur in aged people.¹⁴ In fact, it has been demonstrated that a low lymphocyte count is associated with an increased mortality risk in older persons.¹⁵

There are several limitations to this study. First, the study group might have consisted of elderly who had comorbid diseases, given that they were enrolled from clinical settings. Therefore, our results may apply only to the elderly in ill health. The possibility of an association between TLC and nutritional status in the healthy elderly cannot be excluded. Second, the effect of medication on TLC was not considered in this study, due to the fact that medication data were not available.

In conclusion, we found that TLC is not suitable as a marker of nutritional status in the elderly. TLC appears to be reflective of age rather than of nutritional status.

Acknowledgments

This work was supported by a Grant-in Aid for the Comprehensive Research on Aging and Health from the Ministry of Health and Welfare of Japan.

References

1. Sullivan DH, Sun S, Walls RC. Protein-energy undernutrition among elderly hospitalized patients: a prospective study. *JAMA* 1999;281:2013-9.
2. Edington J, Boorman J, Durrant ER, et al. Prevalence of malnutrition on admission to four hospitals in England. The Malnutrition Prevalence Group. *Clin Nutr* 2000;19:191-5.
3. Sullivan DH, Walls RC. Protein-energy undernutrition and the risk of mortality within six years of hospital discharge. *J Am Coll Nutr* 1998;17:571-8.
4. Keller HH, Ostbye T, Goy R. Nutritional risk predicts quality of life in elderly community-living Canadians. *J Gerontol A Biol Sci Med Sci* 2004;59:68-74.
5. Omran ML, Morley JE. Assessment of protein energy malnutrition in older persons, Part II: Laboratory evaluation. *Nutrition* 2000;16:131-40.
6. Seiler WO. Clinical pictures of malnutrition in ill elderly subjects. *Nutrition* 2001;17:496-8.
7. Vellas B, Guigoz Y, Garry PJ, et al. The Mini Nutritional Assessment (MNA) and its use in grading the nutritional state of elderly patients. *Nutrition* 1999;15:116-22.
8. Kuzuya M, Kanda S, Koike T, Suzuki Y, Satake S, Iguchi A. Evaluation of Mini-Nutritional Assessment for Japanese frail elderly. *Nutrition* 2005 in press.
9. James WP, Francois PJ. The choice of cut-off point for distinguishing normal body weights from underweight or 'chronic energy deficiency' in adults. *Eur J Clin Nutr* 1994;48(Suppl. 3):S179-84.
10. Goodwin JS, Garry PJ. Lack of correlation between indices of nutritional status and immunologic function in elderly humans. *J Gerontol* 1988;43:M46-9.

11. Diaz-Jouamen E, Strickland RG, Williams RC. Studies of human lymphocytes in the newborn and the aged. *Am J Med* 1975;58:620-5.
12. Lesourd BM. Nutrition and immunity in the elderly: modification of immune responses with nutritional treatments. *Am J Clin Nutr* 1997;66:478S-84S.
13. Lesourd BM, Mazari L, Ferry M. The role of nutrition in immunity in the aged. *Nutr Rev* 1998;56(1 Pt 2):S113-25.
14. Lesourd BM, Meaume S. Cell mediated immunity changes in ageing, relative importance of cell subpopulation switches and of nutritional factors. *Immunol Lett* 1994;40:235-42.
15. Izaks GJ, Remarque EJ, Becker SV, Westendorp RG. Lymphocyte count and mortality risk in older persons. The Leiden 85-Plus Study. *J Am Geriatr Soc* 2003;51:1461-5.

Available online at www.sciencedirect.com





ELSEVIER

Nutrition 21 (2005) 498–503

NUTRITION

www.elsevier.com/locate/nut

Applied nutritional investigation

Evaluation of Mini-Nutritional Assessment for Japanese frail elderly

Masafumi Kuzuya, M.D., Ph.D.*, Shigeru Kanda, M.D., Ph.D., Teruhiko Koike, M.D., Ph.D.,
Yusuke Suzuki, M.D., Ph.D., Shosuke Satake, M.D., Ph.D., Akihisa Iguchi, M.D., Ph.D.

Department of Geriatrics, Nagoya University Graduate School of Medicine, Nagoya, Japan

Manuscript received April 18, 2004; accepted August 6, 2004.

Abstract

Objective: We evaluated the Mini-Nutritional Assessment (MNA) test and the short-form MNA as screening tools for malnutrition in the Japanese elderly population.

Methods: A cross-sectional study of 226 elderly Japanese patients (78.6 ± 0.5 y of age, mean \pm standard deviation; 67 men and 159 women) in various settings was carried out. Nutritional assessment included MNA, anthropometric measurements, and biochemical markers.

Results: According to the original cutoff point of the full MNA, 19.9% of those assessed were malnourished, 58.0% were at risk of malnutrition, and 22.1% were well nourished. Significant correlations were found between full MNA scores and age ($r = -0.14$), body mass index ($r = 0.59$), serum albumin ($r = 0.60$), total cholesterol ($r = 0.36$), midarm circumference ($r = 0.50$), and triceps skinfold ($r = 0.37$). The sensitivity and specificity of the full MNA score (<17) for hypoalbuminemia were 0.810 and 0.860, respectively. With a cutoff point lower than 18, sensitivity and specificity hypoalbuminemia were 0.857 and 0.815, respectively. Using a short-form MNA score 12 and higher as normal, its sensitivity and specificity for predicting undernutrition were 0.859 and 0.840, respectively.

Conclusions: The full and short forms of the MNA were useful tools to identify elderly Japanese patients with malnutrition or risk of malnutrition. However, the full MNA cutoff point for malnutrition should be modulated for this population. © 2005 Elsevier Inc. All rights reserved.

Keywords:

Elderly; Malnutrition; Nutritional assessment; Mini-nutritional assessment; Anthropometric measurements

Introduction

Malnutrition is a frequent and serious problem in geriatric patients. Malnutrition in ill elderly subjects is one of the most common and least-heeded problems in hospitals, nursing homes, and home care [1–4]. Different studies have suggested that malnutrition is an important predictor of morbidity and mortality in the elderly [5,6]. In addition, malnutrition has been shown to prolong hospital stays, thereby imposing enormous costs on health services [7,8]. To identify malnourished elderly or subjects at risk of malnutrition, a conventional malnutrition assessment tool is required [9,10]. Anthropometric measurements such as

body mass index (BMI), midarm circumference (MAC), calf circumference (CC), and triceps skinfold (TSF) are essential parts of any nutritional assessment. Biochemical measurements including serum albumin and cholesterol are also frequently used as nutritional parameters, although at present there are no generally accepted criteria for the diagnosis of malnutrition in the elderly. The Mini-Nutritional Assessment (MNA) is a simple clinical scale for the evaluation of the nutritional status of frail elderly subjects. It has been validated in Europe and the United States by comparing its results with a clinical assessment performed by expert geriatric nutritionists [11,12]. Although the MNA was developed specifically for frail older people, it has been validated in a healthy older population [11,12].

The MNA has been demonstrated to be useful in predicting long-term mortality for the institutionalized elderly and acute hospital admission for the elderly living at home [6,13–15]. Recently, the MNA short form (MNA-SF) has been devised as the first step of a two-step process (screen-

This work was supported by a Grant-in Aid for the Comprehensive Research on Aging and Health from the Ministry of Health and Welfare of Japan.

* Corresponding author. Tel.: +81-52-744-2364; fax: +81-52-744-2371.

E-mail address: kuzuya@med.nagoya-u.ac.jp (M. Kuzuya).

ing with the MNA-SF followed by assessment, if needed, by the full MNA) [16]. The MNA has proved to be a simple, noninvasive, well-validated screening tool for malnutrition in elderly persons in Europe and the United States. Despite the fact that Japan has an aging society and ranks first in the world for life expectancy at birth [17], the MNA has not been validated in the Japanese elderly, and whether the MNA and its established cutoff points for the diagnosis of malnutrition and at-risk status are applicable to the Japanese elderly remain unknown.

In the present study we examined whether the MNA can screen and diagnose for malnutrition and risk for malnutrition in the Japanese elderly.

Materials and methods

Subjects

We enrolled 226 elderly (67 men and 159 women; mean age \pm standard deviation = 78.6 ± 0.5 y; age range = 65–95 y) from our geriatric outpatient clinic ($n = 68$), a nursing home ($n = 53$), geriatric hospitals ($n = 72$), and home care patients ($n = 33$). Sixty-eight consecutive outpatients were living at home independently or with mild decline in activities of daily living. Fifty-three residents and 72 inpatients with mild to severe dependency in activities of daily living were randomly chosen from one private nursing home and two geriatric hospitals, respectively. Thirty-three patients living at home and receiving home care services were also eligible for the study. Subjects diagnosed with infection, inflammation, liver disorders, kidney disorders, cancer, or bone marrow proliferative disorders were excluded by physicians. All participants provided written informed consent.

MNA characteristics

The MNA is a two-step procedure: (1) the MNA-SF is used to screen for malnutrition and risk of malnutrition and (2) the full MNA is used to assess nutritional status [16]. The MNA includes 18 items, including anthropometric measurements: BMI, MAC, CC, weight loss, a global assessment (six questions related to lifestyle, medication, and mobility), a dietary questionnaire (eight questions related to number of meals, food, and fluid intake), and a subjective assessment (self-perception of health and nutrition). The MNA-SF comprises 6 of the 18 items. The maximum possible score of the MNA-SF is 14. Scores 12 and above indicate satisfactory nutritional status. A screening score 11 and below suggests possible malnutrition and a need to proceed to the assessment stage of the MNA [16]. The assessment stage has 12 questions, with a maximum possible score of 16 (total = 30 points). The MNA score distinguishes between elderly patients with adequate nutrition

(score ≥ 24), protein-calorie undernutrition (score < 17), and risk of malnutrition (score = 17–23.5) [12].

Anthropometric measurements and biochemical markers

BMI is defined as weight in kilograms divided by height in meters squared. A BMI less than 20 kg/m^2 is widely accepted as underweight [18], particularly in well-developed countries, and 18.5 kg/m^2 is recommended as a practical lower limit for most populations [19]. Therefore, the diagnosis of malnutrition was made when BMI was less than 18.5 kg/m^2 . TSF was measured with Harpenden calipers over the triceps muscle at the midway point between the acromion and the olecranon process. MAC and CC were measured on the left arm and calf, respectively, with a tape measure. Three measurements were taken to the nearest 0.5 mm, with the mean taken as the true value. All anthropometric measurements were taken at least twice by two different investigators, and the reported values are the means of the repeated measurements (interrater reliability with Pearson's correlation coefficient, $r = 0.923$, $P < 0.0001$). Blood samples were collected after an overnight fast. Serum albumin or total cholesterol levels were determined by kinetic immunonephelometry or enzymatically, respectively. Blood was collected into tubes containing ethylene-diaminetetra-acetic acid, and total lymphocyte count was measured with use of a Coulter counter. Serum albumin and total cholesterol levels were used as biochemical markers for undernutrition: levels lower than 3.5 g/dL of albumin or 150 mg/dL of total cholesterol were taken to indicate malnutrition.

Statistical analysis

Differences between groups (MNA total scores < 17 , 17–23.5, and ≥ 24) were determined by analysis of variance or the Kruskal-Wallis test, depending on the distribution of the analyzed variable. Partial rank correlation coefficients adjusted for age were used to measure the relations between MNA total score, MNA-SF, anthropometric measurements, and biochemical markers. To identify optimal threshold values for predicting malnutrition, receiver operating characteristic (ROC) curve analysis was performed by computing the sensitivity and specificity of the different tests at various cutoff levels [20]. The area under the ROC curve was also evaluated. A value of 0.5 under the ROC curve indicates that the variable performs no better than chance, whereas a value of 1.0 indicates perfect discrimination. A larger area under the ROC curve represents a greater reliability and discrimination of the scoring system [21]. Cutoff values can be set depending on the purpose for which the scales are used. For screening purposes, a high sensitivity and a high negative predictive value are required, whereas diagnosis requires a high specificity and a high positive predictive value. Sensitivity, specificity, positive predictive value, and negative predictive value for predicting malnu-

Table 1
MNA score, anthropometric measurements, and clinical chemistry in Japanese elderly

	MNA total score			Analysis of variance (<i>P</i>)
	<17	17–23.5	≥24	
<i>n</i>	45	131	50	
Men/women	16/29	40/91	11/39	0.220*
Age (y) [†]	80.2 (8.0)	78.5 (7.4)	76.9 (7.3)	0.157
BMI (kg/m ²) [†]	18.5 (3.2)	22.2 (3.8)	24.6 (3.0)	<0.0001
MAC (cm) [†]	21.6 (3.4)	24.8 (3.2)	26.2 (2.8)	<0.0001
TSF (mm) [†]	8.1 (6.2)	13.9 (8.2)	15.9 (9.1)	<0.0001
CC (cm) [†]	27.5 (4.7)	31.2 (3.1)	32.0 (3.3)	<0.0001
Albumin (g/dL) [†]	3.6 (0.6)	4.1 (0.3)	4.4 (0.3)	<0.0001
Total cholesterol (mg/dL) [†]	174.0 (42.5)	203.3 (35.4)	217.5 (35.1)	<0.0001
Lymphocyte (/ μ L) [†]	1825.0 (641.1)	1750.5 (725.5)	1744.1 (560.6)	0.805

BMI, body mass index; CC, calf circumference; MAC, midarm circumference; MNA, Mini-Nutritional Assessment; TSF, triceps skinfold

* Kruskal-Wallis test.

[†] Mean (standard deviation).

trition based on the various nutritional markers were also calculated for different cutoff points. The best Youden index (sensitivity + specificity – 1) was used to determine the best cutoff point [22]. The Youden index is used to compare the proportion of cases correctly classified. The higher the Youden index, the more accurate the prediction (higher true positive and true negative and fewer false positive and false negative) at the cutoff point. The Kolmogorov-Smirnov test was used to check the normal distribution of variables. The statistical significance level was set at 0.05. Data evaluation was carried out with SPSS software (SPSS Inc., Chicago, IL, USA).

Results

Subjects' average age was 77.8 ± 13.8 y (mean ± standard deviation). MNA total scores averaged 20.2 ± 4.6 and ranged from a minimum of 4.0 to a maximum of 27.5, with a median at 21.0. Table 1 lists the mean results of variables, which are expressed according to the classification of the MNA. According to the original cutoff point of the full MNA, 19.9% (45 of 226) had an MNA score lower than 17, 58.0% (131 of 226) had an MNA score between 17 and 23.5, and 22.1% (50 of 226) had a score of at least 24. There were significant differences between classes with regard to BMI, MAC, TSF, CC, serum albumin, and total cholesterol levels, but not to age (*P* = 0.157) and lymphocyte count (*P* = 0.805). There was a weak but statistically significant, negative correlation between MNA total score and age (Table 2). There were relations between MNA total score and BMI, MAC, and CC, which are included as anthropometric markers in the MNA. In addition, MNA total score showed good correlation with TSF and serum albumin and total cholesterol, which are not included in the MNA, but no correlation between MNA score and lymphocyte number.

The ROC curves shown in Fig. 1A plot the sensitivity

versus 1-specificity for MNA total score in predicting low serum albumin (<3.5 g/dL), total cholesterol (<150 mg/dL), and low BMI (<18.5 kg/m²) as markers of malnutrition. The area under the ROC curves, which represent the overall accuracy of the MNA total score as a test for malnutrition, was found to be 0.916 (95% confidence interval = 0.846 to 0.985) for albumin (*P* < 0.0001), 0.912 (95% confidence interval = 0.850 to 0.974) for total cholesterol (*P* < 0.0001), and 0.855 (95% confidence interval = 0.801 to 0.908) for BMI (*P* < 0.0001), indicating that the MNA test is relatively accurate. The sensitivity, specificity, Youden index, positive predictive value, and negative predictive value of the MNA total score at the selected threshold MNA score are presented in Table 3. Based on biochemical markers (serum albumin or total cholesterol) or BMI as the indicator of malnutrition, the sensitivity and

Table 2
Correlation between MNA total or MNA-SF score and nutritional parameters in Japanese elderly

	No. of subjects	Correlation with MNA total score*		Correlation with MNA-SF score*	
		<i>r</i>	<i>P</i>	<i>r</i>	<i>P</i>
Age	234	-0.14	0.036 [†]	-0.16	0.012 [†]
BMI	233	0.59	<0.0001	0.57	<0.0001
MAC	227	0.50	<0.0001	0.33	<0.0001
TSF	225	0.37	<0.0001	0.24	0.003
CC	225	0.28	<0.0001	0.31	<0.0001
Albumin	179	0.60	<0.0001	0.56	<0.0001
Total cholesterol	177	0.36	<0.0001	0.30	<0.0001
Lymphocyte	161	0.01	0.930	0.04	0.96
MNA total score	226			0.88	<0.0001

BMI, body mass index; CC, calf circumference; MAC, midarm circumference; MNA, Mini-Nutritional Assessment; MNA-SF, Mini-Nutritional Assessment, Short Form; TSF, triceps skinfold

* Partial rank correlation coefficients adjusted for age were used to measure the association between MNA or MNA-SF score and nutritional parameters except for age.

[†] Spearman's rank correlation.




Fig. 1. Receiver operating characteristics (ROC) curve for the (A) full Mini-Nutritional Assessment (MNA) and (B) the MNA short form (MNA-SF) in the Japanese elderly. (A) ROC for MNA as a predictor of albumin levels lower than 3.5 g/dL, total cholesterol levels lower than 150 mg/dL, or BMI lower than 18.5 kg/m². (B) ROC for MNA-SF as a predictor of an MNA score below 24. BMI, body mass index.

specificity of the MNA total score were 0.810 and 0.860 for albumin, 0.786 and 0.822 for total cholesterol, and 0.558 and 0.839 for BMI, respectively, with a cutoff point lower than 17 indicating malnutrition. This suggests that 19% of elderly persons with hypoalbuminemia would be missed (sensitivity), and that 14% without hypoalbuminemia would be classified as malnourished (specificity).

Maximal discrimination between malnutrition and risk of

malnutrition is by definition reached at the cutoff point that has the highest Youden index (sensitivity + specificity – 1). As presented in Table 3, a cutoff point below 15.5 has the highest Youden index, based on hypoalbuminemia and hypocholesterolemia. With a cutoff point below 15.5, the sensitivity and specificity for hypoalbuminemia (<3.5 g/dL) were 0.762 and 0.936, respectively. The figures for hypocholesterolemia and low BMI (<18.5 kg/m²) showed a

Table 3
Validity values of the MNA total score for malnutritional markers in Japanese elderly

	Threshold for MNA total score							
	<15	<15.5	<16	<16.5	<17	<17.5	<18	<18.5
Albumin (<3.5 g/dL)								
Sensitivity	0.714	0.762	0.762	0.762	0.810	0.810	0.857	0.857
Specificity	0.949	0.936	0.924	0.904	0.860	0.834	0.815	0.777
YI	0.663	0.698	0.686	0.666	0.670	0.644	0.672	0.634
PPV	0.625	0.593	0.552	0.500	0.425	0.386	0.375	0.333
NPV	0.943	0.967	0.967	0.966	0.971	0.970	0.977	0.976
Total cholesterol (<150 mg/dL)								
Sensitivity	0.714	0.786	0.786	0.786	0.786	0.786	0.857	0.857
Specificity	0.914	0.902	0.890	0.871	0.822	0.804	0.785	0.748
YI	0.628	0.688	0.676	0.657	0.608	0.590	0.642	0.605
PPV	0.417	0.407	0.379	0.344	0.275	0.256	0.255	0.226
NPV	0.974	0.980	0.980	0.979	0.978	0.978	0.985	0.984
BMI (<18.5 kg/m ²)								
Sensitivity	0.395	0.419	0.442	0.465	0.558	0.581	0.605	0.674
Specificity	0.902	0.891	0.881	0.870	0.839	0.824	0.798	0.772
YI	0.297	0.310	0.323	0.335	0.397	0.405	0.403	0.446
PPV	0.472	0.462	0.452	0.444	0.436	0.424	0.400	0.397
NPV	0.870	0.873	0.876	0.880	0.895	0.898	0.901	0.914

BMI, body mass index; MNA, Mini-Nutritional Assessment; NPV, negative predictive value; PPV, positive predictive value; YI, Youden index

similar pattern when 15.5 rather than 17 was used as the threshold MNA score; the sensitivities were unchanged or decreased and the specificities increased. With a cutoff point below 18, the sensitivity and specificity for hypoalbuminemia were 0.857 and 0.815, respectively; in this case, the sensitivity of the MNA total score increased but the specificity decreased, as did the positive predictive value.

MNA-SF scores averaged 9.8 ± 0.2 and ranged from a minimum of 1 to a maximum of 14. Although MNA-SF contains only BMI as an anthropometric marker, Table 2 presents the significant correlations between MNA-SF score and age, BMI, MAC, CC, TSF, serum albumin, total cholesterol, or MNA total score. However, the higher degree of correlation existed between these nutritional markers and MNA total score. According to MNA criteria (≥ 24) used to define “well nourished,” only 22.1% of subjects were assessed as such. Thus, 77.9% were malnourished or at risk of malnutrition. The correlation between MNA-SF and MNA total scores was high ($r = 0.88$, $P < 0.0001$). The sensitivity, specificity, Youden index, and positive and negative predictive values for different cutoff points for MNA-SF are presented in Table 4. For MNA-SF, the optimal cutoff point was lower than 12 (sensitivity = 0.861, specificity = 0.840, and Youden index = 0.701). This point can be also determined visually from the ROC curve (Fig. 1B).

Discussion

In the present study, the MNA was validated in the Japanese elderly. We demonstrated that the MNA total score showed a good correlation with anthropometric markers and biochemical markers including serum albumin and total cholesterol. The full MNA contains anthropometric indices including BMI, MAC, and CC. To date, no ethnic-specific anthropometric targets exist; rather, these targets are derived from populations of United States or European origin and are inappropriately applied to men and women of Asian descent. Ethnicity has been recognized as a significant modifier in anthropometric measurements [23]. In addition, MNA contains dietary patterns that may differ across ethnicities [24]. Therefore, the MNA or cutoff point for malnutrition may not be a good fit for the Asian, including Japanese, elderly.

Table 4
Validity values of the MNA-SF score for the risk of malnutrition

	Threshold for MNA-SF score				
	<9	<10	<11	<12	<13
Sensitivity	0.385	0.529	0.679	0.861	0.973
Specificity	1.000	0.980	0.960	0.840	0.540
YI	0.385	0.509	0.639	0.701	0.513
PPV	1.000	0.990	0.984	0.953	0.888
NPV	0.303	0.358	0.444	0.618	0.844

MNA-SF, Mini-Nutritional Assessment, Short Form; NPV, negative predictive value; PPV, positive predictive value; YI, Youden index

Several investigators have dealt with the problem of establishing nutritional parameters for the elderly. In the present study we used anthropometric measurements including BMI, MAC, TSF, and CC and biochemical markers such as serum albumin and total cholesterol as nutritional parameters. Although there are no currently, generally accepted criteria for the diagnosis of malnutrition, these parameters have been widely used to evaluate nutritional status. It should be noted that in the present study cutoff points below 3.5 g/dL for serum albumin and below 150 mg/dL for total cholesterol were considered as undernutrition markers. With aging there may be a small decrease in serum albumin [10]. Total cholesterol levels increase with age in healthy individuals and reach a peak between sixth and ninth decades, only to decrease afterward [10]. The cutoff points used in this study for undernutrition markers are widely accepted even in the elderly [10].

We also showed that the MNA is accurate, based on observation of the ROC curve. These results suggested that the MNA is a useful tool to assess the nutritional status of the Japanese elderly. In the elderly populations in Europe and the United States, an MNA total score cutoff point below 17 as an indicator of protein-calorie undernutrition was found to have a sensitivity of 96%, specificity of 98%, and positive predictive value of 97% [11]. However, the same cutoff point yielded a much lower sensitivity and specificity among the Japanese elderly.

For screening purposes, a malnutrition cutoff point below 18 appears to be better than one below 17 for the Japanese elderly, even though higher cutoff points were associated with lower predictive values. However, if MNA is used as a diagnostic tool, a cutoff point below 15.5 is the best for detecting malnutrition in the Japanese elderly because diagnosis requires a high specificity and a high positive predictive value. MNA is a screening tool mainly for malnutrition. MNA results must be confirmed by other anthropometric, biochemical, and dietary parameters to have a complete nutritional status evaluation and malnutrition diagnosis. Therefore, sensitivity is much more important than specificity, and a cutoff point below 18 appears to be more accurate for the Japanese elderly.

We also demonstrated that there were significant correlations between the MNA-SF score and nutritional parameters in addition to the full MNA score, although these correlations were somewhat stronger in MNA total score than in MNA-SF. When a full MNA score of at least 24 was considered the cutoff point for “normal nutrition” in the Japanese elderly, the optimal MNA-SF cutoff point was at least 12, a finding identical to that in the original report [16]. These results suggest that MNA-SF, which comprises six items, allows a quick screening to determine malnutrition and risk of malnutrition in the Japanese elderly. It has been demonstrated that an MNA score between 17 and 23.5, corresponding to “at risk of malnutrition,” can identify older persons with mild malnutrition [25]. In addition, subjects “at risk of malnutrition” had higher mortality rates than did

subjects with a higher MNA score (≥ 24 , corresponding to “normal nutrition”) [13], suggesting that the identification of the elderly “at risk of malnutrition” is very important for clinical practice. It should be noted that, because the present study was a cross-sectional study, the threshold score between “normal nutrition” and “at risk of malnutrition” could not be determined. The category of “at risk of malnutrition” should be defined by longitudinal observation. In addition, the study group might have consisted of frail Japanese elderly who had comorbid diseases, given that they were enrolled from clinical settings. Therefore, it is hard to consider this sample representative of the elderly Japanese population, and our results may apply only to the elderly in ill health.

We believe this is the first study to examine whether the MNA is useful in identifying undernutrition in a non-Caucasian elderly population. However, because the MNA is used mainly as a screening tool for malnutrition among the Japanese elderly, the MNA total cutoff point shifts to a higher point, namely below 18. Further studies are required to evaluate the cutoff point for the “at risk of malnutrition” group. In addition, further investigation should be conducted to determine whether the MNA can correctly identify those elderly who are likely to benefit from nutritional support.

References

- [1] Sullivan DH, Sun S, Walls RC. Protein-energy undernutrition among elderly hospitalized patients: a prospective study. *JAMA* 1999;281:2013.
- [2] Crogan NL, Pasvogel A. The influence of protein-calorie malnutrition on quality of life in nursing homes. *J Gerontol A Biol Sci Med Sci* 2003;58:159.
- [3] Abbasi AA, Rudman D. Undernutrition in the nursing home: prevalence, consequences, causes and prevention. *Nutr Rev* 1994;52:113.
- [4] Bianchetti A, Rozzini R, Carabellese C, Zanetti O, Trabucchi M. Nutritional intake, socioeconomic conditions, and health status in a large elderly population. *J Am Geriatr Soc* 1990;38:521.
- [5] Sullivan DH, Walls RC. Protein-energy undernutrition and the risk of mortality within six years of hospital discharge. *J Am Coll Nutr* 1998;17:571.
- [6] Persson MD, Brismar KE, Katzarski KS, Nordenstrom J, Cederholm TE. Nutritional status using mini nutritional assessment and subjective global assessment predict mortality in geriatric patients. *J Am Geriatr Soc* 2002;50:1996.
- [7] Thomas DR, Zdrowski CD, Wilson MM, et al. Malnutrition in sub-acute care. *Am J Clin Nutr* 2002;75:308.
- [8] Bickford GR, Brugler LJ, Dolsen S, Vickery CE. Nutrition assessment outcomes: a strategy to improve health care. *Clin Lab Manage Rev* 1999;13:357.
- [9] Omran ML, Morley JE. Assessment of protein energy malnutrition in older persons, part I: history, examination, body composition, and screening tools. *Nutrition* 2000;16:50.
- [10] Omran ML, Morley JE. Assessment of protein energy malnutrition in older persons, Part II: laboratory evaluation. *Nutrition* 2000;16:131.
- [11] Guigoz Y, Vellas B, Garry PJ. Assessing the nutritional status of the elderly: the Mini Nutritional Assessment as part of the geriatric evaluation. *Nutr Rev* 1996;54(1 pt 2):S59.
- [12] Vellas B, Guigoz Y, Garry PJ, et al. The Mini Nutritional Assessment (MNA) and its use in grading the nutritional state of elderly patients. *Nutrition* 1999;15:116.
- [13] Visvanathan R, Macintosh C, Callary M, et al. The nutritional status of 250 older Australian recipients of domiciliary care services and its association with outcomes at 12 months. *J Am Geriatr Soc* 2003;51:1007.
- [14] Donini LM, Savina C, Rosano A, et al. MNA predictive value in the follow-up of geriatric patients. *J Nutr Health Aging* 2003;7:282.
- [15] Van Nes MC, Herrmann FR, Gold G, Michel JP, Rizzoli R. Does the mini nutritional assessment predict hospitalization outcomes in older people? *Age Ageing* 2001;30:221.
- [16] Rubenstein LZ, Harker JO, Salva A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice: developing the short-form mini-nutritional assessment (MNA-SF). *J Gerontol A Biol Sci Med Sci* 2001;56:M366.
- [17] Robine JM, Saito Y, Jagger C. The emergence of extremely old people: the case of Japan. *Exp Gerontol* 2003;38:735.
- [18] Garrow JS, Webster J. Quetelet's index (W/H²) as a measure of fatness. *Int J Obes* 1985;9:147.
- [19] James WP, Franco PJ. The choice of cut-off point for distinguishing normal body weights from underweight or 'chronic energy deficiency' in adults. *Eur J Clin Nutr* 1994;48(suppl 3):S179.
- [20] Murphy JM, Berwick DM, Weinstein MC, et al. Performance of screening and diagnostic tests. Application of receiver operating characteristic analysis. *Arch Gen Psychiatry* 1987;44:550.
- [21] Hanley JA, McNeil BJ. A method of comparing the areas under receiver operating characteristic curves derived from the same cases. *Radiology* 1983;148:839.
- [22] Youden WJ. Index for rating diagnostic test. *Cancer* 1950;3:32035.
- [23] Wang J, Thornton JC, Russell M, et al. Asians have lower body mass index (BMI) but higher percent body fat than do whites: comparisons of anthropometric measurements. *Am J Clin Nutr* 1994;60:23.
- [24] Maskarinec G, Novotny R, Tasaki K. Dietary patterns are associated with body mass index in multiethnic women. *J Nutr* 2000;130:3068.
- [25] Vellas B, Guigoz Y, Baumgartner M, et al. Relationships between nutritional markers and the mini-nutritional assessment in 155 older persons. *J Am Geriatr Soc* 2000;48:1300.

High Morning Home Blood Pressure Is Associated with a Loss of Functional Independence in the Community-Dwelling Elderly Aged 75 Years or Older

Masanori NISHINAGA, Jun TAKATA, Kiyohito OKUMIYA*,
Kozo MATSUBAYASHI**, Toshio OZAWA***, and Yoshinori DOI

To elucidate the relationship between home systolic blood pressure (SBP) and functional impairment in the elderly 75 years or older, 461 community-dwelling subjects (192 men, 269 women, mean age: 80 years) were studied. Home blood pressure was measured twice in the morning and twice in the evening for 5 consecutive days with an automatic cuff-oscillometric device. Total/high-density lipoprotein cholesterol and several functional assessments were evaluated. A subject was determined to exhibit a loss of independence according to the activities of daily living (ADL) score in a study conducted in 2001. Based on the mean home SBPs (mSBP) and morning-evening SBP differences (dSBP), the subjects were classified into 4 groups as follows: hypertensive/morning-dominant (HM; mSBP \geq 135 mmHg, dSBP \geq 15 mmHg), hypertensive/sustained (HS; mSBP \geq 135 mmHg, dSBP $<$ 15 mmHg), normotensive/morning-dominant (NM; mSBP $<$ 135 mmHg, dSBP \geq 15 mmHg), and normotensive/controlled (NC; mSBP $<$ 135 mmHg, dSBP $<$ 15 mmHg). There were no differences in sex, cholesterol levels, history of stroke, other cardiovascular diseases (CVDs), and cognitive function, but there were significant differences in age, antihypertensive medications, the neurobehavioral test scores, and ADL scores. There were no significant differences in terms of mortality and CVD events. In the survivors, HM and HS were independent risk factors for a loss of independence, after adjustments were made for onset of stroke, age, antihypertensive therapy, history of CVD, as well as neurobehavioral test scores and ADL scores (odds ratio [OR]: 12.2 and 3.78, respectively). After the same adjustments as those mentioned above were made, HM and HS were found to be negative determinants of survival and maintenance of independence (OR: 0.082, 0.270, respectively). In conclusion, high home SBP (\geq 135 mmHg) and high dSBP (\geq 15 mmHg) were found to be important in determining the levels of disability for the very elderly. (*Hypertens Res* 2005; 28: 657–663)

Key Words: home blood pressure, elderly, morning hypertension, independence, successful aging

Introduction

Recently there have been rapid increases in both the popula-

tion and life span of the elderly in developed countries, which has resulted in a considerable increase in the number of frail elderly people. "The project to reduce the number of dependent elderly persons" has been promoted as an important

From the Department of Medicine and Geriatrics, Kochi Medical School, Kochi, Japan; *Research Institute for Humanity and Nature, Kyoto, Japan; **South-East Asia Research Center, Kyoto University, Kyoto, Japan; and ***Tokyo Metropolitan Geriatric Hospital, Tokyo, Japan.

This study was supported in part by research grants from the Japan Arteriosclerosis Prevention Fund and the Ministry of Health, Labor and Welfare.

Address for Reprints: Masanori Nishinaga, M.D., Ph.D., Department of Medicine and Geriatrics, Kochi Medical School, Kohasu, Oko-cho, Nankoku 783–8505, Japan. E-mail: nishinam@med.kochi-u.ac.jp

Received November 29, 2004; Accepted in revised form June 15, 2005.




Fig. 1. Classification of four groups according to the distribution of mean home SBP (mSBP) and morning-evening home SBP differences (dSBP). HM, hypertensive/morning-dominant; HS, hypertensive/sustained; NM, normotensive/morning-dominant; NC, normotensive/controlled.




Fig. 2. Outcome after a 9-year follow-up period. CVD, cardiovascular disease, including stroke.

issue not only from the medical, but also from the socioeconomic point of view. This project should be considered a top priority, and appropriate measures should be taken to improve the current situation (1). Although stroke is a major cause of mortality and disability in the elderly (2-4), the management of hypertension in the community has contributed to an outstanding reduction in the incidence of stroke (5, 6).

In many previous epidemiological studies on blood pressure (BP) in the elderly, the mortality and morbidity associated with stroke and other cardiovascular diseases (CVDs) have been selected as the endpoint (7-9). There have been also several epidemiological studies of the relationship between hypertension and dementia (10, 11). However, assessments of functional abilities, the most important factor for elderly persons and their caregivers, have been rarely conducted.

Improving the management of BP for the prevention of stroke and other CVDs has led to the popularization of home BP monitoring devices among the general public (12, 13). However, there have been few reports concerning home BP values in the elderly. Increased BP in the morning is considered as a strong risk factor for stroke and other CVDs (14-17). It remains to be clarified whether such an elevation in BP in the morning is also a risk factor for a loss of functional independence in the elderly.

In the present study, we recruited community-dwelling elderly people 75 years of age or older, and conducted medical and functional assessments. We followed the subjects for 9 years. The purpose of our study was to clarify the relationship between home BP values and functional disabilities, as well as that between home BP and the mortality/incidence of stroke. Furthermore, we studied morning-evening home BP differences in this very elderly sample.

Methods

Subjects

The study subjects were elderly people, aged 75 years or older, who resided in Kahoku Town, Kochi Prefecture, Japan in 1992. All subjects applied to participate in the home BP monitoring program. Subjects with atrial fibrillation were excluded because of the potential inaccuracy of their home BP measurements. A total of 461 people were recruited as the subjects of our study (192 men, 269 women, mean age: 81 years).

Home BP Measurement

Home BP was measured in 1992 with an automatic device (HEM-755C; OMRON Life Science Co., Ltd., Kyoto, Japan) based on the cuff-oscillometric method. The validity of BP measurement according to this method has been reported in several studies (18-20). The subjects and their caregivers were taught by community nurses how to measure the BP at home using this device. According to a previously reported method (21), BP was measured in the non-dominant arm after taking at least a 5-min rest in a sitting position, twice in the morning (6-7 AM) and twice in the evening (8-9 PM), for 5 consecutive weekdays.

We obtained the data regarding the total mean systolic BP (mSBP), morning and evening systolic BP (SBP), morning and evening diastolic BP (DBP) and pulse rates (PR). The mean morning-evening SBP differences (dSBP) were calculated. We defined subjects with mSBP ≥ 135 mmHg as hypertensive or poorly controlled subjects, according to the Japanese Society of Hypertension (JSH) Guidelines for the Self-Monitoring of Blood Pressure at Home (12). We also divided our subjects into two groups by using mean + 1SD of the dSBP. There were no subjects whose mean DBP values alone exceeded 85 mmHg.

Table 1. Basic Characteristics (1992)

	NC (n=159)	NM (n=30)	HS (n=207)	HM (n=65)
Age (years)*	80.3±4.5	81.5±5.1	80.5±5.1	81.4±5.3
Men (n [%])	74 (46.5)	13 (43.3)	79 (38.1)	26 (40.0)
SBP (mmHg)**	125±8	124±8	148±11	150±10
DBP (mmHg)**	72±7	73±9	80±11	81±9
PR(/min)*	68±8	67±7	69±8	68±12
Morning SBP (mmHg)**	126±9	135±9	149±12	162±11
Evening SBP (mmHg)**	123±9	113±9	148±12	138±11
Total cholesterol (mg/dl)	190±35	210±39	186±38	195±42
HDL cholesterol (mg/dl)	48±14	56±10	45±13	44±12
ADL (full score: 21)	20.4±1.9	20.8±0.8	20.2±2.6	20.2±1.9
MMSE (full score: 30)	27.2±4.4	28.4±2.2	27.3±3.2	27.5±3.2
Up and Go test (s)	13.0±3.2	13.0±2.8	14.1±4.6	14.9±5.4
Antihypertensive drugs** (Yes, n [%])	38 (23.9)	10 (33.3)	89 (43.0)	40 (61.5)
History of CVD (Yes, n [%])	16 (10.0)	1 (3.3)	13 (6.3)	4 (8.1)

NC, normotensive/controlled; NM, normotensive/morning-dominant; HS, hypertensive/sustained; HM, hypertensive/morning-dominant; SBP, systolic blood pressure; DBP, diastolic blood pressure; PR, pulse rate; HDL, high-density lipoprotein; ADL, activities of daily living; MMSE, mini-mental state examination; CVD, cardiovascular disease, including stroke. *ANOVA, $p < 0.05$, **ANOVA, $p < 0.01$.

Annual Self-Administered Questionnaire

In the baseline survey (1992), the self-administered questionnaire was addressed to the study subjects to obtain information about characteristics potentially related to their BP, mortality, and disability; the data collected included a history of stroke, heart disease, and bone disease or arthropathy, anti-hypertensive medications, current and past cigarette smoking, current intake of alcohol, and activities of daily living (ADL). All of the response sheets submitted by the subjects were reviewed by community nurses to ascertain their information.

Assessment of ADL, Cognitive Function, Neurobehavioral Function, Mood and Serum Lipid Analysis

The questionnaire regarding the ADL was conducted in 1992 and 2001 in the same manner as was used in our previous study (22). Briefly, ADL were assessed with respect to the following seven items: walking, ascending and descending stairs, feeding, dressing, using the toilet, bathing, and grooming. Each ADL item was scored on a 0–3 scale: 0 = completely dependent, 1 = needs a lot of help, 2 = needs some help, and 3 = completely independent. The scores for these seven items were summarized to obtain a total ADL score ranging from 0 to 21. When a subject did not maintain a score of 21 or 20 points in 2001, he or she was defined as a person that was losing independence.

The mini-mental state examination (MMSE) was used to evaluate each subject's level of cognitive functioning (23). The Up and Go test was used to evaluate neurobehavioral function (24). This latter test measures, in s, the time it takes the subject to stand up from an armchair, walk a distance of 3

m, walk back to the chair, and sit down again. This simple test is a comprehensive evaluation of the subject's balance, gait speed, and functional ability. Since Okumiya and co-workers (25) reported its usefulness in predicting a decline in ADL in the Japanese community-dwelling elderly, the test has been widely accepted for this purpose in many fields. The Geriatric Depression Scale 15 (GDS 15) was also used to evaluate the mood of the subjects (depressive state) (26). Total serum cholesterol and serum high-density lipoprotein (HDL) cholesterol were analyzed in 1992.

During the period between 1992 and 2001, a total of 125 subjects (66 men, 59 women) died. In addition, the information regarding the events of stroke, myocardial infarction, congestive heart failure, and bone/joint diseases as causes of disability were collected by checking the responses provided on the annual questionnaire and the subjects' medical records.

Written informed consent was obtained from each subject at the time of the annual questionnaire. Our study was approved by the Research Ethics Committee of Kochi Medical School, Kochi University, Japan.

Statistical Analysis

All of the values were expressed as mean±SD. Mean values among the groups were compared using ANOVA. A χ^2 test was used to compare the 4 groups with respect to total mortality and incidence of stroke, as well as other CVDs. A logistic multivariate analysis was used to identify the factors that predicted a loss of functional independence or the survival and maintenance of functional independence 9 years after the initial assessment, using Stat View 5.0 for Windows (SAS Institute Inc., Cary, USA).

Table 2. Total/CVD Death and Non-Fatal Stroke

	NC (n=159)	NM (n=30)	HS (n=207)	HM (n=65)
Total death (%)*	36 (22.6)	3 (10.0)	62 (30.0)	24 (36.9)
Non-fatal stroke(%)	16 (10.1)	1 (3.3)	12 (5.8)	4 (6.2)
CVD death (%)	17 (10.7)	1 (3.3)	24 (11.6)	11 (16.9)
CVD events (%)	33 (20.8)	2 (6.7)	36 (17.4)	15 (23.1)

NC, normotensive/controlled; NM, normotensive/morning-dominant; HS, hypertensive/sustained; HM, hypertensive/morning-dominant; CVD, cardiovascular disease, including stroke. *ANOVA $p < 0.1$.

Results

The distribution of mSBP and dSBP are shown in Fig. 1. The subjects were classified into the following 4 groups:

Hypertensive/morning-dominant (HM: mSBP \geq 135 mmHg, dSBP \geq 15 mmHg; $n=65$), hypertensive/sustained (HS: mSBP \geq 135 mmHg, dSBP $<$ 15 mmHg; $n=207$), normotensive/morning-dominant (NM: mSBP $<$ 135 mmHg, dSBP \geq 15 mmHg; $n=30$), and normotensive controlled (NC: mSBP $<$ 135 mmHg, dSBP $<$ 15 mmHg; $n=159$). The NC group, which was expected to be the lowest risk group because both the mSBP and the dSBP were lower than others, was used for reference.

A total of 461 elderly subjects, who were 75 years of age or older in 1992, were followed for 9 years until 2001 (Fig. 2). During that interval, 125 (27%) subjects died; 53 of these subjects had died of stroke and other CVDs. A total of 336 of the subjects were alive 9 years later (2001). One hundred-eighteen subjects had undergone a loss of their functional independence (HM, 17 [41%]; HS, 65 [45%]; NM, 7 [26%]; NC, 29 [24%]).

The basic characteristics of the 4 groups in 1992 are shown in Table 1. The subjects in the NC group were younger than those in the other 3 groups. The percentage of subjects who were taking antihypertensive agents was also the lowest in the NC group. There were no significant differences in terms of sex, PR, total serum cholesterol, HDL cholesterol, the scores of ADL, MMSE, the Up and Go test, or history of CVD among the 4 groups. There were also no differences in the scores on the GDS 15, history of bone/joint diseases, current and past cigarette smoking, and current intake of alcohol (data not shown).

Table 2 shows the total number of deaths, the number and percentage of deaths caused by stroke and other CVDs, and the incidence of non-fatal stroke during the 9-year follow-up period in the 4 groups. Although there was a difference in the total number of deaths among the 4 groups before adjustment for age, the significance of this difference disappeared after adjustment for age. There were no significant differences in the percentage of deaths from stroke or other CVDs. Although 33 subjects suffered from symptomatic strokes, no significant differences were seen in the incidence of strokes among the 4 groups.

The risk factors for loss of functional independence are shown in Table 3. Although a non-fatal event of stroke was one of the most important risk factors for loss of functional independence, HM and HS were also important risk factors, even after adjustment for age, sex, antihypertensive therapy, scores on the Up and Go test in 1992, and the ADL scores in 1992. The adjusted odds ratio (OR) of the HM group (12.2) was significantly higher than that of the HS group (3.78). Therefore, values of mSBP \geq 135 mmHg and dSBP \geq 15 mmHg were independent risk factors for a loss of functional independence (Fig. 3).

The factors associated with successful aging that contributed to the survival and maintenance of functional independence, even among the most elderly (age of 84 or older), are shown in Table 4. Although the non-fatal event of stroke was a significantly negative determinant, the HM and HS also remained as significant independent negative determinants of successful aging, after adjustment for age, sex, antihypertensive therapy, scores on the Up and Go test in 1992, and ADL scores in 1992.

As regards the elderly people aged 75 years or older living in the community, values of mSBP \geq 135 mmHg and dSBP \geq 15 mmHg were independent determinants of a loss of functional independence or successful aging, even when non-fatal stroke and these home SBP variables were simultaneously incorporated into a logistic multivariate analysis model.

In addition, since many of our subjects with morning hypertension had high home SBP, we added dSBP (\geq 15 mmHg) to the same model of multivariate logistic analysis, in order to elucidate whether dSBP was an independent determinant of a loss of functional independence or alive and independence. dSBP remained a significant determinant of a loss of independence (adjusted OR: 3.84, 95% confidence interval [CI]: 1.003–14.73), or alive and independence (adjusted OR: 0.46, 95% CI: 0.183–0.973), in our hypertensive subjects.

Discussion

Our prospective longitudinal study evaluating the maintenance of independence in the elderly aged 75 years or older demonstrated that a mean home SBP of \geq 135 mmHg was a significantly important risk factor for a loss of functional independence. In addition, morning hypertension was an

Table 3. Independent Risk Factors for Loss of Independence in 336 Survivors

Factors	Adjusted odds ratio	95% CI	p
Stroke	17.4	3.67–82.8	0.0003
HM	12.2	3.00–50.0	0.0005
HS	3.78	1.45–9.83	0.0064
Age	1.17	1.05–1.30	0.0036

Data were adjusted for sex, antihypertensive therapy, Up and Go score, and activities of daily living (ADL) score in 1992. CI, confidence interval; HM, hypertensive/morning-dominant; HS, hypertensive/sustained.

Table 4. Independent Negative Factors for Survival and Maintenance of Independence (n=461)

Factors	Adjusted odds ratio	95% CI	p
Stroke	0.058	0.012–0.273	0.0003
HM	0.082	0.020–0.334	0.0005
HS	0.271	0.104–0.704	0.0073
Age	0.855	0.768–0.951	0.0038

Data were adjusted for sex, antihypertensive therapy, Up and Go score, and activities of daily living (ADL) score in 1992. CI, confidence interval; HM, hypertensive/morning-dominant; HS, hypertensive/sustained.




Fig. 3. Adjusted odds ratios for loss of independence among the 4 groups. The abbreviations are the same as those introduced in Fig. 1.

important independent predictor for the functional prognosis of our elderly subjects. The significance of dSBP should be further evaluated.

There were no differences in mortality and incidence of non-fatal stroke among the 4 groups. However, when a loss of functional independence was selected as the endpoint, significant differences were found among these groups. Home BP monitoring in the morning and in the evening was useful in predicting functional prognosis in elderly subjects aged 75 years or older.

Recent large clinical trials have clarified the importance of BP control, even among the elderly (7–9). However, the optimum home BP value for the elderly has not been established, and further prospective studies on the elderly will be necessary to define an adequate home BP value (27).

In the present study, there was no significant difference in the incidence of CVD death, stroke, and other CVD events among the 4 groups. Although the reason for this result is uncertain, further studies with a larger number of subjects may resolve this ambiguity.

Since a relationship between elevation of BP in the morning and stroke and other CVD events (14–17) has been reported, the importance of morning BP has been emphasized in studies of home BP monitoring (28, 29). However, in most of these studies the mortality and morbidity of stroke and

other CVDs, as well as organ damage, were selected as the endpoints. Functional independence, which is important for the elderly as well as for the social economy, is not mentioned in these previous studies. Thus, in our present study, we added a loss of functional independence and successful aging as two new endpoints for geriatric study. Furthermore, we evaluated morning–evening home BP differences in terms of the usefulness of this information for the prognosis of the elderly aged 75 years or older.

Skoog *et al.* (30) reported the relationship between the presence of hypertension at the age of 70 and the development of dementia 10 to 15 years later. In subsequent large studies including SCOPE (10), the association between impaired cognitive function and BP values has been evaluated, although sufficient data on the relationship between BP control and cognitive function have yet to be accumulated. In our previous study (31), a J-curve phenomenon was demonstrated with respect to the profile of the association between BP values and cognitive function 3 years later in an elderly sample. Those findings indicate that BP exerted an effect on cognitive function, not only in the group with high BP, but also in the group with low BP. In our previous study, casual BP was measured twice with the subject in the supine position at the time of physical examination. Here, to avoid the inclusion of various other factors affecting BP measurements, we used 20 home BP measurements in order to calculate the mean value. This method of measurement appeared to have eliminated some of the potential problems with BP monitoring.

Because the follow-up period was so long (9 years), it was difficult to reexamine cognitive function in all of the subjects examined in 2001; some subjects were too old for us to obtain reliable data from them (*i.e.*, among those at least aged 84 years of age and older). Due to our small sample size in the MMSE evaluation ($n=64$, 19%), we did not observe any significant differences between the group with high home BP values (≥ 135 mmHg) and the group with normal home BP values (< 135 mmHg). There were also no significant differences in dSBP (≥ 15 mmHg). Evaluation of the cognitive functions (MMSE) of the most elderly subjects included in the sample was difficult; thus, an appropriate, reliable method

will still need to be developed for the evaluation of this group in future studies.

A variety of factors, including stroke events and other CVDs, as well as the progression of bone/joint diseases and dementia, contributes to disability among the elderly (32). The present study demonstrated for the first time that those elderly persons with a mean home SBP ≥ 135 mmHg were susceptible to a loss of functional independence, even if they did not experience an event of symptomatic stroke. However, the present study did not reveal any direct mechanism to generate this relationship between hypertension reflected by the home BP monitoring value and a loss of independence. The relationships between hypertension or morning BP elevation and pathological conditions such as asymptomatic small infarctions and white matter lesions have previously been reported (33–36). It is thus possible that these lesions are related to BP elevation and may be involved in the impairment of functional abilities required for daily life. To clarify the association between home BP values, asymptomatic brain lesions, and disability among the very elderly, further study is warranted.

References

- Sasaki H, Sekizawa K, Yanai M, Arai H, Yamaya M, Ohru T: Elongation of life expectancy may accompany shift of medical cost to older adults. *J Am Geriatr Soc* 1997; **45**: 254–255 (Letter).
- Carolei A, Sacco S, De Santis F, Marini C: Epidemiology of stroke. *Clin Exp Hypertens* 2002; **24**: 479–483.
- Kobayashi S: Data bank project for acute stroke patients. *Rinsho Shinkeigaku* 2002; **42**: 1176–1178.
- Ikebe T, Ozawa H, Lida M, Shimamoto T, Handa K, Komachi Y: Long-term prognosis after stroke: a community-based study in Japan. *J Epidemiol* 2001; **11**: 8–15.
- Neal B, MacMahon S, Chapman N: Effects of ACE inhibitors, calcium antagonists, and other blood-pressure lowering drugs: results of prospectively designed overviews of randomized trials. *Lancet* 2000; **356**: 1955–1964.
- Sutton-Tyrrell K, Wildman R, Newman A, Kuller LH: Extent of cardiovascular risk reduction associated with treatment of isolated systolic hypertension. *Arch Intern Med* 2003; **163**: 2728–2731.
- Fagard RH, Staessen JA: Treatment of isolated systolic hypertension in the elderly: the Syst-Eur trial. Systolic Hypertension in Europe (Syst-Eur) Trial Investigators. *Clin Exp Hypertens* 1999; **21**: 491–497.
- SHEP Cooperative Research Group: Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program. *JAMA* 1991; **265**: 3255–3264.
- MRC Working Party: Medical Research Council trial of treatment of hypertension in older adults: principal results. *BMJ* 1992; **304**: 405–412.
- Lithell H, Hansson L, Skoog I, *et al*: The study on cognition and prognosis in the elderly (SCOPE): principal results of a randomized double-blind intervention trial. *J Hypertens* 2003; **21**: 875–886.
- Birkenhager WH, Forette F, Seux ML, Wang JG, Staessen JA: Blood pressure, cognitive functions, and prevention of dementias in older persons with hypertension. *Arch Intern Med* 2001; **161**: 152–156.
- Imai Y, Otsuka K, Kawano Y, *et al*: Japanese Society of Hypertension (JSH) Guidelines for Self-Monitoring of Blood Pressure at Home. *Hypertens Res* 2003; **26**: 771–782.
- Pickering TG: How should we take blood pressure in clinical practice? *Br Med J* 2003; **327**: E150–E151.
- Kario K, Pickering TG, Umeda Y, *et al*: Morning surge in blood pressure as a predictor of silent and clinical cerebrovascular disease in elderly hypertensives: a prospective study. *Circulation* 2003; **107**: 1401–1406.
- Imai Y, Hozawa A, Ohkubo T, *et al*: Predictive values of automated blood pressure measurement: what can we learn from the Japanese population—the Ohasama study. *Blood Press Monit* 2001; **6**: 335–339.
- Gosse P, Cipriano C, Bemurat L, *et al*: Prognostic significance of blood pressure measured on rising. *J Hum Hypertens* 2001; **15**: 413–417.
- Kuwajima I, Mitani K, Miyao M, Suzuki Y, Kuramoto K, Ozawa T: Cardiac implications of the morning surge in blood pressure in elderly hypertensive patients: relation to arising time. *Am J Hypertens* 1995; **8**: 29–33.
- Stergiou GS, Voutsas AV, Achimastos AD, Moutokalakis TD: Home self-monitoring of blood pressure: is fully automated oscillometric technique as good as conventional stethoscopic technique? *Am J Hypertens* 1997; **10**: 428–433.
- Yarows SA, Amerena JV: Determination of accuracies of 10 models of home blood pressure monitors using an oscillometric simulator. *Blood Press Monit* 1999; **4**: 45–52.
- Yarows SA, Brooks R: Measurement variation among 12 electronic home blood pressure monitors. *Am J Hypertens* 1999; **13**: 45–52.
- Okumiyama K, Matsubayashi K, Wada T, *et al*: A U-shaped association between home systolic blood pressure and four-year mortality in community-dwelling older men. *J Am Geriatr Soc* 1999; **47**: 1415–1421.
- Matsubayashi K, Okumiyama K, Osaki Y, Fujisawa M, Doi Y: Frailty in elderly Japanese. *Lancet* 1999; **353**: 1445.
- Folstein MF, Folstein SE, McHugh PR: A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; **12**: 189–198.
- Podsiadlo D, Richardson S: The timed “Up & Go”: a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991; **39**: 142–148.
- Okumiyama K, Matsubayashi K, Nakamura T, *et al*: The timed “Up & Go” test and manual button score are useful predictors of functional decline in basic and instrumental ADL in community-dwelling older people. *J Am Geriatr Soc* 1999; **47**: 497–498.
- Brink TL, Yesavage JA, Lum O, *et al*: Screening tests of geriatric depression. *Clin Gerontol* 1982; **1**: 37–43.
- Aoki Y, Asayama K, Ohkubo T, *et al*: Progress report on the HOMED-BP study: hypertension objective treatment based on measurement by electrical devices of blood pressure study. *Clin Exp Hypertens* 2004; **26**: 119–127.

28. Ikeda T, Gomi T, Shibuya Y, et al: Morning rise in blood pressure is a predictor of left ventricular hypertrophy in treated hypertensive patients. *Hypertens Res* 2004; **27**: 939–946.
29. Kamoi K, Kaneko S, Miyakoshi M, Nakagawa O, Soda S: Usefulness of home blood pressure measurement in the morning in type 2 diabetic patients. *Diabetes Care* 2002; **25**: 2218–2223.
30. Skoog I, Lernfelt B, Landahl S, et al: 15-year longitudinal study of blood pressure and dementia. *Lancet* 1996; **347**: 1141–1145.
31. Okumiya K, Matsubayashi K, Wada T, Osaki Y, Doi Y, Ozawa T: J-curve relation between blood pressure and decline in cognitive function in older people living in community, Japan. *J Am Geriatr Soc* 1997; **45**: 1032–1033.
32. Kattainen A, Koskinen S, Reunanen A, Martelin T, Knekt P, Aromaa A: Impact of cardiovascular diseases on activity limitations and need for help among older persons. *J Clin Epidemiol* 2004; **57**: 82–88.
33. Yoshitake T, Kiyohara Y, Kato I, et al: Incidence and risk factors of vascular dementia and Alzheimer's disease in a defined elderly Japanese population: the Hisayama Study. *Neurology* 1995; **45**: 1161–1168.
34. Vermeer SE, Prins ND, den Heijer T, Hofman A, Koudstaal PJ, Breteler MM: Silent brain infarcts and the risk of dementia and cognitive decline. *N Engl J Med* 2003; **348**: 1215–1222.
35. Vermeer SE, Hollander M, van Dijk EJ, Hofman A, Koudstaal PJ, Breteler MM: Silent brain infarcts and white matter lesions increase stroke risk in the general population: the Rotterdam Scan Study. *Stroke* 2003; **34**: 1126–1129.
36. Artero S, Tiemeier H, Prins ND, Sabatier R, Breteler MM, Ritchie K: Neuroanatomical localisation and clinical correlates of white matter lesions in the elderly. *J Neurol Neurosurg Psychiatry* 2004; **75**: 1304–1308.



Arterial stiffness independently predicts cardiovascular events in an elderly community – Longitudinal Investigation for the Longevity and Aging in Hokkaido County (LILAC) study

O. Matsuoka^{a, *}, K. Otsuka^{a,b}, S. Murakami^c, N. Hotta^{a,b},
G. Yamanaka^a, Y. Kubo^a, T. Yamanaka^{a,b}, M. Shinagawa^a, S. Nunoda^a,
Y. Nishimura^a, K. Shibata^{a,b}, H. Saitoh^{a,b}, M. Nishinaga^d, M. Ishine^e,
T. Wada^e, K. Okumiya^f, K. Matsubayashi^g, S. Yano^h, K. Ichiharaⁱ,
G. Cornélissen^j, F. Halberg^j, T. Ozawa^k

^a Department of Medicine, Division of Neurocardiology and Chronoecology, Tokyo Women's Medical University, Medical Center East, Nishiogu 2-1-10, Arakawa, Tokyo, 116-8567, Japan

^b Division of Neurocardiology and Chronoecology, Tokyo Women's Medical University, Medical Center East, Nishiogu 2-1-10, Arakawa, Tokyo, 116-8567, Japan

^c Department of Internal Medicine, Osaka Medical University, Osaka, Japan

^d Department of Gerontology, School of Medicine, Kochi University

^e Department of Field Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan

^f Research Institute for Humanity and Nature, Kyoto, Japan

^g Center for South-east Asian Studies, Kyoto University, Kyoto, Japan

^h Sorachi Health and Welfare Office, Sorachi-godochosha, Iwamizawa, Hokkaido, Japan

ⁱ Division of Clinical Laboratory Sciences, Faculty of Health Sciences, School of Medicine, Yamaguchi University, Ube, Japan

^j Halberg Chronobiology Center, University of Minnesota, Minneapolis, Minnesota, USA

^k Tokyo Metropolitan Geriatric Hospital, Tokyo, Japan

Abstract

We investigated the predictive value of arterial stiffness to assess cardiovascular risk in elderly community-dwelling people by means of a multivariate Cox model. In 298 people older than 75 years (120 men and 178 women, average age: 79.6 years), brachial-ankle pulse wave velocity (baPWV) was measured between the right arm and ankle in a supine position. The LILAC study started on July 25, 2000, consultation was repeated yearly, and the last follow-up ended on November 30, 2004. During this follow-up span of 1227 days, there were nine cardiovascular deaths, the cause of death being myocardial infarction for two men and three women or stroke for two men and two women. In Cox proportional hazard models, baPWV as well as age, Mini-Mental State Examination (MMSE), Hasegawa Dementia Scale Revised (HDSR) and the low-frequency/high-frequency (LF/HF) ratio showed a statistically significant association with the occurrence of cardiovascular death. A two-point increase in MMSE and HDSR score significantly protected against cardiovascular death, the relative risk (RR) being 0.776 ($P = 0.0369$) and 0.753 ($P = 0.0029$), respectively. The LF/HF ratio also was significant ($P = 0.025$), but the other indices of HRV were not. After adjustment for age and HDSR, a 200 cm/s increase in baPWV was associated with a 30.2% increase in risk (RR = 1.302, 95% CI: 1.110–1.525), and a 500 cm/s increase in baPWV with a 93.3% increase in risk (RR = 1.933, 95% CI: 1.300–2.874, $P = 0.0011$), whereas the LF/HF ratio was no longer associated with a statistically significant increase in cardiovascular mortality. In elderly community-dwelling people, arterial stiffness measured by means of baPWV predicted the occurrence of cardiovascular death beyond the prediction provided by age, gender, blood pressure and cognitive functions. baPWV should be added to the cardiovascular assessment in various clinical settings, including field medical surveys and preventive screening. The early detection of risk

* Corresponding author.

E-mail address: otsukagm@dnh.twmu.ac.jp (K. Matsuoka).

by chronomics allows the timely institution of prophylactic measures, thereby shifting the focus from rehabilitation to prehabilitation medicine, as a public service to several Japanese towns.

© 2005 Elsevier SAS. All rights reserved.

Keywords: Pulse wave velocity; Cardiovascular risk; Cognitive function; Elderly community-dwelling people

1. Introduction

As the incidence of myocardial infarction has increased in Japan and since cardiovascular disease, including stroke is the leading cause of mortality and morbidity, the prevention of these conditions is a major goal. Atherosclerosis is an important cause of morbidity and mortality in the elderly, and arterial stiffness may predict cardiovascular events [1]. Several cardiovascular indices, such as blood pressure (BP), heart rate (HR) and heart rate variability (HRV) are predictors of vascular disease risk [2]. This study aimed at assessing their role in association with arterial stiffness in predicting cardiovascular mortality. Arterial stiffness can be assessed non-invasively by measuring pulse wave velocity (PWV), which is a simple and reproducible endpoint.

Recent cohort studies suggested that PWV might be a strong predictor of atherosclerotic cardiovascular events, cardiovascular mortality, and all-cause mortality, independently of age and conventional atherosclerotic risk factors [3–5]. Most studies, however, focused on patients of occidental ancestry, often affected by specific diseases, rather than on Japanese populations. In 2000, we began a community-based study to longitudinally investigate the longevity and aging of a rural population in Hokkaido County (LILAC), and to evaluate this population's neurocardiological function. Our goal was the prevention of cardiovascular events, including strokes and myocardial ischemic events, and to prevent the associated decline in cognitive function of the elderly in this community dwelling.

As the first step of the LILAC study, we focused on the relationship between aortic stiffness measured by PWV and cardiovascular mortality. We assessed the predictive value of arterial stiffness in predicting cardiovascular risk in an elderly population using a multivariate Cox model.

2. Methods

2.1. Subjects and LILAC study design

We examined 298 people older than 75 years (average age, 79.6 years). BP was measured in a sitting position at the beginning of the study, and brachial-ankle PWV (baPWV) was measured between the right arm and ankle in a supine position, using an ABI/Form instrument (Nippon Colin Co., Ltd., Komaki, Japan). The baPWV was measured using a volume-plethysmographic method. ECG electrodes were placed on both wrists, a microphone for detecting heart sounds was placed on the left edge of the sternum, and cuffs

were wrapped on both arms and ankles. The cuffs were connected to a plethysmographic sensor that determines the volume pulse form and oscillometric pressure waveforms were recorded using a semiconductor pressure sensor (the sample acquisition frequency for PWV was set at 1200 Hz). baPWV was measured in duplicate after at least a 5-min rest in each case. Only baPWV measures from participants showing an ankle/brachial pressure index (ABI) value above 0.90 were used for analysis.

We used the first hour of ambulatory ECG recording obtained during routine medical examinations conducted each year in July. The data were processed for HRV using a Fukuda-Denshi Holter analysis system (SCM-280-3). Time-domain measures (SDNN, pNN50, SDANN and Lorenz plot indices: Length (L), Width (W), and L/W ratio) and frequency-domain measures (spectral power in the "very low frequency"—VLF: 0.003–0.04 Hz, "low frequency"—LF: 0.04–0.15 Hz, and "high frequency"—HF: 0.15–0.40 Hz regions, and the LF/HF ratio) were determined. Except for SDNN and HR, calculated over the whole 1-h record, all indices were computed as averages over consecutive 5-min intervals. Spectral indices were obtained by the maximum entropy method (MEM) with the MemCalc/CHIRAM program (Suwa Trust Co., Ltd., Tokyo, Japan).

The Japanese versions of the Mini-Mental State Examination (MMSE) and the Hasegawa Dementia Scale Revised (HDSR) tests were used to measure the overall cognitive function, including verbal orientation, memory, and constructional ability (Kohs' block test). The Up and Go test measured, in seconds, the time it took the subject to stand up from a chair, walk a distance of 3 m, turn, walk back to the chair, and sit down again. This test is a simple measure of physical mobility and demonstrates the subject's balance, gait speed, and functional ability. A lower time score indicates better physical mobility. Functional Reach (FR), used to evaluate balance, represents the maximal distance a subject can reach forward beyond arm's length while maintaining a fixed base of support in the standing position. A higher score indicates better balance. Manual dexterity was assessed using a panel with combinations of 10 hooks, 10 big buttons, and five small buttons. There were three discrete measurements of time recorded for each participant (10 "hook-on"s, 10 big "button-on-and-off"s, and five small "button-on-and-off"s). The total manual dexterity time in seconds, defined as the button score (Button-S), was calculated by adding the average times for one hook-on and one big or small button-on-and-off. A lower button score indicates better manual dexterity.

2.2. Cardiovascular mortality

The LILAC study was started on July 25, 2000 and consultations were repeated every year (end of July, or beginning of August). In addition, one or two doctors of our team visited every 3 months and offered several kinds of health consultation, rehabilitation of disordered function, healthy lifestyle modification by promoting complete cessation of smoking, weight reduction, reduction of salt intake, moderation in the consumption of fruits and vegetables and alcohol intake, as well as providing prescription advice to the local general medical practitioner.

In this investigation, the follow-up ended on November 30, 2004. During this follow-up span, there were nine cardiovascular deaths, the cause of death being myocardial infarction or stroke. Follow-up time was defined as the time elapsed between the first (reference) examination and the time of first cardiovascular event or death.

2.3. Statistical analysis

All data were analyzed with the Statistical Software for Windows (StatFlex Ver.5.0, Artec, Osaka, <http://www.statflex.net>). The effects of classic risk factors on baPWV were assessed by a multivariate regression analysis. We used Cox's regression analysis to calculate the unadjusted and adjusted relative risks (RRs) and 95% confidence intervals (CIs) for cardiovascular death. To identify independent predictors of cardiovascular death in relation to baPWV, we used a multivariate Cox regression analysis with stepwise selection. Variables included in the multivariate models were age, gender, BP, HR, HRV, MMSE, HDSR and Kohs' block test.

An abnormal value of baPWV (>2500 cm/sec) was independently assessed by logistic regression analysis. Kaplan–

Meier event probability curves were computed with two groups, stratified by the abnormal value of baPWV, and the cumulative probability of events of two groups was compared by means of the log-rank test. Statistical significance was considered at a value of $P < 0.05$.

3. Results

The characteristics of the 298 subjects at the start of study (reference) are given in Table 1. The sample comprises 120 men and 178 women. The mean age of participants at entry was 79.6 years. The mean follow-up time was 1227.2 days, during which 9 subjects died (myocardial infarction: 2 men and 3 women; stroke: two men and two women).

Out of the 298 participants, baPWV was measured in 245 subjects, and a baPWV above 2500 cm/s was observed in 33 subjects, five of whom died (three from a myocardial infarction and two from a stroke). Their reference characteristics are given in Table 2, which shows that an increased baPWV is associated with older age, higher systolic and diastolic BP, increased pulse pressure and shorter FR. Subjects with an increased value of baPWV included fewer event-free survivors. Kaplan–Meier curves for event-free survival revealed a significant difference between the two groups stratified by a baPWV of 2500 cm/s ($P < 0.00005$, log-rank test) (Fig. 1).

Among the variables used in Cox proportional hazard models, PWV as well as age, MMSE, HDSR and the LF/HF




Fig. 1. Kaplan–Meier event probability curves for cardiovascular death.

Table 1
Reference characteristics of subjects

Endpoint	Mean	SD
Gender	0.403	
Age	79.0	4.75
BMI	23.4	3.74
SBP	141.2	19.56
DBP	75.5	10.57
PP	65.8	16.41
HR	70.8	11.46
PWV	2097	504.6
ABI	1.02	0.14
Up&Go	16.5	6.54
FR	25.6	7.77
Button	15.7	7.16
MMSE	24.2	4.43
HDSR	23.6	5.30
Kohs	19.1	11.17
GDS	4.8	2.96
VLF	912.3	743.3
LF	207.2	299.0
HF	102.7	250.3
LF/HF	2.84	1.70
pNN50	4.36	8.87
CVRR	5.04	1.95
RMSSD	22.6	12.5
SDNN	37.4	16.0

Gender: Man = 1, Woman = 0 (120 men and 178 women)

Table 2
Reference Characteristics of Subjects with Acceptable or Increased baPWV

Endpoint	n = 212		n = 33		t-value	p-value
	baPWV <2500		baPWV >2500			
	Mean	S.D.	Mean	S.D.		
Gender	0.396		0.424		0.304	N.S.
Age	78.7	4.39	80.4	4.54	2.045	0,0420
BMI	23.5	3.88	23.0	3.41	-0.722	N.S.
SBP	139.2	18.14	154.2	25.81	4.087	0,0001
DBP	74.3	9.82	79.6	14.33	2.635	0,0090
PP	64.9	15.80	74.7	18.66	3.158	0,0018
HR	70.1	11.24	72.5	12.77	1.090	N.S.
PWV	1949	292.0	3045	555.0	17.298	0,0000
ABI	1.03	0.12	0.99	0.17	-1.704	N.S.
Up&Go	16.1	6.36	17.5	6.77	1.128	N.S.
FR	26.2	7.66	22.5	7.64	-2.533	0,0120
Button	15.2	7.05	16.1	4.53	0.682	N.S.
MMSE	24.3	5.18	23.7	5.33	-0.827	N.S.
HDSR	24.0	5.18	22.6	5.13	-1.444	N.S.
Kohs	19.6	11.08	17.1	10.42	-1.177	N.S.
GDS	4.8	2.85	4.8	3.16	0.125	N.S.
VLF	938.7	788.5	738.6	440.9	-1.187	N.S.
LF	211.1	303.7	133.2	113.7	1.216	N.S.
HF	98.8	235.1	63.8	52.1	-0.711	N.S.
LF/HF	2.88	1.70	2.70	1.51	-0.482	N.S.
pNNS50	4.28	8.24	2.70	3.68	-0.908	N.S.
CVRR	4.98	1.55	4.79	1.34	-0.571	N.S.
rMSSD	22.1	9.7	20.8	7.3	-0.624	N.S.
SDNN	37.7	16.2	33.3	9.8	-1.242	N.S.

Gender: Man = 1, Woman = 0.

Statistical significance of difference in PWV between the 2 groups validates classification.

ratio were statistically significantly associated with the occurrence of cardiovascular death (Table 3). In univariate analyses, a 200 or 500 cm/s increase in baPWV was associated with a RR of cardiovascular death of 1.335 or 2.058, respectively ($P < 0.0002$). A two-point increase in MMSE and HDSR score significantly protected against cardiovascular death, being associated with RRs of 0.776 ($P = 0.0369$) and 0.753 ($P = 0.0029$), respectively. The LF/HF ratio also showed significant predictive value ($P = 0.025$), but other HRV indices did not.

In multivariate analyses, when both baPWV and age were used as continuous variables in the same model, baPWV remained statistically significantly associated with the occurrence of cardiovascular death. After adjustment for age and HDSR, a 200 cm/s increase in baPWV was associated with a 30.2% increase in risk (RR = 1.302, 95% CI: 1.110 to 1.525), and a 500 cm/s increase in baPWV with a 93.3% increase in risk (RR = 1.933, 95% CI: 1.300 to 2.874), $P = 0.0011$. In multivariate analyses, when both the LF/HF ratio and age were used as continuous variables in the same model, the LF/HF ratio was no longer statistically significantly associated with cardiovascular mortality.

Table 3
RR of CV Death in relation to PWV, HRV, Cognitive function and classic CV Risk factors

Endpoint	RR	95% CI	p-value
Gender			N.S.
Age (5)	1.859	1.008–3.427	0.0469
BMI			N.S.
SBP			N.S.
DBP			N.S.
PP			N.S.
HR			N.S.
PWV (200)	1.335	1.147–1.553	0.0002
PWV (500)	2.058	1.410–3.005	0.0002
PWV (200)*	1.302	1.110–1.525	0.0011
PWV (500)*	1.933	1.300–2.874	0.0011
ABI			N.S.
Up&Go			N.S.
FR			0.0696
Button			N.S.
MMSE (2)	0.776	0.612–0.985	0.0369
HDSR (2)	0.753	0.624–0.907	0.0029
Kohs			0.0991
GDS			N.S.
VLF			N.S.
LF			N.S.
HF			N.S.
LF/HF (0.20)	0.821	0.690–0.976	0.0255
pNNS50			N.S.
CVRR			N.S.
RMSSD			N.S.
SDNN			N.S.

* After adjustment for age and HDSR

4. Discussion

The main result of the present study is that in elderly community-dwelling people, arterial stiffness measured by means of baPWV predicted the occurrence of cardiovascular death beyond the prediction provided by age, gender, blood pressure and cognitive functions, assessed by a multivariate Cox model. The baPWV measure is a novel noninvasive technique, which has been developed to assess pulse wave transmission between the brachial and tibial arteries [6]. PWV is known to be an indicator of arterial stiffness and a marker of vascular damage [7]. Traditionally, carotid-femoral PWV is an established method for measuring PWV. Contrary to this traditional PWV, baPWV includes peripheral components of the arterial tree. We need to consider the role of this arterial tree because the influence of age changes in different parts of the arterial tree. Although baPWV values are larger compared to those obtained by the traditional method, their validity has been demonstrated by Yamashina et al. [6]. Findings herein suggest the usefulness of baPWV for clinical use. It has also been argued that baPWV is strongly affected by blood pressure [8] and that this effect should be considered in clinical