

|   |   |                          |                     |      |
|---|---|--------------------------|---------------------|------|
| Fujimoto T, Kemppainen J, Kalliokoski KK.   | Skeletal muscle glucose uptake response to exercise in trained and untrained men.   | Med Sci Sports Exerc.    | 35(5):777-83.       | 2003 |
| Fukuyama H, Ouchi Y, Matsuzaki S, Gannon GA, Rhind SG, Shek PN, Shephard RJ.  | Brain functional activity during gait in normal subjects: a SPECT study.  | Neurosci Lett.           | 228(3):183-6.       | 1997 |
| Garcia-Rubi E, Starling RD, Tchernof A, Matthews DE, Walston JD, Shuldiner AR, Silver K, Poehlman ET and Calles-Garza AA, Ha TG, Garcia C, Chen MJ, Russo-Neustadt AA.  | Differential cell adhesion molecule expression and lymphocyte mobilisation during prolonged aerobic Trp64Arg variant of the beta3-adrenoceptor and insulin resistance in obese postmenopausal women.  | Eur J Appl Physiol       | 84(4):272-82        | 2001 |
| Gates PE, Tanaka H, Graves J, Seals DR  | Exercise, antidepressant treatment, and BDNF mRNA expression in the aging brain.  | Pharmacol Biochem Behav. | 77(2):209-20.       | 2004 |
| Giannopoulou I, Fernhall B, Carhart R, Weinstock RS, Baynard T, Figueroa A, Kanaley JA.   | Left ventricular structure and diastolic function with human ageing. Relation to habitual exercise and arterial Effects of diet and/or exercise on the adipocytokine and inflammatory cytokine levels of postmenopausal women with type 2 diabetes.   | Eur Heart J              | 24:2213-20          | 2003 |
| Gill TM, Williams CS, Tinetti ME.   | Assessing risk for the onset of functional dependence among older adults: the role of physical performance.   | J Am Geriatr Soc         | 43(6):603-9         | 1995 |
| Giresi PG, Stevenson EJ, Theilhaber J, Koncarevic A, Parkington J, Fielding RA, Kandarjian SC.  | Identification of a molecular signature of sarcopenia   | Physiol Genomics         | 21:253-63           | 2005 |
| Goldhammer E, Tanchilevitch A, Maor I, Beniamini Y, Rosenschein U, Sagiv M, Gollnick, P.D., Armstrong, R.B., Saltin, B., Saubert, C.W., Sembrowich, W.L., and R.E. Shephard.  | Exercise training modulates cytokines activity in coronary heart disease patients.  | Int J Cardiol            | 100(1):93-9         | 2005 |
| Green D, Walsh JH, Mariorana A, Burke V, Taylor R, O'Driscoll G   | Effect of training on enzyme activity and fiber composition of human skeletal muscle.   | J. Appl. Physiol.        | 34: 107-111         | 1973 |
| Gregg EW, Cauley JA, Stone K, Thompson TJ, Bauer DC, Cummings SR, and Ensrud KE.  | Comparison of resistance and conduit vessel nitric oxide-mediated vascular function in vivo: effects of exercise  | J Appl Physiol           | 97:749-755          | 2004 |
| Gregg EW, Gerzoff RB, Caspersen CJ, Williamson DF, Narayan KM, Gregg, Edward W; Cauley, Jane A; Stone, Katie; Thompson, Theodore J; Bauer, Douglas C; Cummings, Steven R; Ensrud, Greiwe J.S. Holloszy J.O. and C.F. Semenkovich. | Relationship of changes in physical activity and mortality among older women.   | JAMA                     | 289: 2379-2386      | 2003 |
| Greiwe JS, Cheng B, Rubin DC, Yarasheski KE, Semenkovich CF, Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB.  | Relationship of walking to mortality among US adults with diabetes.   | Arch Intern Med.         | 163(12):1440-7.     | 2003 |
| Hagberg JM, Hickson RC, Ehsani AA, et al.   | Relationship of changes in physical activity and mortality among older women  | JAMA                     | 289(18) : 2379-2386 | 2003 |
| Hagberg JM, McCole SD, Brown MD, Ferrell RE, Wilund KR, Huberty A, Douglass LW, and Moore GE  | Exercise induces lipoprotein lipase and GLUT-4 protein in muscle independent of adrenergic-receptor signaling. Resistance exercise decreases skeletal muscle tumor necrosis factor alpha in frail elderly humans Lower-extremity function in persons over age of 70 years as a predictor of subsequent disability. 地域老人の日常生活動作能力に関する追跡的研究 | J. Appl. Physiol.        | 89: 176-181         | 2000 |
|   | Faster adjustment to and recovery from submaximal exercise in the trained state   | FASEB J                  | 15:475-82           | 2001 |
|   | ACE insertion/deletion polymorphism and submaximal exercise hemodynamics in postmenopausal women.   | N Engl J Med             | 332:556-561         | 1995 |
|   |   | 民族衛生                     | 54:217-233          | 1989 |
|   |   | J Appl Physiol           | 48:218-24           | 1980 |
|   |   | J Appl Physiol           | 92: 1083-1088       | 2002 |

|   |  |  |                |      |
|---|--|--|----------------|------|
| Hajduczuk G, Chapleau MW, Johnson SL, Aboud FM  | Increase in sympathetic activity with age. I. Role of impairment of arterial baroreflexes  | Am J Physiol                                 | 260:H1113-20   | 1991 |
| Hakim AA, Petrovitch H, Burchfiel CM, Ross GW, Rodriguez BL, White LR, Yano K, Curb JD, Abbott RD.                            | Effects of walking on mortality among nonsmoking retired men.  | N Engl J Med                                 | 338:94-99      | 1998 |
| Hambrecht R, Niebauer J, Marburger C, Grunze M, Kalberer B, Hauer K, Schlierf G, Kubler W, and Schuler G.                     | Various intensities of leisure time, physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary  | J Am Coll Cardiol                            | 22: 468-477    | 1993 |
| Hambrecht R, Wolf A, Ghelen S, Linke A, Hofer J, Erbs S, Schoene N, Schuler G.  | Effect of exercise on coronary endothelial function in patients with coronary artery disease.  | N Engl J Med.                                | 342(7):454-60. | 2000 |
| Hamed M, Orrell RW, Cobbold M, Goldspink G, Harridge SD.  | Expression of IGF-1 splice variants in young and old human skeletal muscle after high resistance exercise.   | J Physiol                                    | 547:247-54     | 2003 |
| Hannukainen J, Kalliokoski KK, Nuutila P, Fujimoto T,   | In Vivo Measurements of Glucose Uptake in Human Achilles Tendon During Different Exercise Intensities.   | Int J Sports Med.                            | 26(9):727-731. | 2005 |
| Hara Y, Hagihara A, Ikematsu H, Nobutomo K  | Efficacy of influenza vaccine among elderly patients by physical activity status   | Environmental Health and Preventive Medicine | 7: 183-188     | 2002 |
| Haskell WL, Alderman EL, Fair JM, Maron DJ, Mackey SF, Superko HR, Williams PT, Johnstone IM, Champagne MA, Krauss RM, et al. | Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP) | Circulation                                  | 89: 975-90     | 1994 |
| Hass CJ, Gregor RJ, Waddell DE, Oliver A, Smith DW, Fleming RP, Wolf SL.  | The influence of Tai Chi training on the center of pressure trajectory during gait initiation in older adults.   | Arch Phys Med Rehabil                        | 85: 1593-1598  | 2004 |
| Hassmen P, Ceci R, Backman L.   | Exercise for older women: a training method and its influences on physical and cognitive performance.  | Eur J Appl Physiol Occup Physiol.            | 64(5):460-6.   | 1992 |
| Hauer K, Hildebrandt W, Sehl Y, Edler L, Oster P, Droge W.  | Improvement in muscular performance and decrease in tumor necrosis factor level in old age after antioxidant   | J Mol Med.                                   | 81(2):118-25   | 2003 |
| Hayashi T, Tsumura K, Suematsu C, Okada K, Fujii S, Endo G.   | Walking to work and the risk for hypertension in men: the Osaka Health Survey.   | Ann. Intern. Med.                            | 131(1):21-26   | 1999 |
| Hayashi T, Hirshman M.F., Kurth E.J., Winder W.W., and L.J. Goodyear.   | Evidence for 5AMP-activated protein kinase mediation of the effect of muscle contraction on glucose transport.   | Diabetes                                     | 47: 1369-1373  | 1998 |
| Hayes SC, Rowbottom D, Davies PS, Parker TW, Bashford J.  | Immunological changes after cancer treatment and participation in an exercise program  | Med Sci Sports Exerc.                        | 35(1):2-9.     | 2003 |
| Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr.  | Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus.   | N Engl J Med.                                | 325(3):147-52. | 1991 |
| Hess JA, Woollacott M   | Effect of high intensity strength training on functional measures of balance ability in balance-impaired older   | J Manipulative Physiol Ther                  | 28: 582-590    | 2005 |
| Higaki J, Baba S, Katsuya T, Sato N, Ishikawa K, Mannami T, Ogata J, Ogiwara T.   | Deletion allele of angiotensin-converting enzyme gene increases risk of essential hypertension in Japanese men: the Suita Study.   | Circulation.                                 | 101(17):2060-5 | 2000 |
| 樋口博之  | 加速度センサーを内蔵した歩数計による若年者と高齢者も日常身体活動量の比較   | 体力科学   | 52:111-118     | 2003 |
| HIRSCHHORN K, HIRSCHLAND BP, KRAUS H.   | Hypokinetic disease; role of inactivity in production of disease.  | Br J Phys Med.                               | 19(8):180-5    | 1956 |
| Holloszy, J.O.  | Biochemical adaptations in muscle. Effects of exercise on mitochondrial O2 uptake and respiratory enzyme activity in skeletal muscle.  | J. Biol. Chem.                               | 242: 2278-2282 | 1967 |

|  |  |                                 |                                       |              |
|--|--|---------------------------------|---------------------------------------|--------------|
| Holloszy, J.O. and Narahara, H.T.  | Studies of tissue permeability. X. Changes in permeability to 3-methylglucose associated with contraction of isolated frog muscles.  | J. Biol. Chem                   | 240: 3493-3500                        | 1965         |
| Hoppeler H, Howald H, Conley K, Lindstedt SL, Claassen H, Vock P, and Houmard, J. A., Shinebarger, M. H., Dolan, P. L., Leggett-Frazier, N., Bruner, R. K., McCammon, M. R., Israel, R. G., Houmard, J.A., Weidner, M.L., Gavigan, K.E., Thndall, G.L., Hickey, M.S. and Aymen, A. | Endurance training in humans: aerobic capacity and structure of skeletal muscle.<br>Exercise training increases GLUT-4 protein concentration in previously sedentary middle-aged men.  | J Appl Physiol<br>Am J Physiol  | 59: 320-327<br>264 (6 Pt 1), E896-901 | 1985<br>1993 |
| Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, Willett WC, Speizer FE, Manson JE   | Fiber type and citrate synthase activity in the human gastrocnemius and vastus lateralis with aging  | J. Appl. Physiol.               | 85(4):1337-41.                        | 1998         |
| Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, Willett WC, Speizer FE, Manson JE   | Diet, lifestyle, and the risk of type 2 diabetes mellitus in women.  | N Engl J Med.                   | 345(11):790-7.                        | 2001         |
| Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, Willett WC, Speizer FE, Manson JE  | Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study.  | JAMA.                           | 282(15):1433-9.                       | 1999         |
| Hu FB, Stampfer MJ, Solomon C, Liu S, Colditz GA, Speizer FE, Willett WC, Manson JE.   | Walking compared with vigorous physical activity and risk of type 2 diabetes in women: A prospective study.  | JAMA                            | 282:1433-1439                         | 1999         |
| Huang G, Gibson CA, Tran ZV, and Osness WH.  | Physical activity and risk for cardiovascular events in diabetic women.  | Ann Intern Med.                 | 134(2):96-105.                        | 2001         |
| Hue OA, Seynnes O, Ledrole D, Colson SS, Bernard PL  | Controlled endurance exercise training and VO2max changes in older adults: a meta-analysis.  | Prev Cardiol                    | 8: 217-225                            | 2005         |
| Huffman DM, Altene TS, Mawhinney TP, Thomas TR.  | Effects of a physical activity program on postural stability in older people   | Aging Clin Exp Res              | 16: 356-362                           | 2004         |
| Hurley BF, Hagberg JM, Goldberg AP, Seals DR, Ehsani AA, Brennan RE, Hurley BF, Seals DR, Ehsani AA, Cartier L.J, Dalsky GP, Hagberg JM, Holloszy JO   | Effect of n-3 fatty acids on free tryptophan and exercise fatigue.   | Eur J Appl Physiol              | 92(4-5):584-91.                       | 2004         |
| Iemitsu M, Miyauchi T, Maeda S, Tanabe T, Takanashi M, Irukayama-Tomobe Y, Sakai S, Ohmori H, Matsuda M.   | Resistive training can reduce coronary risk factors without altering O2max or percent body fat.  | Med Sci Sports Exerc            | 20:150-154                            | 1988         |
| Iemitsu M, Miyauchi T, Maeda S, Tanabe T, Takanashi M, Matsuda M, Yamaguchi I, Imai K, Sato H, Hori M, Kusuoka H, Ozaki H, Yokoyama H, Takeda H, Inoue M, Kamada T.  | Effects of high-intensity strength training on cardiovascular function.  | Med Sci Sports Exerc            | 16:488-488                            | 1984         |
| Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, Yukawa M, Aiello E, Potter JD, McTiernan A.  | Aging-induced decrease in the PPAR-alpha level in hearts is improved by exercise training  | Am J Physiol Heart Circ Physiol | 283:H1696-H1705                       | 2002         |
| Jackson AS, Wier LT, and Ross RM.  | Exercise training improves cardiac function-related gene levels through thyroid hormone receptor signaling in aged vagally mediated heart rate recovery after exercise is accelerated in athletes but blunted in patients with chronic heart failure | Am J Physiol Heart Circ Physiol | 286:H1750-H1760                       | 2004         |
| Jakicic JM, Marcus BH, Gallagher KI, Napolitano M, Lang W.   | Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomized controlled trial.   | J Am Coll Cardiol               | 24:1529-35                            | 1994         |
| Jackson AS, Wier LT, and Ross RM.  | The validity of non-exercise cardiorespiratory fitness prediction models.  | Med Sci Sports Exerc            | 27: 1096-1098                         | 1995         |
| Jakicic JM, Marcus BH, Gallagher KI, Napolitano M, Lang W.   | Effect of exercise duration and intensity on weight loss in overweight, sedentary women: a randomized trial.   | JAMA.                           | 289(3):323-30.                        | 2003         |
| Jackson AS, Wier LT, and Ross RM.  | The validity of non-exercise cardiorespiratory fitness prediction models.  | Med Sci Sports Exerc            | 27: 1096-1098                         | 1995         |
| Jakicic JM, Marcus BH, Gallagher KI, Napolitano M, Lang W.   | Effect of exercise duration and intensity on weight loss in overweight, sedentary women: a randomized trial.   | JAMA.                           | 290(10):1323-30.                      | 2003         |

|  |   |   |                       |      |
|--|---|---|-----------------------|------|
| Jakicic JM, Winters C, Lang W, Wing RR.  | Effects of intermittent exercise and use of home exercise equipment on adherence, weight loss, and fitness in overweight women: a randomized trial.                   | JAMA.   | 282(16):1554-60.      | 1999 |
| Janssen I, Baumgartner RN, Ross R, Rosenber IH, and Roubenoff R.   | Skeletal muscle cutpoints associated with elevated physical disability risk in older men and women.   | Am J Epidemiol  | 159: 413-421          | 2004 |
| Janssen I, Heymsfield SB, Wang ZM, and Ross R.   | Skeletal muscle mass and distribution in 468 men and women aged 18-88 yr.   | J Appl Physiol  | 89: 81-88             | 2000 |
| Jette AM, Lachman M, Giorgetti MM, Asmann SF, Harris BA, Levenson C, Wernick M, Krebs D.   | Exercise--it's never too late: the strong-for-life program.   | Am J Public Health.   | 89(1): 66-72          | 1999 |
| Jonason, T., Henrikssen, E., Kangro, T., Nilsson, H, Vessby, B. and Ringqvist, I.  | Stiffness of the common carotid artery in healthy 50-years-old subjects   | Clinical Physiology   | 17(6):569-77.         | 1997 |
| Judge JO, Lindsey C, Underwood M, Winsemius D  | Balance improvements in older women: effects of exercise training.  | Phys Ther   | 73: 254-262           | 1993 |
| Judge JO, Underwood M, Gennosa T.  | Exercise to improve gait velocity in older persons.   | Arch Phys Med Rehabil   | 74:400-               | 1993 |
| Kagaya, A. and Homma, S.   | Brachial arterial blood flow during handgrip in young and elderly women.  | J. Exerc. Sci.  | 17: 18-26             | 1997 |
| Kagaya, A., Muraoka, Y., Matsuda, M., Kuno, S., Shimizu, S., Yamamoto, Y., Kera, N. and Kimura, Y.   | Forearm work capacity in association with age and daily physical activity in elderly women.   | In A. Gunji (ed), Inactivity, Health and Aging. Seigakuin University Press. | 61-66                 | 2001 |
| Kahn, S. E., Larson, V. G., Beard, J. C., Cain, K. C., Fellingham, G. W., Schwartz, R. S., Veith, R. C., Stratton, J. R., Cerqueira M D, Abrass, I. B. | Effect of exercise on insulin action, glucose tolerance, and insulin secretion in aging.  | Am J Physiol  | 258 (6 Pt 1), E937-43 | 1990 |
| Kakiyama, T., Matsuda, M. and Koseki, S.   | Effect of physical on the distensibility of the aortic wall in healthy males  | Angiology   | 49(9): 749-57         | 1998 |
| Kalliokoski KK, Knuuti J, Nuutila P.   | Relationship between muscle blood flow and oxygen uptake during exercise in endurance-trained and   | J Appl Physiol.   | 98(1):380-3.          | 2005 |
| Kalliokoski KK, Laaksonen MS, Takala TO, Knuuti J, Nuutila P.  | Muscle oxygen extraction and perfusion heterogeneity during continuous and intermittent static exercise.  | J Appl Physiol.   | 94(3):953-8.          | 2003 |
| Kalliokoski KK, Langberg H, Ryberg AK, Scheede-Bergdahl C,   | The effect of dynamic knee-extension exercise on patellar tendon and quadriceps femoris muscle glucose uptake in humans studied by positron emission tomography.      | J Appl Physiol.   | 99(3):1189-92.        | 2005 |
| Kalliokoski KK, Oikonen V, Takala TO, Sipila H, Knuuti J, Nuutila P.   | Enhanced oxygen extraction and reduced flow heterogeneity in exercising muscle in endurance-trained men.  | Am J Physiol Endocrinol Metab.  | 280(6):E1015-21.      | 2001 |
| Kannel WB, Sorlie P.   | Some health benefits of physical activity. The Framingham Study   | Arch. Intern. Med.  | 139(8):857-861        | 1979 |
| Kaplan GA, Strawbridge WJ, Cohen RD, Hungerford LR.  | Natural history of leisure-time physical activity and its correlates: associations with mortality from all causes and cardiovascular disease over 28 years            | Am. J. Epidemiol.   | 144(8):793-797        | 1996 |
| Katz A. Broberg S, Sahlin K. and J. Wahren.  | Leg glucose uptake during maximal dynamic exercise in humans.   | Am. J. Physiol.   | 251: E65-E70          | 1986 |
| Katzel LI, Bleecker ER, Colman EG, Rogus EM, Sorkin JD, Goldberg AP.   | Effects of weight loss vs aerobic exercise training on risk factors for coronary disease in healthy, obese, middle-aged and older men. A randomized controlled trial. | JAMA.   | 274(24):1915-21.      | 1995 |

|  |   |                                |                   |      |
|--|---|--------------------------------|-------------------|------|
| Katzmarzyk PT, Church TS, Janssen I, Ross R, and Blair SN  | Metabolic syndrome, obesity, and mortality: impact of cardiorespiratory fitness.  | Diabetes Care                  | 28: 391-397       | 2005 |
| Kawakami Y, Akima H, Kubo K, Muraoka Y, Hasegawa H, Kouzaki M, Imai M, Suzuki Y, Gunji A, Kanehisa H, Fukunaga K, Tanaka K, Tabata I, Tanaka A, Higuchi M. | Changes in muscle size, architecture, and neural activation after 20 days of bed rest with and without resistance exercise.<br>Effects of high intensity intermittent swimming on glucose transport in rat epitrochlearis muscle. | Eur J Appl Physiol             | 84: 7-12          | 2001 |
| Keen DA, Yue GH, Enoka RM  | Training-related enhancement in the control of motor output in elderly humans   | J Appl Physiol                 | 77: 2648-2658     | 1994 |
| Kelley G, McClellan P.   | Antihypertensive effects of aerobic exercise. A brief meta-analytic review of randomized controlled trials.   | Am. J. Hypertens.              | 7(2):115-119      | 1994 |
| Kelley G, Tran ZV.   | Aerobic exercise and normotensive adults: a meta-analysis   | Med. Sci. Sports Exer.         | 27(10):1371-1377  | 1995 |
| Kemppainen J, Aalto S, Fujimoto T,   | High intensity exercise decreases global brain glucose uptake in humans.  | J Physiol                      | 568(Pt 1):323-32. | 2005 |
| Kemppainen J, Fujimoto T, Kalliokoski KK.  | Myocardial and skeletal muscle glucose uptake during exercise in humans.  | J Physiol.                     | 542(Pt 2):403-12. | 2002 |
| Kent-Braun JA, Ng AV, and Young K.   | Skeletal muscle contractile and noncontractile components in young and older women and men.   | J Appl Physiol                 | 88: 662-668       | 2000 |
| Kim J, Heshka S, Gallagher D, Kotler DP, Mayer L, Albu J, Shen W, Freda PU, and Hevmsfield SB.   | Intermuscular adipose tissue-free skeletal muscle mass: estimation by dual-energy X-ray absorptiometry in adults.   | J Appl Physiol                 | 97: 655-660       | 2004 |
| King AC, Pruitt LA, Phillips W, Oka R, Rodenburg A, Haskell WL.  | Comparative effects of two physical activity programs on measured and perceived physical functioning and other health-related quality of life outcomes in older adults.   | J Gerontol A Biol Sci Med Sci. | 55(2): M74-83     | 2000 |
| King MC, Marks JH, and Mandell JB.   | Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2.  | Science                        | 302: 643-646      | 2003 |
| Kingwell BA, Sherrard B, Jennings GL, Dart AM.   | Four weeks of cycle training increases basal production of nitric oxide from the forearm  | Am J Physiol                   | 272:H1070-7       | 1997 |
| Kirwan JP, Kohrt WM, Wojta DJ, Bourey RE, Holloszy JO  | Endurance exercise training reduces glucose-stimulated insulin levels in 60- to 70-yr-old men and women.  | J. Gerontol.                   | 48:M84-M90        | 1993 |
| Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM; Diabetes Prevention Program Research Group                            | Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin.   | N Engl J Med.                  | 346(6):393-403.   | 2002 |
| Kohrt WM, Kirwan JP, Staten MA, Bourey RE, King DS, Holloszy JO  | Insulin resistance in aging is related to abdominal obesity.  | Diabetes                       | 42:273-281        | 1993 |
| Kohrt WM, Malley MT, Coggan AR, Spina RJ, Ogawa T, Ehsani AA, Bourey RE, Martin WH, III, Holloszy JO   | Effects of gender, age, and fitness level on the response of O <sub>2</sub> max to training in 60 to 71 yr-olds.  | J Appl Physiol                 | 71:2004-2011      | 1991 |
| Kohrt WM, Malley MT, Dalsky GP, Holloszy JO  | Body composition of healthy sedentary and trained, young and older men and women.   | Med Sci Sports Exer            | 24:832-837        | 1992 |
| Kohut ML, Lee W, Martin A, Arnston B,  | The exercise-induced enhancement of influenza immunity is mediated in part by improvements in psychosocial factors in older adults.   | Brain Behav Immun              | 19(4):357-66      | 2005 |
| Russell DW, Ekkkekakis P, Yoon KJ, Bishon A, Cunnick JE  | Physical training and insulin sensitivity.  | Diabetes Metab Rev             | 1(4), 445-81      | 1986 |
| Koivisto, V. A., Yki-Jarvinen, H., DeFronzo, R. A.   |   |                                |                   |      |

|  |  |  |                    |           |
|--|--|--|--------------------|-----------|
| Kornatz KW, Christou EA, Enoka RM  | Practice reduces motor unit discharge variability in a hand muscle and improves manual dexterity in old adults   | J Appl Physiol                         | 98: 2072-2080      | 2005      |
| Kostek MC, DeImonico MJ, Reichel JB, Roth SM, Douglass L, Ferrell RE, Hurley Kramer AF, Hahn S, Cohen NJ, Banich MT, McAuley E, Harrison CR, Chason J, Vakil E, Bardell L, Boileau RA, Colcombe Kristiansen S, Gade J, Wojtaszewski JF, Kiens B, Richter EA.   | Muscle strength response to strength training is influenced by insulin-like growth factor 1 genotype in<br>Ageing, fitness and neurocognitive function.<br>Glucose uptake is increased in trained vs. untrained muscle during heavy exercise.  | J Appl Physiol                         | 98:2147-54         | 2005      |
| Kritchevsky SB, Nicklas BJ, Visser M, Simonsick EM, Newman AB, Harris TB, Lange EM, Penninx BW, Goodpaster BH, Satterfield S, Colbert I.H., Rubin SM Kubo K, Akima H, Ushiyama J, Tabata I, Fukunaga H, Kanehisa H, Fukunaga T 神崎素樹, 政二 慶, 宮谷昌枝, 村岡哲郎, 白澤葉月, 久野譜也, 金久博昭, 福永哲夫  | Effects of resistance training during bed rest on the viscoelastic properties of tendon structures in the lower limb<br>Angiotensin-converting enzyme insertion/deletion genotype, exercise, and physical decline.   | Nature                                 | 29:400(6743):418-9 | 1999      |
| Laaksonen DE, Lindstrom J, Lakka TA, Eriksson JG, Niskanen L, Wikstrom K, Aunola S, Keinanen-Kiukkaanniemi S, Laakso M, Valle TT, Ilanne-Parikka P, Louheranta A, Hamalainen H, Rastas M, Salminen V, Cepaitis Z, Hakumaki M, Kaikkonen H, Harkonen P, Sundvall J, Tuomilehto J, Uusitupa M: Finnish Laidlaw DH, Kornatz KW, Keen DA, Suzuki S, Enoka RM | Physical activity in the prevention of type 2 diabetes: the Finnish diabetes prevention study.<br>Strength training improves the steadiness of slow lengthening contractions performed by old adults<br>Effect of computerized feedback postural training on posture and attentional demands in older adults | J Appl Physiol                         | 87: 1786-1795      | 1999      |
| Lajoie Y   | Cardiorespiratory fitness and the progression of carotid atherosclerosis in middle-aged men.   | Aging Clin Exp Res                     | 16: 363-368        | 2004      |
| Lakka TA, Laukkanen JA, Rauramaa R, Salonen R, Lakka HM, Kaplan GA, Salonen JT   | The spectrum of physical activity, cardiovascular disease and health: an epidemiologic perspective.  | Ann Intern Med.                        | 134(1):12-20.      | 2001      |
| LaPorte RE, Adams LL, Savage DD, Brenes G, Dearwater S, Cook T, Laukkanen JA, Lakka TA, Rauramaa R, Kuhanen R, Venalainen JM, Salonen R, Salonen JT  | Cardiovascular fitness as a predictor of mortality in men.<br>Physical activity and risk of cognitive impairment and dementia in elderly persons.  | Am J. Epidemiol.                       | 120(4):507-517     | 1984      |
| Laurin D, Verreault R, Lindsay J, MacPherson K, Rockwood K, Lawrenson L, Poole JG, Kim J, Brown C, Patel P and Richardson RS   | Vascular and metabolic response to isolated small muscle mass exercise: effect of age.   | Arch Neurol.                           | 58(3):498-504.     | 2001      |
| Lazarov O, Robinson J, Tang YP, Hairston IS, Korade-Mirnics Z, Lee VM, Hersh LB, Sapolsky RM, Mirnics K, Sisodia SS, Ledin T, Kronhed AC, Moller C, Moller M, Odkvist LM, Olsson B.  | Environmental enrichment reduces Abeta levels and amyloid deposition in transgenic mice.<br>Effects of balance training in elderly evaluated by clinical tests and dynamic posturography.  | Am. J. Physiol. Heart and Circulation. | 285(3):H1023-31.   | 2003      |
|  |  | Cell                                   | 120(5):701-13.     | 2005      |
|  |  | J. Vestib. Res.                        | 1: 129-138         | 1990-1991 |

|   |  |                             |                         |      |
|---|--|-----------------------------|-------------------------|------|
| Lee IM, Hsieh CC, Paffenbarger RS Jr.                               | Exercise intensity and longevity in men. The Harvard Alumni Health Study.  | JAMA.                       | 273(15):1179-84.        | 1995 |
| Lee IM, Manson JE, Hennekens CH, Paffenbarger RS Jr.                | Body weight and mortality. A 27-year follow-up of middle-aged men.   | JAMA.                       | 270(23):2823-8.         | 1993 |
| Lee IM, Paffenbarger RS Jr.   | Associations of light, moderate, and vigorous intensity physical activity with longevity. The Harvard Alumni   | Am J Epidemiol.             | 151(3):293-9.           | 2000 |
| Lee IM, Paffenbarger RS Jr.   | Change in body weight and longevity.   | JAMA.                       | 268(15):2045-9.         | 1992 |
| Lee IM, Paffenbarger RS Jr.   | Physical activity and stroke incidence: the Harvard Alumni Health Study.   | Stroke.                     | 29(10):2049-54.         | 1998 |
| Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr.                      | Relative intensity of physical activity and risk of coronary heart disease.  | Circulation.                | 107(8):1110-6.          | 2003 |
| Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr.                      | The "weekend warrior" and risk of mortality.   | Am J Epidemiol.             | 160(7):636-41.          | 2004 |
| Lee IM, Sesso HD, Paffenbarger RS Jr.                               | Physical activity and coronary heart disease risk in men: does the duration of exercise episodes predict risk?   | Circulation.                | 102(9):981-6.           | 2000 |
| Lee J, Lee S, Shin S, and Kang HS.                                  | Association between the GNB3 polymorphism and blood pressure in young Korean men.  | Med Sci Sports Exerc        | 37: 1138-1143           | 2005 |
| Lee S, Kuk JL, Katzmarzyk PT, Blair SN, Church TS, and Ross R.      | Cardiorespiratory fitness attenuates metabolic risk independent of abdominal subcutaneous and visceral fat   | Diabetes Care               | 28: 895-901             | 2005 |
| Lee, I-Min  | Physical activity in women: how much is good enough?   | JAMA                        | 290(10): 1377-1379      | 2003 |
| Lee, I-Min; Sesso, Howard D; Oguma, Yuko; Paffenbarger. Ralph S. Jr | Relative intensity of physical activity and risk of coronary heart disease   | Circulation                 | 107(8): 1110-1116       | 2003 |
| Levine JA, Eberhardt NL, and Jensen MD.                             | Role of nonexercise activity thermogenesis in resistance to fat gain in humans.  | Science                     | 283: 212-214            | 1999 |
| Levy RL, Linde JA, Feld KA, Crowell MD, Jeffery RW.                 | The association of gastrointestinal symptoms with weight, diet, and exercise in weight-loss program participants.  | Clin Gastroenterol Hepatol. | 3(10):992-6.            | 2005 |
| Levy WC, Cerqueira MD, Abrass IB, Schwartz RS, Stratton JR          | Endurance exercise training augments diastolic filling at rest and during exercise in healthy young and older men. What is the cause of the ageing atrophy? Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. What is the cause of the ageing atrophy? Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. | Circulation                 | 88:116-126              | 1993 |
| Lexell J, Taylor CC, Sjostrom M:                                    | Factors associated with physical activity among older people--a population-based study   | J. Neurol. Sci.             | 84:275-294.             | 1988 |
| Lexell J, Taylor CC, and Sjostrom M.                                | Resistance and agility training reduce fall risk in women aged 75 to 85 with low bone mass: a 6-month randomized, controlled trial.  | J Neurol Sci                | 84: 275-294             | 1988 |
| Lim, Kim; Taylor, Lee   | Balance confidence improves with resistance or agility training. Increase is not correlated with objective changes in fall risk and physical abilities.  | Preventive Medicine         | 40(1): 33-40            | 2005 |
| Liu-Ambrose T, Khan KM, Eng JJ, Janssen PA, Lord SR, McKay HA       | Insulin-like growth factor I is required for vessel remodeling in the adult brain.   | J Am Geriatr Soc            | 52: 657-665             | 2004 |
| Liu-Ambrose T, Khan KM, Eng JJ, Lord SR, McKay HA.                  | The effect of a 12-month exercise trial on balance, strength, and falls in older women: a randomized   | Gerontology                 | 6: 373-382              | 2004 |
| Lopez-Lopez C, LeRoith D, Torres-Aleman L.                          | The effect of a 12-month exercise trial on balance, strength, and falls in older women: a randomized   | Proc Natl Acad Sci U S A.   | 6):9833-8. Epub 2004 Jh | 2004 |
| Lord SR, Ward JA, Williams P, Strudwick M.                          |  | J Am Geriatr Soc.           | 43(11): 1198-206        | 1995 |



|  |  |                                  |                |      |
|--|--|----------------------------------|----------------|------|
| Maeda S, Tanabe T, Miyauchi T, Otsuki T, Sugawara J, Iemitsu M, Kuno S, Ajisaka R, Yamaguchi I, Matsuda M, Magnuson G, Kajiser L, Isberg B, and Saltin B.    | Aerobic exercise training reduces plasma endothelin-1 concentration in older women   | J Appl Physiol                   | 95:336-41      | 2003 |
| Manson JE, Hu FB, Rich-Edwards JW, Colditz GA, Stampfer MJ, Willett WC, Sneizer FE, Hennekens CH.  | Cardiovascular responses during one and two-legged exercise in middle-aged men. A prospective study of walking as compared with vigorous exercise in the prevention of coronary heart disease in women.                | Acta Physiol Scand.              | 150(4):353-62. | 1994 |
| Manson JE, Nathan DM, Krolewski AS, Stampfer MJ, Willett WC, Hennekens CH.   | A prospective study of exercise and incidence of diabetes among US male physicians   | N Engl J Med.                    | 341(9):650-8.  | 1999 |
| Manson JE, Nathan DM, Krolewski AS, Stampfer MJ, Willett WC, Hennekens CH.   | A prospective study of exercise and incidence of diabetes among US male physicians.  | JAMA                             | 268(1):63-67   | 1992 |
| Margaria R, Aghemo P, Rovelli E.   | Indirect determination of maximal O2 consumption in  | JAMA.                            | 268(1):63-7.   | 1992 |
| Mariorana A, O'Driscoll G, Cheetham C, Demo L, Stanton K, Goodman C, Taylor R, Green D.  | The effect of combined aerobic and resistance training on vascular function in type 2 diabetes   | J Appl Physiol.                  | 20(5):1070-3   | 1965 |
| Markel, T.A., Daley III, J.C., Hogeman, C.S. Herr, M.D., Khan, M.H., Gray, K.S., Kunselman & A.R. and Sinoway L.I.   | Ageing and exercise pressor reflex in humans.  | J Am Coll Cardiol                | 38:860-866     | 2001 |
| Matsuda M, Nosaka T, Sato M, Ohshima N.  | Effects of physical exercise on the elasticity and elastic components of the rat aorta   | Circulation                      | 107(5):675-8.  | 2003 |
| Mayer-Davis EJ, D'Agostino R Jr, Karter AJ, Haffner SM, Rewers MJ, Saad M, Bergman RN.   | Intensity and amount of physical activity in relation to insulin sensitivity: the Insulin Resistance Atherosclerosis Study.  | Eur J Appl Physiol Occup Physiol | 66:122-6       | 1993 |
| Mazzeo RS, Rajkumar C, Rolland J, Blaher B, Jennings G, Esler M.   | Immune response to a single bout of exercise in young and elderly subjects   | JAMA.                            | 279(9):669-74. | 1998 |
| McAuley E, Katula J, Mihalko SL, Bliesner B, Duncan TE, Pena M, Dunn E, McGuire DK, Levine BD, Williamson JW, Snell PG, Blomqvist CG, Saltin B, Mitchell JH. | Mode of physical activity and self-efficacy in older adults: a latent growth curve analysis. A 30-year follow-up of the Dallas Bedrest and Training Study: I. Effect of age on the cardiovascular response to exercise | Mech Ageing Dev.                 | 100(2):121-32  | 1998 |
| McMurdo ME, Rennie L.  | A controlled trial of exercise by residents of old people's  | J Gerontol B Psychol Sci Soc.    | 54(5): P283-92 | 1999 |
| McPherron AC et al   | Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member.   | Circulation                      | 104:1350-7     | 2001 |
| Meijer EP, Goris AH, Wouters L & Westertep KR  | Physical inactivity as a determinant of the physical activity level in the elderly.  | Age Ageing                       | 22(1): 11-5    | 1993 |
| Messier SP, Royer TD, Craven TE, O'Toole ML, Burns R, Ettinger WH Jr.  | Long term exercise and its effect on balance in older, osteoarthritic adults: results from the Fitness, Arthritis, and Seniors Trial (FAST).   | Nature                           | 387:83-90      | 1997 |
| Mettter EJ, Talbot LA, Schragger M, Conwit RA.   | Arm-cranking muscle power and arm isometric muscle strength are independent predictors of all-cause mortality in men.  | Int J Obes Relat Metab Disord.   | 25(7): 935-9   | 2001 |
| Miyachi M, Donato AJ, Yamamoto K, Takahashi K, Gates PE, Moreau KL, Tanaka H.  | Greater age-related reductions in central arterial compliance in resistance-trained men  | J Am Geriatr Soc                 | 48: 131-138    | 2000 |
|  |  | J Appl Physiol                   | 96:814-821     | 2004 |
|  |  | Hypertension                     | 41:130-5       | 2003 |



|   |  |   |   |                                      |
|---|--|---|---|--------------------------------------|
| Mizuno M, Kimura Y, Iwakawa T, Oda K, Ishii K, Ishiwata K, Nakamura Y, Muraoka I.   | Regional differences in blood flow and oxygen consumption in resting muscle and their relationship during recovery from exhaustive exercise.   | J Appl Physiol.   | 95(6):2204-10.  | 2003                                 |
| Mizuno M, Kimura Y, Tokizawa K, Ishii K, Mohiaddin RH, Underwood SR, Bogren HG, Firmin DN, Klipstein RH, Rees RS, Longmore DB, Mokdad AH, Bowman BA, Ford ES, Vinicor F, Marks JS, Koplan JP, Monahan KD, Dinunno FA, Seals DR, Clevenger CM, Desouza CA, Tanaka H, Moreau KL, Donato AJ, Seals DR, DeSouza CA, Tanaka H.               | Greater adenosine A(2A) receptor densities in cardiac and skeletal muscle in endurance-trained men: a [(11)C]TMSX PET study.<br>Regional aortic compliance studied by magnetic resonance imaging: the effects of age, training, and coronary artery disease<br>The continuing epidemics of obesity and diabetes in the United States.<br>Age-associated changes in cardiovascular baroreflex sensitivity are related to central arterial compliance<br>Regular exercise, hormone replacement therapy and the age-related decline in carotid arterial compliance in healthy women | Nucl Med Biol.<br>Br Heart J<br>JAMA<br>Am J Physiol Heart Circ Physiol<br>Cardiovasc Res | 32(8):831-6.<br>62:90-96<br>286:1195-1200<br>281:H284-9                   | 2005<br>1989<br>2001<br>2001<br>2003 |
| Morey MC, Pieper CF, Cornoni-Huntley J.   | Is there a threshold between peak oxygen uptake and self-reported physical functioning in older adults?  | Med Sci Sports Exerc  | 30(8):1223-   | 1998                                 |
| Morris RT, Spangenburg EE, Booth FW, Musaro A, McCullagh K, Paul A, Houghton L, Dobrowolny G, Molinaro M, Barton ER, Sweeney HL, Rosenthal N, Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH, Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE, Nagamatsu T, Oida Y, Kitabatake Y, Kohno H, Egawa K, Nezu N, Arao T. | Responsiveness of cell signaling pathways during the failed 15-day regrowth of aged skeletal muscle.<br>Localized Igf-1 transgene expression sustains hypertrophy and regeneration in senescent skeletal muscle.<br>Long-term morbidity and mortality of overweight adolescents. A follow-up of the Harvard Growth Study of exercise capacity and mortality among men referred for exercise testing.<br>A 6-year cohort study on relationship between functional fitness and impairment of ADL in community-dwelling older persons.  | J Appl Physiol<br>Nat Genet<br>New Engl. J. Med.<br>New Engl. J. Med.<br>J Epidemiol.     | 96:398-404<br>27:195-200<br>327:1350-1355<br>346:793-801<br>13(3):142-148 | 2004<br>2001<br>1992<br>2002<br>2003 |
| Nesher, R., Karl, I.E., and Kipnis, D.M.  | Disassociation of effects of insulin and contraction on glucose transport in rat epitrochlearis muscle.  | Am.J.Physiol.   | 249: C226-C232  | 1985                                 |
| Newson, Rachel S; Kemps, Eva B  | General lifestyle activities as a predictor of current cognition and cognitive change in older adults: a cross-sectional and longitudinal examination  | J Gerontol B Psychol Sci Soc Sci.   | 60(3): 113-120  | 2005                                 |
| Nichols JF, Omizo DK, Peterson KK, Nelson KP.   | Efficacy of heavy-resistance training for active women over sixty: muscular strength, body composition, and program adherence.   | J Am Geriatr Soc.   | 41(3): 205-10   | 1993                                 |
| Nieman DC, Henson DA, Gusewitch G, Warren BJ, Dotson RC, Butterworth DE, Nehlsen-Cannarella SL, Nishida, Y., Tokuyama, K., Nagasaka, S., Higaki, Y., Shirai, Y., Kiyonaga, A., Shindo, M., Kusaka, I., Nakamura, T., Ishibashi S, Tanaka H, Nishime EO, Cole CR, Blackstone EH, Pashkow FJ, Lauer MS.                                   | Physical activity and immune function in elderly women.<br>Effect of moderate exercise training on peripheral glucose effectiveness, insulin sensitivity, and endogenous glucose production in healthy humans estimated by a two-compartment-labeled minimal model<br>Heart rate recovery and treadmill exercise score as predictors of mortality in patients referred for exercise  | Med Sci Sports Exerc.<br>Diabetes<br>JAMA.  | 25(7):823-31<br>53, 315-20<br>284(11):1392-8.                             | 1993<br>2004<br>2000                 |

|  |  |                      |                   |      |
|--|--|----------------------|-------------------|------|
| Nuutila P, Peltoniemi P, Oikonen V, Larmola K,   | increases exercise-stimulated glucose uptake in skeletal muscle in humans: studies using [15O]O <sub>2</sub> , [15O]H <sub>2</sub> O, [18F]fluoro-deoxy-glucose, and positron emission tomography. | Diabetes.            | 49(7):1084-91.    | 2000 |
| Ogawa K, Oka J, Yamakawa J, Higuchi M.   | Habitual exercise did not affect the balance of type 1 and type 2 cytokines in elderly people.   | Mech Ageing Dev.     | 124(8-9):951-6    | 2003 |
| Ogawa T, Spina RJ, Martin WH, III, Kohrt WM, Schechtman KB, Holloszy JO, Ehsani AA   | Effects of aging, sex and physical training on cardiovascular responses to exercise.   | Circulation          | 86:494-503        | 1992 |
| Ogawa小川裕, 岩崎清, 安村誠司,   | 地域高齢者の健康度評価に関する追跡的研究-日常生活動作能力の低下と死亡の予知を中心に-  | 日本公衆衛生雑誌             | 40:859-871        | 1993 |
| Oguma Y, Sesso HD, Paffenbarger RS Jr, Lee IM.   | Weight change and risk of developing type 2 diabetes.  | Obes Res.            | 13(5):945-51.     | 2005 |
| Oida Y, Kitabatake Y, Nishijima Y, Nagamatsu T, Kohno H, Egawa K, Arao T.  | programme on mortality and ADL impairment in the elderly.  | Age and Ageing       | 32(6):585-592.    | 2003 |
| Okura T, Nakata Y, Tanaka K.   | Effects of exercise intensity on physical fitness and risk factors for coronary heart disease.   | Obes Res.            | 11(9): 1131-9     | 2003 |
| Orchard TJ, Temprosa M, Goldberg R, Haffner S, Ratner R, Marcovina S, Fowler S; Diabetes Prevention Program Research Group | The effect of metformin and intensive lifestyle intervention on the metabolic syndrome: the Diabetes Prevention Program randomized trial.  | Ann Intern Med.      | 19:142(8):611-9.  | 2005 |
| Owino V, Yang SY, Goldspink G.   | Age-related loss of skeletal muscle function and the inability to express the autocrine form of insulin-like growth factor-1 (IGF) in response to mechanical loading.                              | FEBS Lett            | 505:259-63        | 2001 |
| Paffenbarger RS Jr, Hyde RT, Wing AL, Hsieh CC.  | Physical activity, all-cause mortality, and longevity of college alumni.   | N Engl J Med.        | 314(10):605-13.   | 1986 |
| Paffenbarger RS Jr, Hyde RT, Wing AL, Lee IM, Jung DL, Kampert JB  | The association of changes in physical-activity level and other lifestyle characteristics with mortality among men.  | N. Engl. J. Med.     | 328(8):538-545    | 1993 |
| Paffenbarger RS Jr, Laughlin ME, Gima AS, Black RA.  | Work activity of longshoremen as related to death from coronary heart disease and stroke   | N. Engl. J. Med.     | 282(20):1109-1114 | 1970 |
| Paffenbarger RS Jr, Wing AL, Hyde RT, Jung DL.   | Physical activity and incidence of hypertension in college alumni.   | Am. J. Epidemiol.    | 117(3):245-257    | 1983 |
| Paffenbarger RS, Jr., Hyde RT, Wing AL, and Hsieh CC.  | Physical activity, all-cause mortality, and longevity of college alumni.   | N Engl J Med         | 314:605-613       | 1986 |
| Paffenbarger RS, Jr., Kampert JB, Lee IM, Hyde RT, Leung RW, and Wing AL.  | Changes in physical activity and other lifestyle patterns influencing longevity.   | Med Sci Sports Exerc | 26: 857-865       | 1994 |
| Pappas GP, Olcott EW, Drace JE.  | Imaging of skeletal muscle function using (18)FDG PET: force production, activation, and metabolism.   | J Appl Physiol.      | 90(1):329-37.     | 2001 |
| Parkington JD, LeBrasseur NK, Siebert AP, Fielding RA.   | Contraction-mediated mTOR, p70S6k, and ERK1/2 phosphorylation in aged skeletal muscle.   | J Appl Physiol       | 97:243-8          | 2004 |
| Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, Buchner D, Ettinger W, Heath GW, King AC, et al.            | Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine.   | JAMA                 | 273(5): 402-7     | 1995 |
| Paterson DH, Cunningham DA, Koval JJ, and St Croix CM.   | Aerobic fitness in a population of independently living men and women aged 55-86 years.  | Med Sci Sports Exerc | 31: 1813-1820     | 1999 |
| Paterson DH, Govindasamy D, Vidmar M, Cunningham DA, and Koval JJ.   | Longitudinal study of determinants of dependence in an elderly population.   | J Am Geriatr Soc     | 52: 1632-1638     | 2004 |

|  |   |                                |                  |      |
|--|---|--------------------------------|------------------|------|
| Pattison JS, Folk LC, Madsen RW, Booth FW.   | Selected Contribution: Identification of differentially expressed genes between young and old rat soleus muscle during recovery from immobilization-induced atrophy | J Appl Physiol                 | 95:2171-9        | 2003 |
| Pattison JS, Folk LC, Madsen RW, Childs TE, Booth FW.  | Transcriptional profiling identifies extensive downregulation of extracellular matrix gene expression in sarcopenic rat soleus muscle.                              | Physiol Genomics               | 15:34-43         | 2003 |
| Peltoniemi P, Yki-Jarvinen H, Oikonen V, Oksanen A, Takala TO,   | Resistance to exercise-induced increase in glucose uptake during hyperinsulinemia in insulin-resistant skeletal muscle of patients with type 1 diabetes.            | Diabetes.                      | 50(6):1371-7.    | 2001 |
| Pendergast DR, Fisher NM, Calkins E.   | Cardiovascular, neuromuscular, and metabolic alternations with age leading to frailty.  | J Gerontol                     | 48: 61-7         | 1993 |
| Petrella RJ.   | How effective is exercise training for the treatment of hypertension?   | Clin. J. Sport Med.            | 8(3):224-231     | 1998 |
| Ploutz-Snyder LL, Manini T, Ploutz-Snyder RJ, Wolf DA.   | Functionally relevant thresholds of quadriceps femoris strength.  | J Gerontol A Biol Sci Med Sci. | 57A(4):B144-52   | 2002 |
| Pollock ML, Carroll JF, Graves JE, Leggett SH, Braith RW, Limacher M, Porreca E, Di Febbo C, Pintor S, Baccante G, Gatta V, Moretta V, Nisio MD, Palka C, Cuccurullo F, and Shuppia L. | Injuries and adherence to walk/jog and resistance training programs in the elderly.   | Med Sci Sports Exerc.          | 23(10): 1194-200 | 1991 |
| Posner JD, Gorman KM, Klein HS, and Cline CJ.  | Microsatellite polymorphism of the human leptin gene (LEP) and risk of cardiovascular disease.  | Int J Obes (Lond)              |                  | 2005 |
| Powell KE, Thompson PD, Caspersen CJ, Kendrick JS.   | Ventilatory threshold: measurement and variation with age.  | J Appl Physiol                 | 63: 1519-1525    | 1987 |
| Proctor DN, and Joyner MJ.   | Physical activity and the incidence of coronary heart disease   | Ann. Rev. Public Health        | 8:253-287        | 1987 |
| Proctor, D.N. and Joyner, M.J.   | Skeletal muscle mass and the reduction of VO2max in trained older subjects.   | J Appl Physiol                 | 82: 1411-1415    | 1997 |
| Proctor, D.N., Newcomer, S.C., Koch, D.W., Le, K.U., MacLean, D.A. and Leuenberger, U.A.   | Reduced leg blood flow during dynamic exercise in older endurance-trained men.  | J. Appl. Physiol.              | 85(1):68-75.     | 1998 |
| Rall LC, Roubenoff R, Cannon JG, Abad LW, Dinarello CA, Mevdani SN.  | Leg blood flow during submaximal cycle ergometry is not reduced in healthy older normally active men.   | J. Appl. Physiol.              | 94(5):1859-69    | 2003 |
| Rantanen T and Avela J   | Effects of progressive resistance training on immune response in aging and chronic inflammation.  | Med Sci Sports Exerc.          | 28(11):1356-65   | 1996 |
| Rantanen T, Guralnik JM, Foley D, Masaki K, Leveille S, Curb JD, White L.  | Leg Extension power and waking speed in very old people living independently.   | J Gerontol :Medical Science    | 52A(4):M225-     | 1997 |
| Rantanen T, Era P, Heikkinen E.  | Midlife hand grip strength as a predictor of old age disability.  | JAMA                           | 281(6):558-      | 1999 |
| Ravussin E, Burnand B, Schutz Y, Jequier E.  | Maximal isometric strength and mobility among 75-year old men and women.  | Age and aging                  | 23: 132-         | 1994 |
| Reaven G   | Twenty-four-hour energy expenditure and resting metabolic rate in obese, moderately obese, and control  | Am J Clin Nutr                 | 35(3):566-73     | 1982 |
| Reimers CD, Harder T, and Saxe H.  | Metabolic syndrome: Pathophysiology and implications for management of cardiovascular disease.  | Circulation                    | 106:286-288      | 2002 |
|  | Age-related muscle atrophy does not affect all muscles and can partly be compensated by physical activity: an ultrasound study.                                     | J Neurol Sci                   | 159: 60-66       | 1998 |

|  |   |                                  |                     |      |
|--|---|----------------------------------|---------------------|------|
| Ren, J.M., Semenkovich, C.F., Gulve, E.A., Gao, J., and J.O. Holloszy.   | Exercise induces rapid increases in GLUT4 expression, glucose transport capacity, and insulin-stimulated glycogen storage in muscle.                        | J. Biol. Chem.                   | 269: 14396-14401    | 1994 |
| Rhind SG, Shek PN, Shinkai S, Shephard RJ.   | Effects of moderate endurance exercise and training on in vitro lymphocyte proliferation, interleukin-2 (IL-2) production, and IL-2 receptor expression     | Eur J Appl Physiol Occup Physiol | 74(4):348-60        | 1996 |
| Richter EA., Garetto L.P., Goodman M.N., and N.B. Ruderman.  | Enhanced muscle glucose metabolism after exercise: modulation by local factors.   | Am. J. Physiol.                  | 246: E476-E482      | 1984 |
| Rikli RE, Edwards DJ   | Effects of a three-year exercise program on motor function and cognitive processing speed in older women.   | Res Q Exerc Sport                | 62(1):61-7          | 1991 |
| Rodnick, K.J., Henriksen, E.J., James, D.E., and J.O. Holloszy.  | Exercise training, glucose transporters, and glucose transport in rat skeletal muscles.   | Am. J. Physiol.                  | 262: C9-C14         | 1992 |
| Rodriguez BL, Curb JD, Burchfiel CM, Abbott RD, Petrovitch H, Masaki K, Chiu D   | Physical activity and 23-year incidence of coronary heart disease morbidity and mortality among middle-aged men. The Honolulu Heart Program                 | Circulation                      | 89(6):2540-2544     | 1994 |
| Rogers MA, Hagberg JM, Martin WH, Ehsani AA, Holloszy JO   | Decline in O <sub>2</sub> max with aging in master athletes and sedentary men.  | J Appl Physiol                   | 68:2195-2199        | 1990 |
| Rogers MA, Yamamoto C, King DS, Hagberg JM, Ehsani AA, Holloszy JO   | Improvement in glucose tolerance after one week of exercise in patients with mild NIDDM.  | Diabetes Care                    | 11:613-618          | 1988 |
| Rogers ME, Fernandez JE, Bohlken RM.   | Training to reduce postural sway and increase functional reach in the elderly.  | J Occup Rehabil                  | 11: 291-298         | 2001 |
| Romijn JA, Coyle E.F., Sidossis L.S., Gastaldelli A., Horowitz J.F., Emdert E., and R.R. Wolfe.  | Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration.  | Am. J. Physiol.                  | 265: E380-E391      | 1993 |
| Rooks DS, Kiel DP, Parsons C, Hayes WC.  | Self-paced resistance training and walking exercise in community-dwelling older adults: effects on neuromotor performance.                                  | J Gerontol A Biol Sci Med Sci.   | 52(3): M161-8       | 1997 |
| Ross R, Dagnone D, Jones PJ, Smith H, Paddags A, Hudson R, Janssen I.  | Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. | Ann Intern Med.                  | 133(2):92-103.      | 2000 |
| Roth SM, Martel GF, Ferrell RE, Metter EJ, Hurley BF, Rogers MA.   | Myostatin gene expression is reduced in humans with heavy-resistance strength training: a brief   | Exp Biol Med (Maywood)           | 228:706-9           | 2003 |
| Roth SM, Martel GF, Ivey FM, Lemmer JT, Tracy BL, Metter EJ, Hurley BF, Rubenstein LZ, Josephson KR, Trueblood PR, Loy S, Harker JO, Pietruszka FM, Robbins AS       | Skeletal muscle satellite cell characteristics in young and older men and women after heavy resistance strength   | J Gerontol A Biol Sci Med Sci    | 56:B240-7           | 2001 |
| Safar, M.E., Blacher, J, Mourad, J.J. and London, G.M.   | Effects of a group exercise program on strength, mobility, and falls among fall-prone elderly men.  | J Gerontol A Biol Sci Med Sci.   | 55(6): M317-21      | 2000 |
| Salonen JT, Puska P, Tuomilehto J.   | Stiffness of carotid artery wall material and blood pressure in humans  | Stroke                           | 31(3):782-90.       | 2000 |
| Saltin B, Blomqvist G, Mitchell JH, Johnson RL, Jr, Wildenthal K, Chapman Sandmand M, Bruunsgaard H, Kemp K, Andersen-Ranberg K, Pedersen AN, Skinhoj P, Pedersen BK | Physical activity and risk of myocardial infarction, cerebral stroke and death: a longitudinal study in Eastern   | Am. J. Epidemiol.                | 115(4):526-537      | 1982 |
|  | Response to exercise after bed rest and after training.   | Circulation                      | 38(5 Suppl):VIII-78 | 1968 |
|  | Is ageing associated with a shift in the balance between Type 1 and Type 2 cytokines in humans?   | Clin Exp Immunol                 | 127(1):107-14       | 2002 |

|  |  |   |                              |              |
|--|--|---|------------------------------|--------------|
| Sandri M, Sandri C, Gilbert A, Skurk C, Calabria E, Picard A, Walsh K, Schiaffino S, Lecker SH, Goldberg AL, Sarris WH, Blair SN, van Baak MA, Eaton SB, Davies PS, Di Pietro L, Fogelholm M, Rissanen A, Schoeller D, Swinburn B, Tremblay A, Westerterp KR & Wyatt H Sawada S, Tanaka H, Funakoshi M, Shindo M, Kono S, Ishiko T, Schachter F, Faure-Delanef L, Guenot F, Rouger H, Froguel P, Lesueur-Ginot L, Cohen D. | Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy.<br>How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement.<br>Five year prospective study on blood pressure and maximal oxygen uptake.<br>Genetic associations with human longevity at the APOE and ACE loci. | Cell                                      | 117:399-412                  | 2004         |
| Schlicht J, Camaione DN, Owen SV   | Effect of intense strength training on standing balance, walking speed, and sit-to-stand performance in older  | J Gerontol A Biol Sci Med Sci             | 56:M281-M286                 | 2003         |
| Schnelle JF, MacRae PG, Giacobassi K, MacRae HS, Simmons SF, Ouslander JG, Schuit AJ, Feskens EJ, Launer LJ, Kromhout D.   | Exercise with physically restrained nursing home residents: maximizing benefits of restraint reduction.<br>Physical activity and cognitive decline, the role of the apolipoprotein e4 allele.  | J Am Geriatr Soc.<br>Med Sci Sports Exerc | 44(5): 507-12<br>33(5):772-7 | 1996<br>2001 |
| Seals DR, Hagberg JM, Allen WK, Hurley BF, Dalsky GP, Ehsani AA, Holloszy JO   | Glucose tolerance in young and older athletes and sedentary men  | J Appl Physiol                            | 56:1521-1525                 | 1984         |
| Seals DR, Hagberg JM, Hurley BF, Ehsani AA, Holloszy JO  | Effects of endurance training on glucose tolerance and plasma lipids in older men and women.   | JAMA                                      | 252:645-649                  | 1984         |
| Seals DR, Hagberg JM, Hurley BF, Ehsani AA, Holloszy JO  | Endurance training in older men and women 1. Cardiovascular responses to exercise.   | J Appl Physiol                            | 57:1024-1029                 | 1984         |
| Seals DR, Hagberg JM, Spina RJ, Rogers MA, Schechtman KB, Ehsani AA  | Enhanced left ventricular performance in endurance trained older men   | Circulation                               | 89:198-205                   | 1994         |
| Seals DR, Hagberg JM.  | The effect of exercise training on human hypertension: a review.   | Med. Sci. Sports Exerc.                   | 16(3):207-215                | 1984         |
| Seals DR, Stevenson ET, Jones PP, DeSouza CA, Tanaka H   | Lack of age-associated elevations in 24-h systolic and pulse pressures in women who exercise regularly   | Am J Physiol                              | 277:H947-55                  | 1999         |
| Seals, D. R., Hagberg, J. M., Hurley, B. F., Ehsani, A. A., Holloszy, J. O.  | Effects of endurance training on glucose tolerance and plasma lipid levels in older men and women  | JAMA                                      | 252 (5), 645-9               | 1984         |
| Sesso HD, Paffenbarger RS Jr, Lee IM.  | Physical activity and coronary heart disease in men: The Harvard Alumni Health Study.  | Circulation.                              | 102(9):975-80.               | 2000         |
| Shay KA, Roth DL.  | Association between aerobic fitness and visuospatial performance in healthy older adults.  | Psychol Aging.                            | 7(1):15-24.                  | 1992         |
| Shigematsu R, Chang M, Yabushita N, Sakai T, Nakagaichi M, Nho H, Tanaka K.  | Dance-based aerobic exercise may improve indices of falling risk in older women.   | Age Ageing                                | 31(4): 261-6                 | 2002         |
| Shigematsu R, Tanaka K, Holland G, Nakagaichi M, Chang M, Takeshima N, Noda F, Tanaka Y, Mimura K.   | Validation of the functional fitness age (FFA) index in older Japanese women.  | Aging (Milano).                           | 13(5): 358-390               | 2001         |
| Shigematsu R, Ueno LM, Nakagaichi M, Nho H, Tanaka K.  | Rate of perceived exertion as a tool to monitor cycling exercise intensity in older adults.  | Journal of Aging and Physical Activity    | 12: 3-9                      | 2004         |
| Shinkai S, Kohno H, Kimura K, Komura T, Asai H, Inai R, Oka K, Kurokawa Y, Shephard R.   | Physical activity and immune senescence in men   | Med Sci Sports Med                        | 27 1516-1526                 | 1995         |

|   |   |   |                                |              |
|---|---|---|--------------------------------|--------------|
| Shinkai S, Watanabe S, Kumagai S, Fujiwara Y, Amano H, Yoshida H, Ishizaki T, Yukawa H, Suzuki T, Shibata H, Shinohara M, Yoshitake Y, Kouzaki M, Fukuoka H, Fukunaga T         | Walking speed as a good predictor for the onset of functional dependence in a Japanese rural community population.<br>Strength training counteracts motor performance losses during bed rest.   | Age and Ageing<br>J Appl Physiol          | 29:441-<br>95: 1485-1492       | 2000<br>2003 |
| Sihvonen S, Sipilä S, Taskinen S, Era P.  | Fall incidence in frail older women after individualized visual feedback-based balance training.  | Gerontology                               | 6: 411-416                     | 2004         |
| Sihvonen SE, Sipilä S, Era PA.  | Changes in postural balance in frail elderly women during a 4-week visual feedback training: a randomized   | Gerontology                               | 50: 87-95                      | 2004         |
| Singh MA, Ding W, Manfredi TJ, Solares GS, O'Neill EF, Clements KM, Ryan ND, Kehavias JJ, Fielding RA, Evans WJ.  | Insulin-like growth factor I in skeletal muscle after weight-lifting exercise in frail elders.  | Am J Physiol                              | 277:E185-43                    | 1999         |
| Siscovick DS, LaPorte RE, Newman JM.  | The disease-specific benefits and risks of physical activity and exercise.  | Public Health Reports                     | 100(2):180-188                 | 1985         |
| Slentz CA, Duscha BD, Johnson JL, Ketchum K, Aiken LB, Samsa GP, Howard JA, Bales CW, Kraus WE.   | Effects of the amount of exercise on body weight, body composition, and measures of central obesity: STRRIDE--a randomized controlled study.  | Arch Intern Med.                          | 164(1):31-9.                   | 2004         |
| Sonn U, Frandin K, Grimby G.  | Instrumental activities of daily living related to impairments and functional limitation in 70-year-olds and changes between 70 and 76 years of age.  | Scand J Rehab Med                         | 27:119-128                     | 1995         |
| Spina, R.J., Chi, M.M.-Y., Hopkins, M.G., Nemeth, P.M., Lowry, O.H., and J.O. Holloszy.   | Mitochondrial enzymes increase in muscle in response to 7-10 days of cycle exercise.  | J.Appl.Physiol.                           | J.Appl.Physiol.                | 1996         |
| Spranger J, Kroke A, Mohlig M, Hoffmann K, Bergmann MM, Ristow M, Boeing H, Pfeiffer AF.  | Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and independent actions on fear circuits may lead to therapeutic synergy for anxiety when combining serotonergic and GABAergic agents. | Diabetes                                  | 52(3):812-7.                   | 2003         |
| Stahl SM.   | Myocardial adaptations to endurance exercise in aged rats   | J Clin Psychiatry.                        | 63(10):854-5.                  | 2002         |
| Starnes JW, Beyer RE, Edington DW   | Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol.   | Am J Physiol                              | 245:H560-H566                  | 1983         |
| Stefanick ML, Mackey S, Sheehan M, Ellsworth N, Haskell WL, Wood PD.  | Fitness and fatness as predictors of mortality from all causes and from cardiovascular disease in men and women in the lipid research clinics study.  | N Engl J Med.                             | 339(1):12-20.                  | 1998         |
| Stevens J, Cai J, Evenson KR, and Thomas R.   | Influence of exercise training and age on CD14+ cell-surface expression of toll-like receptor 2 and 4.  | Am J Epidemiol                            | 156: 832-841                   | 2002         |
| Stewart LK, Flynn MG, Campbell WW, Craig BA, Robinson JP, McFarlin BK, Timmerman KL, Coen PM, Felker J, Struder HK, Hollmann W, Platen P, Wostmann R, Weicker H, Molderings GJ. | Effect of acute and chronic exercise on plasma amino acids and prolactin concentrations and on [3H]ketanserin binding to serotonin2A receptors on human platelets.  | Brain Behav Immun                         | 19(5):389-97                   | 2005         |
| 杉浦美穂, 長崎浩, 古名丈人, 奥住英之, Takakura Y, Yoshioka K, Umekawa T, Kogure A, Toda H, Yoshikawa T, and Yoshida T.   | 地域高齢者の歩行能力 4年間の経断変化   | Eur J Appl Physiol Occup Physiol.<br>体力科学 | 79(4):318-24.<br>47(4):443-452 | 1999<br>1998 |
|   | Thr54 allele of the FABP2 gene affects resting metabolic rate and visceral obesity.   | Diabetes Res Clin Pract                   | 67: 86-42                      | 2005         |

|  |   |  |   |  |
|--|---|--|---|--|
| Takala TO, Nuutila P, Knuuti J, Luotolahti M, Yki-Jarvinen H.  | Insulin action on heart and skeletal muscle glucose uptake in weight lifters and endurance athletes.  | Am J Physiol.  | 276(4 Pt 1):E706-11.  | 1999   |
| Takata Y, Ansai T, Awano S, Hamasaki T, Yoshitake Y, Kimura Y, Sonoki K, Wakisaka M, Fukuhara M, Takehara T, Tamai T, Nakai T, Takai H, Fujiwara R, Mivabo S, Higuchi M, Kobayashi S, Tanabe T, Maeda S, Miyauchi T, Iemitsu M, Takanashi M, Irukayama Tomobe Y, Yokota T, Ohmori H, Matsuda M, Tanaka H, Dinanno FA, Monahan KD, Clevenger CM, DeSouza CA, Seals DR, Tanaka K, Okura T, Shigematsu R, Nakata Y, Lee DJ, Wee SW, Yamabuki K, Tanaka K, Sakai T, Nakamura Y, Umeda N, Lee DJ, Nakata Y, Hayashi Y, Akutsu T, Okura T, Yamabuki K, Tanaka K, Shigematsu R, Nakagaichi M, Kim H, Takeshima N. | Relationship of physical fitness to chewing in an 80-year-old population.<br>The effects of physical exercise on plasma lipoprotein and apolipoprotein metabolism in elderly men.<br>Exercise training improves ageing-induced decrease in eNOS expression of the aorta<br>Aging, habitual exercise, and dynamic arterial compliance<br>Target value of intraabdominal fat area for improving coronary heart disease risk factors.<br>Health benefits associated with exercise habituation in older Japanese men.<br>The relationship between functional fitness and coronary heart disease risk factors in older Japanese adults.<br>Potential benefits of exercise on development of coronary collateral during chronic ischemia.<br>Absence of age-related increase in central arterial stiffness in physically active women | Oral Diseases<br>J. Gerontol<br>Acta Physiol Scand<br>Circulation<br>Obes Res<br>Aging Clin Exp Res.<br>Journal of Aging and Physical Activity<br>Advances in Exercise and Sports Physiology<br>Arterioscler Thromb Vas Biol.<br>Circulation | 10(1): 44-9<br>43:M75-79<br>178:3-10<br>102:1270-5<br>12(4): 695-703<br>16(1): 53-59<br>7: 162-174<br>4: 83-91<br>18(1):127-32.<br>102:1270-6 | 2004<br>1988<br>2003<br>2000<br>2004<br>2004<br>2000 |
| Tanaka H, DeSouza, C.A. and Seals, D.R.  | Enhanced calcium uptake of cardiac sarcoplasmic reticulum in exercise-trained old rats  | Am J Physiol   | 258:H431-H435   | 1990   |
| Tanaka, T., Dinanno, F.A., Monahan, K.D., Clevenger, C.M., DeSouza, C.A. and Seals, D.R.   | Effects of high-intensity swimming training on GLUT-4 and glucose transport activity in rat skeletal muscle.  | J.Appl.Physiol   | 90: 2019-2024   | 2001   |
| Tate CA, Taffet GE, Hudson EK, Blaylock SL, McBride RP, Michael LH, Terada S, Yokozeki T, Kawanaka K, Ogawa K, Higuchi M, Ezaki O, and I. Toft AD, Jensen LB, Bruunsgaard H, Ibfelt T, Halkjaer-Kristensen J, Febbraio M, Pedersen BK  | Cytokine response to eccentric exercise in young and elderly humans.<br>Effects of age and exercise on the extent of the myocardial capillary bed<br>Contribution of body composition and physical activity to age related decline in peak VO2 in men and women.<br>Strength training reduces force fluctuations during anisometric contractions of the quadriceps femoris<br>Changes in serum levels of E-selectin correlate to improved glycaemic control and reduced obesity in subjects with the metabolic syndrome.<br>Effect of 4- and 8-wk intensive Tai Chi Training on balance control in the elderly.<br>Exercise training reverses the age-related decline in tyrosine hydroxylase expression in rat hypothalamus.   | Am J Physiol Cell Physiol<br>Anat Rec<br>J Appl Physiol<br>J Appl Physiol<br>Scand J Clin Lab Invest.<br>Med Sci Sports Exerc<br>J Gerontol A Biol Sci Med Sci.  | 283(1):C289-95<br>167:55-62<br>77: 647-652<br>96: 1530-1540<br>65(4):283-90.<br>36: 648-657<br>52(5):B255-9.                                  | 2002<br>1970<br>1994<br>2003<br>2005<br>2004<br>1997 |
| Tomanek RJ   |   |  |   |  |
| Toth MJ, Gardner AW, Ades PA, and Poehlman ET.   |   |  |   |  |
| Tracy BL, Byrnes WC, Enoka RM  |   |  |   |  |
| Troseid M, Lappegard KT, Mollnes TE, Arnesen H, Seljeflot I.   |   |  |   |  |
| Tsang WW, Hui-Chan CW.   |   |  |   |  |
| Tumer N, LaRochelle JS, Yurekli M.   |   |  |   |  |



|  |   |                                     |                         |      |
|--|---|-------------------------------------|-------------------------|------|
| Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukkaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M; Finnish Diabetes Prevention Study Group              | Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance.  | N Engl J Med.                       | 344(18):1343-50.        | 2001 |
| Ullman M, Ullman A, Sommerland H, Skottner A, Oldfors A.   | Effects of growth hormone on muscle regeneration and IGF-I concentration in old rats.   | Acta Physiol Scand                  | 140:521-5               | 1990 |
| Vaitkevicius PV, Fleg JL, Engel JH, O'Connor FC, Wright JG, Lakatta LE, Yin FC, Lakatta EG.  | Effects of age and aerobic capacity on arterial stiffness in healthy adults   | Circulation                         | 88:1456-62              | 1993 |
| van Boxtel MP, Langerak K, Houx PJ, Jolles J.  | Self-reported physical activity, subjective health, and cognitive performance in older adults.  | Exp Aging Res.                      | 22(4):363-79.           | 1996 |
| van Boxtel MP, Paas FG, Houx PJ, Adam JJ, Teeken JC, Jolles J.   | Aerobic capacity and cognitive performance in a cross-sectional aging study.  | Med Sci Sports Exerc.               | 29(10):1357-65.         | 1997 |
| van Heuvelen MJ, Kempen GI, Brouwer WH, de Greef MH.   | Physical fitness related to disability in older persons.  | Gerontology.                        | 46(6):333-41.           | 2000 |
| van Praag H, Kempermann G, Gage FH   | Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus.   | Nature Neuroscience                 | 2(3):266-70.            | 1999 |
| van Praag H, Shubert T, Zhao C, Gage FH.   | Exercise enhances learning and hippocampal neurogenesis in aged mice.   | J Neurosci.                         | 25(38):8680-5           | 2005 |
| van Praag H, Shubert T, Zhao C, Gage FH.   | Exercise enhances learning and hippocampal neurogenesis in aged mice.   | J Neurosci.                         | 25(38):8680-5.          | 2005 |
| van Saase JL, Noteboom WM, and Vandenbroucke JP.   | Longevity of men capable of prolonged vigorous physical exercise: a 32 year follow up of 2259 participants in the Dutch eleven cities ice skating tour. | Br. Med. Jour.                      | 301: 1409-1411          | 1990 |
| Verghese, Joe; Lipton, Richard B; Katz, Mindy J; Hall, Charles B; Derby, Carol A; Kuslansky, Gail; Ambrose, Anne F; Sliwinski, Martin; Buschke, Herman Wadlen TA, Berkowitz RI, Sarwer DB, Prus-Wisniewski R. Steinberg C. | Leisure activities and the risk of dementia in the elderly  | The New England Journal Of Medicine | 348(25) : 2508-2516     | 2003 |
| Wahren, J., Felig, P., Ahlberg, G., and L. Jorfeldt.   | Benefits of lifestyle modification in the pharmacologic treatment of obesity: a randomized trial.   | Arch Intern Med.                    | 161(2):218-27.          | 2001 |
| Walsh JH, Yong G, Cheetham C, Watts G, O'Driscoll G, Taylor R, Green D   | Glucose metabolism during leg exercise in man.  | J.Clin.Invest.                      | 50: 2715-2725           | 1971 |
| Wang G, Dietz WH   | Effect of exercise training on conduit and resistance vessel function in treated and untreated hypercholesterolaemic                                    | Eur Heart J                         | 24:1681-1689            | 2003 |
| Wannamethee G, Shaper AG.  | Economic burden of obesity in youths aged 6 to 17 years:1979-1999.  | Pediatrics                          | 109:E81-91              | 2002 |
| Wasserman K, Whipp BJ, Koyl SN, Beaver WL.   | Physical activity and stroke in British middle aged men   | BMJ                                 | 304(6827):597-601       | 1992 |
| Wei M, Gibbons LW, Mitchell TL, Kampert JB, Lee CD, Blair SN.  | Anaerobic threshold and respiratory gas exchange during exercise.   | J Appl Physiol.                     | 35(2):236-43            | 1973 |
| Wei M, Kampert JB, Barlow CE, Nichaman MZ, Gibbons LW, Paffenbarger RS Jr, Blair SN.   | The association between cardiorespiratory fitness and impaired fasting glucose and type 2 diabetes mellitus in  | Ann Intern Med.                     | 39-96. Erratum in: 131( | 1999 |
|  | Relationship between low cardiorespiratory fitness and mortality in normal-weight, overweight, and obese men.   | JAMA.                               | 282(16):1547-53.        | 1999 |

|   |   |   |  |              |
|---|---|---|--|--------------|
| Welle S, Brooks AI, Delehanty JM, Needler N, Thornton CA, Westertep KR.   | Gene expression profile of aging in human muscle. Pattern and intensity of physical activity.   | Physiol Genomics                                | 14:149-59                                | 2003         |
| Weuve J, Kang JH, Manson JE, Breteler MM, Ware JH, Grodstein F, Whittemore LA, Song K, Li X, Aghajanian J, Davies M, Kirgenrath S, Hill JJ, Jalenak M, Kelley P, Knight A, Maylor R, O'Hara D, Pearson A, Quazi A, Ryerson S, Tan XY, Tomkinson KN, Veldman GM, Widom A, Wright JF, Wujakka S, Zhao J, Wildman RP, Farhat GN, Patel AS, Mackey RH, Brockwell S, Thompson T, Sutton-Tyrrell K. | Physical activity, including walking, and cognitive function in older women. Inhibition of myostatin in adult mice increases skeletal muscle mass and strength. Weight Change Is Associated With Change in Arterial Stiffness Among Healthy Young Adults  | Nature<br>JAMA                                  | 410: 539<br>292:1454-                    | 2001<br>2004 |
| Woo KS, Chook P, Yu CW, Sung RY, Qiao M, Leung SS, Lam CW, Metreweli C, Celermaier DS, Woods JA, Ceddia MA, Wolters BW, Evans JK, Lu Q, McAuley E.  | Effects of diet and exercise on obesity-related vascular dysfunction in children Effects of 6 months of moderate aerobic exercise training on immune function in the elderly. Effects of maximal exercise on natural killer (NK) cell cytotoxicity and responsiveness to interferon-alpha in the young and old. | Circulation<br>Mech Ageing Dev                  | 109: 1981-6<br>109(1):1-19               | 2004<br>1999 |
| Woods JA, Evans JK, Wolters BW, Ceddia MA, McAuley E.   | Effects of maximal exercise on natural killer (NK) cell cytotoxicity and responsiveness to interferon-alpha in the young and old.   | J Gerontol A Biol Sci Med Sci.                  | 53(6):B430-7                             | 1998         |
| Yaffe K, Barnes D, Nevitt M, Lui LY, Covinsky K, Yamaga T, Yoshihara A, Ando Y, Yoshitake Y, Kimura Y, Shimada M, Nishimuta M, Mizazaki H.  | A prospective study of physical activity and cognitive decline in elderly women: women who walk. Relationship between dental occlusion and physical fitness in an elderly population.   | Arch Intern Med.<br>J Gerontol..Medical Science | 161(14):1703-8.<br>57A(9):M616- 10:44-49 | 2001<br>2002 |
| Yamamoto T, Castell LM, Botella J, Powell H, Yamanouchi K, Shinozaki T, Chikada K, Nishikawa T, Ito K, Shimizu S, Ozawa N, Suzuki Y, Maeno H, Kato K, et al.  | Changes in the albumin binding of tryptophan during postoperative recovery: a possible link with central Daily walking combined with diet therapy is a useful means for obese NIDDM patients not only to reduce body weight but also to improve insulin sensitivity.  | Brain Res Bull.<br>Diabetes Care                | 43(1):43-6.<br>18:775-778.               | 1997<br>1995 |
| Yamanouchi K, Nakajima H, Shinozaki T, Chikada K, Oshida Y, Sato Y, Higuchi M, Kobavashi S.   | Effects of daily physical activity on insulin action in the elderly.  | J. Appl. Physiol.                               | 73:2241-2245                             | 1992         |
| Yamanouchi K, Shinozaki T, Chikada K, Nishikawa T, Ito K, Shimizu S, Ozawa N, Suzuki Y, Maeno H, Kato K, et al.   | Daily walking combined with diet therapy is a useful means for obese NIDDM patients not only to reduce body weight but also to improve insulin sensitivity.   | Diabetes Care.                                  | 18(6): 775-8                             | 1995         |
| Yan H, Kuroiwa A, Tanaka H, Shindo M, Kivonaga A, Nagavama A.   | Effect of moderate exercise on immune senescence in men.  | Eur J Appl Physiol                              | 86(2):105-11                             | 2001         |
| Yarasheski KE, Zachwieja JJ, Bier DM.   | Acute effects of resistance exercise on muscle protein synthesis rate in young and elderly men and women.   | Am J Physiol                                    | 265:E210-4                               | 1993         |
| Yarasheski KE, Pak-Loduca J, Hasten DL, Obert KA, Brown MB, Sinacore DR.  | Resistance exercise training increases mixed muscle protein synthesis rate in frail women and men >=76 yr   | Am J Physiol                                    | 277:E118-25                              | 1999         |
| Yarasheski KE, Bhasin S, Sinha-Hikim I, Pak-Loduca J, Gonzalez-Cadavid NF.  | Serum myostatin immunoreactive protein is increased in 60-92 year old women and men with muscle wasting.  | J Nutr Health Aging                             | 6:343-8                                  | 2002         |
| Yoshida T, Sakane N, Umekawa T, Sakai M, Takahashi T, and Kondo M.  | Mutation of beta 3-adrenergic-receptor gene and response to treatment of obesity.   | Lancet  | 346: 1433-1434                           | 1995         |

|  |  |   |                      |      |
|--|--|---|----------------------|------|
| Yoshiga CC, Higuchi M, Oka J   | Serum lipoprotein cholesterol in older oarsmen.  | Eur. J. Appl. Physiol.  | 87:228-232           | 2002 |
| Yoshiga CC, Yashiro K, Higuchi M, Oka J  | Rowing prevents muscle wasting in older men.   | Eur. J. Appl. Physiol.  | 88:1-4               | 2002 |
| Yoshioka M, Ayabe M, Yahiro T, Higuchi H, Higaki Y, St-Amand J, Miyazaki H, Yoshitake Y, Shindo M, Tanaka H. | Long-term accelerometer monitoring shows the role of physical activity in overweight and obesity.  | Int J Obes  | 29:502-508           | 2005 |
| Yoshitake Y., et al.   | Relation between physical fitness and functional performance in 80-year-old men and women residing in a community for the elderly.   | In Exercise for preventing common diseases, (eds) Tanaka H. and Shindo M, Springer-Verlag, Tokyo              | pp. 147-153          | 1999 |
| Yoshitake Y., et al.   | Relationship between physical fitness and functional performances in older women.  | In Recent Advances in Physiological Anthropology, (eds) Sato M, Tokura H, Watanuki S, kyushu University Press | pp. 299-308          | 1999 |
| Young A  | Exercise physiology in geriatric practice  | Acta Med Scand  | Suppl. 711:227-      | 1986 |
| Young A, Skelton DA  | Applied physiology of strength and power in old age.   | Int J Sport Med   | 15:149-151           | 1994 |
| Zhang B, Sakai T, Miura S, Kiyonaga A, Tanaka H, Shindo M, Saku K.   | Association of angiotensin-converting-enzyme gene polymorphism with the depressor response to mild exercise therapy in patients with mild to moderate effect of exercise testing protocol on parameters of aerobic function. | Clin Genet.   | 62(4):328-33         | 2002 |
| Zhang YY, Johnson MC 2nd, Chow N, Wasserman K  | Effect of exercise testing protocol on parameters of aerobic function.   | Med Sci Sports Exerc  | 23(5):625-30         | 1991 |
| Zinman, B., Ruderman, N., Campaigne, B. N., Devlin, J. T., Schneider, S. H.                                  | Physical activity/exercise and diabetes.   | Diabetes Care   | 27 (Suppl 1), S58-62 | 2004 |



利用者側 文献検索フロー

