

表23 血清LDL-Cコレステロールの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						mg/dl	
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル					
			5	25	50	75			95	5	25	50		75
29歳以下	83	102.1 (33.5)	52.2	83.0	98.0	115.0	167.8	179	95.1 (26.7)	57.0	77.0	92.0	111.0	145.0
30-39	211	116.7 (32.7)	70.0	90.0	113.0	141.0	178.2	618	100.8 (26.1)	62.0	83.0	99.0	116.0	147.0
40-44	343	117.2 (32.7)	68.2	94.0	116.0	137.0	175.0	789	107.5 (26.2)	69.0	89.0	106.0	123.0	152.5
45-49	456	117.4 (32.5)	68.7	95.0	115.5	137.0	172.3	1,163	116.8 (28.9)	73.2	96.0	115.0	135.0	164.0
50-54	730	118.2 (29.6)	73.0	98.0	116.0	137.0	169.0	1,852	124.7 (30.5)	79.0	104.0	123.0	143.0	180.0
55-59	770	114.4 (29.2)	69.6	95.0	113.5	132.0	161.9	2,103	127.3 (28.9)	82.0	108.0	126.0	145.0	176.0
60-64	1,345	114.5 (30.2)	70.0	93.0	113.0	134.0	166.0	2,830	128.1 (27.8)	85.0	109.0	127.0	145.0	177.0
65-69	1,874	112.7 (28.7)	67.0	93.0	112.0	132.0	162.0	3,148	126.0 (27.8)	83.0	107.0	125.0	143.0	173.0
70-74	1,749	111.6 (27.8)	67.0	93.0	110.0	130.0	158.0	2,557	125.0 (26.4)	85.0	107.0	124.0	141.0	171.0
75-79	1,027	112.4 (27.3)	71.0	94.0	110.0	129.0	157.0	1,336	124.4 (28.2)	81.0	104.0	123.0	142.0	172.0
80歳以上	371	109.7 (27.5)	66.0	91.0	110.0	129.0	154.4	393	121.5 (28.1)	78.0	104.0	120.0	138.0	170.6
計	8,959	113.6 (29.3)	69.0	94.0	112.0	132.0	163.0	16,968	123.3 (28.9)	78.0	104.0	122.0	141.0	172.0

表24 血清中性脂肪の状況 (性・年齢階級別)

年齢階級	総数						人 (%)	
	-39	40-99	100-149	150-199	200-239	240mg/dl以上		
男								
-29歳	86 (100.0)	38 (44.2)	23 (26.7)	14 (16.3)	3 (3.5)	5 (5.8)		
30-39	214 (100.0)	80 (37.4)	54 (25.2)	29 (13.6)	20 (9.3)	26 (12.1)		
40-44	348 (100.0)	105 (30.2)	81 (23.3)	76 (21.8)	22 (6.3)	58 (16.7)		
45-49	465 (100.0)	158 (34.0)	132 (28.4)	73 (15.7)	37 (8.0)	61 (13.1)		
50-54	741 (100.0)	271 (36.6)	230 (31.0)	97 (13.1)	36 (4.9)	95 (12.8)		
55-59	778 (100.0)	324 (41.6)	216 (27.8)	105 (13.5)	34 (4.4)	80 (10.3)		
60-64	1,368 (100.0)	577 (42.2)	379 (27.7)	180 (13.2)	91 (6.7)	114 (8.3)		
65-69	1,912 (100.0)	909 (47.5)	534 (27.9)	228 (11.9)	81 (4.2)	124 (6.5)		
70-74	1,803 (100.0)	927 (51.4)	456 (25.3)	202 (11.2)	86 (4.8)	86 (4.8)		
75-79	1,060 (100.0)	547 (51.6)	283 (26.7)	111 (10.5)	40 (3.8)	49 (4.6)		
80歳以上	385 (100.0)	222 (57.7)	102 (26.5)	34 (8.8)	12 (3.1)	11 (2.9)		
計	9,160 (100.0)	4,158 (45.4)	2,490 (27.2)	1,149 (12.5)	462 (5.0)	709 (7.7)		

年齢階級	総数						人 (%)	
	-39	40-99	100-149	150-199	200-239	240mg/dl以上		
女								
-29歳	180 (100.0)	125 (69.4)	16 (8.9)	4 (2.2)	1 (0.6)	6 (3.3)		
30-39	620 (100.0)	403 (65.0)	89 (14.4)	37 (6.0)	16 (2.6)	20 (3.2)		
40-44	795 (100.0)	482 (60.6)	145 (18.2)	53 (6.7)	23 (2.9)	26 (3.3)		
45-49	1,185 (100.0)	703 (59.3)	241 (20.3)	109 (9.2)	36 (3.0)	40 (3.4)		
50-54	1,878 (100.0)	995 (53.0)	494 (26.3)	174 (9.3)	66 (3.5)	91 (4.8)		
55-59	2,138 (100.0)	1,045 (48.9)	575 (26.9)	260 (12.2)	99 (4.6)	104 (4.9)		
60-64	2,878 (100.0)	1,319 (45.8)	853 (29.6)	383 (13.3)	123 (4.3)	135 (4.7)		
65-69	3,215 (100.0)	1,465 (45.6)	972 (30.2)	410 (12.8)	139 (4.3)	173 (5.4)		
70-74	2,621 (100.0)	1,235 (47.1)	798 (30.4)	340 (13.0)	92 (3.5)	133 (5.1)		
75-79	1,383 (100.0)	641 (46.3)	441 (31.9)	169 (12.2)	54 (3.9)	63 (4.6)		
80歳以上	412 (100.0)	201 (48.8)	127 (30.8)	48 (11.7)	22 (5.3)	12 (2.9)		
計	17,305 (100.0)	8,614 (49.8)	4,751 (27.5)	1,987 (11.5)	671 (3.9)	803 (4.6)		

表25 血清中性脂肪の平均値、標準偏差および分布 (性・年齢階級別)

年齢階級	男						女						mg/dl		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	86	122.4 (85.6)	46.4	76.8	103.5	150.3	313.3	180	75.7 (69.3)	31.0	44.0	56.0	83.0	166.7	
30-39	214	144.0 (97.1)	49.0	78.0	116.0	184.3	341.8	620	89.3 (62.5)	34.0	51.0	70.5	103.0	208.8	
40-44	348	160.8 (106.6)	50.5	89.0	129.5	194.0	383.7	795	92.8 (61.9)	37.0	55.0	74.0	109.0	218.0	
45-49	465	149.7 (106.4)	52.3	86.0	123.0	176.0	345.2	1,185	101.8 (86.1)	40.0	58.5	81.0	124.5	216.0	
50-54	741	141.1 (99.6)	49.0	78.0	112.0	166.0	337.9	1,878	109.1 (69.8)	43.0	66.0	92.0	128.3	236.1	
55-59	778	130.7 (87.0)	48.0	75.0	107.5	157.0	299.1	2,138	114.8 (66.7)	45.0	70.0	97.0	141.0	237.1	
60-64	1,368	130.7 (87.1)	47.0	75.0	108.0	158.0	279.6	2,878	116.3 (63.5)	48.0	74.0	102.0	143.0	232.0	
65-69	1,912	120.3 (80.2)	47.0	71.0	101.0	144.0	260.0	3,215	118.5 (65.5)	48.0	74.0	103.0	143.0	244.2	
70-74	1,803	114.1 (69.7)	46.0	70.0	95.0	139.0	235.8	2,621	118.1 (63.9)	51.0	76.0	102.0	142.0	241.0	
75-79	1,060	111.2 (67.2)	45.0	66.0	94.0	135.8	236.0	1,383	116.6 (60.5)	53.0	77.0	102.0	139.0	231.8	
80歳以上	385	104.3 (54.1)	48.3	69.0	89.0	124.5	209.0	412	113.2 (54.5)	51.0	76.0	100.0	136.5	210.4	
計	9,160	125.1 (83.6)	47.0	72.0	103.0	150.0	278.0	17,305	112.5 (66.9)	44.0	69.0	96.0	137.0	233.0	

表26 血清ASTの状況(性・年齢階級別)

年齢階級	人 (%)				
	総数	-19	20-39	40-59	60U/L以上
男					
-29歳	86 (100.0)	32 (37.2)	51 (59.3)	2 (2.3)	1 (1.2)
30-39	214 (100.0)	47 (22.0)	152 (71.0)	9 (4.2)	6 (2.8)
40-44	348 (100.0)	57 (16.4)	258 (74.1)	25 (7.2)	8 (2.3)
45-49	465 (100.0)	90 (19.4)	336 (72.3)	30 (6.5)	9 (1.9)
50-54	741 (100.0)	113 (15.2)	562 (75.8)	44 (5.9)	22 (3.0)
55-59	778 (100.0)	119 (15.3)	585 (75.2)	51 (6.6)	23 (3.0)
60-64	1,368 (100.0)	200 (14.6)	1,051 (76.8)	82 (6.0)	35 (2.6)
65-69	1,912 (100.0)	232 (12.1)	1,485 (77.7)	154 (8.1)	41 (2.1)
70-74	1,803 (100.0)	220 (12.2)	1,433 (79.5)	113 (6.3)	37 (2.1)
75-79	1,060 (100.0)	151 (14.2)	821 (77.5)	70 (6.6)	18 (1.7)
80歳以上	385 (100.0)	48 (12.5)	310 (80.5)	22 (5.7)	5 (1.3)
計	9,160 (100.0)	1,309 (14.3)	7,044 (76.9)	602 (6.6)	205 (2.2)

年齢階級	人 (%)				
	総数	-19	20-39	40-59	60U/L以上
女					
-29歳	180 (100.0)	140 (77.8)	37 (20.6)	3 (1.7)	0 (0.0)
30-39	620 (100.0)	439 (70.8)	172 (27.7)	8 (1.3)	1 (0.2)
40-44	795 (100.0)	517 (65.0)	269 (33.8)	5 (0.6)	4 (0.5)
45-49	1,185 (100.0)	617 (52.1)	534 (45.1)	28 (2.4)	6 (0.5)
50-54	1,878 (100.0)	666 (35.5)	1,163 (61.9)	38 (2.0)	11 (0.6)
55-59	2,138 (100.0)	506 (23.7)	1,559 (72.9)	58 (2.7)	15 (0.7)
60-64	2,878 (100.0)	528 (18.3)	2,240 (77.8)	85 (3.0)	25 (0.9)
65-69	3,215 (100.0)	585 (18.2)	2,501 (77.8)	97 (3.0)	32 (1.0)
70-74	2,621 (100.0)	415 (15.8)	2,111 (80.5)	79 (3.0)	16 (0.6)
75-79	1,383 (100.0)	220 (15.9)	1,108 (80.1)	51 (3.7)	4 (0.3)
80歳以上	412 (100.0)	79 (19.2)	322 (78.2)	9 (2.2)	2 (0.5)
計	17,305 (100.0)	4,712 (27.2)	12,016 (69.4)	461 (2.7)	116 (0.7)

表27 血清ASTの平均値、標準偏差および分布（性・年齢階級別）

U/L

年齢階級	男						女							
	人数	平均値(SD)	5	25	50	75	95	人数	平均値(SD)	5	25	50	75	95
29歳以下	86	23.1 (8.7)	14.0	18.0	21.0	25.0	37.3	180	17.5 (5.2)	12.1	15.0	16.5	19.0	25.0
30-39	214	26.7 (11.3)	16.0	20.0	24.0	29.0	47.3	620	18.4 (6.0)	13.0	15.0	17.0	20.0	27.0
40-44	348	27.6 (11.2)	17.0	21.0	25.0	31.0	44.6	795	19.4 (9.0)	13.0	16.0	18.0	21.0	28.0
45-49	465	27.9 (19.6)	16.3	21.0	25.0	30.0	43.7	1,185	20.7 (7.1)	14.0	17.0	19.0	22.0	33.0
50-54	741	28.0 (13.0)	17.0	21.0	25.0	30.0	49.0	1,878	22.9 (8.1)	15.0	18.0	22.0	25.0	34.1
55-59	778	28.7 (18.6)	17.0	21.0	25.0	31.0	48.1	2,138	24.0 (7.6)	16.0	20.0	23.0	26.0	36.0
60-64	1,368	28.5 (17.4)	17.0	21.0	25.0	31.0	45.6	2,878	25.1 (13.8)	17.0	20.0	23.0	27.0	37.0
65-69	1,912	28.4 (13.3)	18.0	22.0	25.0	31.0	47.0	3,215	25.3 (8.5)	17.0	21.0	24.0	28.0	38.0
70-74	1,803	27.9 (12.0)	17.0	22.0	26.0	31.0	44.0	2,621	25.2 (8.0)	17.0	21.0	24.0	28.0	37.0
75-79	1,060	27.5 (11.7)	17.0	21.0	25.0	31.0	42.0	1,383	25.2 (7.0)	17.0	21.0	24.0	28.0	37.0
80歳以上	385	27.6 (12.6)	18.0	22.0	26.0	30.0	43.0	412	24.6 (6.6)	17.0	21.0	24.0	27.0	36.4
計	9,160	28.0 (14.3)	17.0	21.0	25.0	31.0	45.0	17,305	23.9 (9.3)	15.0	19.0	22.0	27.0	36.0

表28 血清ALTの状況 (性・年齢階級別)

年齢階級	人 (%)				
	総数	-19	20-39	40-59	60U/L以上
-29歳	86 (100.0)	39 (45.3)	34 (39.5)	7 (8.1)	6 (7.0)
30-39	214 (100.0)	44 (20.6)	111 (51.9)	31 (14.5)	28 (13.1)
40-44	348 (100.0)	62 (17.8)	188 (54.0)	60 (17.2)	38 (10.9)
45-49	465 (100.0)	105 (22.6)	262 (56.3)	73 (15.7)	25 (5.4)
50-54	741 (100.0)	167 (22.5)	443 (59.8)	86 (11.6)	45 (6.1)
55-59	778 (100.0)	183 (23.5)	462 (59.4)	87 (11.2)	46 (5.9)
60-64	1,368 (100.0)	380 (27.8)	812 (59.4)	119 (8.7)	57 (4.2)
65-69	1,912 (100.0)	588 (30.8)	1,106 (57.8)	162 (8.5)	56 (2.9)
70-74	1,803 (100.0)	654 (36.3)	997 (55.3)	111 (6.2)	41 (2.3)
75-79	1,060 (100.0)	454 (42.8)	530 (50.0)	58 (5.5)	18 (1.7)
80歳以上	385 (100.0)	204 (53.0)	165 (42.9)	10 (2.6)	6 (1.6)
計	9,160 (100.0)	2,880 (31.4)	5,110 (55.8)	804 (8.8)	366 (4.0)

男

年齢階級	人 (%)				
	総数	-19	20-39	40-59	60U/L以上
-29歳	180 (100.0)	147 (81.7)	28 (15.6)	4 (2.2)	1 (0.6)
30-39	620 (100.0)	492 (79.4)	107 (17.3)	11 (1.8)	10 (1.6)
40-44	795 (100.0)	586 (73.7)	184 (23.1)	18 (2.3)	7 (0.9)
45-49	1,185 (100.0)	781 (65.9)	331 (27.9)	52 (4.4)	21 (1.8)
50-54	1,878 (100.0)	969 (51.6)	778 (41.4)	101 (5.4)	30 (1.6)
55-59	2,138 (100.0)	1,006 (47.1)	992 (46.4)	99 (4.6)	41 (1.9)
60-64	2,878 (100.0)	1,345 (46.7)	1,328 (46.1)	141 (4.9)	64 (2.2)
65-69	3,215 (100.0)	1,515 (47.1)	1,519 (47.2)	130 (4.0)	51 (1.6)
70-74	2,621 (100.0)	1,417 (54.1)	1,098 (41.9)	75 (2.9)	31 (1.2)
75-79	1,383 (100.0)	853 (61.7)	487 (35.2)	36 (2.6)	7 (0.5)
80歳以上	412 (100.0)	292 (70.9)	112 (27.2)	8 (1.9)	0 (0.0)
計	17,305 (100.0)	9,403 (54.3)	6,964 (40.2)	675 (3.9)	263 (1.5)

女

表29 血清ALTの平均値、標準偏差および分布（性・年齢階級別）

U/L

年齢階級	男						女							
	人数	平均値(SD)	5	25	50	75	95	人数	平均値(SD)	5	25	50	75	95
29歳以下	86	27.9 (22.2)	11.0	16.0	21.0	29.0	78.6	180	15.3 (9.9)	8.0	10.0	13.0	16.0	35.0
30-39	214	37.8 (27.3)	13.0	21.0	30.0	44.0	104.0	620	17.2 (12.3)	9.0	12.0	14.0	18.0	35.0
40-44	348	35.0 (20.5)	15.0	22.0	29.0	43.0	73.0	795	17.9 (11.6)	9.0	12.0	15.0	20.0	35.0
45-49	465	32.1 (28.4)	14.0	20.0	27.0	37.0	64.7	1,185	20.0 (13.4)	10.0	13.0	17.0	22.0	44.7
50-54	741	31.2 (20.3)	14.1	20.0	27.0	36.0	64.0	1,878	22.3 (13.5)	11.0	15.0	19.0	25.0	43.0
55-59	778	30.6 (19.1)	14.0	20.0	26.0	35.0	63.0	2,138	23.2 (14.0)	12.0	16.0	20.0	26.3	43.0
60-64	1,368	28.2 (17.0)	14.0	19.0	24.0	32.0	56.0	2,878	23.7 (22.6)	12.0	16.0	20.0	26.0	46.0
65-69	1,912	26.7 (14.7)	13.0	18.0	23.0	30.0	51.3	3,215	22.5 (11.9)	12.0	16.0	20.0	25.0	41.0
70-74	1,803	25.0 (13.3)	12.0	17.0	22.0	28.0	46.0	2,621	21.2 (10.0)	12.0	16.0	19.0	24.0	37.0
75-79	1,060	23.7 (12.8)	12.0	16.3	21.0	27.0	43.0	1,383	19.7 (8.5)	11.0	15.0	18.0	22.0	35.0
80歳以上	385	21.6 (12.6)	11.0	15.0	19.0	25.0	36.0	412	17.7 (7.1)	9.7	13.0	17.0	20.0	30.0
計	9,160	27.6 (17.4)	13.0	18.0	23.0	31.0	55.0	17,305	21.6 (14.3)	11.0	15.0	19.0	24.0	40.0

表30 γ-GTPの状況(性・年齢階級別)

年齢階級	総数					人(%)	
	-24	25-49	50-74	75-99	100U/L以上		
男							
-29歳	86 (100.0)	27 (31.4)	9 (10.5)	2 (2.3)	5 (5.8)		
30-39	214 (100.0)	87 (40.7)	30 (14.0)	16 (7.5)	31 (14.5)		
40-44	348 (100.0)	129 (37.1)	56 (16.1)	32 (9.2)	67 (19.3)		
45-49	465 (100.0)	176 (37.8)	74 (15.9)	44 (9.5)	72 (15.5)		
50-54	741 (100.0)	271 (36.6)	130 (17.5)	65 (8.8)	107 (14.4)		
55-59	778 (100.0)	286 (36.8)	129 (16.6)	70 (9.0)	101 (13.0)		
60-64	1,368 (100.0)	552 (40.4)	197 (14.4)	75 (5.5)	167 (12.2)		
65-69	1,912 (100.0)	743 (38.9)	237 (12.4)	109 (5.7)	193 (10.1)		
70-74	1,803 (100.0)	719 (39.9)	199 (11.0)	82 (4.5)	115 (6.4)		
75-79	1,060 (100.0)	389 (36.7)	102 (9.6)	37 (3.5)	53 (5.0)		
80歳以上	385 (100.0)	122 (31.7)	36 (9.4)	9 (2.3)	14 (3.6)		
計	9,160 (100.0)	3,501 (38.2)	1,199 (13.1)	541 (5.9)	925 (10.1)		

年齢階級	総数					人(%)	
	-24	25-49	50-74	75-99	100U/L以上		
女							
-29歳	180 (100.0)	13 (7.2)	2 (1.1)	1 (0.6)	0 (0.0)		
30-39	620 (100.0)	88 (14.2)	12 (1.9)	6 (1.0)	5 (0.8)		
40-44	795 (100.0)	115 (14.5)	24 (3.0)	8 (1.0)	9 (1.1)		
45-49	1,185 (100.0)	202 (17.0)	50 (4.2)	15 (1.3)	27 (2.3)		
50-54	1,878 (100.0)	437 (23.3)	114 (6.1)	36 (1.9)	43 (2.3)		
55-59	2,138 (100.0)	592 (27.7)	122 (5.7)	32 (1.5)	44 (2.1)		
60-64	2,878 (100.0)	696 (24.2)	135 (4.7)	47 (1.6)	52 (1.8)		
65-69	3,215 (100.0)	782 (24.3)	141 (4.4)	50 (1.6)	41 (1.3)		
70-74	2,621 (100.0)	552 (21.1)	82 (3.1)	33 (1.3)	19 (0.7)		
75-79	1,383 (100.0)	279 (20.2)	49 (3.5)	18 (1.3)	14 (1.0)		
80歳以上	412 (100.0)	75 (18.2)	9 (2.2)	2 (0.5)	3 (0.7)		
計	17,305 (100.0)	3,831 (22.1)	740 (4.3)	248 (1.4)	257 (1.5)		

表31 血清γ-GTPの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	U/L													
	男					女								
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル					
5			25	50	75	5			25	50	75	95		
29歳以下	86	35.5 (30.2)	14.0	18.0	24.5	43.3	109.7	180	16.1 (9.7)	9.0	11.0	14.0	17.0	33.0
30-39	214	58.7 (64.1)	14.0	25.0	39.0	66.0	161.3	620	20.0 (17.6)	9.0	12.0	15.0	20.0	44.0
40-44	348	71.5 (78.9)	17.0	28.0	44.0	82.8	242.1	795	21.2 (20.0)	10.0	13.0	16.0	22.0	53.0
45-49	465	63.7 (74.9)	16.3	25.5	42.0	74.0	178.4	1,185	25.7 (36.4)	10.0	13.0	17.0	24.0	65.0
50-54	741	66.7 (91.9)	17.0	26.0	40.0	72.0	192.8	1,878	27.3 (26.7)	11.0	14.0	19.0	29.0	70.0
55-59	778	61.7 (86.5)	16.0	25.0	39.0	68.0	164.1	2,138	27.4 (24.3)	11.0	15.0	20.0	30.0	66.0
60-64	1,368	59.4 (94.3)	15.0	24.0	34.0	60.0	168.6	2,878	26.5 (26.5)	11.0	15.0	19.5	28.0	63.1
65-69	1,912	53.5 (87.7)	15.0	22.0	32.0	54.0	156.1	3,215	25.5 (28.0)	11.0	15.0	19.0	27.0	58.2
70-74	1,803	45.6 (75.9)	14.0	20.0	29.0	46.0	115.8	2,621	23.0 (16.7)	11.0	14.0	18.0	25.0	50.0
75-79	1,060	39.1 (48.0)	13.0	18.0	26.0	41.0	100.9	1,383	23.7 (30.2)	10.2	14.0	18.0	25.0	53.0
80歳以上	385	40.8 (96.5)	13.0	17.0	24.0	36.5	81.4	412	21.2 (16.2)	11.0	13.0	17.0	23.0	44.4
計	9,160	53.6 (82.2)	15.0	22.0	32.0	55.0	152.0	17,305	25.0 (25.7)	11.0	14.0	18.0	27.0	59.0

表32 血清クレアチニンの状況 (性・年齢階級別)

年齢階級	総数	人 (%)			
		-0.7	0.8-1.0	1.1-1.4	1.5mg/dl以上
男					
-29歳	86 (100.0)	23 (26.7)	62 (72.1)	1 (1.2)	0 (0.0)
30-39	214 (100.0)	69 (32.2)	144 (67.3)	1 (0.5)	0 (0.0)
40-44	348 (100.0)	106 (30.5)	232 (66.7)	10 (2.9)	0 (0.0)
45-49	465 (100.0)	183 (39.4)	266 (57.2)	16 (3.4)	0 (0.0)
50-54	741 (100.0)	287 (38.7)	437 (59.0)	17 (2.3)	0 (0.0)
55-59	778 (100.0)	288 (37.0)	454 (58.4)	34 (4.4)	2 (0.3)
60-64	1,368 (100.0)	474 (34.6)	823 (60.2)	64 (4.7)	7 (0.5)
65-69	1,912 (100.0)	660 (34.5)	1,151 (60.2)	89 (4.7)	12 (0.6)
70-74	1,803 (100.0)	536 (29.7)	1,109 (61.5)	141 (7.8)	17 (0.9)
75-79	1,060 (100.0)	236 (22.3)	681 (64.2)	128 (12.1)	15 (1.4)
80歳以上	385 (100.0)	73 (19.0)	241 (62.6)	62 (16.1)	9 (2.3)
計	9,160 (100.0)	2,935 (32.0)	5,600 (61.1)	563 (6.1)	62 (0.7)

年齢階級	総数	人 (%)			
		-0.7	0.8-1.0	1.1-1.4	1.5mg/dl以上
女					
-29歳	153 (100.0)	146 (95.4)	7 (4.6)	0 (0.0)	0 (0.0)
30-39	514 (100.0)	485 (94.4)	28 (5.4)	1 (0.2)	0 (0.0)
40-44	795 (100.0)	743 (93.5)	51 (6.4)	1 (0.1)	0 (0.0)
45-49	1,185 (100.0)	1,122 (94.7)	60 (5.1)	2 (0.2)	1 (0.1)
50-54	1,878 (100.0)	1,750 (93.2)	126 (6.7)	0 (0.0)	2 (0.1)
55-59	2,138 (100.0)	1,950 (91.2)	185 (8.7)	2 (0.1)	1 (0.0)
60-64	2,878 (100.0)	2,558 (88.9)	314 (10.9)	6 (0.2)	0 (0.0)
65-69	3,215 (100.0)	2,778 (86.4)	424 (13.2)	9 (0.3)	4 (0.1)
70-74	2,621 (100.0)	2,132 (81.3)	470 (17.9)	16 (0.6)	3 (0.1)
75-79	1,383 (100.0)	1,063 (76.9)	293 (21.2)	20 (1.4)	7 (0.5)
80歳以上	412 (100.0)	292 (70.9)	108 (26.2)	8 (1.9)	4 (1.0)
計	17,172 (100.0)	15,019 (87.5)	2,066 (12.0)	65 (0.4)	22 (0.1)

表33 血清クレアチニンの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	mg/dl													
	男					女								
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル					
5			25	50	75	95			5	25	50	75	95	
29歳以下	86	0.8 (0.1)	0.6	0.7	0.8	0.9	1.0	153	0.6 (0.1)	0.5	0.6	0.6	0.7	0.7
30-39	214	0.8 (0.1)	0.6	0.7	0.8	0.9	0.9	514	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
40-44	348	0.8 (0.1)	0.6	0.7	0.8	0.9	1.0	795	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
45-49	465	0.8 (0.1)	0.6	0.7	0.8	0.9	1.0	1,185	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
50-54	741	0.8 (0.1)	0.6	0.7	0.8	0.9	1.0	1,878	0.6 (0.2)	0.5	0.6	0.6	0.7	0.8
55-59	778	0.8 (0.1)	0.6	0.7	0.8	0.9	1.0	2,138	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
60-64	1,368	0.8 (0.3)	0.6	0.7	0.8	0.9	1.1	2,878	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
65-69	1,912	0.8 (0.2)	0.6	0.7	0.8	0.9	1.1	3,215	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
70-74	1,803	0.8 (0.2)	0.6	0.7	0.8	0.9	1.1	2,621	0.7 (0.1)	0.5	0.6	0.6	0.7	0.9
75-79	1,060	0.9 (0.2)	0.6	0.8	0.8	1.0	1.2	1,383	0.7 (0.2)	0.5	0.6	0.7	0.7	0.9
80歳以上	385	0.9 (0.2)	0.7	0.8	0.9	1.0	1.2	412	0.7 (0.2)	0.5	0.6	0.7	0.8	1.0
計	9,160	0.8 (0.2)	0.6	0.7	0.8	0.9	1.1	17,172	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8

表34 血糖の状況（空腹時採血者のみ、性・年齢階級別）

年齢階級	男						女							
	総数	-89	90-109	110-125	126-139	140-199	200mg/dl以上	総数	-89	90-109	110-125	126-139	140-199	200mg/dl以上
-29歳	14 (100.0)	8 (57.1)	6 (42.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	27 (100.0)	18 (66.7)	9 (33.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	39 (100.0)	20 (51.3)	17 (43.6)	2 (5.1)	0 (0.0)	0 (0.0)	0 (0.0)	86 (100.0)	45 (52.3)	40 (46.5)	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)
40-44	46 (100.0)	13 (28.3)	26 (56.5)	3 (6.5)	2 (4.3)	1 (2.2)	1 (2.2)	95 (100.0)	43 (45.3)	48 (50.5)	2 (2.1)	2 (2.1)	0 (0.0)	0 (0.0)
45-49	80 (100.0)	16 (20.0)	53 (66.3)	5 (6.3)	1 (1.3)	4 (5.0)	1 (1.3)	173 (100.0)	61 (35.3)	92 (53.2)	9 (5.2)	5 (2.9)	5 (2.9)	1 (0.6)
50-54	125 (100.0)	21 (16.8)	67 (53.6)	22 (17.6)	9 (7.2)	4 (3.2)	2 (1.6)	297 (100.0)	103 (34.7)	164 (55.2)	17 (5.7)	7 (2.4)	5 (1.7)	1 (0.3)
55-59	142 (100.0)	12 (8.5)	87 (61.3)	26 (18.3)	10 (7.0)	5 (3.5)	2 (1.4)	323 (100.0)	88 (27.2)	197 (61.0)	23 (7.1)	7 (2.2)	7 (2.2)	1 (0.3)
60-64	227 (100.0)	40 (17.6)	130 (57.3)	41 (18.1)	8 (3.5)	8 (3.5)	0 (0.0)	527 (100.0)	120 (22.8)	319 (60.5)	48 (9.1)	18 (3.4)	17 (3.2)	5 (0.9)
65-69	361 (100.0)	55 (15.2)	206 (57.1)	62 (17.2)	16 (4.4)	16 (4.4)	6 (1.7)	567 (100.0)	131 (23.1)	344 (60.7)	66 (11.6)	15 (2.6)	10 (1.8)	1 (0.2)
70-74	363 (100.0)	54 (14.9)	211 (58.1)	69 (19.0)	16 (4.4)	11 (3.0)	2 (0.6)	508 (100.0)	112 (22.0)	300 (59.1)	53 (10.4)	20 (3.9)	20 (3.9)	3 (0.6)
75-79	214 (100.0)	37 (17.3)	140 (65.4)	23 (10.7)	4 (1.9)	7 (3.3)	3 (1.4)	255 (100.0)	61 (23.9)	139 (54.5)	32 (12.5)	9 (3.5)	11 (4.3)	3 (1.2)
80歳以上	66 (100.0)	11 (16.7)	33 (50.0)	10 (15.2)	6 (9.1)	5 (7.6)	1 (1.5)	68 (100.0)	9 (13.2)	41 (60.3)	9 (13.2)	5 (7.4)	4 (5.9)	0 (0.0)
計	1,677 (100.0)	287 (17.1)	976 (58.2)	263 (15.7)	72 (4.3)	61 (3.6)	18 (1.1)	2,926 (100.0)	791 (27.0)	1,693 (57.9)	260 (8.9)	88 (3.0)	79 (2.7)	15 (0.5)

表35 血糖の平均値、標準偏差および分布（空腹時採血者のみ、性・年齢階級別）

年齢階級	mg/dl													
	男					女								
	人数	パーセンタイル				人数	パーセンタイル							
平均値 (SD)		5	25	50	75		95	平均値 (SD)	5	25	50	75	95	
29歳以下	14	88.2 (5.1)	78.0	84.8	88.0	92.3	95.0	27	86.9 (6.9)	73.0	82.0	87.0	90.0	100.2
30-39	39	92.0 (9.0)	82.0	84.0	89.0	97.0	111.0	86	89.4 (6.8)	78.4	85.0	89.0	93.0	101.7
40-44	46	99.9 (22.9)	79.7	88.8	94.5	102.8	150.0	95	92.2 (9.8)	80.0	86.0	90.0	95.0	109.6
45-49	80	102.0 (24.6)	82.0	91.0	96.0	104.0	154.7	173	99.0 (32.8)	80.7	87.0	93.0	100.5	134.3
50-54	125	105.9 (23.7)	85.3	93.0	100.0	110.5	142.2	297	96.2 (17.5)	80.0	87.5	93.0	101.0	119.1
55-59	142	106.7 (22.2)	87.0	94.0	100.0	113.0	140.6	323	98.0 (16.7)	83.0	89.0	95.0	102.0	123.8
60-64	227	102.8 (15.8)	83.0	92.0	99.0	110.0	132.6	527	100.8 (19.9)	83.0	91.0	97.0	104.0	133.6
65-69	361	106.8 (27.5)	85.0	92.0	101.0	111.0	145.8	567	98.9 (14.2)	83.0	90.0	96.0	104.0	125.0
70-74	363	104.0 (21.4)	83.2	93.0	100.0	111.0	132.8	508	101.3 (20.5)	81.5	90.0	96.0	105.0	136.7
75-79	214	103.2 (26.8)	83.0	92.0	99.0	107.0	135.8	255	101.9 (22.2)	81.0	90.0	97.0	107.0	144.0
80歳以上	66	108.0 (25.4)	82.4	94.0	99.5	117.0	166.6	68	106.4 (22.5)	82.8	92.0	101.0	110.0	168.5
計	1,677	104.2 (23.4)	83.0	92.0	99.0	109.0	138.0	2,926	99.1 (19.4)	82.0	89.0	95.0	103.0	129.0

表36 血糖の状況 (食後4時間以上, 性・年齢階級別)

年齢階級	総数					人 (%)				
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
男										
-29歳	25 (100.0)	10 (40.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
30-39	64 (100.0)	29 (45.3)	5 (7.8)	0 (0.0)	0 (0.0)	0 (0.0)				
40-44	107 (100.0)	62 (57.9)	10 (9.3)	2 (1.9)	2 (1.9)	2 (1.9)				
45-49	149 (100.0)	85 (57.0)	17 (11.4)	3 (2.0)	5 (3.4)	3 (2.0)				
50-54	243 (100.0)	142 (58.4)	33 (13.6)	17 (7.0)	6 (2.5)	3 (1.2)				
55-59	257 (100.0)	151 (58.8)	44 (17.1)	14 (5.4)	6 (2.3)	5 (1.9)				
60-64	448 (100.0)	248 (55.4)	80 (17.9)	20 (4.5)	20 (4.5)	3 (0.7)				
65-69	635 (100.0)	358 (56.4)	106 (16.7)	28 (4.4)	33 (5.2)	9 (1.4)				
70-74	632 (100.0)	358 (56.6)	109 (17.2)	33 (5.2)	26 (4.1)	9 (1.4)				
75-79	373 (100.0)	222 (59.5)	55 (14.7)	15 (4.0)	13 (3.5)	4 (1.1)				
80歳以上	119 (100.0)	62 (52.1)	17 (14.3)	9 (7.6)	9 (7.6)	2 (1.7)				
計	3,052 (100.0)	1,727 (56.6)	476 (15.6)	141 (4.6)	120 (3.9)	40 (1.3)				

年齢階級	総数					人 (%)				
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
女										
-29歳	48 (100.0)	19 (39.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
30-39	195 (100.0)	99 (50.8)	3 (1.5)	0 (0.0)	1 (0.5)	1 (0.5)				
40-44	241 (100.0)	141 (58.5)	5 (2.1)	2 (0.8)	2 (0.8)	0 (0.0)				
45-49	382 (100.0)	228 (59.7)	17 (4.5)	7 (1.8)	6 (1.6)	2 (0.5)				
50-54	645 (100.0)	377 (58.4)	49 (7.6)	14 (2.2)	9 (1.4)	3 (0.5)				
55-59	684 (100.0)	433 (63.3)	52 (7.6)	16 (2.3)	13 (1.9)	1 (0.1)				
60-64	973 (100.0)	608 (62.5)	103 (10.6)	34 (3.5)	23 (2.4)	9 (0.9)				
65-69	1,051 (100.0)	651 (61.9)	126 (12.0)	29 (2.8)	19 (1.8)	3 (0.3)				
70-74	899 (100.0)	543 (60.4)	102 (11.3)	36 (4.0)	36 (4.0)	7 (0.8)				
75-79	456 (100.0)	256 (56.1)	68 (14.9)	23 (5.0)	17 (3.7)	7 (1.5)				
80歳以上	129 (100.0)	77 (59.7)	20 (15.5)	7 (5.4)	10 (7.8)	0 (0.0)				
計	5,703 (100.0)	3,432 (60.2)	545 (9.6)	168 (2.9)	136 (2.4)	33 (0.6)				

表37 血糖の平均値、標準偏差および分布（食後4時間以上，性・年齢階級別）

年齢階級	mg/dl													
	男					女								
	人数	平均値 (SD)	パーセンタイル				人数	平均値 (SD)	パーセンタイル					
5			25	50	75	95			5	25	50	75	95	
29歳以下	25	87.6 (8.5)	66.1	84.0	87.0	92.5	104.7	48	88.5 (7.3)	77.4	83.3	88.0	93.8	103.2
30-39	64	93.2 (9.2)	82.0	86.0	90.0	101.0	110.8	195	91.3 (13.9)	78.0	85.0	90.0	95.0	103.2
40-44	107	101.8 (41.2)	79.4	89.0	95.0	102.0	133.4	241	92.6 (11.2)	80.0	87.0	91.0	96.0	106.9
45-49	149	104.0 (40.2)	82.0	90.0	95.0	104.5	150.5	382	97.0 (24.0)	82.0	88.0	93.0	100.0	118.9
50-54	243	103.1 (20.9)	84.0	91.0	97.0	109.0	132.8	645	97.4 (17.3)	81.0	89.0	95.0	102.0	119.0
55-59	257	105.2 (25.5)	84.0	92.0	99.0	111.0	137.1	684	98.0 (14.9)	83.0	90.0	95.0	103.0	122.0
60-64	448	104.8 (20.2)	83.5	93.0	100.5	111.0	142.1	973	101.3 (20.7)	84.0	91.0	97.0	105.0	130.0
65-69	635	106.4 (25.7)	84.0	93.0	101.0	111.0	149.4	1,051	99.9 (15.9)	83.0	91.0	97.0	105.0	125.0
70-74	632	105.6 (24.1)	83.0	93.0	100.0	111.0	143.4	899	102.3 (20.8)	84.0	91.0	97.0	106.0	139.0
75-79	373	103.8 (24.7)	83.0	92.0	99.0	108.0	134.0	456	104.4 (25.0)	83.0	91.0	99.0	110.0	143.0
80歳以上	119	108.6 (30.7)	81.0	94.0	102.0	114.0	162.0	129	107.3 (22.1)	83.0	93.0	102.0	110.5	154.0
計	3,052	104.7 (25.8)	83.0	92.0	99.0	110.0	142.4	5,703	99.6 (19.3)	82.0	90.0	96.0	104.0	128.0

表38 HbA1cの状況(性・年齢階級別)

年齢階級	総数					人 (%)		
	-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上			
男								
-29歳	86 (100.0)	69 (80.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	214 (100.0)	174 (81.3)	6 (2.8)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)
40-44	348 (100.0)	291 (83.6)	12 (3.4)	7 (2.0)	5 (1.4)	5 (1.4)	5 (1.4)	5 (1.4)
45-49	465 (100.0)	380 (81.7)	33 (7.1)	2 (0.4)	12 (2.6)	12 (2.6)	12 (2.6)	12 (2.6)
50-54	741 (100.0)	579 (78.1)	74 (10.0)	16 (2.2)	13 (1.8)	13 (1.8)	13 (1.8)	13 (1.8)
55-59	778 (100.0)	598 (76.9)	97 (12.5)	25 (3.2)	15 (1.9)	15 (1.9)	15 (1.9)	15 (1.9)
60-64	1,368 (100.0)	1,067 (78.0)	169 (12.4)	42 (3.1)	27 (2.0)	27 (2.0)	27 (2.0)	27 (2.0)
65-69	1,912 (100.0)	1,423 (74.4)	281 (14.7)	64 (3.3)	42 (2.2)	42 (2.2)	42 (2.2)	42 (2.2)
70-74	1,802 (100.0)	1,345 (74.6)	276 (15.3)	65 (3.6)	36 (2.0)	36 (2.0)	36 (2.0)	36 (2.0)
75-79	1,060 (100.0)	797 (75.2)	158 (14.9)	30 (2.8)	18 (1.7)	18 (1.7)	18 (1.7)	18 (1.7)
80歳以上	385 (100.0)	287 (74.5)	57 (14.8)	16 (4.2)	5 (1.3)	5 (1.3)	5 (1.3)	5 (1.3)
計	9,159 (100.0)	7,010 (76.5)	1,163 (12.7)	268 (2.9)	174 (1.9)	174 (1.9)	174 (1.9)	174 (1.9)

年齢階級	総数					人 (%)		
	-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上			
女								
-29歳	180 (100.0)	137 (76.1)	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	620 (100.0)	510 (82.3)	7 (1.1)	0 (0.0)	1 (0.2)	1 (0.2)	1 (0.2)	1 (0.2)
40-44	795 (100.0)	646 (81.3)	23 (2.9)	1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)
45-49	1,185 (100.0)	984 (83.0)	67 (5.7)	9 (0.8)	13 (1.1)	13 (1.1)	13 (1.1)	13 (1.1)
50-54	1,878 (100.0)	1,556 (82.9)	175 (9.3)	22 (1.2)	18 (1.0)	18 (1.0)	18 (1.0)	18 (1.0)
55-59	2,138 (100.0)	1,777 (83.1)	230 (10.8)	22 (1.0)	33 (1.5)	33 (1.5)	33 (1.5)	33 (1.5)
60-64	2,877 (100.0)	2,321 (80.7)	372 (12.9)	49 (1.7)	42 (1.5)	42 (1.5)	42 (1.5)	42 (1.5)
65-69	3,213 (100.0)	2,541 (79.1)	463 (14.4)	71 (2.2)	41 (1.3)	41 (1.3)	41 (1.3)	41 (1.3)
70-74	2,621 (100.0)	2,039 (77.8)	421 (16.1)	70 (2.7)	29 (1.1)	29 (1.1)	29 (1.1)	29 (1.1)
75-79	1,383 (100.0)	1,054 (76.2)	240 (17.4)	40 (2.9)	19 (1.4)	19 (1.4)	19 (1.4)	19 (1.4)
80歳以上	411 (100.0)	323 (78.6)	58 (14.1)	15 (3.6)	7 (1.7)	7 (1.7)	7 (1.7)	7 (1.7)
計	17,301 (100.0)	13,888 (80.3)	2,057 (11.9)	299 (1.7)	204 (1.2)	204 (1.2)	204 (1.2)	204 (1.2)

表39 HbA1cの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女							
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル					
			5	25	50	75			95	5	25	50	75	95
29歳以下	86	4.7 (0.3)	4.2	4.5	4.7	4.9	5.2	180	4.7 (0.3)	4.2	4.5	4.6	4.8	5.1
30-39	214	4.8 (0.5)	4.3	4.6	4.8	5.0	5.4	620	4.8 (0.4)	4.2	4.5	4.7	4.9	5.3
40-44	348	4.9 (0.8)	4.3	4.7	4.9	5.0	5.8	795	4.8 (0.4)	4.3	4.5	4.8	5.0	5.3
45-49	465	5.0 (0.8)	4.4	4.7	4.9	5.1	5.9	1,185	4.9 (0.6)	4.3	4.7	4.9	5.1	5.6
50-54	741	5.1 (0.7)	4.4	4.7	5.0	5.2	6.1	1,878	5.0 (0.7)	4.4	4.8	5.0	5.2	5.7
55-59	778	5.2 (0.8)	4.4	4.8	5.0	5.3	6.5	2,138	5.1 (0.6)	4.5	4.8	5.0	5.3	5.9
60-64	1,368	5.2 (0.7)	4.5	4.8	5.0	5.3	6.5	2,877	5.2 (0.7)	4.5	4.8	5.0	5.3	6.0
65-69	1,912	5.2 (0.8)	4.4	4.8	5.0	5.4	6.6	3,213	5.2 (0.6)	4.5	4.8	5.1	5.3	6.0
70-74	1,802	5.2 (0.8)	4.5	4.8	5.0	5.4	6.6	2,621	5.2 (0.6)	4.5	4.9	5.1	5.4	6.2
75-79	1,060	5.2 (0.7)	4.4	4.8	5.1	5.3	6.4	1,383	5.2 (0.6)	4.6	4.9	5.1	5.4	6.3
80歳以上	385	5.2 (0.6)	4.4	4.8	5.1	5.4	6.6	411	5.2 (0.7)	4.5	4.8	5.1	5.3	6.5
計	9,159	5.1 (0.7)	4.4	4.8	5.0	5.3	6.4	17,301	5.1 (0.6)	4.5	4.8	5.0	5.3	6.0

表40 赤血球数の状況 (性・年齢階級別)

年齢階級	人 (%)					
	総数	-399	400-449	450-499	500-549	550×10 <sup>4</sup> /μ以上
-29歳	86 (100.0)	0 (0.0)	5 (5.8)	27 (31.4)	46 (53.5)	8 (9.3)
30-39	214 (100.0)	0 (0.0)	17 (7.9)	97 (45.3)	81 (37.9)	19 (8.9)
40-44	348 (100.0)	4 (1.1)	52 (14.9)	171 (49.1)	109 (31.3)	12 (3.4)
45-49	465 (100.0)	7 (1.5)	91 (19.6)	213 (45.8)	141 (30.3)	13 (2.8)
50-54	741 (100.0)	16 (2.2)	138 (18.6)	380 (51.3)	189 (25.5)	18 (2.4)
55-59	778 (100.0)	26 (3.3)	180 (23.1)	421 (54.1)	136 (17.5)	15 (1.9)
60-64	1,368 (100.0)	69 (5.0)	388 (28.4)	637 (46.6)	259 (18.9)	15 (1.1)
65-69	1,912 (100.0)	127 (6.6)	591 (30.9)	881 (46.1)	287 (15.0)	26 (1.4)
70-74	1,803 (100.0)	186 (10.3)	677 (37.5)	720 (39.9)	204 (11.3)	16 (0.9)
75-79	1,060 (100.0)	140 (13.2)	394 (37.2)	403 (38.0)	113 (10.7)	10 (0.9)
80歳以上	385 (100.0)	86 (22.3)	171 (44.4)	101 (26.2)	25 (6.5)	2 (0.5)
計	9,160 (100.0)	661 (7.2)	2,704 (29.5)	4,051 (44.2)	1,590 (17.4)	154 (1.7)

年齢階級	人 (%)					
	総数	-399	400-449	450-499	500-549	550×10 <sup>4</sup> /μ以上
-29歳	180 (100.0)	22 (12.2)	101 (56.1)	51 (28.3)	6 (3.3)	0 (0.0)
30-39	620 (100.0)	123 (19.8)	340 (54.8)	145 (23.4)	11 (1.8)	1 (0.2)
40-44	795 (100.0)	138 (17.4)	467 (58.7)	177 (22.3)	13 (1.6)	0 (0.0)
45-49	1,185 (100.0)	224 (18.9)	640 (54.0)	283 (23.9)	37 (3.1)	1 (0.1)
50-54	1,878 (100.0)	300 (16.0)	982 (52.3)	524 (27.9)	66 (3.5)	6 (0.3)
55-59	2,138 (100.0)	334 (15.6)	1,184 (55.4)	560 (26.2)	58 (2.7)	2 (0.1)
60-64	2,878 (100.0)	595 (20.7)	1,540 (53.5)	682 (23.7)	60 (2.1)	1 (0.0)
65-69	3,215 (100.0)	761 (23.7)	1,690 (52.6)	709 (22.1)	54 (1.7)	1 (0.0)
70-74	2,621 (100.0)	759 (29.0)	1,301 (49.6)	509 (19.4)	51 (1.9)	1 (0.0)
75-79	1,383 (100.0)	481 (34.8)	637 (46.1)	242 (17.5)	21 (1.5)	2 (0.1)
80歳以上	412 (100.0)	160 (38.8)	188 (45.6)	61 (14.8)	3 (0.7)	0 (0.0)
計	17,305 (100.0)	3,897 (22.5)	9,070 (52.4)	3,943 (22.8)	380 (2.2)	15 (0.1)

表41 赤血球数の平均値、標準偏差および分布 (性・年齢階級別)

10<sup>4</sup>/μl

年齢階級	男					女								
	人数	平均値 (SD)	5	25	50	75	95	人数	平均値 (SD)	5	25	50	75	95
29歳以下	86	506.4 (35.5)	442.4	481.0	507.0	534.0	560.7	180	434.4 (31.6)	386.1	412.3	435.0	454.0	494.6
30-39	214	498.9 (36.0)	434.8	475.0	497.5	525.3	558.8	620	428.3 (33.2)	377.0	404.3	425.0	450.0	486.0
40-44	348	486.6 (36.8)	426.5	460.3	488.0	509.0	546.6	795	428.2 (31.7)	375.0	407.0	429.0	448.0	481.2
45-49	465	481.9 (37.4)	420.3	457.5	485.0	507.0	541.7	1,185	430.3 (36.0)	376.0	406.0	429.0	452.0	490.0
50-54	741	477.7 (36.8)	416.0	454.0	479.0	503.0	537.9	1,878	434.1 (35.5)	378.0	410.0	433.0	457.0	493.0
55-59	778	470.1 (36.1)	410.0	447.0	470.0	493.0	530.1	2,138	432.6 (33.6)	378.0	410.0	432.0	454.0	491.0
60-64	1,368	465.5 (39.3)	399.0	440.3	466.0	493.0	524.0	2,878	428.2 (35.3)	372.0	405.0	428.0	451.0	485.0
65-69	1,912	460.2 (41.6)	393.7	434.0	462.0	489.0	525.0	3,215	425.0 (34.9)	370.0	401.0	424.0	448.0	483.0
70-74	1,803	451.1 (43.7)	382.0	425.0	452.0	480.0	520.0	2,621	420.2 (36.7)	361.0	396.0	420.0	445.0	480.0
75-79	1,060	447.6 (43.5)	375.1	421.0	449.0	475.8	518.0	1,383	415.4 (39.1)	353.0	388.0	415.0	442.0	481.0
80歳以上	385	430.1 (46.3)	350.3	402.0	430.0	462.0	507.7	412	407.6 (39.8)	340.3	382.0	409.0	434.0	472.0
計	9,160	462.2 (43.3)	391.0	435.0	464.0	492.0	530.0	17,305	426.3 (36.0)	369.0	402.0	426.0	450.0	486.0

表42 ヘモグロビン値の状況 (性・年齢階級別)

年齢階級	総数	人 (%)						
		-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上	
男								
-29歳	86 (100.0)	0 (0.0)	0 (0.0)	4 (4.7)	24 (27.9)	31 (36.0)	27 (31.4)	
30-39	214 (100.0)	0 (0.0)	2 (0.9)	13 (6.1)	37 (17.3)	98 (45.8)	64 (29.9)	
40-44	348 (100.0)	0 (0.0)	6 (1.7)	26 (7.5)	92 (26.4)	149 (42.8)	75 (21.6)	
45-49	465 (100.0)	3 (0.6)	11 (2.4)	36 (7.7)	157 (33.8)	167 (35.9)	91 (19.6)	
50-54	741 (100.0)	6 (0.8)	12 (1.6)	76 (10.3)	239 (32.3)	255 (34.4)	153 (20.6)	
55-59	778 (100.0)	7 (0.9)	23 (3.0)	103 (13.2)	277 (35.6)	250 (32.1)	118 (15.2)	
60-64	1,368 (100.0)	14 (1.0)	57 (4.2)	203 (14.8)	467 (34.1)	436 (31.9)	191 (14.0)	
65-69	1,912 (100.0)	50 (2.6)	111 (5.8)	331 (17.3)	621 (32.5)	540 (28.2)	259 (13.5)	
70-74	1,803 (100.0)	68 (3.8)	132 (7.3)	399 (22.1)	592 (32.8)	428 (23.7)	184 (10.2)	
75-79	1,060 (100.0)	43 (4.1)	115 (10.8)	262 (24.7)	327 (30.8)	223 (21.0)	90 (8.5)	
80歳以上	385 (100.0)	44 (11.4)	54 (14.0)	117 (30.4)	104 (27.0)	49 (12.7)	17 (4.4)	
計	9,160 (100.0)	235 (2.6)	523 (5.7)	1,570 (17.1)	2,937 (32.1)	2,626 (28.7)	1,269 (13.9)	

年齢階級	総数	人 (%)					
		-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上
女							
-29歳	180 (100.0)	31 (17.2)	61 (33.9)	65 (36.1)	20 (11.1)	3 (1.7)	0 (0.0)
30-39	620 (100.0)	133 (21.5)	213 (34.4)	196 (31.6)	65 (10.5)	13 (2.1)	0 (0.0)
40-44	795 (100.0)	205 (25.8)	236 (29.7)	244 (30.7)	103 (13.0)	7 (0.9)	0 (0.0)
45-49	1,185 (100.0)	285 (24.1)	329 (27.8)	357 (30.1)	186 (15.7)	25 (2.1)	3 (0.3)
50-54	1,878 (100.0)	242 (12.9)	541 (28.8)	695 (37.0)	348 (18.5)	48 (2.6)	4 (0.2)
55-59	2,138 (100.0)	177 (8.3)	636 (29.7)	860 (40.2)	389 (18.2)	73 (3.4)	3 (0.1)
60-64	2,878 (100.0)	312 (10.8)	856 (29.7)	1,110 (38.6)	514 (17.9)	82 (2.8)	4 (0.1)
65-69	3,215 (100.0)	372 (11.6)	1,007 (31.3)	1,217 (37.9)	527 (16.4)	88 (2.7)	4 (0.1)
70-74	2,621 (100.0)	389 (14.8)	845 (32.2)	907 (34.6)	411 (15.7)	65 (2.5)	4 (0.2)
75-79	1,383 (100.0)	283 (20.5)	435 (31.5)	437 (31.6)	184 (13.3)	42 (3.0)	2 (0.1)
80歳以上	412 (100.0)	111 (26.9)	127 (30.8)	125 (30.3)	41 (10.0)	8 (1.9)	0 (0.0)
計	17,305 (100.0)	2,540 (14.7)	5,286 (30.5)	6,213 (35.9)	2,788 (16.1)	454 (2.6)	24 (0.1)