

表33 血糖の状況（空腹時採血者のみ、性・年齢階級別）

男 年齢階級	総数					人 (%)				
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
-29歳	3 (100.0)	1 (33.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
30-39	4 (66.7)	1 (16.7)	1 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)				
40-44	7 (46.7)	6 (40.0)	0 (0.0)	1 (6.7)	1 (6.7)	0 (0.0)				
45-49	7 (29.2)	14 (58.3)	0 (0.0)	1 (4.2)	1 (4.2)	1 (4.2)				
50-54	8 (26.7)	18 (60.0)	2 (6.7)	2 (6.7)	0 (0.0)	0 (0.0)				
55-59	5 (17.2)	15 (51.7)	4 (13.8)	3 (10.3)	1 (3.4)	1 (3.4)				
60-64	17 (28.3)	30 (50.0)	8 (13.3)	3 (5.0)	2 (3.3)	0 (0.0)				
65-69	22 (24.2)	47 (51.6)	15 (16.5)	4 (4.4)	2 (2.2)	1 (1.1)				
70-74	31 (30.4)	54 (52.9)	9 (8.8)	6 (5.9)	2 (2.0)	0 (0.0)				
75-79	15 (23.1)	39 (60.0)	8 (12.3)	0 (0.0)	1 (1.5)	2 (3.1)				
80歳以上	5 (31.3)	8 (50.0)	0 (0.0)	2 (12.5)	1 (6.3)	0 (0.0)				
計	123 (27.9)	233 (52.8)	47 (10.7)	22 (5.0)	11 (2.5)	5 (1.1)				

女 年齢階級	総数					人 (%)				
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
-29歳	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
30-39	5 (55.6)	4 (44.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
40-44	20 (69.0)	9 (31.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
45-49	21 (42.9)	19 (38.8)	3 (6.1)	1 (2.0)	4 (8.2)	1 (2.0)				
50-54	59 (54.6)	45 (41.7)	1 (0.9)	2 (1.9)	1 (0.9)	0 (0.0)				
55-59	47 (42.7)	58 (52.7)	2 (1.8)	1 (0.9)	2 (1.8)	0 (0.0)				
60-64	44 (33.6)	79 (60.3)	2 (1.5)	3 (2.3)	2 (1.5)	1 (0.8)				
65-69	68 (37.4)	96 (52.7)	15 (8.2)	2 (1.1)	1 (0.5)	0 (0.0)				
70-74	61 (33.3)	98 (53.6)	9 (4.9)	8 (4.4)	6 (3.3)	1 (0.5)				
75-79	38 (39.2)	47 (48.5)	7 (7.2)	2 (2.1)	2 (2.1)	1 (1.0)				
80歳以上	6 (25.0)	15 (62.5)	2 (8.3)	1 (4.2)	0 (0.0)	0 (0.0)				
計	370 (40.1)	470 (50.9)	41 (4.4)	20 (2.2)	18 (2.0)	4 (0.4)				

表34 血糖の平均値、標準偏差および分布 (空腹時採血者のみ、性・年齢階級別)

年齢階級	男						女								
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50	75	95	
29歳以下	3	88.3 (3.2)	86.0	87.0	87.0	87.0	92.0	1	83.0						
30-39	6	90.3 (13.1)	81.0	81.8	84.5	100.0	115.0	9	90.6 (7.1)	82.0	86.0	89.0	94.0	106.0	
40-44	15	96.3 (22.1)	69.0	83.0	91.0	102.0	131.0	29	87.3 (5.4)	79.0	83.0	87.0	91.5	98.5	
45-49	24	103.3 (36.5)	81.3	87.0	92.0	99.3	229.8	49	106.2 (58.2)	77.0	85.0	92.0	101.0	178.0	
50-54	30	96.8 (11.5)	82.7	89.0	94.0	101.3	128.7	108	90.7 (10.0)	79.5	85.0	89.0	94.0	108.6	
55-59	29	107.0 (28.7)	80.5	90.5	100.0	117.0	193.5	110	93.6 (12.8)	81.0	87.0	90.0	98.0	114.4	
60-64	60	100.0 (15.5)	82.0	88.3	96.0	107.0	136.7	131	95.3 (15.0)	80.6	88.0	93.0	99.0	117.0	
65-69	91	102.2 (22.6)	82.8	90.0	96.0	109.0	138.0	182	94.7 (12.0)	80.2	87.0	93.0	100.0	117.6	
70-74	102	98.1 (14.9)	78.3	89.0	96.0	103.0	129.7	183	98.0 (20.2)	79.0	87.0	93.0	101.0	134.8	
75-79	65	104.2 (41.8)	79.6	90.5	96.0	104.0	143.5	97	96.1 (22.0)	80.0	86.0	91.0	99.0	130.8	
80歳以上	16	102.1 (26.8)	76.0	85.5	96.0	103.5	132.0	24	98.5 (13.7)	79.5	89.3	94.5	106.8	132.5	
計	441	100.8 (25.0)	81.0	89.0	95.0	105.0	135.0	923	95.4 (20.4)	80.0	87.0	92.0	99.0	123.8	

表35 血糖の状況 (食後4時間以上, 性・年齢階級別)

年齢階級	総数						人 (%)		
	-89	90-109	110-125	126-139	140-199	200mg/dl以上			
男									
-29歳	6 (100.0)	2 (33.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	15 (100.0)	4 (26.7)	2 (13.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
40-44	49 (100.0)	27 (55.1)	3 (6.1)	1 (2.0)	2 (4.1)	1 (2.0)	1 (2.0)	1 (1.7)	1 (1.7)
45-49	60 (100.0)	29 (48.3)	5 (8.3)	1 (1.7)	2 (3.3)	1 (1.7)	1 (1.7)	0 (0.0)	0 (0.0)
50-54	84 (100.0)	55 (65.5)	4 (4.8)	6 (7.1)	0 (0.0)	0 (0.0)	0 (0.0)	2 (2.8)	2 (2.8)
55-59	71 (100.0)	35 (49.3)	10 (14.1)	4 (5.6)	1 (1.4)	1 (1.4)	1 (1.4)	0 (0.0)	0 (0.0)
60-64	142 (100.0)	75 (52.8)	20 (14.1)	7 (4.9)	4 (2.8)	0 (0.0)	0 (0.0)	3 (1.4)	3 (1.4)
65-69	211 (100.0)	113 (53.6)	32 (15.2)	6 (2.8)	10 (4.7)	2 (0.9)	2 (0.9)	3 (2.2)	3 (2.2)
70-74	224 (100.0)	125 (55.8)	23 (10.3)	9 (4.0)	8 (3.6)	0 (0.0)	0 (0.0)	2 (5.6)	2 (5.6)
75-79	136 (100.0)	78 (57.4)	16 (11.8)	3 (2.2)	5 (3.7)	0 (0.0)	0 (0.0)	12 (1.2)	12 (1.2)
80歳以上	36 (100.0)	20 (55.6)	3 (8.3)	2 (5.6)	2 (5.6)	0 (0.0)	0 (0.0)		
計	1,034 (100.0)	563 (54.4)	118 (11.4)	39 (3.8)	34 (3.3)	12 (1.2)	12 (1.2)		

年齢階級	総数						人 (%)		
	-89	90-109	110-125	126-139	140-199	200mg/dl以上			
女									
-29歳	3 (100.0)	2 (66.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	36 (100.0)	16 (44.4)	0 (0.0)	0 (0.0)	1 (2.8)	1 (2.8)	1 (2.8)	0 (0.0)	0 (0.0)
40-44	85 (100.0)	40 (47.1)	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.8)	1 (0.8)
45-49	123 (100.0)	63 (51.2)	5 (4.1)	2 (1.6)	4 (3.3)	1 (0.5)	1 (0.5)	0 (0.0)	0 (0.0)
50-54	217 (100.0)	97 (44.7)	11 (5.1)	3 (1.4)	2 (0.9)	1 (0.5)	1 (0.5)	2 (0.7)	2 (0.7)
55-59	239 (100.0)	132 (55.2)	8 (3.3)	4 (1.7)	5 (2.1)	0 (0.0)	0 (0.0)	4 (1.1)	4 (1.1)
60-64	303 (100.0)	185 (61.1)	16 (5.3)	6 (2.0)	3 (1.0)	2 (0.7)	2 (0.7)	8 (2.2)	8 (2.2)
65-69	392 (100.0)	231 (58.9)	35 (8.9)	6 (1.5)	4 (1.0)	1 (0.3)	1 (0.3)	3 (1.6)	3 (1.6)
70-74	357 (100.0)	211 (59.1)	24 (6.7)	11 (3.1)	8 (2.2)	2 (1.1)	2 (1.1)	40 (2.0)	40 (2.0)
75-79	183 (100.0)	102 (55.7)	14 (7.7)	7 (3.8)	3 (1.6)	0 (0.0)	0 (0.0)	12 (1.2)	12 (1.2)
80歳以上	54 (100.0)	36 (66.7)	6 (11.1)	1 (1.9)	0 (0.0)	0 (0.0)	0 (0.0)		
計	1,992 (100.0)	1,115 (56.0)	120 (6.0)	40 (2.0)	30 (1.5)	12 (0.6)	12 (0.6)		

表36 血糖の平均値、標準偏差および分布（食後4時間以上、性・年齢階級別）

年齢階級	男						女							
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル				
			5	25	50	75	95			5	25	50	75	95
29歳以下	6	88.0 (3.3)	84.0	85.5	87.0	92.0	92.0	3	91.7 (7.6)	83.0	83.0	95.0	96.0	97.0
30-39	15	91.5 (10.9)	80.0	83.0	87.0	99.0	112.0	36	95.6 (28.6)	76.9	84.0	89.5	97.8	157.9
40-44	49	105.7 (57.3)	79.0	86.5	95.0	102.0	158.5	85	89.0 (7.2)	78.3	83.5	89.0	93.5	101.7
45-49	60	98.4 (25.9)	81.0	87.0	91.0	98.5	151.2	123	98.1 (37.7)	80.2	86.0	92.0	99.0	137.0
50-54	84	96.6 (12.3)	78.8	90.0	94.0	100.8	126.8	217	92.9 (14.5)	79.0	85.0	90.0	96.0	116.0
55-59	71	102.1 (24.6)	78.0	89.0	95.0	108.0	146.8	239	94.7 (13.2)	81.0	87.0	91.0	99.0	120.0
60-64	142	100.1 (14.9)	82.2	89.0	97.0	107.3	131.9	303	96.5 (15.3)	81.0	88.0	94.0	101.0	120.4
65-69	211	103.6 (24.7)	81.6	91.0	97.0	109.0	153.8	392	97.0 (15.4)	81.7	88.0	94.0	102.0	120.0
70-74	224	100.7 (19.9)	81.0	89.0	97.0	104.0	133.8	357	99.1 (21.5)	81.0	89.0	95.0	102.0	131.3
75-79	136	103.7 (34.8)	80.7	90.0	96.0	104.0	145.5	183	98.6 (23.6)	81.0	88.0	93.0	101.0	129.8
80歳以上	36	101.1 (21.1)	76.9	88.0	97.0	104.0	155.1	54	98.6 (12.3)	79.0	90.0	97.0	107.0	124.3
計	1,034	101.3 (25.8)	81.0	89.0	96.0	105.0	136.3	1,992	96.4 (19.3)	80.7	87.3	93.0	100.0	122.0

表37 HbA1cの状況 (性・年齢階級別)

男 年齢階級	総数					人 (%)		
	-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上			
-29歳	41 (100.0)	32 (78.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	96 (100.0)	76 (79.2)	2 (2.1)	1 (1.0)	1 (1.0)	1 (1.0)	1 (1.0)	1 (1.0)
40-44	155 (100.0)	128 (82.6)	4 (2.6)	5 (3.2)	2 (1.3)	2 (1.3)	2 (1.3)	2 (1.3)
45-49	224 (100.0)	183 (81.7)	13 (5.8)	0 (0.0)	5 (2.2)	5 (2.2)	5 (2.2)	5 (2.2)
50-54	296 (100.0)	231 (78.0)	27 (9.1)	6 (2.0)	7 (2.4)	7 (2.4)	7 (2.4)	7 (2.4)
55-59	253 (100.0)	184 (72.7)	27 (10.7)	9 (3.6)	8 (3.2)	8 (3.2)	8 (3.2)	8 (3.2)
60-64	443 (100.0)	339 (76.5)	49 (11.1)	16 (3.6)	10 (2.3)	10 (2.3)	10 (2.3)	10 (2.3)
65-69	657 (100.0)	473 (72.0)	94 (14.3)	23 (3.5)	11 (1.7)	11 (1.7)	11 (1.7)	11 (1.7)
70-74	676 (100.0)	511 (75.6)	94 (13.9)	21 (3.1)	12 (1.8)	12 (1.8)	12 (1.8)	12 (1.8)
75-79	371 (100.0)	279 (75.2)	50 (13.5)	7 (1.9)	9 (2.4)	9 (2.4)	9 (2.4)	9 (2.4)
80歳以上	150 (100.0)	115 (76.7)	16 (10.7)	7 (4.7)	1 (0.7)	1 (0.7)	1 (0.7)	1 (0.7)
計	3,362 (100.0)	2,551 (75.9)	376 (11.2)	95 (2.8)	66 (2.0)	66 (2.0)	66 (2.0)	66 (2.0)

女 年齢階級	総数					人 (%)		
	-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上			
-29歳	41 (100.0)	27 (65.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	182 (100.0)	138 (75.8)	1 (0.5)	0 (0.0)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)
40-44	305 (100.0)	227 (74.4)	8 (2.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
45-49	416 (100.0)	341 (82.0)	19 (4.6)	4 (1.0)	9 (2.2)	9 (2.2)	9 (2.2)	9 (2.2)
50-54	622 (100.0)	513 (82.5)	44 (7.1)	8 (1.3)	9 (1.4)	9 (1.4)	9 (1.4)	9 (1.4)
55-59	689 (100.0)	561 (81.4)	67 (9.7)	10 (1.5)	14 (2.0)	14 (2.0)	14 (2.0)	14 (2.0)
60-64	936 (100.0)	779 (83.2)	86 (9.2)	18 (1.9)	10 (1.1)	10 (1.1)	10 (1.1)	10 (1.1)
65-69	1,138 (100.0)	902 (79.3)	140 (12.3)	29 (2.5)	18 (1.6)	18 (1.6)	18 (1.6)	18 (1.6)
70-74	986 (100.0)	783 (79.4)	122 (12.4)	33 (3.3)	13 (1.3)	13 (1.3)	13 (1.3)	13 (1.3)
75-79	560 (100.0)	441 (78.8)	74 (13.2)	17 (3.0)	7 (1.3)	7 (1.3)	7 (1.3)	7 (1.3)
80歳以上	171 (100.0)	137 (80.1)	22 (12.9)	6 (3.5)	2 (1.2)	2 (1.2)	2 (1.2)	2 (1.2)
計	6,046 (100.0)	4,849 (80.2)	583 (9.6)	125 (2.1)	83 (1.4)	83 (1.4)	83 (1.4)	83 (1.4)

表38 HbA1cの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女							
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル				
			5	25	50	75	95			5	25	50	75	95
29歳以下	41	4.6 (0.3)	4.2	4.5	4.6	4.8	5.1	41	4.6 (0.3)	4.1	4.4	4.6	4.8	5.1
30-39	96	4.8 (0.6)	4.3	4.5	4.8	5.0	5.4	182	4.7 (0.6)	4.2	4.5	4.7	4.9	5.2
40-44	155	4.9 (1.0)	4.3	4.6	4.8	5.0	6.2	305	4.7 (0.3)	4.2	4.5	4.7	4.9	5.3
45-49	224	5.0 (0.7)	4.3	4.7	4.9	5.1	5.9	416	5.0 (0.8)	4.3	4.6	4.9	5.1	5.8
50-54	296	5.1 (0.8)	4.4	4.7	4.9	5.2	6.1	622	5.0 (0.8)	4.4	4.7	4.9	5.1	5.9
55-59	253	5.1 (0.9)	4.4	4.7	4.9	5.3	6.8	689	5.1 (0.7)	4.4	4.8	5.0	5.2	6.0
60-64	443	5.1 (0.8)	4.4	4.7	4.9	5.3	6.7	936	5.1 (0.6)	4.5	4.8	5.0	5.2	5.8
65-69	657	5.1 (0.8)	4.4	4.7	5.0	5.3	6.5	1,138	5.1 (0.7)	4.5	4.8	5.0	5.3	6.1
70-74	676	5.1 (0.7)	4.4	4.7	5.0	5.3	6.4	986	5.2 (0.7)	4.5	4.8	5.0	5.3	6.4
75-79	371	5.1 (0.8)	4.4	4.7	5.0	5.3	6.2	560	5.2 (0.7)	4.5	4.8	5.0	5.3	6.3
80歳以上	150	5.1 (0.6)	4.3	4.7	5.0	5.3	6.6	171	5.2 (0.6)	4.5	4.8	5.1	5.3	6.5
計	3,362	5.1 (0.8)	4.4	4.7	4.9	5.2	6.4	6,046	5.1 (0.7)	4.4	4.7	5.0	5.2	6.1

表39 赤血球数の状況 (性・年齢階級別)

年齢階級	人 (%)					
	総数	-399	400-449	450-499	500-549	550 × 10 ⁴ / μl 以上
男						
-29歳	41 (100.0)	0 (0.0)	3 (7.3)	15 (36.6)	19 (46.3)	4 (9.8)
30-39	96 (100.0)	0 (0.0)	12 (12.5)	42 (43.8)	36 (37.5)	6 (6.3)
40-44	155 (100.0)	1 (0.6)	25 (16.1)	74 (47.7)	52 (33.5)	3 (1.9)
45-49	224 (100.0)	4 (1.8)	44 (19.6)	101 (45.1)	66 (29.5)	9 (4.0)
50-54	296 (100.0)	6 (2.0)	56 (18.9)	145 (49.0)	79 (26.7)	10 (3.4)
55-59	253 (100.0)	12 (4.7)	70 (27.7)	127 (50.2)	41 (16.2)	3 (1.2)
60-64	443 (100.0)	33 (7.4)	124 (28.0)	201 (45.4)	83 (18.7)	2 (0.5)
65-69	657 (100.0)	56 (8.5)	209 (31.8)	277 (42.2)	108 (16.4)	7 (1.1)
70-74	676 (100.0)	75 (11.1)	255 (37.7)	260 (38.5)	79 (11.7)	7 (1.0)
75-79	371 (100.0)	55 (14.8)	143 (38.5)	129 (34.8)	38 (10.2)	6 (1.6)
80歳以上	150 (100.0)	29 (19.3)	74 (49.3)	37 (24.7)	9 (6.0)	1 (0.7)
計	3,362 (100.0)	271 (8.1)	1,015 (30.2)	1,408 (41.9)	610 (18.1)	58 (1.7)

年齢階級	人 (%)					
	総数	-399	400-449	450-499	500-549	550 × 10 ⁴ / μl 以上
女						
-29歳	41 (100.0)	4 (9.8)	18 (43.9)	17 (41.5)	2 (4.9)	0 (0.0)
30-39	182 (100.0)	41 (22.5)	88 (48.4)	52 (28.6)	1 (0.5)	0 (0.0)
40-44	305 (100.0)	54 (17.7)	180 (59.0)	64 (21.0)	7 (2.3)	0 (0.0)
45-49	416 (100.0)	64 (15.4)	215 (51.7)	117 (28.1)	20 (4.8)	0 (0.0)
50-54	622 (100.0)	94 (15.1)	320 (51.4)	182 (29.3)	23 (3.7)	3 (0.5)
55-59	689 (100.0)	101 (14.7)	382 (55.4)	185 (26.9)	21 (3.0)	0 (0.0)
60-64	936 (100.0)	198 (21.2)	499 (53.3)	228 (24.4)	11 (1.2)	0 (0.0)
65-69	1,138 (100.0)	242 (21.3)	607 (53.3)	266 (23.4)	23 (2.0)	0 (0.0)
70-74	986 (100.0)	266 (27.0)	484 (49.1)	213 (21.6)	23 (2.3)	0 (0.0)
75-79	560 (100.0)	196 (35.0)	247 (44.1)	101 (18.0)	15 (2.7)	1 (0.2)
80歳以上	171 (100.0)	60 (35.1)	81 (47.4)	30 (17.5)	0 (0.0)	0 (0.0)
計	6,046 (100.0)	1,320 (21.8)	3,121 (51.6)	1,455 (24.1)	146 (2.4)	4 (0.1)

表40 赤血球数の平均値、標準偏差および分布 (性・年齢階級別)

年齢階級	男										女									
	パーセントイル					平均値 (SD)	パーセントイル					平均値 (SD)	パーセントイル							
	人数	5	25	50	75		95	人数	5	25	50		75	95						
29歳以下	41	501.1 (36.4)	429.7	475.5	504.0	522.5	573.0	41	443.7 (31.8)	395.0	413.0	440.0	465.5	503.4						
30-39	96	495.7 (37.3)	431.6	474.3	496.5	525.8	552.6	182	427.2 (34.4)	374.2	401.0	423.0	455.0	483.7						
40-44	155	486.5 (37.7)	421.6	460.0	490.0	509.0	544.0	305	429.6 (32.7)	375.0	408.0	429.0	448.0	486.7						
45-49	224	483.8 (38.9)	419.8	455.0	487.0	507.8	547.3	416	435.7 (36.1)	378.0	413.3	434.5	456.0	498.3						
50-54	296	479.0 (37.8)	417.9	455.0	480.0	504.0	542.2	622	435.3 (36.7)	376.3	411.0	434.0	459.0	495.9						
55-59	253	464.8 (36.3)	402.6	442.5	466.0	490.0	520.9	689	433.5 (33.7)	379.0	409.0	432.0	456.5	494.0						
60-64	443	463.1 (41.2)	393.0	439.0	466.0	492.0	523.8	936	427.4 (34.4)	370.0	404.0	427.5	450.0	485.0						
65-69	657	458.8 (42.6)	387.7	431.0	460.0	489.0	526.0	1,138	426.7 (34.8)	372.0	403.0	425.0	450.0	484.0						
70-74	676	451.0 (44.6)	379.9	424.0	450.5	480.8	524.0	986	422.4 (36.8)	363.0	397.0	422.0	448.0	487.7						
75-79	371	446.2 (44.5)	373.0	418.0	445.0	474.0	520.0	560	416.2 (41.6)	352.0	389.0	415.0	443.0	488.0						
80歳以上	150	430.8 (42.5)	359.6	405.0	431.0	463.0	502.5	171	410.2 (40.3)	332.6	383.0	413.0	438.0	472.8						
計	3,362	461.9 (44.2)	388.0	434.0	463.0	492.0	531.0	6,046	427.2 (36.5)	369.0	403.0	427.0	451.0	488.0						

10⁴/μl

表41 ヘモグロビン値の状況 (性・年齢階級別)

年齢階級	総数	人 (%)					
		-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上
男							
-29歳	41 (100.0)	0 (0.0)	0 (0.0)	3 (7.3)	15 (36.6)	14 (34.1)	9 (22.0)
30-39	96 (100.0)	0 (0.0)	1 (1.0)	10 (10.4)	13 (13.5)	46 (47.9)	26 (27.1)
40-44	155 (100.0)	0 (0.0)	4 (2.6)	13 (8.4)	38 (24.5)	63 (40.6)	37 (23.9)
45-49	224 (100.0)	3 (1.3)	7 (3.1)	22 (9.8)	69 (30.8)	80 (35.7)	43 (19.2)
50-54	296 (100.0)	3 (1.0)	6 (2.0)	36 (12.2)	90 (30.4)	106 (35.8)	55 (18.6)
55-59	253 (100.0)	4 (1.6)	11 (4.3)	41 (16.2)	99 (39.1)	70 (27.7)	28 (11.1)
60-64	443 (100.0)	9 (2.0)	26 (5.9)	72 (16.3)	155 (35.0)	128 (28.9)	53 (12.0)
65-69	657 (100.0)	20 (3.0)	55 (8.4)	114 (17.4)	227 (34.6)	151 (23.0)	90 (13.7)
70-74	676 (100.0)	30 (4.4)	56 (8.3)	165 (24.4)	215 (31.8)	148 (21.9)	62 (9.2)
75-79	371 (100.0)	16 (4.3)	57 (15.4)	103 (27.8)	110 (29.6)	59 (15.9)	26 (7.0)
80歳以上	150 (100.0)	14 (9.3)	19 (12.7)	53 (35.3)	45 (30.0)	15 (10.0)	4 (2.7)
計	3,362 (100.0)	99 (2.9)	242 (7.2)	632 (18.8)	1,076 (32.0)	880 (26.2)	433 (12.9)

年齢階級	総数	人 (%)					
		-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上
女							
-29歳	41 (100.0)	6 (14.6)	16 (39.0)	13 (31.7)	5 (12.2)	1 (2.4)	0 (0.0)
30-39	182 (100.0)	38 (20.9)	75 (41.2)	53 (29.1)	14 (7.7)	2 (1.1)	0 (0.0)
40-44	305 (100.0)	74 (24.3)	94 (30.8)	89 (29.2)	44 (14.4)	4 (1.3)	0 (0.0)
45-49	416 (100.0)	91 (21.9)	104 (25.0)	144 (34.6)	63 (15.1)	13 (3.1)	1 (0.2)
50-54	622 (100.0)	86 (13.8)	192 (30.9)	209 (33.6)	114 (18.3)	19 (3.1)	2 (0.3)
55-59	689 (100.0)	62 (9.0)	222 (32.2)	252 (36.6)	128 (18.6)	24 (3.5)	1 (0.1)
60-64	936 (100.0)	108 (11.5)	319 (34.1)	345 (36.9)	149 (15.9)	15 (1.6)	0 (0.0)
65-69	1,138 (100.0)	148 (13.0)	365 (32.1)	421 (37.0)	172 (15.1)	29 (2.5)	3 (0.3)
70-74	986 (100.0)	151 (15.3)	341 (34.6)	322 (32.7)	143 (14.5)	28 (2.8)	1 (0.1)
75-79	560 (100.0)	124 (22.1)	183 (32.7)	162 (28.9)	74 (13.2)	16 (2.9)	1 (0.2)
80歳以上	171 (100.0)	46 (26.9)	54 (31.6)	50 (29.2)	17 (9.9)	4 (2.3)	0 (0.0)
計	6,046 (100.0)	934 (15.4)	1,965 (32.5)	2,060 (34.1)	923 (15.3)	155 (2.6)	9 (0.1)

表42 ヘモグロビン値の平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						g/dl	
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル					
			5	25	50	75			95	5	25	50		75
29歳以下	41	15.2 (1.0)	13.3	14.7	15.2	15.9	16.9	41	12.9 (1.1)	10.8	12.2	12.8	13.7	14.8
30-39	96	15.4 (1.0)	13.6	14.9	15.5	16.1	16.8	182	12.6 (1.2)	10.5	12.1	12.7	13.4	14.6
40-44	155	15.3 (1.1)	13.6	14.6	15.2	15.9	16.9	305	12.6 (1.4)	10.0	12.0	12.7	13.6	14.5
45-49	224	15.0 (1.3)	13.1	14.3	15.1	15.8	16.9	416	12.7 (1.5)	9.8	12.1	13.0	13.8	14.7
50-54	296	15.0 (1.2)	13.2	14.4	15.1	15.7	16.9	622	13.0 (1.2)	11.0	12.3	13.1	13.8	14.8
55-59	253	14.7 (1.2)	12.8	14.1	14.6	15.4	16.5	689	13.2 (1.0)	11.5	12.5	13.1	13.8	14.7
60-64	443	14.7 (1.2)	12.6	14.0	14.7	15.4	16.5	936	13.0 (1.0)	11.4	12.4	13.0	13.7	14.6
65-69	657	14.5 (1.3)	12.3	13.8	14.6	15.3	16.5	1,138	13.0 (1.0)	11.3	12.3	13.1	13.7	14.7
70-74	676	14.3 (1.4)	12.0	13.5	14.3	15.2	16.3	986	13.0 (1.1)	11.2	12.2	13.0	13.7	14.7
75-79	371	14.0 (1.3)	12.0	13.2	14.0	14.8	16.1	560	12.8 (1.3)	10.7	12.0	12.8	13.6	14.7
80歳以上	150	13.7 (1.4)	11.5	13.1	13.8	14.6	15.5	171	12.5 (1.4)	10.4	11.8	12.7	13.4	14.7
計	3,362	14.6 (1.3)	12.3	13.8	14.6	15.4	16.5	6,046	12.9 (1.2)	11.0	12.3	13.0	13.7	14.7

表43 ヘマトクリット値の状況 (性・年齢階級別)

年齢階級	総数	-29.9	30.0-34.9	35.0-39.9	40.0-44.9	45.0-49.9	50.0%以上
男							
-29歳	41 (100.0)	0 (0.0)	0 (0.0)	3 (7.3)	17 (41.5)	20 (48.8)	1 (2.4)
30-39	96 (100.0)	0 (0.0)	0 (0.0)	2 (2.1)	30 (31.3)	61 (63.5)	3 (3.1)
40-44	155 (100.0)	0 (0.0)	0 (0.0)	5 (3.2)	67 (43.2)	75 (48.4)	8 (5.2)
45-49	224 (100.0)	1 (0.4)	1 (0.4)	13 (5.8)	98 (43.8)	102 (45.5)	9 (4.0)
50-54	296 (100.0)	1 (0.3)	2 (0.7)	13 (4.4)	136 (45.9)	129 (43.6)	15 (5.1)
55-59	253 (100.0)	1 (0.4)	2 (0.8)	22 (8.7)	147 (58.1)	76 (30.0)	5 (2.0)
60-64	443 (100.0)	1 (0.2)	3 (0.7)	54 (12.2)	221 (49.9)	148 (33.4)	16 (3.6)
65-69	657 (100.0)	0 (0.0)	12 (1.8)	99 (15.1)	327 (49.8)	206 (31.4)	13 (2.0)
70-74	676 (100.0)	4 (0.6)	18 (2.7)	128 (18.9)	343 (50.7)	169 (25.0)	14 (2.1)
75-79	371 (100.0)	0 (0.0)	9 (2.4)	94 (25.3)	187 (50.4)	77 (20.8)	4 (1.1)
80歳以上	150 (100.0)	2 (1.3)	4 (2.7)	44 (29.3)	76 (50.7)	23 (15.3)	1 (0.7)
計	3,362 (100.0)	10 (0.3)	51 (1.5)	477 (14.2)	1,649 (49.0)	1,086 (32.3)	89 (2.6)

年齢階級	総数	-29.9	30.0-34.9	35.0-39.9	40.0-44.9	45.0-49.9	50.0%以上
女							
-29歳	41 (100.0)	0 (0.0)	3 (7.3)	22 (53.7)	16 (39.0)	0 (0.0)	0 (0.0)
30-39	182 (100.0)	2 (1.1)	15 (8.2)	117 (64.3)	44 (24.2)	4 (2.2)	0 (0.0)
40-44	305 (100.0)	6 (2.0)	40 (13.1)	157 (51.5)	100 (32.8)	2 (0.7)	0 (0.0)
45-49	416 (100.0)	10 (2.4)	44 (10.6)	196 (47.1)	154 (37.0)	12 (2.9)	0 (0.0)
50-54	622 (100.0)	1 (0.2)	47 (7.6)	291 (46.8)	263 (42.3)	20 (3.2)	0 (0.0)
55-59	689 (100.0)	1 (0.1)	24 (3.5)	324 (47.0)	314 (45.6)	26 (3.8)	0 (0.0)
60-64	936 (100.0)	2 (0.2)	49 (5.2)	475 (50.7)	387 (41.3)	23 (2.5)	0 (0.0)
65-69	1,138 (100.0)	4 (0.4)	63 (5.5)	575 (50.5)	458 (40.2)	38 (3.3)	0 (0.0)
70-74	986 (100.0)	3 (0.3)	77 (7.8)	490 (49.7)	378 (38.3)	38 (3.9)	0 (0.0)
75-79	560 (100.0)	6 (1.1)	68 (12.1)	279 (49.8)	185 (33.0)	22 (3.9)	0 (0.0)
80歳以上	171 (100.0)	4 (2.3)	26 (15.2)	86 (50.3)	50 (29.2)	5 (2.9)	0 (0.0)
計	6,046 (100.0)	39 (0.6)	456 (7.5)	3,012 (49.8)	2,349 (38.9)	190 (3.1)	0 (0.0)

表44 ハマトクリット値の平均値、標準偏差および分布 (性・年齢階級別)

年齢階級	男						女							
	人数	平均値 (SD)	パーセンタイル					人数	平均値 (SD)	パーセンタイル				
			5	25	50	75	95			5	25	50	75	95
29歳以下	41	45.0 (2.8)	39.7	43.1	45.2	46.9	49.3	41	39.0 (2.8)	34.1	37.0	39.2	41.2	44.2
30-39	96	45.8 (2.8)	40.4	44.3	46.1	47.5	49.5	182	38.3 (3.0)	32.7	36.7	38.1	40.0	43.5
40-44	155	45.2 (3.0)	40.3	43.2	45.3	47.3	50.2	305	38.3 (3.4)	32.6	36.6	38.7	40.5	43.4
45-49	224	44.7 (3.5)	39.2	42.8	44.9	46.8	49.7	416	38.8 (3.9)	31.7	37.0	39.3	41.3	44.1
50-54	296	44.7 (3.5)	39.4	42.6	44.8	46.8	50.1	622	39.6 (3.3)	34.0	37.6	39.6	41.8	44.7
55-59	253	43.6 (3.3)	38.2	41.8	43.6	45.6	48.3	689	39.8 (2.8)	35.4	37.9	39.9	41.8	44.5
60-64	443	43.7 (3.6)	37.9	41.6	43.8	46.0	49.6	936	39.5 (2.9)	34.9	37.6	39.4	41.3	44.2
65-69	657	43.4 (3.7)	37.0	41.1	43.5	45.9	49.0	1,138	39.5 (3.0)	34.6	37.5	39.5	41.5	44.3
70-74	676	42.6 (3.9)	35.7	40.3	42.8	45.2	48.5	986	39.3 (3.2)	34.2	37.3	39.3	41.5	44.7
75-79	371	42.1 (3.7)	36.2	39.7	42.0	44.6	48.2	560	38.8 (3.6)	33.2	36.5	38.9	41.3	44.7
80歳以上	150	41.1 (3.9)	35.2	39.0	41.2	43.6	46.6	171	38.2 (3.7)	32.2	36.1	38.5	40.7	44.4
計	3,362	43.4 (3.8)	37.1	41.2	43.6	45.9	49.1	6,046	39.3 (3.2)	33.9	37.3	39.3	41.4	44.4

表45 血清ALPの状況(性・年齢階級別)

年齢階級	総数							人 (%)
	-199	200-249	250-299	300-349	350-399	400U/L以上		
男								
-29歳	41 (100.0)	10 (24.4)	10 (24.4)	4 (9.8)	3 (7.3)	1 (2.4)		
30-39	96 (100.0)	26 (27.1)	26 (27.1)	8 (8.3)	7 (7.3)	1 (1.0)		
40-44	144 (100.0)	47 (32.6)	32 (22.2)	26 (18.1)	6 (4.2)	3 (2.1)		
45-49	211 (100.0)	70 (33.2)	47 (22.3)	25 (11.8)	11 (5.2)	7 (3.3)		
50-54	269 (100.0)	80 (29.7)	68 (25.3)	26 (9.7)	15 (5.6)	10 (3.7)		
55-59	230 (100.0)	87 (37.8)	50 (21.7)	36 (15.7)	5 (2.2)	6 (2.6)		
60-64	396 (100.0)	115 (29.0)	110 (27.8)	38 (9.6)	24 (6.1)	17 (4.3)		
65-69	583 (100.0)	167 (28.6)	156 (26.8)	74 (12.7)	30 (5.1)	36 (6.2)		
70-74	608 (100.0)	169 (27.8)	141 (23.2)	89 (14.6)	36 (5.9)	32 (5.3)		
75-79	328 (100.0)	70 (21.3)	91 (27.7)	46 (14.0)	26 (7.9)	18 (5.5)		
80歳以上	131 (100.0)	33 (25.2)	42 (32.1)	18 (13.7)	6 (4.6)	9 (6.9)		
計	3,037 (100.0)	874 (28.8)	773 (25.5)	390 (12.8)	169 (5.6)	140 (4.6)		

年齢階級	総数							人 (%)
	-199	200-249	250-299	300-349	350-399	400U/L以上		
女								
-29歳	41 (100.0)	10 (24.4)	8 (19.5)	0 (0.0)	1 (2.4)	1 (2.4)		
30-39	182 (100.0)	48 (26.4)	15 (8.2)	4 (2.2)	1 (0.5)	4 (2.2)		
40-44	292 (100.0)	78 (26.7)	28 (9.6)	7 (2.4)	3 (1.0)	3 (1.0)		
45-49	383 (100.0)	121 (31.6)	48 (12.5)	24 (6.3)	5 (1.3)	8 (2.1)		
50-54	570 (100.0)	176 (30.9)	111 (19.5)	86 (15.1)	39 (6.8)	30 (5.3)		
55-59	618 (100.0)	151 (24.4)	194 (31.4)	106 (17.2)	49 (7.9)	41 (6.6)		
60-64	840 (100.0)	225 (26.8)	217 (25.8)	174 (20.7)	68 (8.1)	73 (8.7)		
65-69	998 (100.0)	257 (25.8)	243 (24.3)	182 (18.2)	117 (11.7)	87 (8.7)		
70-74	888 (100.0)	233 (26.2)	223 (25.1)	178 (20.0)	86 (9.7)	89 (10.0)		
75-79	500 (100.0)	114 (22.8)	114 (22.8)	84 (16.8)	57 (11.4)	68 (13.6)		
80歳以上	146 (100.0)	25 (17.1)	38 (26.0)	31 (21.2)	20 (13.7)	19 (13.0)		
計	5,458 (100.0)	1,438 (26.3)	1,239 (22.7)	876 (16.0)	446 (8.2)	423 (7.8)		

表46 血清ALPの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						U/L		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	41	246.5 (69.1)	157.1	193.0	239.0	283.5	375.8	41	209.2 (80.9)	115.9	160.0	198.0	249.5	351.1	
30-39	96	243.6 (66.6)	150.7	186.8	240.0	281.8	366.1	182	192.2 (64.8)	107.6	150.0	184.0	215.5	305.7	
40-44	144	254.0 (86.5)	162.8	208.0	245.0	297.5	353.0	292	196.3 (73.9)	118.0	154.0	188.0	220.5	290.4	
45-49	211	252.2 (84.6)	156.0	200.0	238.0	289.0	381.0	383	216.3 (79.1)	131.0	172.0	206.0	240.0	328.8	
50-54	269	248.6 (80.4)	138.5	196.5	238.0	289.0	389.0	570	261.3 (82.9)	161.0	203.0	245.5	305.0	406.9	
55-59	230	251.7 (81.9)	164.2	205.8	235.5	287.3	351.1	618	277.7 (74.6)	176.0	224.0	268.5	314.0	413.2	
60-64	396	253.1 (77.1)	153.9	201.0	244.0	286.0	394.0	840	285.0 (79.8)	180.1	230.0	272.0	325.0	439.9	
65-69	583	260.2 (81.4)	157.0	205.0	250.0	298.0	415.0	998	288.1 (89.5)	178.0	228.0	275.0	336.3	429.1	
70-74	608	262.1 (93.2)	156.5	201.0	248.5	302.0	407.7	888	290.0 (86.3)	180.5	229.0	279.0	332.8	439.0	
75-79	328	264.8 (84.9)	156.4	204.3	256.0	311.0	412.9	500	295.5 (104.5)	165.1	229.0	278.5	349.8	459.0	
80歳以上	131	266.9 (72.4)	174.0	211.0	259.0	300.0	437.6	146	307.7 (89.7)	178.8	247.3	293.0	354.8	451.7	
計	3,037	257.2 (83.4)	156.9	203.0	247.0	294.0	394.0	5,458	271.4 (90.1)	154.0	211.0	259.0	317.0	424.0	

表47 血清コレステロール値の状況 (性・年齢階級別)

年齢階級	人数							人 (%)
	総数	-119	120-159	160-199	200-239	240-279	280U/L以上	
男								
-29歳	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	18 (100.0)	0 (0.0)	2 (11.1)	11 (61.1)	2 (11.1)	3 (16.7)	0 (0.0)	0 (0.0)
40-44	84 (100.0)	2 (2.4)	7 (8.3)	36 (42.9)	28 (33.3)	9 (10.7)	2 (2.4)	2 (2.4)
45-49	128 (100.0)	2 (1.6)	12 (9.4)	64 (50.0)	39 (30.5)	11 (8.6)	0 (0.0)	0 (0.0)
50-54	167 (100.0)	4 (2.4)	22 (13.2)	82 (49.1)	48 (28.7)	9 (5.4)	2 (1.2)	2 (1.2)
55-59	144 (100.0)	5 (3.5)	31 (21.5)	65 (45.1)	35 (24.3)	8 (5.6)	0 (0.0)	0 (0.0)
60-64	295 (100.0)	13 (4.4)	71 (24.1)	121 (41.0)	74 (25.1)	14 (4.7)	2 (0.7)	2 (0.7)
65-69	452 (100.0)	18 (4.0)	134 (29.6)	198 (43.8)	87 (19.2)	11 (2.4)	4 (0.9)	4 (0.9)
70-74	475 (100.0)	31 (6.5)	172 (36.2)	207 (43.6)	54 (11.4)	10 (2.1)	1 (0.2)	1 (0.2)
75-79	266 (100.0)	21 (7.9)	117 (44.0)	97 (36.5)	25 (9.4)	6 (2.3)	0 (0.0)	0 (0.0)
80歳以上	108 (100.0)	11 (10.2)	56 (51.9)	31 (28.7)	6 (5.6)	3 (2.8)	1 (0.9)	1 (0.9)
計	2,137 (100.0)	107 (5.0)	624 (29.2)	912 (42.7)	398 (18.6)	84 (3.9)	12 (0.6)	12 (0.6)

年齢階級	人数							人 (%)
	総数	-119	120-159	160-199	200-239	240-279	280U/L以上	
女								
-29歳	1 (100.0)	0 (0.0)	0 (0.0)	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	47 (100.0)	1 (2.1)	23 (48.9)	20 (42.6)	3 (6.4)	0 (0.0)	0 (0.0)	0 (0.0)
40-44	199 (100.0)	8 (4.0)	80 (40.2)	76 (38.2)	33 (16.6)	2 (1.0)	0 (0.0)	0 (0.0)
45-49	283 (100.0)	7 (2.5)	89 (31.4)	114 (40.3)	55 (19.4)	15 (5.3)	3 (1.1)	3 (1.1)
50-54	443 (100.0)	9 (2.0)	78 (17.6)	209 (47.2)	108 (24.4)	31 (7.0)	8 (1.8)	8 (1.8)
55-59	471 (100.0)	4 (0.8)	56 (11.9)	195 (41.4)	151 (32.1)	53 (11.3)	12 (2.5)	12 (2.5)
60-64	640 (100.0)	1 (0.2)	85 (13.3)	307 (48.0)	179 (28.0)	50 (7.8)	18 (2.8)	18 (2.8)
65-69	796 (100.0)	11 (1.4)	134 (16.8)	344 (43.2)	219 (27.5)	67 (8.4)	21 (2.6)	21 (2.6)
70-74	731 (100.0)	12 (1.6)	174 (23.8)	339 (46.4)	150 (20.5)	42 (5.7)	14 (1.9)	14 (1.9)
75-79	431 (100.0)	9 (2.1)	128 (29.7)	192 (44.5)	84 (19.5)	15 (3.5)	3 (0.7)	3 (0.7)
80歳以上	128 (100.0)	8 (6.3)	49 (38.3)	50 (39.1)	16 (12.5)	2 (1.6)	3 (2.3)	3 (2.3)
計	4,170 (100.0)	70 (1.7)	896 (21.5)	1,847 (44.3)	998 (23.9)	277 (6.6)	82 (2.0)	82 (2.0)

表48 血清コリンエステラーゼの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	U/L													
	男					女								
	パーセントाइル					パーセントाइル								
	人数	平均値(SD)	5	25	50	75	95	人数	平均値(SD)	5	25	50	75	95
29歳以下	0							1	180.0					
30-39	18	194.0 (33.2)	147.0	167.5	188.0	219.8	248.0	47	160.6 (25.1)	127.8	139.0	158.0	181.0	209.0
40-44	84	198.4 (37.9)	141.0	172.8	197.0	218.8	272.5	199	167.1 (30.7)	120.0	146.0	162.0	189.0	220.0
45-49	128	193.6 (33.6)	139.0	172.0	192.5	213.8	256.6	283	177.8 (35.6)	127.0	154.0	172.0	200.0	245.8
50-54	167	188.2 (34.9)	131.0	166.0	186.0	216.0	244.6	443	188.7 (37.4)	130.0	165.0	186.0	208.0	253.4
55-59	144	179.9 (35.0)	120.0	159.3	177.0	204.0	241.8	471	198.5 (37.5)	141.0	173.0	196.0	222.0	264.0
60-64	295	181.6 (36.6)	120.8	156.0	181.0	207.0	243.6	640	195.3 (37.0)	144.0	170.0	190.0	216.8	265.0
65-69	452	175.4 (34.4)	123.0	153.0	172.0	196.8	231.0	796	192.9 (39.0)	139.7	167.0	189.0	215.0	259.2
70-74	475	166.6 (33.1)	116.6	143.0	164.0	187.0	225.0	731	184.0 (37.2)	133.6	159.0	180.0	204.0	257.0
75-79	266	161.0 (33.7)	110.4	140.0	158.0	180.0	222.0	431	178.4 (34.8)	129.6	154.0	175.0	198.0	239.0
80歳以上	108	154.8 (35.7)	107.5	136.3	150.0	173.8	225.2	128	170.1 (37.3)	114.2	148.3	165.0	188.8	235.8
計	2,137	174.9 (36.3)	119.0	151.0	173.0	197.0	236.0	4,170	187.1 (37.9)	133.0	161.0	183.0	208.0	255.0

表49 総ビリルビンの状況 (性・年齢階級別)

年齢階級	人 (%)					
	総数	-0.5	0.6-0.8	0.9-1.1	1.2-1.4	1.5mg/dl以上
男						
-29歳	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	18 (100.0)	8 (44.4)	7 (38.9)	3 (16.7)	0 (0.0)	0 (0.0)
40-44	84 (100.0)	32 (38.1)	31 (36.9)	15 (17.9)	5 (6.0)	1 (1.2)
45-49	128 (100.0)	60 (46.9)	46 (35.9)	14 (10.9)	2 (1.6)	6 (4.7)
50-54	167 (100.0)	61 (36.5)	71 (42.5)	26 (15.6)	9 (5.4)	0 (0.0)
55-59	144 (100.0)	56 (38.9)	60 (41.7)	19 (13.2)	5 (3.5)	4 (2.8)
60-64	295 (100.0)	106 (35.9)	122 (41.4)	54 (18.3)	9 (3.1)	4 (1.4)
65-69	452 (100.0)	158 (35.0)	201 (44.5)	64 (14.2)	20 (4.4)	9 (2.0)
70-74	475 (100.0)	167 (35.2)	201 (42.3)	82 (17.3)	19 (4.0)	6 (1.3)
75-79	266 (100.0)	124 (46.6)	94 (35.3)	37 (13.9)	9 (3.4)	2 (0.8)
80歳以上	108 (100.0)	51 (47.2)	43 (39.8)	8 (7.4)	4 (3.7)	2 (1.9)
計	2,137 (100.0)	823 (38.5)	876 (41.0)	322 (15.1)	82 (3.8)	34 (1.6)

年齢階級	人 (%)					
	総数	-0.5	0.6-0.8	0.9-1.1	1.2-1.4	1.5mg/dl以上
女						
-29歳	1 (100.0)	0 (0.0)	1 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	47 (100.0)	34 (72.3)	10 (21.3)	1 (2.1)	1 (2.1)	1 (2.1)
40-44	199 (100.0)	96 (48.2)	78 (39.2)	16 (8.0)	5 (2.5)	4 (2.0)
45-49	283 (100.0)	142 (50.2)	108 (38.2)	26 (9.2)	3 (1.1)	4 (1.4)
50-54	443 (100.0)	195 (44.0)	193 (43.6)	46 (10.4)	5 (1.1)	4 (0.9)
55-59	471 (100.0)	205 (43.5)	206 (43.7)	46 (9.8)	11 (2.3)	3 (0.6)
60-64	640 (100.0)	309 (48.3)	253 (39.5)	54 (8.4)	15 (2.3)	9 (1.4)
65-69	796 (100.0)	357 (44.8)	343 (43.1)	71 (8.9)	15 (1.9)	10 (1.3)
70-74	731 (100.0)	348 (47.6)	291 (39.8)	68 (9.3)	16 (2.2)	8 (1.1)
75-79	431 (100.0)	244 (56.6)	149 (34.6)	30 (7.0)	7 (1.6)	1 (0.2)
80歳以上	128 (100.0)	74 (57.8)	46 (35.9)	6 (4.7)	2 (1.6)	0 (0.0)
計	4,170 (100.0)	2,004 (48.1)	1,678 (40.2)	364 (8.7)	80 (1.9)	44 (1.1)

表50 総ビリルビンの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男										女										mg/dl
	人数	パーセンタイル					平均値 (SD)	人数	パーセンタイル					平均値 (SD)							
		5	25	50	75	95			5	25	50	75	95								
29歳以下	0						1	0.7													
30-39	18	0.6 (0.2)	0.3	0.5	0.6	0.7	1.0	47	0.5 (0.3)	0.3	0.4	0.5	0.6	1.1	199	0.6 (0.3)	0.3	0.5	0.6	0.7	1.1
40-44	84	0.7 (0.3)	0.3	0.5	0.7	0.9	1.3	283	0.6 (0.2)	0.3	0.4	0.5	0.7	1.0	443	0.6 (0.2)	0.3	0.5	0.6	0.7	1.0
45-49	128	0.7 (0.3)	0.4	0.5	0.6	0.8	1.2	471	0.6 (0.2)	0.3	0.5	0.6	0.7	1.0	640	0.6 (0.3)	0.3	0.4	0.6	0.7	1.1
50-54	167	0.7 (0.2)	0.4	0.5	0.6	0.8	1.2	796	0.6 (0.2)	0.3	0.5	0.6	0.7	1.0	731	0.6 (0.2)	0.3	0.5	0.6	0.7	1.1
55-59	144	0.7 (0.3)	0.3	0.5	0.6	0.8	1.3	431	0.6 (0.2)	0.3	0.4	0.5	0.6	1.0	128	0.5 (0.2)	0.3	0.4	0.5	0.6	0.9
60-64	295	0.7 (0.3)	0.4	0.5	0.6	0.8	1.1	4,170	0.6 (0.2)	0.3	0.5	0.6	0.7	1.0							
65-69	452	0.7 (0.3)	0.4	0.5	0.6	0.8	1.2														
70-74	475	0.7 (0.3)	0.3	0.5	0.6	0.8	1.2														
75-79	266	0.6 (0.3)	0.3	0.5	0.6	0.8	1.1														
80歳以上	108	0.6 (0.3)	0.3	0.5	0.6	0.7	1.2														
計	2,137	0.7 (0.3)	0.3	0.5	0.6	0.8	1.2														

表51 血清総蛋白の状況 (性・年齢階級別)

年齢階級	人 (%)							
	総数	-6.3	6.4-6.7	6.8-7.1	7.2-7.5	7.6-7.9	8.0g/dl以上	
男								
-29歳	41 (100.0)	1 (2.4)	1 (2.4)	9 (22.0)	17 (41.5)	12 (29.3)	1 (2.4)	
30-39	96 (100.0)	1 (1.0)	6 (6.3)	31 (32.3)	36 (37.5)	20 (20.8)	2 (2.1)	
40-44	144 (100.0)	0 (0.0)	11 (7.6)	34 (23.6)	41 (28.5)	44 (30.6)	14 (9.7)	
45-49	211 (100.0)	1 (0.5)	15 (7.1)	59 (28.0)	77 (36.5)	48 (22.7)	11 (5.2)	
50-54	269 (100.0)	6 (2.2)	18 (6.7)	74 (27.5)	89 (33.1)	61 (22.7)	21 (7.8)	
55-59	230 (100.0)	2 (0.9)	18 (7.8)	71 (30.9)	77 (33.5)	53 (23.0)	9 (3.9)	
60-64	396 (100.0)	3 (0.8)	33 (8.3)	115 (29.0)	125 (31.6)	90 (22.7)	30 (7.6)	
65-69	583 (100.0)	6 (1.0)	56 (9.6)	165 (28.3)	219 (37.6)	113 (19.4)	24 (4.1)	
70-74	608 (100.0)	13 (2.1)	66 (10.9)	176 (28.9)	202 (33.2)	118 (19.4)	33 (5.4)	
75-79	328 (100.0)	2 (0.6)	42 (12.8)	99 (30.2)	124 (37.8)	43 (13.1)	18 (5.5)	
80歳以上	131 (100.0)	1 (0.8)	30 (22.9)	33 (25.2)	46 (35.1)	14 (10.7)	7 (5.3)	
計	3,037 (100.0)	36 (1.2)	296 (9.7)	866 (28.5)	1,053 (34.7)	616 (20.3)	170 (5.6)	

年齢階級	人 (%)							
	総数	-6.3	6.4-6.7	6.8-7.1	7.2-7.5	7.6-7.9	8.0g/dl以上	
女								
-29歳	41 (100.0)	0 (0.0)	2 (4.9)	8 (19.5)	19 (46.3)	11 (26.8)	1 (2.4)	
30-39	182 (100.0)	0 (0.0)	13 (7.1)	54 (29.7)	66 (36.3)	35 (19.2)	14 (7.7)	
40-44	292 (100.0)	1 (0.3)	23 (7.9)	72 (24.7)	118 (40.4)	67 (22.9)	11 (3.8)	
45-49	383 (100.0)	0 (0.0)	28 (7.3)	85 (22.2)	135 (35.2)	87 (22.7)	48 (12.5)	
50-54	570 (100.0)	1 (0.2)	23 (4.0)	125 (21.9)	203 (35.6)	160 (28.1)	58 (10.2)	
55-59	618 (100.0)	6 (1.0)	18 (2.9)	131 (21.2)	248 (40.1)	158 (25.6)	57 (9.2)	
60-64	840 (100.0)	1 (0.1)	43 (5.1)	185 (22.0)	348 (41.4)	202 (24.0)	61 (7.3)	
65-69	998 (100.0)	4 (0.4)	69 (6.9)	242 (24.2)	410 (41.1)	197 (19.7)	76 (7.6)	
70-74	888 (100.0)	7 (0.8)	54 (6.1)	229 (25.8)	346 (39.0)	196 (22.1)	56 (6.3)	
75-79	500 (100.0)	3 (0.6)	44 (8.8)	131 (26.2)	186 (37.2)	104 (20.8)	32 (6.4)	
80歳以上	146 (100.0)	5 (3.4)	14 (9.6)	39 (26.7)	52 (35.6)	25 (17.1)	11 (7.5)	
計	5,458 (100.0)	28 (0.5)	331 (6.1)	1,301 (23.8)	2,131 (39.0)	1,242 (22.8)	425 (7.8)	

表52 血清総蛋白の平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						g/dl	
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル					
			5	25	50	75			95	5	25	50		75
29歳以下	41	7.3 (0.4)	6.7	7.1	7.4	7.6	7.8	41	7.4 (0.3)	6.7	7.2	7.4	7.6	7.9
30-39	96	7.3 (0.4)	6.7	7.0	7.2	7.5	7.9	182	7.3 (0.4)	6.7	7.0	7.3	7.6	8.0
40-44	144	7.4 (0.5)	6.5	7.0	7.3	7.7	8.2	292	7.3 (0.4)	6.6	7.0	7.3	7.6	7.9
45-49	211	7.3 (0.4)	6.7	7.0	7.3	7.6	8.0	383	7.4 (0.5)	6.7	7.1	7.4	7.7	8.1
50-54	269	7.3 (0.4)	6.7	7.0	7.3	7.6	8.0	570	7.4 (0.4)	6.8	7.1	7.4	7.7	8.2
55-59	230	7.3 (0.4)	6.7	7.0	7.3	7.6	7.9	618	7.4 (0.4)	6.8	7.1	7.4	7.7	8.1
60-64	396	7.3 (0.4)	6.6	7.0	7.3	7.6	8.0	840	7.4 (0.4)	6.7	7.1	7.4	7.6	8.0
65-69	583	7.3 (0.4)	6.6	7.0	7.3	7.5	7.9	998	7.3 (0.4)	6.7	7.1	7.3	7.6	8.0
70-74	608	7.2 (0.4)	6.5	7.0	7.2	7.5	8.0	888	7.3 (0.4)	6.7	7.0	7.3	7.6	8.0
75-79	328	7.2 (0.4)	6.6	6.9	7.2	7.5	8.0	500	7.3 (0.4)	6.6	7.0	7.3	7.6	8.0
80歳以上	131	7.2 (0.5)	6.5	6.8	7.2	7.4	8.0	146	7.3 (0.5)	6.4	7.0	7.3	7.5	8.1
計	3,037	7.3 (0.4)	6.6	7.0	7.3	7.6	8.0	5,458	7.4 (0.4)	6.7	7.1	7.4	7.6	8.0