

表24 血清中性脂肪の平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						mg/dl		
	人数	平均値 (SD)	パーセンタイル					人数	平均値 (SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	38	132.6 (93.3)	45.2	78.0	111.5	168.0	363.1	100	75.5 (77.7)	29.1	42.0	56.0	81.5	160.4	
30-39	80	126.5 (99.7)	39.4	70.0	97.0	152.0	336.6	299	85.4 (62.0)	33.0	50.0	67.0	97.0	210.0	
40-44	114	148.0 (96.9)	52.0	81.5	119.5	186.3	325.3	279	88.8 (57.0)	36.0	53.0	72.0	106.0	219.0	
45-49	145	131.4 (89.3)	47.9	69.5	111.0	161.5	333.1	434	92.3 (53.2)	40.0	55.0	78.0	110.3	190.5	
50-54	286	131.3 (81.5)	50.0	74.8	107.0	156.3	291.6	731	100.2 (55.4)	41.0	65.0	87.0	119.0	208.4	
55-59	317	127.3 (81.2)	45.9	73.0	108.0	153.5	299.8	860	110.7 (65.0)	44.1	68.0	94.0	135.0	235.0	
60-64	585	119.6 (73.4)	45.0	71.0	99.0	149.0	251.5	1,220	110.5 (59.0)	46.0	70.0	97.0	136.0	213.0	
65-69	793	116.4 (77.4)	47.0	71.0	97.0	137.0	241.3	1,258	113.3 (58.4)	46.0	72.0	100.0	137.0	232.0	
70-74	705	106.1 (63.3)	46.0	67.0	90.0	127.0	227.4	998	109.8 (55.6)	50.0	73.0	97.0	129.0	202.2	
75-79	470	103.2 (56.6)	42.0	64.8	90.0	125.0	210.9	512	111.1 (56.9)	48.0	73.0	97.0	133.8	216.4	
80歳以上	143	96.6 (49.3)	44.2	68.0	84.0	114.0	208.2	173	106.9 (54.2)	51.0	73.5	92.0	124.5	207.3	
計	3,676	116.5 (74.4)	46.0	69.0	97.0	140.0	254.0	6,864	106.1 (59.2)	43.0	66.0	92.0	129.0	216.0	

表25 血清ASTの状況(性・年齢階級別)

年齢階級	人(%)				
	総数	-19	20-39	40-59	60U/L以上
-29歳	38 (100.0)	16 (42.1)	20 (52.6)	1 (2.6)	1 (2.6)
30-39	80 (100.0)	22 (27.5)	51 (63.8)	4 (5.0)	3 (3.8)
40-44	114 (100.0)	28 (24.6)	79 (69.3)	6 (5.3)	1 (0.9)
45-49	145 (100.0)	40 (27.6)	97 (66.9)	6 (4.1)	2 (1.4)
50-54	286 (100.0)	47 (16.4)	216 (75.5)	16 (5.6)	7 (2.4)
55-59	317 (100.0)	54 (17.0)	239 (75.4)	19 (6.0)	5 (1.6)
60-64	585 (100.0)	108 (18.5)	429 (73.3)	31 (5.3)	17 (2.9)
65-69	793 (100.0)	117 (14.8)	596 (75.2)	62 (7.8)	18 (2.3)
70-74	705 (100.0)	103 (14.6)	552 (78.3)	38 (5.4)	12 (1.7)
75-79	470 (100.0)	68 (14.5)	369 (78.5)	29 (6.2)	4 (0.9)
80歳以上	143 (100.0)	17 (11.9)	118 (82.5)	7 (4.9)	1 (0.7)
計	3,676 (100.0)	620 (16.9)	2,766 (75.2)	219 (6.0)	71 (1.9)

年齢階級	人(%)				
	総数	-19	20-39	40-59	60U/L以上
-29歳	100 (100.0)	80 (80.0)	20 (20.0)	0 (0.0)	0 (0.0)
30-39	299 (100.0)	226 (75.6)	72 (24.1)	1 (0.3)	0 (0.0)
40-44	279 (100.0)	198 (71.0)	77 (27.6)	1 (0.4)	3 (1.1)
45-49	434 (100.0)	267 (61.5)	156 (35.9)	9 (2.1)	2 (0.5)
50-54	731 (100.0)	341 (46.6)	381 (52.1)	6 (0.8)	3 (0.4)
55-59	860 (100.0)	268 (31.2)	564 (65.6)	22 (2.6)	6 (0.7)
60-64	1,220 (100.0)	283 (23.2)	893 (73.2)	32 (2.6)	12 (1.0)
65-69	1,258 (100.0)	279 (22.2)	931 (74.0)	35 (2.8)	13 (1.0)
70-74	998 (100.0)	206 (20.6)	756 (75.8)	29 (2.9)	7 (0.7)
75-79	512 (100.0)	102 (19.9)	392 (76.6)	17 (3.3)	1 (0.2)
80歳以上	173 (100.0)	41 (23.7)	128 (74.0)	3 (1.7)	1 (0.6)
計	6,864 (100.0)	2,291 (33.4)	4,370 (63.7)	155 (2.3)	48 (0.7)

表26 血清ASTの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						U/L		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	38	22.6 (10.6)	11.9	17.0	21.0	24.0	51.0	100	16.8 (3.6)	12.0	14.0	16.0	19.0	24.0	
30-39	80	26.9 (12.8)	15.1	19.0	24.0	30.0	48.0	299	17.4 (4.0)	12.0	15.0	17.0	19.0	24.0	
40-44	114	25.5 (8.8)	16.0	19.8	24.0	30.0	41.5	279	19.0 (12.6)	12.0	15.0	17.0	20.0	29.0	
45-49	145	26.0 (13.3)	15.0	19.0	24.0	29.5	40.7	434	19.6 (6.9)	13.0	16.0	18.0	21.0	31.0	
50-54	286	27.8 (15.1)	16.0	21.0	25.0	30.0	48.7	731	21.3 (6.4)	14.0	17.0	20.0	24.0	32.4	
55-59	317	26.6 (10.8)	16.0	21.0	24.0	29.0	44.0	860	22.9 (7.3)	15.0	19.0	22.0	25.0	35.0	
60-64	585	28.1 (18.1)	16.0	21.0	24.0	30.0	46.7	1,220	24.7 (18.9)	16.0	20.0	23.0	26.0	36.0	
65-69	793	28.1 (13.5)	17.0	21.0	25.0	30.0	48.0	1,258	24.6 (9.2)	16.0	20.0	23.0	27.0	37.0	
70-74	705	27.2 (9.7)	17.0	21.0	25.0	31.0	42.0	998	24.6 (7.8)	17.0	20.0	23.0	27.0	37.0	
75-79	470	26.7 (10.7)	17.0	21.0	25.0	30.0	41.5	512	24.5 (7.3)	17.0	20.0	23.0	27.0	36.0	
80歳以上	143	27.0 (12.7)	18.0	21.0	25.0	29.0	42.6	173	23.7 (7.0)	16.0	20.0	23.0	26.0	34.3	
計	3,676	27.3 (13.1)	16.0	21.0	25.0	30.0	44.0	6,864	23.0 (10.9)	15.0	18.0	22.0	25.0	35.0	

表27 血清ALTの状況 (性・年齢階級別)

年齢階級	人 (%)				
	総数	-19	20-39	40-59	60U/L以上
男					
-29歳	38 (100.0)	17 (44.7)	14 (36.8)	2 (5.3)	5 (13.2)
30-39	80 (100.0)	19 (23.8)	38 (47.5)	8 (10.0)	15 (18.8)
40-44	114 (100.0)	26 (22.8)	59 (51.8)	19 (16.7)	10 (8.8)
45-49	145 (100.0)	36 (24.8)	76 (52.4)	26 (17.9)	7 (4.8)
50-54	286 (100.0)	51 (17.8)	185 (64.7)	33 (11.5)	17 (5.9)
55-59	317 (100.0)	93 (29.3)	175 (55.2)	35 (11.0)	14 (4.4)
60-64	585 (100.0)	188 (32.1)	327 (55.9)	45 (7.7)	25 (4.3)
65-69	793 (100.0)	244 (30.8)	456 (57.5)	71 (9.0)	22 (2.8)
70-74	705 (100.0)	263 (37.3)	384 (54.5)	44 (6.2)	14 (2.0)
75-79	470 (100.0)	206 (43.8)	231 (49.1)	30 (6.4)	3 (0.6)
80歳以上	143 (100.0)	77 (53.8)	62 (43.4)	1 (0.7)	3 (2.1)
計	3,676 (100.0)	1,220 (33.2)	2,007 (54.6)	314 (8.5)	135 (3.7)
女					
-29歳	100 (100.0)	79 (79.0)	19 (19.0)	2 (2.0)	0 (0.0)
30-39	299 (100.0)	244 (81.6)	50 (16.7)	3 (1.0)	2 (0.7)
40-44	279 (100.0)	204 (73.1)	65 (23.3)	7 (2.5)	3 (1.1)
45-49	434 (100.0)	308 (71.0)	103 (23.7)	17 (3.9)	6 (1.4)
50-54	731 (100.0)	432 (59.1)	262 (35.8)	29 (4.0)	8 (1.1)
55-59	860 (100.0)	438 (50.9)	376 (43.7)	33 (3.8)	13 (1.5)
60-64	1,220 (100.0)	577 (47.3)	556 (45.6)	59 (4.8)	28 (2.3)
65-69	1,258 (100.0)	641 (51.0)	554 (44.0)	42 (3.3)	21 (1.7)
70-74	998 (100.0)	558 (55.9)	397 (39.8)	31 (3.1)	12 (1.2)
75-79	512 (100.0)	334 (65.2)	163 (31.8)	11 (2.1)	4 (0.8)
80歳以上	173 (100.0)	127 (73.4)	43 (24.9)	3 (1.7)	0 (0.0)
計	6,864 (100.0)	3,942 (57.4)	2,588 (37.7)	237 (3.5)	97 (1.4)

表28 血清ALPの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						U/L		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	38	31.8 (27.7)	9.9	16.8	20.5	32.0	114.0	100	14.6 (7.1)	7.0	10.0	13.0	17.8	27.0	
30-39	80	39.5 (29.2)	12.1	20.3	27.5	47.5	109.7	299	15.6 (8.8)	8.0	11.0	13.0	17.0	30.0	
40-44	114	32.7 (19.1)	12.0	20.0	25.0	40.0	73.0	279	18.2 (13.9)	9.0	12.0	15.0	20.0	37.0	
45-49	145	30.4 (15.1)	13.0	19.5	27.0	37.5	63.2	434	18.7 (12.2)	9.0	12.0	15.0	21.0	43.3	
50-54	286	32.1 (24.5)	15.0	21.0	27.0	35.0	64.0	731	20.4 (10.9)	10.0	14.0	18.0	23.0	40.0	
55-59	317	28.7 (16.0)	13.0	19.0	25.0	33.0	57.1	860	22.3 (13.9)	11.0	15.0	19.0	25.0	40.0	
60-64	585	27.7 (19.8)	13.0	18.0	23.0	31.0	57.7	1,220	23.9 (31.3)	12.0	16.0	20.0	26.0	48.0	
65-69	793	26.4 (13.5)	13.0	18.0	23.0	31.0	51.0	1,258	22.1 (12.7)	12.0	16.0	19.0	25.0	40.0	
70-74	705	24.6 (12.2)	12.0	17.0	22.0	28.0	44.7	998	20.9 (9.7)	12.0	15.0	19.0	24.0	38.0	
75-79	470	22.8 (10.2)	12.0	16.0	20.0	26.0	43.5	512	19.1 (8.5)	11.0	14.0	17.0	21.0	31.4	
80歳以上	143	21.3 (11.2)	12.0	15.0	19.0	24.0	33.8	173	17.3 (6.6)	9.0	13.0	16.0	20.0	29.0	
計	3,676	26.9 (16.6)	13.0	18.0	23.0	31.0	54.2	6,864	21.0 (16.9)	10.0	14.3	18.0	24.0	39.0	

表29 γ-GTPの状況(性・年齢階級別)

年齢階級	人(%)					
	総数	-24	25-49	50-74	75-99	100U/L以上
男						
-29歳	38 (100.0)	20 (52.6)	10 (26.3)	6 (15.8)	0 (0.0)	2 (5.3)
30-39	80 (100.0)	21 (26.3)	38 (47.5)	11 (13.8)	3 (3.8)	7 (8.8)
40-44	114 (100.0)	22 (19.3)	42 (36.8)	17 (14.9)	14 (12.3)	19 (16.7)
45-49	145 (100.0)	37 (25.5)	51 (35.2)	19 (13.1)	17 (11.7)	21 (14.5)
50-54	286 (100.0)	64 (22.4)	96 (33.6)	54 (18.9)	26 (9.1)	46 (16.1)
55-59	317 (100.0)	80 (25.2)	118 (37.2)	62 (19.6)	18 (5.7)	39 (12.3)
60-64	585 (100.0)	173 (29.6)	238 (40.7)	76 (13.0)	27 (4.6)	71 (12.1)
65-69	793 (100.0)	254 (32.0)	309 (39.0)	90 (11.3)	50 (6.3)	90 (11.3)
70-74	705 (100.0)	272 (38.6)	279 (39.6)	77 (10.9)	39 (5.5)	38 (5.4)
75-79	470 (100.0)	209 (44.5)	177 (37.7)	45 (9.6)	15 (3.2)	24 (5.1)
80歳以上	143 (100.0)	78 (54.5)	45 (31.5)	13 (9.1)	2 (1.4)	5 (3.5)
計	3,676 (100.0)	1,230 (33.5)	1,403 (38.2)	470 (12.8)	211 (5.7)	362 (9.8)

年齢階級	人(%)					
	総数	-24	25-49	50-74	75-99	100U/L以上
女						
-29歳	100 (100.0)	93 (93.0)	6 (6.0)	1 (1.0)	0 (0.0)	0 (0.0)
30-39	299 (100.0)	255 (85.3)	36 (12.0)	4 (1.3)	2 (0.7)	2 (0.7)
40-44	279 (100.0)	231 (82.8)	36 (12.9)	6 (2.2)	2 (0.7)	4 (1.4)
45-49	434 (100.0)	337 (77.6)	70 (16.1)	18 (4.1)	5 (1.2)	4 (0.9)
50-54	731 (100.0)	519 (71.0)	149 (20.4)	34 (4.7)	16 (2.2)	13 (1.8)
55-59	860 (100.0)	545 (63.4)	245 (28.5)	41 (4.8)	16 (1.9)	13 (1.5)
60-64	1,220 (100.0)	824 (67.5)	290 (23.8)	62 (5.1)	19 (1.6)	25 (2.0)
65-69	1,258 (100.0)	882 (70.1)	297 (23.6)	46 (3.7)	16 (1.3)	17 (1.4)
70-74	998 (100.0)	734 (73.5)	216 (21.6)	29 (2.9)	10 (1.0)	9 (0.9)
75-79	512 (100.0)	382 (74.6)	97 (18.9)	22 (4.3)	7 (1.4)	4 (0.8)
80歳以上	173 (100.0)	142 (82.1)	26 (15.0)	5 (2.9)	0 (0.0)	0 (0.0)
計	6,864 (100.0)	4,944 (72.0)	1,468 (21.4)	268 (3.9)	93 (1.4)	91 (1.3)

表30 血清γ-GTPの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						U/L		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	38	34.0 (27.5)	13.0	16.8	22.5	45.3	111.0	100	14.9 (7.4)	9.0	11.0	13.0	16.0	29.8	
30-39	80	48.5 (51.5)	13.1	24.0	31.5	50.8	152.4	299	18.2 (13.4)	8.0	12.0	15.0	19.0	43.0	
40-44	114	64.2 (61.4)	13.8	27.0	43.0	84.3	158.8	279	20.8 (24.0)	10.0	12.0	15.0	20.0	45.0	
45-49	145	59.0 (49.4)	16.0	24.0	40.0	79.5	176.3	434	22.1 (20.0)	9.0	12.0	16.0	23.0	53.3	
50-54	286	73.8 (116.6)	15.0	27.0	41.0	75.0	207.3	731	25.3 (23.7)	10.0	14.0	18.0	27.0	69.4	
55-59	317	60.6 (76.5)	15.0	24.0	40.0	64.0	220.1	860	26.3 (19.6)	11.0	15.0	20.0	30.0	63.0	
60-64	585	57.3 (86.6)	14.0	22.0	33.0	55.0	164.0	1,220	27.1 (28.5)	11.0	15.0	19.5	29.0	65.0	
65-69	793	58.1 (106.7)	15.0	22.0	33.0	56.0	159.0	1,258	24.6 (21.5)	11.0	15.0	19.0	26.0	53.1	
70-74	705	41.8 (44.7)	13.0	20.0	29.0	46.0	110.1	998	22.8 (16.7)	11.0	14.0	18.0	25.0	49.0	
75-79	470	40.2 (55.3)	14.0	18.0	26.0	41.0	101.5	512	24.4 (43.5)	10.0	14.0	18.0	25.0	54.7	
80歳以上	143	42.0 (109.6)	13.0	17.0	24.0	36.0	77.2	173	19.3 (10.4)	10.0	13.0	17.0	21.0	46.3	
計	3,676	53.1 (82.6)	14.0	21.0	32.0	54.0	148.0	6,864	24.2 (24.2)	10.0	14.0	18.0	26.0	56.0	

表31 血清クレアチニンの状況（性・年齢階級別）

年齢階級	総数	人 (%)			
		-0.7	0.8-1.0	1.1-1.4	1.5mg/dl以上
-29歳	38 (100.0)	10 (26.3)	27 (71.1)	1 (2.6)	0 (0.0)
30-39	80 (100.0)	26 (32.5)	53 (66.3)	1 (1.3)	0 (0.0)
40-44	114 (100.0)	31 (27.2)	79 (69.3)	4 (3.5)	0 (0.0)
45-49	145 (100.0)	51 (35.2)	88 (60.7)	6 (4.1)	0 (0.0)
50-54	286 (100.0)	104 (36.4)	171 (59.8)	11 (3.8)	0 (0.0)
55-59	317 (100.0)	104 (32.8)	197 (62.1)	15 (4.7)	1 (0.3)
60-64	585 (100.0)	182 (31.1)	370 (63.2)	31 (5.3)	2 (0.3)
65-69	793 (100.0)	243 (30.6)	498 (62.8)	45 (5.7)	7 (0.9)
70-74	705 (100.0)	195 (27.7)	444 (63.0)	63 (8.9)	3 (0.4)
75-79	470 (100.0)	100 (21.3)	306 (65.1)	54 (11.5)	10 (2.1)
80歳以上	143 (100.0)	25 (17.5)	92 (64.3)	23 (16.1)	3 (2.1)
計	3,676 (100.0)	1,071 (29.1)	2,325 (63.2)	254 (6.9)	26 (0.7)

年齢階級	総数	人 (%)			
		-0.7	0.8-1.0	1.1-1.4	1.5mg/dl以上
-29歳	100 (100.0)	95 (95.0)	5 (5.0)	0 (0.0)	0 (0.0)
30-39	299 (100.0)	278 (93.0)	20 (6.7)	1 (0.3)	0 (0.0)
40-44	279 (100.0)	256 (91.8)	23 (8.2)	0 (0.0)	0 (0.0)
45-49	434 (100.0)	400 (92.2)	33 (7.6)	1 (0.2)	0 (0.0)
50-54	731 (100.0)	678 (92.7)	52 (7.1)	0 (0.0)	1 (0.1)
55-59	860 (100.0)	778 (90.5)	80 (9.3)	2 (0.2)	0 (0.0)
60-64	1,220 (100.0)	1,067 (87.5)	150 (12.3)	3 (0.2)	0 (0.0)
65-69	1,258 (100.0)	1,061 (84.3)	193 (15.3)	3 (0.2)	1 (0.1)
70-74	998 (100.0)	797 (79.9)	194 (19.4)	5 (0.5)	2 (0.2)
75-79	512 (100.0)	400 (78.1)	104 (20.3)	5 (1.0)	3 (0.6)
80歳以上	173 (100.0)	122 (70.5)	46 (26.6)	3 (1.7)	2 (1.2)
計	6,864 (100.0)	5,932 (86.4)	900 (13.1)	23 (0.3)	9 (0.1)

表32 血清クレアチニンの平均値、標準偏差および分布 (性・年齢階級別)

年齢階級	mg/dl													
	男						女							
	人数	平均値 (SD)	パーセンタイル					人数	平均値 (SD)	パーセンタイル				
5			25	50	75	95	5			25	50	75	95	
29歳以下	38	0.8 (0.1)	0.7	0.7	0.8	0.9	1.0	100	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
30-39	80	0.8 (0.1)	0.7	0.7	0.8	0.9	0.9	299	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
40-44	114	0.8 (0.1)	0.7	0.7	0.8	0.9	1.0	279	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
45-49	145	0.8 (0.1)	0.6	0.7	0.8	0.9	1.0	434	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
50-54	286	0.8 (0.1)	0.6	0.7	0.8	0.9	1.0	731	0.6 (0.3)	0.5	0.6	0.6	0.7	0.8
55-59	317	0.8 (0.1)	0.6	0.7	0.8	0.9	1.1	860	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
60-64	585	0.8 (0.4)	0.6	0.7	0.8	0.9	1.1	1,220	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
65-69	793	0.8 (0.2)	0.6	0.7	0.8	0.9	1.1	1,258	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8
70-74	705	0.8 (0.2)	0.6	0.7	0.8	0.9	1.1	998	0.7 (0.2)	0.5	0.6	0.6	0.7	0.9
75-79	470	0.9 (0.2)	0.6	0.8	0.9	1.0	1.2	512	0.7 (0.1)	0.5	0.6	0.7	0.7	0.9
80歳以上	143	0.9 (0.2)	0.7	0.8	0.9	1.0	1.3	173	0.7 (0.2)	0.5	0.6	0.7	0.8	1.0
計	3,676	0.8 (0.2)	0.6	0.7	0.8	0.9	1.1	6,864	0.6 (0.1)	0.5	0.6	0.6	0.7	0.8

表33 血糖の状況（空腹時採血者のみ、性・年齢階級別）

年齢階級	総数					人 (%)				
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
男										
-29歳	7 (100.0)	3 (42.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
30-39	22 (100.0)	11 (50.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
40-44	19 (100.0)	15 (78.9)	2 (10.5)	0 (0.0)	0 (0.0)	0 (0.0)				
45-49	35 (100.0)	21 (60.0)	5 (14.3)	0 (0.0)	3 (8.6)	0 (0.0)				
50-54	59 (100.0)	28 (47.5)	14 (23.7)	5 (8.5)	4 (6.8)	1 (1.7)				
55-59	71 (100.0)	45 (63.4)	16 (22.5)	3 (4.2)	2 (2.8)	0 (0.0)				
60-64	108 (100.0)	61 (56.5)	27 (25.0)	5 (4.6)	2 (1.9)	0 (0.0)				
65-69	183 (100.0)	110 (60.1)	26 (14.2)	9 (4.9)	12 (6.6)	5 (2.7)				
70-74	194 (100.0)	117 (60.3)	46 (23.7)	9 (4.6)	7 (3.6)	1 (0.5)				
75-79	130 (100.0)	87 (66.9)	13 (10.0)	4 (3.1)	5 (3.8)	1 (0.8)				
80歳以上	45 (100.0)	22 (48.9)	10 (22.2)	4 (8.9)	4 (8.9)	1 (2.2)				
計	873 (100.0)	520 (59.6)	159 (18.2)	39 (4.5)	39 (4.5)	9 (1.0)				

年齢階級	総数					人 (%)				
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
女										
-29歳	23 (100.0)	9 (39.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)				
30-39	66 (100.0)	32 (48.5)	1 (1.5)	0 (0.0)	0 (0.0)	0 (0.0)				
40-44	44 (100.0)	25 (56.8)	1 (2.3)	2 (4.5)	0 (0.0)	0 (0.0)				
45-49	71 (100.0)	41 (57.7)	2 (2.8)	3 (4.2)	1 (1.4)	0 (0.0)				
50-54	126 (100.0)	88 (69.8)	11 (8.7)	3 (2.4)	2 (1.6)	0 (0.0)				
55-59	142 (100.0)	95 (66.9)	18 (12.7)	5 (3.5)	4 (2.8)	1 (0.7)				
60-64	266 (100.0)	159 (59.8)	33 (12.4)	12 (4.5)	13 (4.9)	3 (1.1)				
65-69	264 (100.0)	171 (64.8)	35 (13.3)	9 (3.4)	4 (1.5)	1 (0.4)				
70-74	248 (100.0)	155 (62.5)	34 (13.7)	9 (3.6)	9 (3.6)	2 (0.8)				
75-79	125 (100.0)	73 (58.4)	22 (17.6)	6 (4.8)	5 (4.0)	2 (1.6)				
80歳以上	41 (100.0)	24 (58.5)	6 (14.6)	4 (9.8)	4 (9.8)	0 (0.0)				
計	1,416 (100.0)	872 (61.6)	163 (11.5)	53 (3.7)	42 (3.0)	9 (0.6)				

表34 血糖の平均値、標準偏差および分布（空腹時採血者のみ、性・年齢階級別）

年齢階級	男										女									
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル										
			5	25	50	75	95			5	25	50	75	95						
29歳以下	7	88.7 (5.4)	82.0	84.0	89.0	97.0	97.0	23	87.5 (7.3)	72.0	82.0	88.0	92.0	99.0						
30-39	22	91.4 (7.7)	82.0	84.0	89.5	97.8	106.7	66	89.5 (6.8)	79.0	85.0	89.5	93.0	101.7						
40-44	19	98.9 (7.6)	88.0	94.0	98.0	105.0	107.0	44	94.7 (11.7)	81.0	88.0	92.0	101.8	124.8						
45-49	35	104.4 (21.2)	84.4	94.0	100.0	109.0	161.8	71	96.5 (13.6)	82.6	88.0	93.0	101.0	131.6						
50-54	59	111.3 (26.9)	87.0	97.0	104.0	117.0	165.0	126	98.9 (13.4)	84.0	91.0	96.0	103.0	119.7						
55-59	71	104.9 (13.4)	87.6	95.0	101.0	115.0	135.8	142	102.3 (18.8)	85.0	92.0	98.0	106.0	130.9						
60-64	108	104.4 (14.4)	88.0	93.3	101.0	111.8	129.8	266	104.1 (21.7)	85.4	92.0	98.5	109.0	143.0						
65-69	183	110.4 (32.9)	86.2	95.0	103.0	112.0	164.2	264	100.7 (14.5)	84.0	91.3	98.0	106.0	126.0						
70-74	194	106.5 (19.7)	88.0	96.0	101.0	113.0	136.0	248	102.8 (20.3)	84.5	92.0	98.0	107.0	137.2						
75-79	130	102.6 (16.6)	86.0	93.0	99.5	107.0	137.2	125	105.7 (21.9)	84.3	93.5	101.0	111.0	150.7						
80歳以上	45	111.9 (25.4)	83.6	96.0	104.0	119.5	166.9	41	110.7 (26.0)	85.3	94.5	102.0	113.0	186.5						
計	873	106.2 (22.7)	86.0	94.0	101.0	111.0	143.0	1,416	101.3 (18.6)	84.0	91.0	97.0	105.8	132.0						

表35 血糖の状況 (食後4時間以上, 性・年齢階級別)

年齢階級	人 (%)						
	総数	-89	90-109	110-125	126-139	140-199	200mg/dl以上
-29歳	15 (100.0)	9 (60.0)	6 (40.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	31 (100.0)	13 (41.9)	17 (54.8)	1 (3.2)	0 (0.0)	0 (0.0)	0 (0.0)
40-44	30 (100.0)	3 (10.0)	24 (80.0)	3 (10.0)	0 (0.0)	0 (0.0)	0 (0.0)
45-49	48 (100.0)	8 (16.7)	30 (62.5)	5 (10.4)	1 (2.1)	3 (6.3)	1 (2.1)
50-54	96 (100.0)	12 (12.5)	53 (55.2)	19 (19.8)	6 (6.3)	5 (5.2)	1 (1.0)
55-59	103 (100.0)	10 (9.7)	59 (57.3)	23 (22.3)	6 (5.8)	3 (2.9)	2 (1.9)
60-64	173 (100.0)	24 (13.9)	94 (54.3)	41 (23.7)	6 (3.5)	7 (4.0)	1 (0.6)
65-69	251 (100.0)	27 (10.8)	151 (60.2)	39 (15.5)	13 (5.2)	15 (6.0)	6 (2.4)
70-74	250 (100.0)	18 (7.2)	148 (59.2)	57 (22.8)	13 (5.2)	11 (4.4)	3 (1.2)
75-79	169 (100.0)	24 (14.2)	109 (64.5)	21 (12.4)	8 (4.7)	6 (3.6)	1 (0.6)
80歳以上	58 (100.0)	5 (8.6)	29 (50.0)	12 (20.7)	6 (10.3)	5 (8.6)	1 (1.7)
計	1,224 (100.0)	153 (12.5)	720 (58.8)	221 (18.1)	59 (4.8)	55 (4.5)	16 (1.3)

年齢階級	人 (%)						
	総数	-89	90-109	110-125	126-139	140-199	200mg/dl以上
-29歳	34 (100.0)	21 (61.8)	13 (38.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	115 (100.0)	54 (47.0)	59 (51.3)	2 (1.7)	0 (0.0)	0 (0.0)	0 (0.0)
40-44	86 (100.0)	28 (32.6)	55 (64.0)	1 (1.2)	2 (2.3)	0 (0.0)	0 (0.0)
45-49	123 (100.0)	40 (32.5)	73 (59.3)	5 (4.1)	4 (3.3)	1 (0.8)	0 (0.0)
50-54	222 (100.0)	42 (18.9)	154 (69.4)	18 (8.1)	5 (2.3)	3 (1.4)	0 (0.0)
55-59	251 (100.0)	33 (13.1)	172 (68.5)	32 (12.7)	7 (2.8)	6 (2.4)	1 (0.4)
60-64	396 (100.0)	56 (14.1)	244 (61.6)	56 (14.1)	18 (4.5)	16 (4.0)	6 (1.5)
65-69	390 (100.0)	57 (14.6)	257 (65.9)	54 (13.8)	13 (3.3)	8 (2.1)	1 (0.3)
70-74	339 (100.0)	47 (13.9)	209 (61.7)	49 (14.5)	16 (4.7)	16 (4.7)	2 (0.6)
75-79	178 (100.0)	20 (11.2)	97 (54.5)	38 (21.3)	11 (6.2)	9 (5.1)	3 (1.7)
80歳以上	59 (100.0)	4 (6.8)	33 (55.9)	12 (20.3)	4 (6.8)	6 (10.2)	0 (0.0)
計	2,193 (100.0)	402 (18.3)	1,366 (62.3)	267 (12.2)	80 (3.6)	65 (3.0)	13 (0.6)

表36 血糖の平均値、標準偏差および分布 (食後4時間以上, 性・年齢階級別)

年齢階級	男						女							
	パーセントाइル						パーセントाइル							
	人数	平均値(SD)	5	25	50	75	95	人数	平均値(SD)	5	25	50	75	95
29歳以下	15	87.6 (10.5)	61.0	84.0	86.0	95.0	97.0	34	88.3 (8.0)	74.8	82.8	88.0	92.3	106.0
30-39	31	92.7 (8.3)	82.0	87.0	90.0	100.0	109.8	115	90.1 (7.0)	79.0	86.0	90.0	95.0	101.2
40-44	30	97.8 (7.9)	83.5	92.8	96.5	102.0	113.9	86	93.6 (9.5)	81.0	88.8	92.0	98.3	108.7
45-49	48	106.9 (31.9)	83.9	94.0	100.0	108.0	173.7	123	95.8 (12.4)	80.4	88.0	94.0	101.0	122.6
50-54	96	107.3 (23.1)	86.0	94.0	102.0	115.8	146.4	222	98.9 (12.8)	85.0	91.0	97.0	103.0	119.0
55-59	103	108.1 (28.5)	86.2	94.0	101.0	117.0	139.4	251	101.6 (16.8)	85.0	92.0	98.0	106.0	129.4
60-64	173	105.9 (19.1)	86.0	94.0	102.0	113.5	140.2	396	105.4 (25.8)	86.0	93.0	99.0	109.0	143.0
65-69	251	109.8 (30.4)	86.0	95.0	103.0	113.0	160.6	390	101.5 (14.6)	85.0	92.0	99.0	107.0	127.0
70-74	250	108.0 (22.9)	88.0	96.0	102.0	114.0	143.0	339	104.1 (19.9)	85.0	93.0	99.0	109.0	141.0
75-79	169	103.6 (16.9)	86.0	94.0	100.0	108.0	133.5	178	108.2 (23.0)	85.0	95.0	103.0	113.3	152.4
80歳以上	58	111.3 (23.7)	83.0	96.8	104.5	119.3	162.4	59	111.0 (23.9)	85.0	99.0	104.0	114.0	182.0
計	1,224	106.6 (24.1)	86.0	94.0	101.0	112.0	143.0	2,193	101.7 (19.1)	84.0	91.0	98.0	106.0	131.0

表37 HbA1cの状況 (性・年齢階級別)

年齢階級	総数	人 (%)				
		-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上
男						
-29歳	38 (100.0)	8 (21.1)	30 (78.9)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	80 (100.0)	15 (18.8)	62 (77.5)	3 (3.8)	0 (0.0)	0 (0.0)
40-44	114 (100.0)	13 (11.4)	95 (83.3)	4 (3.5)	1 (0.9)	1 (0.9)
45-49	145 (100.0)	8 (5.5)	120 (82.8)	12 (8.3)	1 (0.7)	4 (2.8)
50-54	286 (100.0)	24 (8.4)	227 (79.4)	25 (8.7)	6 (2.1)	4 (1.4)
55-59	317 (100.0)	10 (3.2)	255 (80.4)	38 (12.0)	10 (3.2)	4 (1.3)
60-64	585 (100.0)	22 (3.8)	467 (79.8)	70 (12.0)	16 (2.7)	10 (1.7)
65-69	793 (100.0)	34 (4.3)	607 (76.5)	105 (13.2)	26 (3.3)	21 (2.6)
70-74	704 (100.0)	25 (3.6)	533 (75.7)	107 (15.2)	24 (3.4)	15 (2.1)
75-79	470 (100.0)	23 (4.9)	362 (77.0)	63 (13.4)	16 (3.4)	6 (1.3)
80歳以上	143 (100.0)	5 (3.5)	109 (76.2)	23 (16.1)	5 (3.5)	1 (0.7)
計	3,675 (100.0)	187 (5.1)	2,867 (78.0)	450 (12.2)	105 (2.9)	66 (1.8)

年齢階級	総数	人 (%)				
		-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上
女						
-29歳	100 (100.0)	18 (18.0)	81 (81.0)	1 (1.0)	0 (0.0)	0 (0.0)
30-39	299 (100.0)	46 (15.4)	248 (82.9)	5 (1.7)	0 (0.0)	0 (0.0)
40-44	279 (100.0)	29 (10.4)	241 (86.4)	8 (2.9)	1 (0.4)	0 (0.0)
45-49	434 (100.0)	45 (10.4)	364 (83.9)	19 (4.4)	4 (0.9)	2 (0.5)
50-54	731 (100.0)	42 (5.7)	603 (82.5)	73 (10.0)	9 (1.2)	4 (0.5)
55-59	860 (100.0)	26 (3.0)	724 (84.2)	90 (10.5)	8 (0.9)	12 (1.4)
60-64	1,220 (100.0)	29 (2.4)	974 (79.8)	174 (14.3)	20 (1.6)	23 (1.9)
65-69	1,257 (100.0)	30 (2.4)	1,003 (79.8)	186 (14.8)	25 (2.0)	13 (1.0)
70-74	998 (100.0)	18 (1.8)	779 (78.1)	170 (17.0)	21 (2.1)	10 (1.0)
75-79	512 (100.0)	5 (1.0)	395 (77.1)	94 (18.4)	14 (2.7)	4 (0.8)
80歳以上	173 (100.0)	2 (1.2)	139 (80.3)	23 (13.3)	5 (2.9)	4 (2.3)
計	6,863 (100.0)	290 (4.2)	5,551 (80.9)	843 (12.3)	107 (1.6)	72 (1.0)

表38 HbA1cの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男					女							
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル				
			5	25	50	75			95	5	25	50	75
29歳以下	38	4.7 (0.3)	4.0	4.5	4.8	5.0	100	4.7 (0.3)	4.3	4.5	4.7	4.9	5.2
30-39	80	4.7 (0.3)	4.2	4.5	4.8	4.9	299	4.7 (0.3)	4.2	4.5	4.7	4.9	5.4
40-44	114	4.9 (0.5)	4.3	4.6	4.8	5.0	279	4.8 (0.3)	4.4	4.6	4.8	5.0	5.5
45-49	145	5.1 (0.8)	4.4	4.7	4.9	5.1	434	4.9 (0.5)	4.3	4.7	4.9	5.1	6.0
50-54	286	5.1 (0.6)	4.4	4.7	5.0	5.2	731	5.0 (0.5)	4.4	4.8	5.0	5.2	6.0
55-59	317	5.1 (0.7)	4.5	4.8	5.0	5.3	860	5.1 (0.6)	4.5	4.8	5.0	5.3	6.3
60-64	585	5.2 (0.6)	4.5	4.8	5.0	5.3	1,220	5.2 (0.7)	4.5	4.9	5.1	5.3	6.3
65-69	793	5.2 (0.7)	4.5	4.8	5.1	5.4	1,257	5.2 (0.6)	4.6	4.9	5.1	5.3	6.6
70-74	704	5.2 (0.7)	4.5	4.8	5.0	5.4	998	5.2 (0.6)	4.6	4.9	5.1	5.4	6.6
75-79	470	5.2 (0.6)	4.5	4.9	5.1	5.3	512	5.2 (0.5)	4.7	4.9	5.1	5.4	6.3
80歳以上	143	5.1 (0.6)	4.5	4.8	5.0	5.3	173	5.2 (0.8)	4.6	4.9	5.1	5.3	6.2
計	3,675	5.1 (0.7)	4.4	4.8	5.0	5.3	6,863	5.1 (0.6)	4.5	4.8	5.0	5.3	6.4

表39 赤血球数の状況 (性・年齢階級別)

年齢階級	総数					人 (%)	
	-399	400-449	450-499	500-549	550×10 ⁴ /μl以上		
男							
-29歳	38 (100.0)	2 (5.3)	9 (23.7)	23 (60.5)	4 (10.5)		
30-39	80 (100.0)	5 (6.3)	37 (46.3)	29 (36.3)	9 (11.3)		
40-44	114 (100.0)	20 (17.5)	55 (48.2)	36 (31.6)	3 (2.6)		
45-49	145 (100.0)	32 (22.1)	65 (44.8)	47 (32.4)	0 (0.0)		
50-54	286 (100.0)	52 (18.2)	151 (52.8)	74 (25.9)	4 (1.4)		
55-59	317 (100.0)	63 (19.9)	179 (56.5)	61 (19.2)	5 (1.6)		
60-64	585 (100.0)	176 (30.1)	260 (44.4)	120 (20.5)	8 (1.4)		
65-69	793 (100.0)	223 (28.1)	383 (48.3)	129 (16.3)	15 (1.9)		
70-74	705 (100.0)	263 (37.3)	296 (42.0)	78 (11.1)	9 (1.3)		
75-79	470 (100.0)	167 (35.5)	189 (40.2)	58 (12.3)	3 (0.6)		
80歳以上	143 (100.0)	59 (41.3)	42 (29.4)	9 (6.3)	1 (0.7)		
計	3,676 (100.0)	1,062 (28.9)	1,666 (45.3)	664 (18.1)	61 (1.7)		

年齢階級	総数					人 (%)	
	-399	400-449	450-499	500-549	550×10 ⁴ /μl以上		
女							
-29歳	100 (100.0)	60 (60.0)	25 (25.0)	2 (-2.0)	0 (0.0)		
30-39	299 (100.0)	178 (59.5)	57 (19.1)	8 (2.7)	1 (0.3)		
40-44	279 (100.0)	159 (57.0)	64 (22.9)	3 (1.1)	0 (0.0)		
45-49	434 (100.0)	239 (55.1)	95 (21.9)	10 (2.3)	0 (0.0)		
50-54	731 (100.0)	386 (52.8)	188 (25.7)	22 (3.0)	2 (0.3)		
55-59	860 (100.0)	469 (54.5)	219 (25.5)	26 (3.0)	2 (0.2)		
60-64	1,220 (100.0)	651 (53.4)	294 (24.1)	29 (2.4)	0 (0.0)		
65-69	1,258 (100.0)	648 (51.5)	286 (22.7)	23 (1.8)	1 (0.1)		
70-74	998 (100.0)	487 (48.8)	195 (19.5)	17 (1.7)	1 (0.1)		
75-79	512 (100.0)	238 (46.5)	90 (17.6)	5 (1.0)	1 (0.2)		
80歳以上	173 (100.0)	76 (43.9)	23 (13.3)	1 (0.6)	0 (0.0)		
計	6,864 (100.0)	3,591 (52.3)	1,536 (22.4)	146 (2.1)	8 (0.1)		

表40 赤血球数の平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女							
	人数	平均値 (SD)	パーセンタイル					人数	平均値 (SD)	パーセンタイル				
			5	25	50	75	95			5	25	50	75	95
29歳以下	38	512.9 (34.7)	440.3	487.5	517.5	538.5	561.2	100	432.2 (28.8)	384.1	412.8	430.0	450.0	483.0
30-39	80	500.8 (35.1)	449.0	477.5	498.0	524.8	558.0	299	428.9 (32.1)	383.0	409.0	426.0	447.0	482.0
40-44	114	486.1 (32.1)	431.5	463.3	490.0	506.0	541.3	279	427.0 (31.9)	374.0	406.0	429.0	449.0	479.0
45-49	145	478.7 (35.6)	415.8	456.0	483.0	506.0	541.7	434	426.9 (34.3)	372.8	403.0	425.0	449.0	488.3
50-54	286	478.5 (36.1)	416.4	455.0	479.5	502.3	537.0	731	431.6 (35.1)	378.0	407.0	430.0	453.0	492.0
55-59	317	473.4 (35.4)	410.9	453.0	475.0	496.0	533.0	860	432.7 (34.2)	378.0	411.0	432.0	454.0	491.0
60-64	585	466.6 (38.3)	404.3	441.0	466.0	495.0	523.7	1,220	428.8 (34.4)	373.1	406.0	428.0	452.0	485.0
65-69	793	463.1 (42.8)	395.7	438.0	465.0	491.0	526.0	1,258	425.9 (35.5)	370.0	401.0	425.0	449.0	484.0
70-74	705	453.5 (43.2)	386.0	427.0	455.0	482.0	520.0	998	419.6 (37.3)	360.0	395.0	419.0	445.3	479.1
75-79	470	450.5 (43.6)	381.0	424.8	452.0	478.0	518.9	512	414.9 (38.7)	350.7	389.0	413.5	442.8	481.0
80歳以上	143	433.0 (47.5)	353.6	402.0	435.0	465.0	520.6	173	404.6 (39.2)	339.7	380.5	406.0	428.0	467.6
計	3,676	463.8 (42.8)	395.0	437.0	465.0	493.0	530.0	6,864	425.9 (35.8)	368.0	402.0	425.0	449.0	484.0

表41 ヘモグロビン値の状況 (性・年齢階級別)

年齢階級	人 (%)						
	総数	-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上
男							
-29歳	38 (100.0)	0 (0.0)	0 (0.0)	1 (2.6)	7 (18.4)	14 (36.8)	16 (42.1)
30-39	80 (100.0)	0 (0.0)	1 (1.3)	2 (2.5)	15 (18.8)	39 (48.8)	23 (28.8)
40-44	114 (100.0)	0 (0.0)	2 (1.8)	8 (7.0)	34 (29.8)	48 (42.1)	22 (19.3)
45-49	145 (100.0)	0 (0.0)	3 (2.1)	9 (6.2)	56 (38.6)	50 (34.5)	27 (18.6)
50-54	286 (100.0)	2 (0.7)	4 (1.4)	23 (8.0)	95 (33.2)	93 (32.5)	69 (24.1)
55-59	317 (100.0)	2 (0.6)	9 (2.8)	34 (10.7)	114 (36.0)	106 (33.4)	52 (16.4)
60-64	585 (100.0)	2 (0.3)	25 (4.3)	77 (13.2)	193 (33.0)	191 (32.6)	97 (16.6)
65-69	793 (100.0)	22 (2.8)	36 (4.5)	125 (15.8)	244 (30.8)	248 (31.3)	118 (14.9)
70-74	705 (100.0)	24 (3.4)	45 (6.4)	135 (19.1)	233 (33.0)	181 (25.7)	87 (12.3)
75-79	470 (100.0)	17 (3.6)	37 (7.9)	107 (22.8)	147 (31.3)	116 (24.7)	46 (9.8)
80歳以上	143 (100.0)	14 (9.8)	19 (13.3)	43 (30.1)	37 (25.9)	21 (14.7)	9 (6.3)
計	3,676 (100.0)	83 (2.3)	181 (4.9)	564 (15.3)	1,175 (32.0)	1,107 (30.1)	566 (15.4)

年齢階級	人 (%)						
	総数	-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上
女							
-29歳	100 (100.0)	18 (18.0)	34 (34.0)	36 (36.0)	10 (10.0)	2 (2.0)	0 (0.0)
30-39	299 (100.0)	64 (21.4)	89 (29.8)	104 (34.8)	35 (11.7)	7 (2.3)	0 (0.0)
40-44	279 (100.0)	81 (29.0)	77 (27.6)	90 (32.3)	29 (10.4)	2 (0.7)	0 (0.0)
45-49	434 (100.0)	124 (28.6)	125 (28.8)	108 (24.9)	71 (16.4)	5 (1.2)	1 (0.2)
50-54	731 (100.0)	107 (14.6)	208 (28.5)	285 (39.0)	116 (15.9)	15 (2.1)	0 (0.0)
55-59	860 (100.0)	71 (8.3)	242 (28.1)	356 (41.4)	163 (19.0)	27 (3.1)	1 (0.1)
60-64	1,220 (100.0)	131 (10.7)	337 (27.6)	475 (38.9)	235 (19.3)	39 (3.2)	3 (0.2)
65-69	1,258 (100.0)	126 (10.0)	382 (30.4)	475 (37.8)	231 (18.4)	43 (3.4)	1 (0.1)
70-74	998 (100.0)	153 (15.3)	313 (31.4)	335 (33.6)	167 (16.7)	28 (2.8)	2 (0.2)
75-79	512 (100.0)	106 (20.7)	151 (29.5)	172 (33.6)	65 (12.7)	17 (3.3)	1 (0.2)
80歳以上	173 (100.0)	47 (27.2)	55 (31.8)	51 (29.5)	18 (10.4)	2 (1.2)	0 (0.0)
計	6,864 (100.0)	1,028 (15.0)	2,013 (29.3)	2,487 (36.2)	1,140 (16.6)	187 (2.7)	9 (0.1)

表42 ヘモグロビン値の平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						g/dl		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	38	15.7 (1.0)	14.0	15.2	15.9	16.3	17.6	100	12.8 (1.2)	10.7	12.3	12.9	13.5	14.4	
30-39	80	15.6 (0.8)	14.1	15.1	15.6	16.0	17.1	299	12.7 (1.4)	9.8	12.2	12.9	13.5	14.7	
40-44	114	15.2 (0.9)	13.8	14.6	15.3	15.7	16.6	279	12.4 (1.5)	9.5	11.8	12.7	13.5	14.4	
45-49	145	15.1 (0.9)	13.7	14.5	15.0	15.8	16.7	434	12.5 (1.5)	9.4	11.7	12.7	13.7	14.6	
50-54	286	15.1 (1.1)	13.4	14.5	15.1	15.9	16.9	731	13.0 (1.2)	10.9	12.3	13.1	13.7	14.7	
55-59	317	15.0 (1.1)	13.1	14.3	14.9	15.6	16.7	860	13.2 (1.0)	11.7	12.6	13.3	13.9	14.7	
60-64	585	14.9 (1.1)	13.0	14.2	14.9	15.6	16.7	1,220	13.2 (1.0)	11.6	12.6	13.2	13.9	14.8	
65-69	793	14.7 (1.3)	12.7	14.0	14.8	15.6	16.5	1,258	13.2 (1.0)	11.5	12.5	13.2	13.8	14.9	
70-74	705	14.5 (1.4)	12.3	13.8	14.6	15.5	16.5	998	13.0 (1.1)	11.2	12.3	13.1	13.8	14.7	
75-79	470	14.4 (1.3)	12.3	13.6	14.4	15.3	16.3	512	12.9 (1.2)	10.9	12.1	12.9	13.7	14.7	
80歳以上	143	13.8 (1.5)	11.1	13.0	13.8	14.8	16.1	173	12.6 (1.2)	10.4	11.9	12.7	13.5	14.5	
計	3,676	14.7 (1.3)	12.6	14.0	14.8	15.6	16.6	6,864	13.0 (1.2)	11.1	12.4	13.1	13.8	14.7	

表43 ハマトクレット値の状況 (性・年齢階級別)

男	人 (%)						
年齢階級	総数	-29.9	30.0-34.9	35.0-39.9	40.0-44.9	45.0-49.9	50.0%以上
-29歳	38 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	10 (26.3)	23 (60.5)	5 (13.2)
30-39	80 (100.0)	0 (0.0)	0 (0.0)	1 (1.3)	21 (26.3)	52 (65.0)	6 (7.5)
40-44	114 (100.0)	0 (0.0)	0 (0.0)	1 (0.9)	48 (42.1)	62 (54.4)	3 (2.6)
45-49	145 (100.0)	0 (0.0)	0 (0.0)	4 (2.8)	75 (51.7)	63 (43.4)	3 (2.1)
50-54	286 (100.0)	0 (0.0)	0 (0.0)	9 (3.1)	131 (45.8)	130 (45.5)	16 (5.6)
55-59	317 (100.0)	1 (0.3)	0 (0.0)	25 (7.9)	141 (44.5)	141 (44.5)	9 (2.8)
60-64	585 (100.0)	1 (0.2)	1 (0.2)	46 (7.9)	270 (46.2)	247 (42.2)	20 (3.4)
65-69	793 (100.0)	1 (0.1)	14 (1.8)	76 (9.6)	355 (44.8)	317 (40.0)	30 (3.8)
70-74	705 (100.0)	4 (0.6)	13 (1.8)	91 (12.9)	346 (49.1)	231 (32.8)	20 (2.8)
75-79	470 (100.0)	4 (0.9)	9 (1.9)	73 (15.5)	232 (49.4)	141 (30.0)	11 (2.3)
80歳以上	143 (100.0)	1 (0.7)	8 (5.6)	41 (28.7)	66 (46.2)	25 (17.5)	2 (1.4)
計	3,676 (100.0)	12 (0.3)	45 (1.2)	367 (10.0)	1,695 (46.1)	1,432 (39.0)	125 (3.4)

女	人 (%)						
年齢階級	総数	-29.9	30.0-34.9	35.0-39.9	40.0-44.9	45.0-49.9	50.0%以上
-29歳	100 (100.0)	2 (2.0)	6 (6.0)	59 (59.0)	31 (31.0)	2 (2.0)	0 (0.0)
30-39	299 (100.0)	7 (2.3)	25 (8.4)	157 (52.5)	104 (34.8)	6 (2.0)	0 (0.0)
40-44	279 (100.0)	9 (3.2)	41 (14.7)	139 (49.8)	89 (31.9)	1 (0.4)	0 (0.0)
45-49	434 (100.0)	11 (2.5)	64 (14.7)	205 (47.2)	147 (33.9)	6 (1.4)	1 (0.2)
50-54	731 (100.0)	4 (0.5)	56 (7.7)	338 (46.2)	309 (42.3)	24 (3.3)	0 (0.0)
55-59	860 (100.0)	1 (0.1)	22 (2.6)	380 (44.2)	421 (49.0)	36 (4.2)	0 (0.0)
60-64	1,220 (100.0)	3 (0.2)	43 (3.5)	555 (45.5)	573 (47.0)	46 (3.8)	0 (0.0)
65-69	1,258 (100.0)	2 (0.2)	52 (4.1)	599 (47.6)	547 (43.5)	58 (4.6)	0 (0.0)
70-74	998 (100.0)	4 (0.4)	67 (6.7)	461 (46.2)	429 (43.0)	36 (3.6)	1 (0.1)
75-79	512 (100.0)	1 (0.2)	53 (10.4)	241 (47.1)	197 (38.5)	19 (3.7)	1 (0.2)
80歳以上	173 (100.0)	2 (1.2)	27 (15.6)	86 (49.7)	54 (31.2)	4 (2.3)	0 (0.0)
計	6,864 (100.0)	46 (0.7)	456 (6.6)	3,220 (46.9)	2,901 (42.3)	238 (3.5)	3 (0.0)