

表35 血糖の状況（食後4時間以上，性・年齢階級別）

年齢階級	総数							人 (%)		
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
男										
-29歳	4 (100.0)	2 (50.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	18 (100.0)	8 (44.4)	2 (11.1)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
40-44	28 (100.0)	11 (39.3)	4 (14.3)	1 (3.6)	0 (0.0)	0 (0.0)	1 (3.6)	0 (0.0)	1 (3.6)	1 (3.6)
45-49	41 (100.0)	6 (14.6)	7 (17.1)	1 (2.4)	0 (0.0)	0 (0.0)	1 (2.4)	0 (0.0)	1 (2.4)	1 (2.4)
50-54	63 (100.0)	11 (17.5)	10 (15.9)	5 (7.9)	1 (1.6)	1 (1.6)	2 (3.2)	1 (1.6)	2 (3.2)	2 (3.2)
55-59	83 (100.0)	8 (9.6)	11 (13.3)	4 (4.8)	2 (2.4)	2 (2.4)	1 (1.2)	1 (1.2)	1 (1.2)	1 (1.2)
60-64	133 (100.0)	17 (12.8)	19 (14.3)	7 (5.3)	9 (6.8)	9 (6.8)	2 (1.5)	2 (1.5)	2 (1.5)	2 (1.5)
65-69	173 (100.0)	27 (15.6)	35 (20.2)	9 (5.2)	8 (4.6)	8 (4.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
70-74	158 (100.0)	22 (13.9)	29 (18.4)	11 (7.0)	7 (4.4)	7 (4.4)	4 (2.5)	4 (2.5)	4 (2.5)	4 (2.5)
75-79	68 (100.0)	9 (13.2)	18 (26.5)	4 (5.9)	2 (2.9)	2 (2.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
80歳以上	25 (100.0)	6 (24.0)	2 (8.0)	1 (4.0)	2 (8.0)	2 (8.0)	1 (4.0)	1 (4.0)	1 (4.0)	1 (4.0)
計	794 (100.0)	127 (16.0)	137 (17.3)	43 (5.4)	31 (3.9)	31 (3.9)	12 (1.5)	12 (1.5)	12 (1.5)	12 (1.5)

年齢階級	総数							200mg/dl以上		
	-89	90-109	110-125	126-139	140-199	200mg/dl以上				
女										
-29歳	11 (100.0)	4 (36.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	44 (100.0)	24 (54.5)	1 (2.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
40-44	70 (100.0)	46 (65.7)	3 (4.3)	0 (0.0)	2 (2.9)	2 (2.9)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
45-49	136 (100.0)	92 (67.6)	7 (5.1)	1 (0.7)	1 (0.7)	1 (0.7)	1 (0.7)	1 (0.7)	1 (0.7)	1 (0.7)
50-54	206 (100.0)	126 (61.2)	20 (9.7)	6 (2.9)	4 (1.9)	4 (1.9)	2 (1.0)	2 (1.0)	2 (1.0)	2 (1.0)
55-59	194 (100.0)	129 (66.5)	12 (6.2)	5 (2.6)	2 (1.0)	2 (1.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
60-64	274 (100.0)	179 (65.3)	31 (11.3)	10 (3.6)	4 (1.5)	4 (1.5)	1 (0.4)	1 (0.4)	1 (0.4)	1 (0.4)
65-69	269 (100.0)	163 (60.6)	37 (13.8)	10 (3.7)	7 (2.6)	7 (2.6)	1 (0.4)	1 (0.4)	1 (0.4)	1 (0.4)
70-74	203 (100.0)	123 (60.6)	29 (14.3)	9 (4.4)	12 (5.9)	12 (5.9)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)
75-79	95 (100.0)	57 (60.0)	16 (16.8)	5 (5.3)	5 (5.3)	5 (5.3)	2 (2.1)	2 (2.1)	2 (2.1)	2 (2.1)
80歳以上	16 (100.0)	8 (50.0)	2 (12.5)	2 (12.5)	4 (25.0)	4 (25.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
計	1,518 (100.0)	951 (62.6)	158 (10.4)	48 (3.2)	41 (2.7)	41 (2.7)	8 (0.5)	8 (0.5)	8 (0.5)	8 (0.5)

表36 血糖の平均値、標準偏差および分布（食後4時間以上，性・年齢階級別）

年齢階級	男						女						mg/dl		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	4	87.3 (6.9)	78.0	80.0	89.0	92.8	93.0	11	88.4 (5.4)	80.0	84.0	88.0	94.0	95.0	
30-39	18	95.4 (9.1)	83.0	87.8	94.5	102.5	110.0	44	91.0 (8.0)	77.3	85.0	90.0	96.0	105.8	
40-44	28	99.4 (26.6)	75.7	84.5	92.5	101.0	177.9	70	95.8 (15.2)	84.0	88.8	93.0	98.0	121.4	
45-49	41	109.0 (61.0)	81.3	91.5	100.0	108.0	131.2	136	97.0 (13.9)	82.0	89.3	95.0	102.0	115.0	
50-54	63	105.6 (24.4)	86.0	91.0	99.0	114.0	143.2	206	100.4 (22.6)	81.0	90.0	97.0	105.0	130.7	
55-59	83	104.1 (22.0)	84.2	94.0	99.0	107.0	137.0	194	97.5 (13.1)	83.0	90.0	95.0	102.0	120.3	
60-64	133	108.5 (25.1)	83.0	94.0	102.0	114.0	160.6	274	100.7 (16.0)	84.0	92.0	98.0	106.0	126.0	
65-69	173	105.0 (17.7)	81.0	93.0	102.0	112.0	140.5	269	101.8 (17.7)	84.0	92.0	98.0	107.0	133.0	
70-74	158	108.6 (29.8)	83.0	93.0	102.0	115.0	156.4	203	105.0 (20.4)	86.0	92.0	100.0	110.0	144.8	
75-79	68	104.8 (15.3)	85.5	94.0	103.5	115.0	130.9	95	108.7 (29.0)	86.8	96.0	102.0	111.0	145.2	
80歳以上	25	113.4 (50.1)	80.5	90.0	99.0	109.0	283.8	16	123.0 (28.9)	93.0	100.0	111.0	147.5	178.5	
計	794	106.2 (27.7)	83.0	92.0	101.0	111.0	144.0	1,518	100.8 (18.9)	84.0	91.0	97.0	105.0	131.0	

表37 HbA1cの状況 (性・年齢階級別)

年齢階級	総数	人 (%)				
		-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上
-29歳	7 (100.0)	0 (0.0)	7 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	38 (100.0)	1 (2.6)	36 (94.7)	1 (2.6)	0 (0.0)	0 (0.0)
40-44	79 (100.0)	4 (5.1)	68 (86.1)	4 (5.1)	1 (1.3)	2 (2.5)
45-49	96 (100.0)	7 (7.3)	77 (80.2)	8 (8.3)	1 (1.0)	3 (3.1)
50-54	159 (100.0)	10 (6.3)	121 (76.1)	22 (13.8)	4 (2.5)	2 (1.3)
55-59	208 (100.0)	8 (3.8)	159 (76.4)	32 (15.4)	6 (2.9)	3 (1.4)
60-64	340 (100.0)	12 (3.5)	261 (76.8)	50 (14.7)	10 (2.9)	7 (2.1)
65-69	462 (100.0)	12 (2.6)	343 (74.2)	82 (17.7)	15 (3.2)	10 (2.2)
70-74	422 (100.0)	17 (4.0)	301 (71.3)	75 (17.8)	20 (4.7)	9 (2.1)
75-79	219 (100.0)	8 (3.7)	156 (71.2)	45 (20.5)	7 (3.2)	3 (1.4)
80歳以上	92 (100.0)	4 (4.3)	63 (68.5)	18 (19.6)	4 (4.3)	3 (3.3)
計	2,122 (100.0)	83 (3.9)	1,592 (75.0)	337 (15.9)	68 (3.2)	42 (2.0)

年齢階級	総数	人 (%)				
		-4.4	4.5-5.4	5.5-6.4	6.5-7.4	7.5%以上
-29歳	39 (100.0)	10 (25.6)	29 (74.4)	0 (0.0)	0 (0.0)	0 (0.0)
30-39	139 (100.0)	14 (10.1)	124 (89.2)	1 (0.7)	0 (0.0)	0 (0.0)
40-44	211 (100.0)	25 (11.8)	178 (84.4)	7 (3.3)	0 (0.0)	1 (0.5)
45-49	335 (100.0)	24 (7.2)	279 (83.3)	29 (8.7)	1 (0.3)	2 (0.6)
50-54	525 (100.0)	17 (3.2)	440 (83.8)	58 (11.0)	5 (1.0)	5 (1.0)
55-59	589 (100.0)	13 (2.2)	492 (83.5)	73 (12.4)	4 (0.7)	7 (1.2)
60-64	721 (100.0)	21 (2.9)	568 (78.8)	112 (15.5)	11 (1.5)	9 (1.2)
65-69	818 (100.0)	18 (2.2)	636 (77.8)	137 (16.7)	17 (2.1)	10 (1.2)
70-74	637 (100.0)	9 (1.4)	477 (74.9)	129 (20.3)	16 (2.5)	6 (0.9)
75-79	311 (100.0)	4 (1.3)	218 (70.1)	72 (23.2)	9 (2.9)	8 (2.6)
80歳以上	67 (100.0)	2 (3.0)	47 (70.1)	13 (19.4)	4 (6.0)	1 (1.5)
計	4,392 (100.0)	157 (3.6)	3,488 (79.4)	631 (14.4)	67 (1.5)	49 (1.1)

表38 HbA1cの平均値、標準偏差および分布 (性・年齢階級別)

年齢階級	%													
	男					女								
	パーセンタイル					パーセンタイル								
	人数	平均値(SD)	5	25	50	75	95	人数	平均値(SD)	5	25	50	75	95
29歳以下	7	4.9 (0.3)	4.6	4.6	5.0	5.2	5.2	39	4.6 (0.3)	4.1	4.4	4.6	4.8	5.1
30-39	38	4.8 (0.2)	4.5	4.7	4.8	4.9	5.3	139	4.8 (0.3)	4.3	4.6	4.8	5.0	5.3
40-44	79	5.1 (0.7)	4.4	4.8	4.9	5.2	6.1	211	4.8 (0.4)	4.3	4.6	4.8	5.1	5.3
45-49	96	5.1 (1.0)	4.3	4.7	4.9	5.1	6.5	335	5.0 (0.5)	4.4	4.7	4.9	5.2	5.6
50-54	159	5.1 (0.7)	4.4	4.8	5.0	5.3	6.3	525	5.1 (0.6)	4.5	4.8	5.0	5.3	5.7
55-59	208	5.2 (0.8)	4.5	4.8	5.0	5.4	6.4	589	5.1 (0.5)	4.6	4.9	5.1	5.3	5.9
60-64	340	5.2 (0.7)	4.5	4.8	5.1	5.4	6.5	721	5.2 (0.6)	4.5	4.9	5.1	5.3	6.0
65-69	462	5.2 (0.7)	4.6	4.8	5.1	5.4	6.6	818	5.2 (0.6)	4.6	4.9	5.1	5.4	6.0
70-74	422	5.3 (0.8)	4.5	4.9	5.1	5.4	6.8	637	5.2 (0.6)	4.6	4.9	5.1	5.4	6.2
75-79	219	5.3 (0.7)	4.5	4.9	5.1	5.5	6.4	311	5.3 (0.7)	4.6	4.9	5.2	5.5	6.6
80歳以上	92	5.3 (0.7)	4.5	5.0	5.2	5.5	6.8	67	5.3 (0.6)	4.5	4.9	5.2	5.5	6.7
計	2,122	5.2 (0.7)	4.5	4.8	5.1	5.4	6.5	4,392	5.1 (0.6)	4.5	4.8	5.1	5.3	6.0

表39 赤血球数の状況 (性・年齢階級別)

年齢階級	人 (%)					
	総数	-399	400-449	450-499	500-549	550×10 ⁴ /μl以上
男						
-29歳	7 (100.0)	0 (0.0)	0 (0.0)	3 (42.9)	4 (57.1)	0 (0.0)
30-39	38 (100.0)	0 (0.0)	0 (0.0)	18 (47.4)	16 (42.1)	4 (10.5)
40-44	79 (100.0)	3 (3.8)	7 (8.9)	42 (53.2)	21 (26.6)	6 (7.6)
45-49	96 (100.0)	2 (2.1)	15 (15.6)	47 (49.0)	28 (29.2)	4 (4.2)
50-54	159 (100.0)	5 (3.1)	30 (18.9)	84 (52.8)	36 (22.6)	4 (2.5)
55-59	208 (100.0)	5 (2.4)	47 (22.6)	115 (55.3)	34 (16.3)	7 (3.4)
60-64	340 (100.0)	15 (4.4)	88 (25.9)	176 (51.8)	56 (16.5)	5 (1.5)
65-69	462 (100.0)	28 (6.1)	159 (34.4)	221 (47.8)	50 (10.8)	4 (0.9)
70-74	422 (100.0)	52 (12.3)	159 (37.7)	164 (38.9)	47 (11.1)	0 (0.0)
75-79	219 (100.0)	32 (14.6)	84 (38.4)	85 (38.8)	17 (7.8)	1 (0.5)
80歳以上	92 (100.0)	25 (27.2)	38 (41.3)	22 (23.9)	7 (7.6)	0 (0.0)
計	2,122 (100.0)	167 (7.9)	627 (29.5)	977 (46.0)	316 (14.9)	35 (1.6)

年齢階級	人 (%)					
	総数	-399	400-449	450-499	500-549	550×10 ⁴ /μl以上
女						
-29歳	39 (100.0)	5 (12.8)	23 (59.0)	9 (23.1)	2 (5.1)	0 (0.0)
30-39	139 (100.0)	27 (19.4)	74 (53.2)	36 (25.9)	2 (1.4)	0 (0.0)
40-44	211 (100.0)	31 (14.7)	128 (60.7)	49 (23.2)	3 (1.4)	0 (0.0)
45-49	335 (100.0)	70 (20.9)	186 (55.5)	71 (21.2)	7 (2.1)	1 (0.3)
50-54	525 (100.0)	73 (13.9)	276 (52.6)	154 (29.3)	21 (4.0)	1 (0.2)
55-59	589 (100.0)	89 (15.1)	333 (56.5)	156 (26.5)	11 (1.9)	0 (0.0)
60-64	722 (100.0)	151 (20.9)	390 (54.0)	160 (22.2)	20 (2.8)	1 (0.1)
65-69	819 (100.0)	219 (26.7)	435 (53.1)	157 (19.2)	8 (1.0)	0 (0.0)
70-74	637 (100.0)	195 (30.6)	330 (51.8)	101 (15.9)	11 (1.7)	0 (0.0)
75-79	311 (100.0)	107 (34.4)	152 (48.9)	51 (16.4)	1 (0.3)	0 (0.0)
80歳以上	68 (100.0)	27 (39.7)	31 (45.6)	8 (11.8)	2 (2.9)	0 (0.0)
計	4,395 (100.0)	994 (22.6)	2,358 (53.7)	952 (21.7)	88 (2.0)	3 (0.1)

表40 赤血球数の平均値、標準偏差および分布 (性・年齢階級別)

年齢階級	男						女						10 ⁴ /μl		
	人数	平均値 (SD)	パーセンタイル					人数	平均値 (SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	7	502.4 (33.0)	464.0	465.0	500.0	534.0	540.5	39	430.4 (36.7)	372.0	410.0	429.0	454.0	503.0	
30-39	38	502.8 (34.4)	458.7	470.5	503.0	527.5	565.7	139	428.2 (34.0)	375.0	404.0	425.0	451.0	494.0	
40-44	79	487.7 (41.6)	420.0	462.0	482.0	513.0	561.0	211	427.7 (29.7)	377.2	408.0	427.0	449.0	477.8	
45-49	96	482.3 (36.8)	422.2	463.0	480.0	509.5	541.5	335	427.8 (37.2)	374.8	403.0	423.0	448.0	485.0	
50-54	159	473.7 (36.0)	406.0	452.0	475.0	500.0	530.0	525	436.1 (34.4)	379.3	413.0	436.0	458.5	493.7	
55-59	208	471.5 (36.2)	410.0	447.8	471.0	494.8	537.7	589	431.5 (32.6)	376.5	410.0	432.0	453.0	487.0	
60-64	340	466.6 (38.3)	402.0	443.0	464.5	491.8	527.0	722	428.3 (37.9)	374.0	405.0	428.5	450.0	488.9	
65-69	462	457.3 (37.8)	398.0	432.8	458.0	483.0	518.0	819	421.1 (34.0)	367.0	397.0	420.0	445.0	478.0	
70-74	422	447.5 (42.7)	382.2	422.0	449.5	475.0	515.9	637	418.0 (35.3)	359.9	395.0	418.0	441.0	476.1	
75-79	219	443.7 (41.6)	372.0	418.0	445.0	471.0	508.0	311	414.8 (35.2)	360.0	387.0	415.0	440.0	473.8	
80歳以上	92	424.4 (50.2)	327.0	396.0	425.0	458.5	508.0	68	408.5 (40.0)	349.4	376.8	407.0	429.5	487.1	
計	2,122	459.9 (42.5)	389.0	434.0	461.0	487.3	525.9	4,395	425.5 (35.6)	369.0	402.0	425.0	448.0	484.0	

表41 ヘモグロビン値の状況 (性・年齢階級別)

年齢階級	人 (%)						
	総数	-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上
男							
-29歳	7 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (28.6)	3 (42.9)	2 (28.6)
30-39	38 (100.0)	0 (0.0)	0 (0.0)	1 (2.6)	9 (23.7)	13 (34.2)	15 (39.5)
40-44	79 (100.0)	0 (0.0)	0 (0.0)	5 (6.3)	20 (25.3)	38 (48.1)	16 (20.3)
45-49	96 (100.0)	0 (0.0)	1 (1.0)	5 (5.2)	32 (33.3)	37 (38.5)	21 (21.9)
50-54	159 (100.0)	1 (0.6)	2 (1.3)	17 (10.7)	54 (34.0)	56 (35.2)	29 (18.2)
55-59	208 (100.0)	1 (0.5)	3 (1.4)	28 (13.5)	64 (30.8)	74 (35.6)	38 (18.3)
60-64	340 (100.0)	3 (0.9)	6 (1.8)	54 (15.9)	119 (35.0)	117 (34.4)	41 (12.1)
65-69	462 (100.0)	8 (1.7)	20 (4.3)	92 (19.9)	150 (32.5)	141 (30.5)	51 (11.0)
70-74	422 (100.0)	14 (3.3)	31 (7.3)	99 (23.5)	144 (34.1)	99 (23.5)	35 (8.3)
75-79	219 (100.0)	10 (4.6)	21 (9.6)	52 (23.7)	70 (32.0)	48 (21.9)	18 (8.2)
80歳以上	92 (100.0)	16 (17.4)	16 (17.4)	21 (22.8)	22 (23.9)	13 (14.1)	4 (4.3)
計	2,122 (100.0)	53 (2.5)	100 (4.7)	374 (17.6)	686 (32.3)	639 (30.1)	270 (12.7)

年齢階級	人 (%)						
	総数	-11.9	12.0-12.9	13.0-13.9	14.0-14.9	15.0-15.9	16.0g/dl以上
女							
-29歳	39 (100.0)	7 (17.9)	11 (28.2)	16 (41.0)	5 (12.8)	0 (0.0)	0 (0.0)
30-39	139 (100.0)	31 (22.3)	49 (35.3)	39 (28.1)	16 (11.5)	4 (2.9)	0 (0.0)
40-44	211 (100.0)	50 (23.7)	65 (30.8)	65 (30.8)	30 (14.2)	1 (0.5)	0 (0.0)
45-49	335 (100.0)	70 (20.9)	100 (29.9)	105 (31.3)	52 (15.5)	7 (2.1)	1 (0.3)
50-54	525 (100.0)	49 (9.3)	141 (26.9)	201 (38.3)	118 (22.5)	14 (2.7)	2 (0.4)
55-59	589 (100.0)	44 (7.5)	172 (29.2)	252 (42.8)	98 (16.6)	22 (3.7)	1 (0.2)
60-64	722 (100.0)	73 (10.1)	200 (27.7)	290 (40.2)	130 (18.0)	28 (3.9)	1 (0.1)
65-69	819 (100.0)	98 (12.0)	260 (31.7)	321 (39.2)	124 (15.1)	16 (2.0)	0 (0.0)
70-74	637 (100.0)	85 (13.3)	191 (30.0)	250 (39.2)	101 (15.9)	9 (1.4)	1 (0.2)
75-79	311 (100.0)	53 (17.0)	101 (32.5)	103 (33.1)	45 (14.5)	9 (2.9)	0 (0.0)
80歳以上	68 (100.0)	18 (26.5)	18 (26.5)	24 (35.3)	6 (8.8)	2 (2.9)	0 (0.0)
計	4,395 (100.0)	578 (13.2)	1,308 (29.8)	1,666 (37.9)	725 (16.5)	112 (2.5)	6 (0.1)

表42 ヘモグロビン値の平均値、標準偏差および分布（性・年齢階級別）

年齢階級	g/dl													
	男					女								
	人数	平均値(SD)	パーセンタイル				人数	平均値(SD)	パーセンタイル					
5			25	50	75	95			5	25	50	75	95	
29歳以下	7	15.5 (0.7)	14.4	14.5	15.7	16.2	16.2	39	13.0 (0.9)	11.3	12.2	13.1	13.6	14.7
30-39	38	15.7 (0.9)	14.3	14.9	15.9	16.4	17.3	139	12.7 (1.4)	10.1	12.0	12.8	13.6	14.9
40-44	79	15.3 (1.0)	13.7	14.8	15.2	15.9	17.5	211	12.7 (1.3)	10.1	12.0	12.8	13.6	14.5
45-49	96	15.3 (1.0)	13.7	14.7	15.2	15.9	16.9	335	12.7 (1.6)	9.6	12.1	12.9	13.6	14.6
50-54	159	15.1 (1.0)	13.3	14.4	15.1	15.8	16.6	525	13.2 (1.1)	11.3	12.7	13.3	14.0	14.7
55-59	208	15.1 (1.1)	13.4	14.3	15.0	15.7	16.7	589	13.2 (0.9)	11.8	12.7	13.3	13.8	14.8
60-64	340	14.9 (1.0)	13.2	14.2	14.9	15.6	16.4	722	13.2 (1.0)	11.5	12.6	13.3	13.9	14.9
65-69	462	14.6 (1.2)	12.8	13.9	14.7	15.4	16.4	819	13.1 (1.0)	11.4	12.5	13.1	13.7	14.6
70-74	422	14.4 (1.3)	12.2	13.6	14.5	15.1	16.3	637	13.0 (1.0)	11.3	12.4	13.1	13.7	14.5
75-79	219	14.2 (1.4)	12.1	13.5	14.3	15.2	16.5	311	13.0 (1.1)	11.3	12.2	13.0	13.7	14.7
80歳以上	92	13.6 (1.5)	10.9	12.7	13.6	14.7	15.9	68	12.8 (1.0)	11.1	11.9	12.9	13.5	14.8
計	2,122	14.7 (1.3)	12.7	14.0	14.8	15.5	16.5	4,395	13.1 (1.1)	11.3	12.5	13.1	13.8	14.7

表43 ヘマトクリット値の状況 (性・年齢階級別)

年齢階級	総数	人 (%)						
		-29.9	30.0-34.9	35.0-39.9	40.0-44.9	45.0-49.9	50.0%以上	
男								
-29歳	7 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (28.6)	5 (71.4)	0 (0.0)	
30-39	38 (100.0)	0 (0.0)	0 (0.0)	0 (0.0)	13 (34.2)	22 (57.9)	3 (7.9)	
40-44	79 (100.0)	0 (0.0)	0 (0.0)	3 (3.8)	36 (45.6)	35 (44.3)	5 (6.3)	
45-49	96 (100.0)	0 (0.0)	0 (0.0)	4 (4.2)	46 (47.9)	43 (44.8)	3 (3.1)	
50-54	159 (100.0)	0 (0.0)	0 (0.0)	11 (6.9)	79 (49.7)	64 (40.3)	5 (3.1)	
55-59	208 (100.0)	0 (0.0)	0 (0.0)	11 (5.3)	109 (52.4)	82 (39.4)	6 (2.9)	
60-64	340 (100.0)	0 (0.0)	0 (0.0)	28 (8.2)	190 (55.9)	112 (32.9)	10 (2.9)	
65-69	462 (100.0)	0 (0.0)	5 (1.1)	60 (13.0)	252 (54.5)	135 (29.2)	10 (2.2)	
70-74	422 (100.0)	3 (0.7)	7 (1.7)	81 (19.2)	220 (52.1)	106 (25.1)	5 (1.2)	
75-79	219 (100.0)	1 (0.5)	7 (3.2)	42 (19.2)	115 (52.5)	48 (21.9)	6 (2.7)	
80歳以上	92 (100.0)	0 (0.0)	9 (9.8)	32 (34.8)	37 (40.2)	12 (13.0)	2 (2.2)	
計	2,122 (100.0)	4 (0.2)	28 (1.3)	272 (12.8)	1,099 (51.8)	664 (31.3)	55 (2.6)	

年齢階級	総数	人 (%)						
		-29.9	30.0-34.9	35.0-39.9	40.0-44.9	45.0-49.9	50.0%以上	
女								
-29歳	39 (100.0)	0 (0.0)	3 (7.7)	22 (56.4)	14 (35.9)	0 (0.0)	0 (0.0)	
30-39	139 (100.0)	3 (2.2)	14 (10.1)	78 (56.1)	39 (28.1)	5 (3.6)	0 (0.0)	
40-44	211 (100.0)	1 (0.5)	26 (12.3)	116 (55.0)	67 (31.8)	1 (0.5)	0 (0.0)	
45-49	335 (100.0)	11 (3.3)	35 (10.4)	173 (51.6)	110 (32.8)	5 (1.5)	1 (0.3)	
50-54	525 (100.0)	5 (1.0)	21 (4.0)	247 (47.0)	243 (46.3)	9 (1.7)	0 (0.0)	
55-59	589 (100.0)	0 (0.0)	20 (3.4)	285 (48.4)	267 (45.3)	17 (2.9)	0 (0.0)	
60-64	722 (100.0)	2 (0.3)	34 (4.7)	352 (48.8)	306 (42.4)	27 (3.7)	1 (0.1)	
65-69	819 (100.0)	1 (0.1)	51 (6.2)	425 (51.9)	321 (39.2)	21 (2.6)	0 (0.0)	
70-74	637 (100.0)	1 (0.2)	41 (6.4)	337 (52.9)	242 (38.0)	16 (2.5)	0 (0.0)	
75-79	311 (100.0)	0 (0.0)	26 (8.4)	162 (52.1)	114 (36.7)	9 (2.9)	0 (0.0)	
80歳以上	68 (100.0)	0 (0.0)	10 (14.7)	38 (55.9)	17 (25.0)	3 (4.4)	0 (0.0)	
計	4,395 (100.0)	24 (0.5)	281 (6.4)	2,235 (50.9)	1,740 (39.6)	113 (2.6)	2 (0.0)	

表44 ハマトクレット値の平均値、標準偏差および分布 (性・年齢階級別)

年齢階級	男						女							
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル				
			5	25	50	75	95			5	25	50	75	95
29歳以下	7	46.3 (2.8)	42.3	42.5	47.7	48.4	48.7	39	38.9 (2.6)	34.6	36.9	39.4	40.5	43.1
30-39	38	46.5 (2.5)	42.6	44.1	46.8	48.3	50.3	139	38.4 (3.4)	32.6	36.7	38.6	40.5	44.2
40-44	79	45.2 (2.8)	41.0	43.3	45.0	46.7	51.0	211	38.3 (3.1)	32.7	36.5	38.4	40.5	42.9
45-49	96	44.9 (2.7)	40.0	43.3	44.9	46.5	49.8	335	38.5 (3.9)	31.4	36.5	38.9	40.8	43.5
50-54	159	44.4 (2.9)	39.2	42.5	44.5	46.4	49.3	525	39.8 (3.0)	34.9	38.2	39.8	41.9	44.1
55-59	208	44.5 (3.0)	39.7	42.8	44.5	46.5	49.6	589	39.8 (2.6)	35.4	38.2	39.8	41.5	44.1
60-64	340	44.0 (3.0)	39.1	42.1	44.0	45.9	49.2	722	39.7 (3.1)	34.9	37.9	39.7	41.7	44.7
65-69	462	43.3 (3.2)	37.8	41.2	43.7	45.4	48.4	819	39.3 (2.9)	34.5	37.5	39.3	41.2	44.0
70-74	422	42.7 (3.7)	36.8	40.5	42.8	45.1	48.7	637	39.3 (2.9)	34.5	37.4	39.3	41.3	43.9
75-79	219	42.4 (3.9)	36.0	40.1	42.5	44.9	48.5	311	39.1 (3.1)	34.1	37.1	38.8	41.3	44.7
80歳以上	92	40.5 (4.4)	33.0	37.6	40.4	43.4	47.8	68	38.6 (3.1)	34.0	36.1	38.6	40.4	45.2
計	2,122	43.5 (3.5)	37.6	41.4	43.7	45.7	49.1	4,395	39.3 (3.1)	34.3	37.4	39.4	41.4	44.0

表45 血清ALPの状況(性・年齢階級別)

年齢階級	総数							人 (%)
	-199	200-249	250-299	300-349	350-399	400U/L以上		
男								
-29歳	7 (100.0)	2 (28.6)	2 (28.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
30-39	17 (100.0)	6 (35.3)	5 (29.4)	1 (5.9)	0 (0.0)	0 (0.0)	0 (0.0)	
40-44	54 (100.0)	14 (25.9)	16 (29.6)	6 (11.1)	1 (1.9)	1 (1.9)	1 (1.9)	
45-49	76 (100.0)	33 (43.4)	13 (17.1)	5 (6.6)	0 (0.0)	2 (2.6)	2 (2.6)	
50-54	129 (100.0)	55 (42.6)	26 (20.2)	10 (7.8)	9 (7.0)	4 (3.1)	4 (3.1)	
55-59	156 (100.0)	47 (30.1)	34 (21.8)	19 (12.2)	4 (2.6)	6 (3.8)	6 (3.8)	
60-64	268 (100.0)	81 (30.2)	61 (22.8)	30 (11.2)	10 (3.7)	3 (1.1)	3 (1.1)	
65-69	357 (100.0)	108 (30.3)	84 (23.5)	36 (10.1)	15 (4.2)	7 (2.0)	7 (2.0)	
70-74	333 (100.0)	111 (33.3)	61 (18.3)	47 (14.1)	16 (4.8)	14 (4.2)	14 (4.2)	
75-79	170 (100.0)	61 (35.9)	36 (21.2)	17 (10.0)	15 (8.8)	5 (2.9)	5 (2.9)	
80歳以上	68 (100.0)	16 (23.5)	17 (25.0)	12 (17.6)	4 (5.9)	5 (7.4)	5 (7.4)	
計	1,635 (100.0)	534 (32.7)	355 (21.7)	183 (11.2)	74 (4.5)	47 (2.9)	47 (2.9)	

年齢階級	総数							人 (%)
	-199	200-249	250-299	300-349	350-399	400U/L以上		
女								
-29歳	12 (100.0)	3 (25.0)	2 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
30-39	31 (100.0)	6 (19.4)	1 (3.2)	0 (0.0)	1 (3.2)	0 (0.0)	0 (0.0)	
40-44	162 (100.0)	32 (19.8)	11 (6.8)	2 (1.2)	3 (1.9)	1 (0.6)	1 (0.6)	
45-49	243 (100.0)	62 (25.5)	33 (13.6)	13 (5.3)	3 (1.2)	1 (0.4)	1 (0.4)	
50-54	381 (100.0)	101 (26.5)	89 (23.4)	47 (12.3)	29 (7.6)	13 (3.4)	13 (3.4)	
55-59	440 (100.0)	116 (26.4)	132 (30.0)	65 (14.8)	38 (8.6)	23 (5.2)	23 (5.2)	
60-64	546 (100.0)	157 (28.8)	147 (26.9)	103 (18.9)	27 (4.9)	26 (4.8)	26 (4.8)	
65-69	621 (100.0)	152 (24.5)	199 (32.0)	97 (15.6)	49 (7.9)	32 (5.2)	32 (5.2)	
70-74	510 (100.0)	140 (27.5)	142 (27.8)	66 (12.9)	43 (8.4)	34 (6.7)	34 (6.7)	
75-79	250 (100.0)	50 (20.0)	68 (27.2)	47 (18.8)	23 (9.2)	10 (4.0)	10 (4.0)	
80歳以上	58 (100.0)	13 (22.4)	21 (36.2)	10 (17.2)	3 (5.2)	3 (5.2)	3 (5.2)	
計	3,254 (100.0)	832 (25.6)	845 (26.0)	450 (13.8)	219 (6.7)	143 (4.4)	143 (4.4)	

表46 血清ALPの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						U/L		
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	7	218.9 (43.9)	177.0	177.0	213.0	267.0	273.0	12	189.2 (44.9)	121.0	153.3	177.0	222.8	256.0	
30-39	17	229.6 (44.2)	144.0	193.5	223.0	262.0	291.0	31	183.5 (55.6)	118.8	148.0	174.0	206.0	336.2	
40-44	54	240.8 (59.3)	152.0	193.0	230.0	286.5	348.3	162	185.5 (53.3)	126.0	148.8	171.5	209.3	288.1	
45-49	76	230.8 (72.3)	153.1	189.0	223.5	250.8	345.2	243	208.6 (74.4)	124.2	166.0	195.0	241.0	327.6	
50-54	129	248.0 (65.0)	160.5	205.5	239.0	278.5	375.0	381	252.0 (76.0)	146.0	195.5	243.0	297.0	389.6	
55-59	156	247.6 (92.3)	147.1	188.0	236.5	280.0	363.9	440	274.5 (85.5)	174.1	219.0	260.5	307.0	422.9	
60-64	268	237.3 (68.2)	151.9	188.0	227.0	274.8	354.5	546	265.5 (69.9)	164.0	217.8	259.0	307.0	396.6	
65-69	357	242.3 (68.7)	152.0	194.0	235.0	279.5	353.2	621	272.4 (80.8)	171.2	219.5	265.0	310.0	400.0	
70-74	333	257.5 (188.2)	145.7	199.0	234.0	292.5	390.3	510	270.6 (85.4)	159.0	217.8	258.0	310.3	416.0	
75-79	170	255.6 (86.3)	161.1	204.0	240.5	290.5	382.0	250	272.0 (89.6)	155.7	205.8	264.0	321.0	394.0	
80歳以上	68	268.4 (78.7)	152.4	214.8	257.0	316.0	421.8	58	271.8 (76.4)	157.5	222.3	275.0	302.5	412.7	
計	1,635	247.2 (107.8)	153.0	197.0	234.0	281.0	365.0	3,254	258.6 (82.6)	149.0	203.0	251.0	299.0	394.0	

表47 血清コリンエステラーゼの状況 (性・年齢階級別)

年齢階級	総数							人 (%)	
	-119	120-159	160-199	200-239	240-279	280U/L以上			
-29歳	7 (100.0)	2 (28.6)	5 (71.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
30-39	17 (100.0)	1 (5.9)	10 (58.8)	5 (29.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
40-44	54 (100.0)	0 (0.0)	23 (42.6)	14 (25.9)	1 (1.9)	2 (3.7)	2 (3.7)	2 (3.7)	
45-49	76 (100.0)	3 (3.9)	30 (39.5)	22 (28.9)	7 (9.2)	0 (0.0)	0 (0.0)	0 (0.0)	
50-54	129 (100.0)	3 (2.3)	68 (52.7)	35 (27.1)	6 (4.7)	1 (0.8)	1 (0.8)	1 (0.8)	
55-59	156 (100.0)	3 (1.9)	79 (50.6)	31 (19.9)	5 (3.2)	1 (0.6)	1 (0.6)	1 (0.6)	
60-64	268 (100.0)	9 (3.4)	126 (47.0)	52 (19.4)	9 (3.4)	1 (0.4)	1 (0.4)	1 (0.4)	
65-69	357 (100.0)	16 (4.5)	182 (51.0)	40 (11.2)	9 (2.5)	1 (0.3)	1 (0.3)	1 (0.3)	
70-74	333 (100.0)	23 (6.9)	145 (43.5)	34 (10.2)	0 (0.0)	2 (0.6)	2 (0.6)	2 (0.6)	
75-79	170 (100.0)	20 (11.8)	57 (33.5)	18 (10.6)	5 (2.9)	0 (0.0)	0 (0.0)	0 (0.0)	
80歳以上	68 (100.0)	8 (11.8)	17 (25.0)	5 (7.4)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
計	1,635 (100.0)	86 (5.3)	742 (45.4)	256 (15.7)	42 (2.6)	8 (0.5)	8 (0.5)	8 (0.5)	

年齢階級	総数							人 (%)	
	-119	120-159	160-199	200-239	240-279	280U/L以上			
-29歳	12 (100.0)	8 (66.7)	3 (25.0)	1 (8.3)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
30-39	31 (100.0)	13 (41.9)	12 (38.7)	3 (9.7)	1 (3.2)	0 (0.0)	0 (0.0)	0 (0.0)	
40-44	162 (100.0)	9 (5.6)	62 (38.3)	15 (9.3)	3 (1.9)	1 (0.6)	1 (0.6)	1 (0.6)	
45-49	243 (100.0)	12 (4.9)	80 (32.9)	39 (16.0)	7 (2.9)	3 (1.2)	3 (1.2)	3 (1.2)	
50-54	381 (100.0)	6 (1.6)	167 (43.8)	81 (21.3)	28 (7.3)	8 (2.1)	8 (2.1)	8 (2.1)	
55-59	440 (100.0)	7 (1.6)	208 (47.3)	106 (24.1)	36 (8.2)	9 (2.0)	9 (2.0)	9 (2.0)	
60-64	546 (100.0)	10 (1.8)	259 (47.4)	140 (25.6)	27 (4.9)	11 (2.0)	11 (2.0)	11 (2.0)	
65-69	621 (100.0)	15 (2.4)	284 (45.7)	143 (23.0)	42 (6.8)	12 (1.9)	12 (1.9)	12 (1.9)	
70-74	510 (100.0)	8 (1.6)	255 (50.0)	93 (18.2)	22 (4.3)	5 (1.0)	5 (1.0)	5 (1.0)	
75-79	250 (100.0)	8 (3.2)	125 (50.0)	33 (13.2)	11 (4.4)	4 (1.6)	4 (1.6)	4 (1.6)	
80歳以上	58 (100.0)	0 (0.0)	29 (50.0)	3 (5.2)	2 (3.4)	2 (3.4)	2 (3.4)	2 (3.4)	
計	3,254 (100.0)	77 (2.4)	1,484 (45.6)	657 (20.2)	179 (5.5)	55 (1.7)	55 (1.7)	55 (1.7)	

表48 血清コリンエステラーゼの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	男						女						U/L		
	人数	平均値 (SD)	パーセンタイル					人数	平均値 (SD)	パーセンタイル					
			5	25	50	75	95			5	25	50		75	95
29歳以下	7	162.9 (14.3)	147.0	147.0	164.0	167.0	178.0	12	160.2 (28.6)	134.0	142.3	148.0	173.5	212.0	
30-39	17	187.9 (28.4)	117.0	175.0	192.0	214.0	218.0	31	161.4 (30.5)	107.6	144.0	161.0	177.0	228.2	
40-44	54	187.2 (37.6)	141.0	158.0	183.0	205.5	269.8	162	163.3 (33.0)	116.0	139.8	159.5	182.0	213.0	
45-49	76	184.5 (40.4)	117.9	163.0	179.5	211.0	250.4	243	168.2 (37.9)	119.4	142.0	162.0	188.0	236.6	
50-54	129	190.4 (33.6)	139.5	169.5	187.0	212.5	241.0	381	185.6 (38.6)	135.0	158.5	181.0	208.0	257.8	
55-59	156	179.7 (34.7)	128.7	157.5	177.5	197.0	236.3	440	190.6 (39.2)	138.0	165.3	186.0	212.0	257.0	
60-64	268	177.5 (32.5)	126.9	154.3	176.5	198.0	235.1	546	187.7 (35.1)	134.0	164.0	183.0	208.0	245.7	
65-69	357	171.7 (36.0)	120.9	151.5	170.0	187.0	220.1	621	186.0 (35.9)	133.1	162.0	183.0	206.0	250.9	
70-74	333	162.4 (32.4)	115.4	138.5	163.0	179.0	217.6	510	180.3 (33.1)	132.0	158.0	176.0	198.3	241.0	
75-79	170	160.6 (35.1)	111.6	139.0	155.0	178.0	224.9	250	175.9 (37.6)	126.0	152.0	173.0	191.3	243.9	
80歳以上	68	149.2 (27.1)	106.9	127.5	146.0	168.0	206.6	58	173.7 (33.4)	137.0	153.8	166.5	184.0	267.2	
計	1,635	172.2 (35.6)	119.0	149.0	171.0	192.0	228.0	3,254	182.2 (37.0)	130.0	157.0	178.0	203.0	247.0	

表49 総ビリルビンの状況 (性・年齢階級別)

年齢階級	総数	人 (%)				
		-0.5	0.6-0.8	0.9-1.1	1.2-1.4	1.5mg/dl以上
-29歳	7 (100.0)	3 (42.9)	1 (14.3)	1 (14.3)	1 (14.3)	1 (14.3)
30-39	17 (100.0)	7 (41.2)	4 (23.5)	5 (29.4)	1 (5.9)	0 (0.0)
40-44	54 (100.0)	18 (33.3)	20 (37.0)	10 (18.5)	4 (7.4)	2 (3.7)
45-49	76 (100.0)	24 (31.6)	34 (44.7)	11 (14.5)	6 (7.9)	1 (1.3)
50-54	129 (100.0)	29 (22.5)	63 (48.8)	26 (20.2)	6 (4.7)	5 (3.9)
55-59	156 (100.0)	29 (18.6)	86 (55.1)	29 (18.6)	10 (6.4)	2 (1.3)
60-64	268 (100.0)	53 (19.8)	120 (44.8)	74 (27.6)	19 (7.1)	2 (0.7)
65-69	357 (100.0)	56 (15.7)	154 (43.1)	104 (29.1)	29 (8.1)	14 (3.9)
70-74	333 (100.0)	65 (19.5)	177 (53.2)	69 (20.7)	13 (3.9)	9 (2.7)
75-79	170 (100.0)	35 (20.6)	75 (44.1)	46 (27.1)	9 (5.3)	5 (2.9)
80歳以上	68 (100.0)	18 (26.5)	33 (48.5)	16 (23.5)	0 (0.0)	1 (1.5)
計	1,635 (100.0)	337 (20.6)	767 (46.9)	391 (23.9)	98 (6.0)	42 (2.6)

年齢階級	総数	人 (%)				
		-0.5	0.6-0.8	0.9-1.1	1.2-1.4	1.5mg/dl以上
-29歳	12 (100.0)	1 (8.3)	6 (50.0)	3 (25.0)	2 (16.7)	0 (0.0)
30-39	31 (100.0)	10 (32.3)	15 (48.4)	4 (12.9)	1 (3.2)	1 (3.2)
40-44	162 (100.0)	67 (41.4)	61 (37.7)	24 (14.8)	7 (4.3)	3 (1.9)
45-49	243 (100.0)	73 (30.0)	125 (51.4)	34 (14.0)	8 (3.3)	3 (1.2)
50-54	381 (100.0)	95 (24.9)	200 (52.5)	66 (17.3)	13 (3.4)	7 (1.8)
55-59	440 (100.0)	97 (22.0)	239 (54.3)	79 (18.0)	17 (3.9)	8 (1.8)
60-64	546 (100.0)	120 (22.0)	274 (50.2)	126 (23.1)	18 (3.3)	8 (1.5)
65-69	621 (100.0)	146 (23.5)	314 (50.6)	122 (19.6)	27 (4.3)	12 (1.9)
70-74	510 (100.0)	127 (24.9)	259 (50.8)	88 (17.3)	29 (5.7)	7 (1.4)
75-79	250 (100.0)	74 (29.6)	118 (47.2)	43 (17.2)	10 (4.0)	5 (2.0)
80歳以上	58 (100.0)	14 (24.1)	33 (56.9)	11 (19.0)	0 (0.0)	0 (0.0)
計	3,254 (100.0)	824 (25.3)	1,644 (50.5)	600 (18.4)	132 (4.1)	54 (1.7)

表50 総ビリルビンの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	mg/dl.													
	男						女							
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル				
5			25	50	75	95	5			25	50	75	95	
29歳以下	7	0.8 (0.4)	0.4	0.5	0.7	1.2	1.4	12	0.8 (0.3)	0.4	0.6	0.8	1.1	1.4
30-39	17	0.7 (0.3)	0.4	0.5	0.6	0.9	1.1	31	0.7 (0.3)	0.3	0.5	0.6	0.8	1.4
40-44	54	0.8 (0.4)	0.4	0.5	0.7	0.9	1.5	162	0.7 (0.3)	0.4	0.5	0.6	0.8	1.3
45-49	76	0.7 (0.3)	0.4	0.5	0.7	0.8	1.2	243	0.7 (0.2)	0.4	0.5	0.6	0.8	1.1
50-54	129	0.8 (0.3)	0.4	0.6	0.7	0.9	1.4	381	0.7 (0.3)	0.4	0.6	0.7	0.8	1.2
55-59	156	0.8 (0.3)	0.5	0.6	0.7	0.9	1.3	440	0.7 (0.3)	0.4	0.6	0.7	0.8	1.2
60-64	268	0.8 (0.3)	0.4	0.6	0.7	0.9	1.3	546	0.7 (0.2)	0.4	0.6	0.7	0.9	1.1
65-69	357	0.8 (0.3)	0.5	0.6	0.8	0.9	1.4	621	0.7 (0.3)	0.4	0.6	0.7	0.9	1.2
70-74	333	0.8 (0.3)	0.4	0.6	0.7	0.9	1.3	510	0.7 (0.3)	0.4	0.6	0.7	0.8	1.2
75-79	170	0.8 (0.3)	0.4	0.6	0.7	0.9	1.3	250	0.7 (0.3)	0.4	0.5	0.7	0.8	1.2
80歳以上	68	0.7 (0.2)	0.4	0.5	0.7	0.9	1.0	58	0.7 (0.2)	0.3	0.6	0.6	0.8	1.1
計	1,635	0.8 (0.3)	0.4	0.6	0.7	0.9	1.3	3,254	0.7 (0.3)	0.4	0.5	0.7	0.8	1.2

表51 血清総蛋白の状況 (性・年齢階級別)

男	年齢階級	総数	-6.3	6.4-6.7	6.8-7.1	7.2-7.5	7.6-7.9	8.0g/dl以上	人 (%)
	-29歳	7 (100.0)	0 (0.0)	0 (0.0)	2 (28.6)	2 (28.6)	3 (42.9)	0 (0.0)	0 (0.0)
	30-39	17 (100.0)	0 (0.0)	1 (5.9)	4 (23.5)	7 (41.2)	5 (29.4)	0 (0.0)	0 (0.0)
	40-44	54 (100.0)	0 (0.0)	2 (3.7)	20 (37.0)	25 (46.3)	6 (11.1)	1 (1.9)	1 (1.9)
	45-49	76 (100.0)	0 (0.0)	8 (10.5)	19 (25.0)	28 (36.8)	20 (26.3)	1 (1.3)	1 (1.3)
	50-54	129 (100.0)	1 (0.8)	9 (7.0)	40 (31.0)	49 (38.0)	25 (19.4)	5 (3.9)	5 (3.9)
	55-59	156 (100.0)	0 (0.0)	15 (9.6)	51 (32.7)	54 (34.6)	29 (18.6)	7 (4.5)	7 (4.5)
	60-64	268 (100.0)	6 (2.2)	20 (7.5)	70 (26.1)	111 (41.4)	48 (17.9)	13 (4.9)	13 (4.9)
	65-69	357 (100.0)	5 (1.4)	36 (10.1)	119 (33.3)	128 (35.9)	53 (14.8)	16 (4.5)	16 (4.5)
	70-74	333 (100.0)	7 (2.1)	33 (9.9)	124 (37.2)	106 (31.8)	54 (16.2)	9 (2.7)	9 (2.7)
	75-79	170 (100.0)	3 (1.8)	26 (15.3)	60 (35.3)	57 (33.5)	18 (10.6)	6 (3.5)	6 (3.5)
	80歳以上	68 (100.0)	2 (2.9)	11 (16.2)	23 (33.8)	21 (30.9)	9 (13.2)	2 (2.9)	2 (2.9)
	計	1,635 (100.0)	24 (1.5)	161 (9.8)	532 (32.5)	588 (36.0)	270 (16.5)	60 (3.7)	60 (3.7)

女	年齢階級	総数	-6.3	6.4-6.7	6.8-7.1	7.2-7.5	7.6-7.9	8.0g/dl以上	人 (%)
	-29歳	12 (100.0)	0 (0.0)	0 (0.0)	1 (8.3)	5 (41.7)	3 (25.0)	3 (25.0)	3 (25.0)
	30-39	31 (100.0)	0 (0.0)	1 (3.2)	7 (22.6)	14 (45.2)	8 (25.8)	1 (3.2)	1 (3.2)
	40-44	162 (100.0)	1 (0.6)	10 (6.2)	44 (27.2)	64 (39.5)	30 (18.5)	13 (8.0)	13 (8.0)
	45-49	243 (100.0)	2 (0.8)	16 (6.6)	70 (28.8)	102 (42.0)	42 (17.3)	11 (4.5)	11 (4.5)
	50-54	381 (100.0)	1 (0.3)	20 (5.2)	88 (23.1)	149 (39.1)	95 (24.9)	28 (7.3)	28 (7.3)
	55-59	440 (100.0)	1 (0.2)	31 (7.0)	110 (25.0)	173 (39.3)	99 (22.5)	26 (5.9)	26 (5.9)
	60-64	546 (100.0)	4 (0.7)	31 (5.7)	168 (30.8)	220 (40.3)	106 (19.4)	17 (3.1)	17 (3.1)
	65-69	621 (100.0)	10 (1.6)	54 (8.7)	197 (31.7)	235 (37.8)	101 (16.3)	24 (3.9)	24 (3.9)
	70-74	510 (100.0)	8 (1.6)	49 (9.6)	184 (36.1)	174 (34.1)	80 (15.7)	15 (2.9)	15 (2.9)
	75-79	250 (100.0)	3 (1.2)	33 (13.2)	92 (36.8)	81 (32.4)	31 (12.4)	10 (4.0)	10 (4.0)
	80歳以上	58 (100.0)	1 (1.7)	5 (8.6)	29 (50.0)	17 (29.3)	6 (10.3)	0 (0.0)	0 (0.0)
	計	3,254 (100.0)	31 (1.0)	250 (7.7)	990 (30.4)	1,234 (37.9)	601 (18.5)	148 (4.5)	148 (4.5)

表52 血清総蛋白の平均値、標準偏差および分布（性・年齢階級別）

年齢階級	g/dl													
	男						女							
	人数	平均値(SD)	パーセンタイル					人数	平均値(SD)	パーセンタイル				
5			25	50	75	95	5			25	50	75	95	
29歳以下	7	7.4 (0.3)	6.9	7.1	7.5	7.6	7.7	12	7.7 (0.5)	7.0	7.3	7.6	7.9	8.5
30-39	17	7.3 (0.3)	6.7	7.1	7.3	7.7	7.7	31	7.4 (0.3)	6.8	7.1	7.4	7.6	7.9
40-44	54	7.2 (0.3)	6.8	7.0	7.2	7.5	7.8	162	7.3 (0.4)	6.7	7.0	7.3	7.6	8.0
45-49	76	7.3 (0.4)	6.7	7.0	7.3	7.6	7.8	243	7.3 (0.4)	6.6	7.0	7.3	7.5	7.9
50-54	129	7.3 (0.4)	6.6	7.0	7.2	7.5	7.9	381	7.4 (0.4)	6.7	7.1	7.4	7.6	8.0
55-59	156	7.3 (0.4)	6.7	7.0	7.2	7.5	7.9	440	7.3 (0.4)	6.7	7.1	7.3	7.6	8.0
60-64	268	7.3 (0.4)	6.5	7.0	7.3	7.5	8.0	546	7.3 (0.4)	6.7	7.0	7.3	7.5	7.9
65-69	357	7.2 (0.4)	6.5	7.0	7.2	7.5	7.9	621	7.2 (0.4)	6.6	7.0	7.2	7.5	7.9
70-74	333	7.2 (0.4)	6.5	6.9	7.2	7.5	7.8	510	7.2 (0.4)	6.6	7.0	7.2	7.4	7.8
75-79	170	7.1 (0.4)	6.5	6.9	7.1	7.4	7.8	250	7.2 (0.4)	6.4	6.9	7.1	7.4	7.9
80歳以上	68	7.1 (0.4)	6.4	6.8	7.1	7.3	7.9	58	7.1 (0.3)	6.6	6.9	7.1	7.3	7.7
計	1,635	7.2 (0.4)	6.5	6.9	7.2	7.5	7.9	3,254	7.3 (0.4)	6.6	7.0	7.3	7.5	7.9

表53 血清アルブミンの状況 (性・年齢階級別)

年齢階級	総数	人 (%)						
		-3.5	3.6-3.9	4.0-4.3	4.4-4.7	4.8-5.1	5.2g/dl以上	
男								
-29歳	7 (100.0)	0 (0.0)	0 (0.0)	1 (14.3)	6 (85.7)	0 (0.0)	0 (0.0)	
30-39	17 (100.0)	0 (0.0)	0 (0.0)	3 (17.6)	9 (52.9)	5 (29.4)	0 (0.0)	
40-44	54 (100.0)	0 (0.0)	1 (1.9)	11 (20.4)	32 (59.3)	10 (18.5)	0 (0.0)	
45-49	76 (100.0)	1 (1.3)	1 (1.3)	17 (22.4)	45 (59.2)	11 (14.5)	1 (1.3)	
50-54	129 (100.0)	0 (0.0)	0 (0.0)	41 (31.8)	73 (56.6)	14 (10.9)	1 (0.8)	
55-59	156 (100.0)	1 (0.6)	1 (0.6)	52 (33.3)	89 (57.1)	12 (7.7)	1 (0.6)	
60-64	268 (100.0)	0 (0.0)	8 (3.0)	113 (42.2)	122 (45.5)	23 (8.6)	2 (0.7)	
65-69	357 (100.0)	1 (0.3)	18 (5.0)	162 (45.4)	160 (44.8)	16 (4.5)	0 (0.0)	
70-74	333 (100.0)	1 (0.3)	25 (7.5)	173 (52.0)	120 (36.0)	14 (4.2)	0 (0.0)	
75-79	170 (100.0)	0 (0.0)	23 (13.5)	91 (53.5)	55 (32.4)	1 (0.6)	0 (0.0)	
80歳以上	68 (100.0)	0 (0.0)	16 (23.5)	42 (61.8)	9 (13.2)	1 (1.5)	0 (0.0)	
計	1,635 (100.0)	4 (0.2)	93 (5.7)	706 (43.2)	720 (44.0)	107 (6.5)	5 (0.3)	

年齢階級	総数	人 (%)					
		-3.5	3.6-3.9	4.0-4.3	4.4-4.7	4.8-5.1	5.2g/dl以上
女							
-29歳	12 (100.0)	0 (0.0)	0 (0.0)	2 (16.7)	8 (66.7)	2 (16.7)	0 (0.0)
30-39	31 (100.0)	0 (0.0)	0 (0.0)	9 (29.0)	20 (64.5)	2 (6.5)	0 (0.0)
40-44	162 (100.0)	0 (0.0)	3 (1.9)	59 (36.4)	94 (58.0)	6 (3.7)	0 (0.0)
45-49	243 (100.0)	0 (0.0)	17 (7.0)	103 (42.4)	111 (45.7)	12 (4.9)	0 (0.0)
50-54	381 (100.0)	1 (0.3)	6 (1.6)	147 (38.6)	201 (52.8)	26 (6.8)	0 (0.0)
55-59	440 (100.0)	0 (0.0)	13 (3.0)	166 (37.7)	231 (52.5)	29 (6.6)	1 (0.2)
60-64	546 (100.0)	1 (0.2)	15 (2.7)	220 (40.3)	275 (50.4)	35 (6.4)	0 (0.0)
65-69	621 (100.0)	4 (0.6)	24 (3.9)	273 (44.0)	302 (48.6)	16 (2.6)	2 (0.3)
70-74	510 (100.0)	2 (0.4)	26 (5.1)	237 (46.5)	230 (45.1)	15 (2.9)	0 (0.0)
75-79	250 (100.0)	3 (1.2)	23 (9.2)	128 (51.2)	88 (35.2)	8 (3.2)	0 (0.0)
80歳以上	58 (100.0)	0 (0.0)	6 (10.3)	29 (50.0)	21 (36.2)	2 (3.4)	0 (0.0)
計	3,254 (100.0)	11 (0.3)	133 (4.1)	1,373 (42.2)	1,581 (48.6)	153 (4.7)	3 (0.1)

表54 血清アルブミンの平均値、標準偏差および分布（性・年齢階級別）

年齢階級	g/dl													
	男					女								
	パーセンタイル					パーセンタイル								
	人数	平均値(SD)	5	25	50	75	95	人数	平均値(SD)	5	25	50	75	95
29歳以下	7	4.5 (0.2)	4.3	4.4	4.5	4.7	4.7	12	4.5 (0.2)	4.3	4.4	4.4	4.7	4.9
30-39	17	4.6 (0.2)	4.3	4.5	4.7	4.8	4.8	31	4.4 (0.2)	4.1	4.3	4.4	4.6	4.8
40-44	54	4.5 (0.2)	4.2	4.4	4.5	4.7	4.9	162	4.4 (0.2)	4.0	4.3	4.4	4.5	4.7
45-49	76	4.5 (0.3)	4.1	4.3	4.5	4.6	5.0	243	4.3 (0.2)	3.9	4.2	4.4	4.5	4.8
50-54	129	4.5 (0.2)	4.1	4.3	4.5	4.6	4.9	381	4.4 (0.2)	4.0	4.3	4.4	4.6	4.8
55-59	156	4.4 (0.3)	4.1	4.3	4.4	4.6	4.8	440	4.4 (0.2)	4.0	4.3	4.4	4.6	4.8
60-64	268	4.4 (0.3)	4.0	4.2	4.4	4.6	4.8	546	4.4 (0.2)	4.0	4.2	4.4	4.5	4.8
65-69	357	4.3 (0.2)	3.9	4.2	4.3	4.5	4.7	621	4.3 (0.2)	4.0	4.2	4.4	4.5	4.7
70-74	333	4.3 (0.3)	3.9	4.1	4.3	4.5	4.7	510	4.3 (0.2)	3.9	4.2	4.3	4.5	4.7
75-79	170	4.2 (0.2)	3.8	4.1	4.2	4.4	4.6	250	4.3 (0.3)	3.9	4.1	4.3	4.4	4.7
80歳以上	68	4.1 (0.3)	3.6	4.0	4.1	4.3	4.6	58	4.3 (0.3)	3.9	4.1	4.3	4.4	4.6
計	1,635	4.4 (0.3)	3.9	4.2	4.4	4.5	4.8	3,254	4.4 (0.2)	4.0	4.2	4.4	4.5	4.7