

## Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Loop diuretic	Female	0	0.0	2	1.5	1	0.7	2	1.8	2	10.5	7	1.3
	Total	0	0.0	2	0.9	2	0.7	4	1.7	3	6.7	11	1.0
Potassium sparing diuretic	Male	0	0.0	0	0.0	1	0.7	2	1.6	0	0.0	3	0.6
	Female	0	0.0	0	0.0	0	0.0	2	1.8	0	0.0	2	0.4
	Total	0	0.0	0	0.0	1	0.4	4	1.7	0	0.0	5	0.5
Carbonate dehydratase inhibitor	Male	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	1	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
Other diuretic	Male	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	1	0.2
	Female	0	0.0	0	0.0	0	0.0	2	1.8	0	0.0	2	0.4
	Total	0	0.0	0	0.0	0	0.0	3	1.3	0	0.0	3	0.3
ACE inhibitor	Male	0	0.0	5	4.9	18	12.1	23	18.3	5	19.2	51	9.6
	Female	1	0.7	5	3.8	11	8.1	24	21.4	5	26.3	46	8.6
	Total	1	0.4	10	4.3	29	10.2	47	19.7	10	22.2	97	9.1
Ca antagonist	Male	2	1.6	9	8.7	34	22.8	47	37.3	6	23.1	98	18.4
	Female	2	1.5	13	9.9	19	14.1	38	33.9	6	31.6	78	14.6
	Total	4	1.5	22	9.4	53	18.7	85	35.7	12	26.7	176	16.5
Other antihypertensive agent	Male	1	0.8	2	1.9	6	4.0	3	2.4	0	0.0	12	2.3
	Female	0	0.0	0	0.0	1	0.7	3	2.7	1	5.3	5	0.9
	Total	1	0.4	2	0.9	7	2.5	6	2.5	1	2.2	17	1.6
Vascular reinforcing agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Vasoconstrictor	Male	1	0.8	0	0.0	0	0.0	1	0.8	0	0.0	2	0.4
	Female	0	0.0	0	0.0	0	0.0	2	1.8	0	0.0	2	0.4

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Vasoconstrictor	Total	1	0.4	0	0.0	0	0.0	3	1.3	0	0.0	4	0.4
Vasodilator	Male	0	0.0	2	1.9	6	4.0	8	6.3	3	11.5	19	3.6
	Female	0	0.0	0	0.0	1	0.7	5	4.5	1	5.3	7	1.3
	Total	0	0.0	2	0.9	7	2.5	13	5.5	4	8.9	26	2.4
Antihyperlipidemic agent	Male	0	0.0	6	5.8	15	10.1	16	12.7	2	7.7	39	7.3
	Female	0	0.0	8	6.1	25	18.5	28	25.0	6	31.6	67	12.6
	Total	0	0.0	14	6.0	40	14.1	44	18.5	8	17.8	106	9.9
Other agent on cardiovascular system	Male	0	0.0	2	1.9	3	2.0	10	7.9	4	15.4	19	3.6
	Female	0	0.0	4	3.1	1	0.7	11	9.8	3	15.8	19	3.6
	Total	0	0.0	6	2.6	4	1.4	21	8.8	7	15.6	38	3.6
Respiratory stimulator	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antitussive agent	Male	0	0.0	1	1.0	1	0.7	1	0.8	0	0.0	3	0.6
	Female	1	0.7	1	0.8	3	2.2	4	3.6	0	0.0	9	1.7
	Total	1	0.4	2	0.9	4	1.4	5	2.1	0	0.0	12	1.1
Expectorant	Male	1	0.8	1	1.0	4	2.7	5	4.0	2	7.7	13	2.4
	Female	1	0.7	1	0.8	2	1.5	4	3.6	1	5.3	9	1.7
	Total	2	0.8	2	0.9	6	2.1	9	3.8	3	6.7	22	2.1
Antitussive agent and expectorant	Male	1	0.8	0	0.0	2	1.3	3	2.4	0	0.0	6	1.1
	Female	2	1.5	2	1.5	2	1.5	0	0.0	0	0.0	6	1.1
	Total	3	1.1	2	0.9	4	1.4	3	1.3	0	0.0	12	1.1
Bronchodilator	Male	2	1.6	2	1.9	1	0.7	6	4.8	2	7.7	13	2.4
	Female	0	0.0	2	1.5	2	1.5	1	0.9	0	0.0	5	0.9
	Total	2	0.8	4	1.7	3	1.1	7	2.9	2	4.4	18	1.7

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Gargle	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent acting on respiratory system	Male	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2
	Female	0	0.0	1	0.8	0	0.0	1	0.9	0	0.0	2	0.4
	Total	1	0.4	1	0.4	0	0.0	1	0.4	0	0.0	3	0.3
Antidiarrheal drug	Male	1	0.8	2	1.9	4	2.7	3	2.4	1	3.8	11	2.1
	Female	1	0.7	4	3.1	4	3.0	5	4.5	0	0.0	14	2.6
	Total	2	0.8	6	2.6	8	2.8	8	3.4	1	2.2	25	2.3
Antiulcerative agent	Male	4	3.1	6	5.8	21	14.1	30	23.8	5	19.2	66	12.4
	Female	6	4.4	11	8.4	16	11.9	33	29.5	7	36.8	73	13.7
	Total	10	3.8	17	7.3	37	13.0	63	26.5	12	26.7	139	13.0
Stomachic	Male	1	0.8	1	1.0	4	2.7	4	3.2	4	15.4	14	2.6
	Female	0	0.0	3	2.3	1	0.7	8	7.1	2	10.5	14	2.6
	Total	1	0.4	4	1.7	5	1.8	12	5.0	6	13.3	28	2.6
Antacid	Male	0	0.0	0	0.0	1	0.7	1	0.8	0	0.0	2	0.4
	Female	1	0.7	1	0.8	0	0.0	0	0.0	0	0.0	2	0.4
	Total	1	0.4	1	0.4	1	0.4	1	0.4	0	0.0	4	0.4
Laxative	Male	1	0.8	3	2.9	3	2.0	12	9.5	5	19.2	24	4.5
	Female	5	3.7	5	3.8	6	4.4	12	10.7	2	10.5	30	5.6
	Total	6	2.3	8	3.4	9	3.2	24	10.1	7	15.6	54	5.1
Cholagogue	Male	1	0.8	1	1.0	2	1.3	1	0.8	0	0.0	5	0.9
	Female	0	0.0	2	1.5	1	0.7	1	0.9	0	0.0	4	0.8
	Total	1	0.4	3	1.3	3	1.1	2	0.8	0	0.0	9	0.8
Combined digestive drug	Male	8	6.2	4	3.9	6	4.0	2	1.6	0	0.0	20	3.8

(Continued)

## Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Combined digestive drug	Female	7	5.1	6	4.6	1	0.7	4	3.6	0	0.0	18	3.4
	Total	15	5.7	10	4.3	7	2.5	6	2.5	0	0.0	38	3.6
Other agent acting on alimentary system	Male	1	0.8	1	1.0	3	2.0	6	4.8	2	7.7	13	2.4
	Female	1	0.7	3	2.3	0	0.0	9	8.0	3	15.8	16	3.0
	Total	2	0.8	4	1.7	3	1.1	15	6.3	5	11.1	29	2.7
Pituitary hormone drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Hormone drug of salivary gland	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Thyroid parathyroid hormone drug	Male	1	0.8	1	1.0	0	0.0	1	0.8	0	0.0	3	0.6
	Female	4	2.9	3	2.3	2	1.5	4	3.6	1	5.3	14	2.6
	Total	5	1.9	4	1.7	2	0.7	5	2.1	1	2.2	17	1.6
Anabolic steroid drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Adrenocortical hormone drug	Male	3	2.3	1	1.0	6	4.0	5	4.0	0	0.0	15	2.8
	Female	1	0.7	3	2.3	5	3.7	0	0.0	1	5.3	10	1.9
	Total	4	1.5	4	1.7	11	3.9	5	2.1	1	2.2	25	2.3
Androgen drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Estrogen or progesterone drug	Male	0	0.0	0	0.0	0	0.0	3	2.4	0	0.0	3	0.6
	Female	0	0.0	4	3.1	0	0.0	0	0.0	0	0.0	4	0.8

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Estrogen or progesterone drug	Total	0	0.0	4	1.7	0	0.0	3	1.3	0	0.0	7	0.7
Mixed hormone drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other hormone drug	Male	0	0.0	0	0.0	0	0.0	2	1.6	1	3.8	3	0.6
	Female	0	0.0	0	0.0	1	0.7	2	1.8	1	5.3	4	0.8
	Total	0	0.0	0	0.0	1	0.4	4	1.7	2	4.4	7	0.7
Agent acting on urinary system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent acting on genital system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Uterine constrictor	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Contraceptive	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	2	1.5	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4
	Total	2	0.8	0	0.0	0	0.0	0	0.0	0	0.0	2	0.2
Drugs for hemorrhoids	Male	0	0.0	0	0.0	2	1.3	0	0.0	0	0.0	2	0.4
	Female	0	0.0	0	0.0	1	0.7	0	0.0	0	0.0	1	0.2
	Total	0	0.0	0	0.0	3	1.1	0	0.0	0	0.0	3	0.3
Other agent acting on urogenital or anus	Male	0	0.0	0	0.0	5	3.4	10	7.9	2	7.7	17	3.2
	Female	0	0.0	0	0.0	0	0.0	3	2.7	1	5.3	4	0.8
	Total	0	0.0	0	0.0	5	1.8	13	5.5	3	6.7	21	2.0

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Integumentary microbiosite	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Protective agent for wound	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pyostatic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Analgenic (external application)	Male	1	0.8	0	0.0	0	0.0	1	0.8	0	0.0	2	0.4
	Female	1	0.7	0	0.0	1	0.7	0	0.0	0	0.0	2	0.4
	Total	2	0.8	0	0.0	1	0.4	1	0.4	0	0.0	4	0.4
Integumentary Parasiticide	Male	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
Emollient	Male	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
Hair. tonic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Bath liquid	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other surface-acting agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other surface-acting agent	Female	1	0.7	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2
	Total	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
Dental local anesthetic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for devitalization of the pulp	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Analgesic or sedative for dental use	Male	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	1	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
Agent for mummification of the pulp	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for pulp capping	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics for dental use	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent for dental use	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Vitamin A or D	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

## Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Vitamin A or D	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Vitamin B1	Male	0	0.0	1	1.0	1	0.7	2	1.6	2	7.7	6	1.1
	Female	0	0.0	0	0.0	1	0.7	2	1.8	1	5.3	4	0.8
	Total	0	0.0	1	0.4	2	0.7	4	1.7	3	6.7	10	0.9
Vitamin B complex	Male	2	1.6	1	1.0	7	4.7	9	7.1	7	26.9	26	4.9
	Female	3	2.2	4	3.1	1	0.7	11	9.8	4	21.1	23	4.3
	Total	5	1.9	5	2.1	8	2.8	20	8.4	11	24.4	49	4.6
Vitamin C	Male	0	0.0	1	1.0	0	0.0	1	0.8	0	0.0	2	0.4
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	1	0.4	0	0.0	1	0.4	0	0.0	2	0.2
Vitamin E	Male	0	0.0	0	0.0	1	0.7	2	1.6	0	0.0	3	0.6
	Female	0	0.0	1	0.8	0	0.0	0	0.0	0	0.0	1	0.2
	Total	0	0.0	1	0.4	1	0.4	2	0.8	0	0.0	4	0.4
Vitamin K	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	2	1.8	0	0.0	2	0.4
	Total	0	0.0	0	0.0	0	0.0	2	0.8	0	0.0	2	0.2
Mixed vitamin	Male	1	0.8	0	0.0	0	0.0	3	2.4	0	0.0	4	0.8
	Female	1	0.7	1	0.8	2	1.5	3	2.7	1	5.3	8	1.5
	Total	2	0.8	1	0.4	2	0.7	6	2.5	1	2.2	12	1.1
Other vitamin	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Calcium	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	1	0.8	4	3.0	2	1.8	1	5.3	8	1.5
	Total	0	0.0	1	0.4	4	1.4	2	0.8	1	2.2	8	0.8

(Continued)



Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Mineral preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	1	0.9	1	5.3	2	0.4
	Total	0	0.0	0	0.0	0	0.0	1	0.4	1	2.2	2	0.2
Saccaride preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Organic acid preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
protein or amino acid preparation	Male	0	0.0	1	1.0	0	0.0	0	0.0	0	0.0	1	0.2
	Female	0	0.0	0	0.0	1	0.7	0	0.0	1	5.3	2	0.4
	Total	0	0.0	1	0.4	1	0.4	0	0.0	1	2.2	3	0.3
Hormone drugs made from internal organ	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Drugs for infant	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other tonic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Blood substitute	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Styptic	Male	0	0.0	0	0.0	1	0.7	2	1.6	0	0.0	3	0.6

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Styptic	Female	3	2.2	1	0.8	2	1.5	1	0.9	0	0.0	7	1.3
	Total	3	1.1	1	0.4	3	1.1	3	1.3	0	0.0	10	0.9
Anticoagulant	Male	0	0.0	1	1.0	0	0.0	0	0.0	0	0.0	1	0.2
	Female	0	0.0	0	0.0	2	1.5	0	0.0	0	0.0	2	0.4
	Total	0	0.0	1	0.4	2	0.7	0	0.0	0	0.0	3	0.3
Other blood substitute	Male	0	0.0	0	0.0	7	4.7	12	9.5	4	15.4	23	4.3
	Female	0	0.0	0	0.0	3	2.2	9	8.0	2	10.5	14	2.6
	Total	0	0.0	0	0.0	10	3.5	21	8.8	6	13.3	37	3.5
Agent for hemodialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for peritoneal dialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent for dialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for liver disease	Male	1	0.8	1	1.0	2	1.3	2	1.6	0	0.0	6	1.1
	Female	0	0.0	0	0.0	2	1.5	1	0.9	0	0.0	3	0.6
	Total	1	0.4	1	0.4	4	1.4	3	1.3	0	0.0	9	0.8
Antidote	Male	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	1	0.2
	Female	0	0.0	0	0.0	1	0.7	1	0.9	0	0.0	2	0.4
	Total	0	0.0	0	0.0	1	0.4	2	0.8	0	0.0	3	0.3
Agent for habitual intoxication	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

## Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Agent for habitual intoxication	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Drugs for gout	Male	3	2.3	6	5.8	3	2.0	9	7.1	2	7.7	23	4.3
	Female	0	0.0	0	0.0	1	0.7	1	0.9	0	0.0	2	0.4
	Total	3	1.1	6	2.6	4	1.4	10	4.2	2	4.4	25	2.3
Enzyme preparation	Male	1	0.8	1	1.0	2	1.3	1	0.8	0	0.0	5	0.9
	Female	0	0.0	2	1.5	4	3.0	4	3.6	0	0.0	10	1.9
	Total	1	0.4	3	1.3	6	2.1	5	2.1	0	0.0	15	1.4
Antidiabetics	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.7	0	0.0	0	0.0	1	0.2
	Total	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	1	0.1
Other agent on metabolic system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Unclassified agent on metabolic system	Male	2	1.6	1	1.0	2	1.3	1	0.8	2	7.7	8	1.5
	Female	1	0.7	3	2.3	2	1.5	3	2.7	0	0.0	9	1.7
	Total	3	1.1	4	1.7	4	1.4	4	1.7	2	4.4	17	1.6
Chlorophyll agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pigment agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other cell activation agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	1	0.9	0	0.0	1	0.2
	Total	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Alkylating agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antimetabolite	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.7	1	0.9	0	0.0	2	0.4
	Total	0	0.0	0	0.0	1	0.4	1	0.4	0	0.0	2	0.2
Antibiotics for antitumor agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Plant component for antitumor agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other antitumor agent	Male	1	0.8	0	0.0	1	0.7	1	0.8	1	3.8	4	0.8
	Female	1	0.7	0	0.0	1	0.7	3	2.7	0	0.0	5	0.9
	Total	2	0.8	0	0.0	2	0.7	4	1.7	1	2.2	9	0.8
Antihistaminic drug	Male	3	2.3	2	1.9	4	2.7	1	0.8	1	3.8	11	2.1
	Female	10	7.4	3	2.3	3	2.2	3	2.7	0	0.0	19	3.6
	Total	13	4.9	5	2.1	7	2.5	4	1.7	1	2.2	30	2.8
Agent for stimulus therapy	Male	0	0.0	0	0.0	0	0.0	2	1.6	0	0.0	2	0.4
	Female	0	0.0	1	0.8	0	0.0	0	0.0	0	0.0	1	0.2
	Total	0	0.0	1	0.4	0	0.0	2	0.8	0	0.0	3	0.3
Agent for nonspecific immunogenicity	Male	0	0.0	0	0.0	0	0.0	0	0.0	1	3.8	1	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	1	2.2	1	0.1
Other antiallergic drug	Male	5	3.9	5	4.9	10	6.7	9	7.1	1	3.8	30	5.6

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other antiallergic drug	Female	8	5.9	10	7.6	8	5.9	3	2.7	0	0.0	29	5.4
	Total	13	4.9	15	6.4	18	6.3	12	5.0	1	2.2	59	5.5
Crude drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.7	1	0.9	0	0.0	2	0.4
	Total	0	0.0	0	0.0	1	0.4	1	0.4	0	0.0	2	0.2
Herb medicine	Male	12	9.3	4	3.9	22	14.8	16	12.7	2	7.7	56	10.5
	Female	16	11.8	18	13.7	20	14.8	15	13.4	6	31.6	75	14.1
	Total	28	10.6	22	9.4	42	14.8	31	13.0	8	17.8	131	12.3
Other drugs by a herb medicine	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(+))	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(-))	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(+), G(-))	Male	0	0.0	1	1.0	3	2.0	2	1.6	0	0.0	6	1.1
	Female	4	2.9	1	0.8	6	4.4	1	0.9	0	0.0	12	2.3
	Total	4	1.5	2	0.9	9	3.2	3	1.3	0	0.0	18	1.7
Antibiotics (G(+), G(-), Mycoplasma)	Male	0	0.0	1	1.0	3	2.0	2	1.6	1	3.8	7	1.3
	Female	2	1.5	0	0.0	0	0.0	1	0.9	0	0.0	3	0.6
	Total	2	0.8	1	0.4	3	1.1	3	1.3	1	2.2	10	0.9
Antibiotics (Rickettsia, Chlamidia)	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Antibiotics (Rickettsia, Chlamidia)	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics against acid- fastbacterium	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics against fungus	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other antibiotics	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.7	0	0.0	0	0.0	1	0.2
	Total	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	1	0.1
Sulfa drug	Male	0	0.0	0	0.0	1	0.7	0	0.0	0	0.0	1	0.2
	Female	1	0.7	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2
	Total	1	0.4	0	0.0	1	0.4	0	0.0	0	0.0	2	0.2
Antituberculous agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Drugs for leprosy	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Synthetic antibiotics	Male	1	0.8	0	0.0	2	1.3	0	0.0	0	0.0	3	0.6
	Female	1	0.7	1	0.8	1	0.7	4	3.6	0	0.0	7	1.3
	Total	2	0.8	1	0.4	3	1.1	4	1.7	0	0.0	10	0.9
Antiviral agent	Male	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	1	0.2
	Female	0	0.0	1	0.8	1	0.7	0	0.0	0	0.0	2	0.4
	Total	0	0.0	1	0.4	1	0.4	1	0.4	0	0.0	3	0.3

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other agent for chemotherapy	Male	1	0.8	1	1.0	1	0.7	1	0.8	0	0.0	4	0.8
	Female	2	1.5	1	0.8	0	0.0	1	0.9	0	0.0	4	0.8
	Total	3	1.1	2	0.9	1	0.4	2	0.8	0	0.0	8	0.8
Vaccine	Male	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0	1	0.2
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
Poison or toxoid	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antitoxic serum or leptospira serum	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Blood supplements	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for biological test	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Mixed biological agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other biological agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Protozoicide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Protozoicide	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Parasiticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other drug against parasitic disease	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Diluent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Ointment base	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Solution	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Corrective, tinction	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Emulsifier	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other dispensing drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)



Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other dispensing drug	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
X-ray contrast medium	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for functional examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other test	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antiseptic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pesticide for prevention of epidemics	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Insecticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Insecticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Rodenticide	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other drug used for public health	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for general examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for blood analysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for chemical analysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for immunologic test	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for microbiological examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Reagent for pathological examination	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Radiopharmaceutical for diagnosis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other reagent for diagnosis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks.

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other reagent for diagnosis	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Strap	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other drug not for treatment	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Opiate alkaloid agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Coca alkaloid agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other alkaloid agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Synthetic narcotic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Unclassifiable drug	Male	0	0.0	1	1.0	0	0.0	1	0.8	0	0.0	2	0.4
	Female	0	0.0	0	0.0	0	0.0	3	2.7	0	0.0	3	0.6
	Total	0	0.0	1	0.4	0	0.0	4	1.7	0	0.0	5	0.5
No use of drug	Male	60	46.5	44	42.7	41	27.5	24	19.0	4	15.4	173	32.5
	Female	53	39.0	41	31.3	30	22.2	19	17.0	1	5.3	144	27.0
	Total	113	42.6	85	36.3	71	25.0	43	18.1	5	11.1	317	29.7
Total	Male	129	100.0	103	100.0	149	100.0	126	100.0	26	100.0	533	100.0
	Female	136	100.0	131	100.0	135	100.0	112	100.0	19	100.0	533	100.0
	Total	265	100.0	234	100.0	284	100.0	238	100.0	45	100.0	1066	100.0

## IV. Nutritional Examinations