

Figure 3. Immunohistochemical analysis of transplanted MSCs. **(a)**, Native NP cells in the normal control disc do not stain with GFP. Staining for **(b)** keratin sulfate, **(c)** chondroitin-4-sulfate, **(d)** chondroitin-6-sulfate, and **(e)** Type II collagen showed positive expressions in both native NP cells marked by arrow and GFP-positive MSCs marked by asterisk. On the other hand, there was no expression of **(f)** Type I collagen in either native cells or MSCs. Bars = 25 μm .

demonstrated that some of GFP-positive MSCs costain together with keratan sulfate, chondroitin sulfate, and Type II collagen, the main matrix components of intervertebral disc cells. Furthermore, these cells stained positively for HIF-1 α , GLUT-1, and MMP-2. Oil-red-O and Von Kossa staining showed that there was no apparent differentiation of MSCs toward adipogenesis or osteogenesis.

These results suggest that MSCs were induced by the microenvironment of the nucleus pulposus to undergo differentiation toward the nucleus pulposus cell phenotype *in vivo*. It is unlikely that atelocollagen scaffold alone could induce such a change in MSC phenotype. In our preliminary experiment with culturing MSCs in atelocollagen *in vitro* failed to induce chondrogenesis without chondrogenic induction medium containing TGF- β .




Figure 4. Some of the MSCs transplanted to NP area of degenerative discs showed expression of hypoxia-responsive proteins. Triple-colored image of cells on the nucleus region showing that some of the NP cells including GFP-positive MSCs (shown in higher magnification in the right panels) stain positively for **(a)** HIF-1 α , **(b)** GLUT-1, and **(c)** MMP-2. Bars = 25 μm (left), 5 μm (right).

Therefore, it is more likely that the strong expression of nucleus pulposus chondrocyte-like phenotype of the transplanted MSCs was mediated by stimulation and interaction from the surrounding microenvironment. However, much remains to be investigated regarding the differentiation conditions required for definitive induction. It was not possible to determine whether any of the MSCs differentiated into nucleus pulposus chondrocyte-like cells, as the complete profile characteristic of nucleus pulposus cells is unknown.³⁸ Further investigation is needed on nucleus pulposus cell development and profiles of final cell characteristics in the etiology of disc degeneration.

We used atelocollagen as a cell-delivering scaffold. This collagen gel is safe, as the antigenic telopeptide region is removed by pepsin digestion and differential salt




Figure 5. MSC-transplanted discs do not show apparent differentiation toward adipogenesis or osteogenesis: a, Oil-Red O staining; and b, Von Kossa staining. Bars = 1 mm.

precipitation during purification. Atelocollagen is a liquid at 4 C but gels after incubation at 37 C. Atelocollagen is composed of Type I collagen purified from calf dermis and is used by physicians for subcutaneous injection during cosmetic and plastic surgery. Atelocollagen dissolves in water after 2 weeks, another evidence indicating that it would have minimal effects on differentiation of MSCs when used as scaffold for cell delivery.

Biochemical analysis and RT-PCR showed that the total glycosaminoglycan content of the MSC-transplanted discs recovered significantly and the expression of collagen and proteoglycan genes was restored compared with nontreated discs. This supports the proposal that transplanted MSCs differentiate into cells, presenting a nucleus pulposus chondrocyte-like phenotype.

The lack of an ideal animal model that directly reflects symptomatic disc degeneration is a problem. The rabbit disc degeneration model used in this study mimics nucleotomy, a popular surgical procedure used in the treatment of herniated lumbar discs. Nucleus pulposus tissue is evacuated in this procedure, and it causes partial degeneration evidenced as nucleus pulposus cell loss, re-




Figure 6. Changes in proteoglycan content in the three groups. Bar graph showing significant decrease of PG content in discs from degenerative disc model (DG) compared with normal control (NC), which is delayed in MSC-transplanted discs (MSC).

sulting to a decrease in proteoglycan content and disc height. This model has been used in many cell transplantation studies.^{18–21} It is necessary to confirm the effect of this procedure in a more appropriate large-animal model, possibly in primates, before any of the newly developed techniques can be considered clinically applicable. The fact that only about 5 mg of nucleus tissue (about 40%–50% of the total nucleus pulposus tissue in L4–L5 discs) was evacuated raises the question of whether the microenvironment after nucleus pulposus




Figure 7. RT-PCR results showed that while increase in Type I collagen mRNA expression is seen in discs from degenerative disc model (DG), Type II collagen and aggrecan mRNA are markedly decreased. In MSC-transplanted discs (MSC), increase in Type I collagen expression is suspended with maintaining expression of Type II collagen and aggrecan mRNA.

evacuation promotes appropriate differentiation in transplanted MSCs. It is important to observe MSC differentiation in an undamaged nucleus pulposus microenvironment. However, if one were to apply this technique clinically, target intervertebral discs would be damaged to some extent. Furthermore, many obstacles remain to be overcome before application to humans can be considered; these include the appropriate age of patients, extent of disc degeneration, and the efficacy of treatment of degenerative damage caused by aging *versus* postspinal surgery. With regard to the effects of age and extent of disc degeneration, it is suggested that it would be best to apply this technique to mildly to moderately damaged intervertebral discs, as it is impossible to restore highly damaged discs by altered nutrition, and a ruptured annular structure makes cell survival and retention in the closed cavity crucial.

■ Conclusion

We have shown that MSCs can differentiate into cells expressing some of the major phenotypes of nucleus pulposus chondrocyte-like cells. Because it is easier to obtain MSCs from bone marrow than from chondrocytes, there is a good chance that the former may become an alternative source of cells for cell therapy in intervertebral disc regeneration. Although there is a need for further research to define protocols for obtaining the maximum effect of this treatment and to develop patient selection methods, our data indicate that transplantation of autologous MSCs may become an effective option for treating the relentlessly progressive process of intervertebral disc degeneration.

■ Key Points

- Differentiation of mesenchymal stem cells was assessed after transplantation to the nucleus pulposus area of degenerative disc model in rabbits.
- Transplanted mesenchymal stem cells survived and proliferated after transplantation to degenerative discs.
- In phenotype comparison performed before and after transplantation, mesenchymal stem cells expressed apparently different character, suggesting differentiation toward nucleus pulposus cells after transplantation.
- Proteoglycan content and matrix-related gene expression recovered close to normal after mesenchymal stem cell transplantation.
- Despite many obstacles needed to overcome in concerning clinical application, mesenchymal stem cells may become an alternative cell source for cell transplantation therapy for treatment of disc degeneration.

Acknowledgments

The authors thank Drs. Y. Oka, I. Saito, T. Kutsuna, and M. Kamimoto for their generous assistance.

References

1. Deyo RA, Weinstein JN. Low back pain. *N Engl J Med* 2001;344:363–70.
2. Frymoyer JW, Cats-Baril WL. An overview of the incidence and costs of low back pain. *Orthop Clin North Am* 1991;22:263–271.
3. Nishida K, Kang JD, Gilbertson LG, et al. 1999 Volvo Award in basic science. Modulation of the biologic activity of the rabbit intervertebral disc by gene therapy: an in vivo study of adenovirus-mediated transfer of the human transforming growth factor beta1 encoding gene. *Spine* 1999;24:2419–25.
4. Anderson JAD. Back pain and occupation. In: *The Lumbar Spine and Back Pain*, 3rd ed. London: Churchill Livingstone, 1987:2–36.
5. An HS, Thonar EJ, Masuda K. Biological repair of intervertebral disc. *Spine* 2003;28(suppl):86–92.
6. Alini M, Roughley PJ, Antoniou J, et al. A biological approach to treating disc degeneration: not for today, but maybe for tomorrow. *Eur Spine J* 2002;11(suppl):215–20.
7. Nishida K, Kang JD, Suh JK, et al. Adenovirus-mediated gene transfer to nucleus pulposus cells: implications for the treatment of intervertebral disc degeneration. *Spine* 1998;23:2437–42.
8. Walsh AJ, Bradford DS, Lotz JC, et al. In vivo growth factor treatment of degenerated intervertebral discs. *Spine* 2004;29:156–63.
9. Yoon ST, Kim SK, Li J, et al. The effect of bone morphogenetic protein-2 on rat intervertebral disc cells in vitro. *Spine* 2003;28:1773–80.
10. Masuda K, Takegami K, An H, et al. Recombinant osteogenic protein-1 upregulates extracellular matrix metabolism by rabbit annulus fibrosus and nucleus pulposus cells cultured in alginate beads. *J Orthop Res* 2003;21:922–30.
11. Mizuno H, Roy AK, Vacanti CA, et al. Tissue-engineered composites of annulus fibrosus and nucleus pulposus for intervertebral disc replacement. *Spine* 2004;29:1290–7.
12. Seguin CA, Grynbas MD, Pilliar RM, et al. Tissue engineered nucleus pulposus tissue formed on a porous calcium polyphosphate substrate. *Spine* 2004;29:1299–306.
13. Alini M, Li W, Markovic P, et al. The potential and limitations of a cell-seeded collagen/hyaluronan scaffold to engineer an intervertebral disc-like matrix. *Spine* 2003;28:446–54.
14. Brisby H, Tao H, Ma DD, et al. Cell therapy for disc degeneration—potentials and pitfalls. *Orthop Clin North Am* 2004;35:85–93.
15. Ganey T, Libera J, Moos V, et al. Disc chondrocyte transplantation in a canine model: a treatment for degenerated or damaged intervertebral disc. *Spine* 2003;28:2609–20.
16. Gruber HE, Johnson TL, Leslie K, et al. Autologous intervertebral disc cell implantation: a model using Psammomys obesus, the sand rat. *Spine* 2002;27:1626–33.
17. Gorenssek M, Jaksimovic C, Kregar-Velikonja N, et al. Nucleus pulposus repair with cultured autologous elastic cartilage derived chondrocytes. *Cell Mol Biol Lett* 2004;9:363–73.
18. Nishimura K, Mochida J. Percutaneous reinsertion of the nucleus pulposus: an experimental study. *Spine* 1998;22:1531–9.
19. Okuma M, Mochida J, Nishimura K, et al. Reinsertion of activated nucleus pulposus cells retards intervertebral disc degeneration: an in vitro and in vivo experimental study. *J Orthop Res* 2000;18:988–97.
20. Nomura T, Mochida J, Okuma M, et al. Nucleus pulposus allograft retards intervertebral disc degeneration. *Clin Orthop* 2001;389:94–101.
21. Sakai D, Mochida J, Yamamoto Y, et al. Transplantation of mesenchymal stem cells embedded in Atelocollagen gel to the intervertebral disc: a potential therapeutic model for disc degeneration. *Biomaterials* 2003;24:3531–41.
22. Sobajima S, Kim JS, Niyibizi C, et al. Stem cell therapy for intervertebral disc diseases in vitro and in vivo feasibility study. *Orthop Res Soc Trans* 2004;29:137.
23. Crevensten G, Walsh AJ, Ananthakrishnan D, et al. Intervertebral disc cell therapy for regeneration: mesenchymal stem cell implantation in rat intervertebral discs. *Ann Biomed Eng* 2004;32:430–4.
24. Horner HA, Roberts S, Bielby RC, et al. Cells from different regions of the intervertebral disc. *Spine* 2002;27:1018–28.
25. Gruber HE, Hanley EN Jr. Human disc cells in monolayer vs 3D culture: cell shape, division and matrix formation. *BMC Musculoskeletal Disord* 2000;1:1.
26. Gruber HE, Stasky AA, Hanley EN Jr. Characterization and phenotypic stability of human disc cells in vitro. *Matrix Biol* 1997;16:285–8.
27. Risbud MV, Albert TJ, Guttapalli A, et al. Differentiation of mesenchymal stem cells towards a nucleus pulposus-like phenotype in vitro: implications for cell-based transplantation therapy. *Spine* 2004;29:2627–32.
28. Deans RJ, Moseley AB. Mesenchymal stem cells: biology and potential clinical uses. *Exp Hematol* 2000;28:875–84.
29. Caplan AI. Mesenchymal stem cells. *J Orthop Res* 1991;9:641–50.
30. Hawley RG, Lieu FH, Fong AZ, et al. Versatile retroviral vectors for potential use in gene therapy. *Gene Ther* 1994;1:136.

31. Oki M, et al. Efficient lentiviral transduction of human cord blood CD34⁺ cells followed by their expansion and differentiation into dendritic cells. *Exp Hematol* 2001;29:1210–7.
32. Gerson SL. Mesenchymal stem cells: no longer second class marrow citizens. *Nat Med* 1999;5:262–4.
33. Horwitz EM, Prockop DJ, Fitzpatrick LA, et al. Transplantability and therapeutic effects of bone marrow-derived mesenchymal cells in children with osteogenesis imperfecta. *Nat Med* 1999;5:309–13.
34. Liechty KW, MacKenzie TC, Shaaban AF, et al. Human mesenchymal stem cells engraft and demonstrate site-specific differentiation after in utero transplantation in sheep. *Nat Med* 2000;6:1282–6.
35. Toma C, Pittenger MF, Cahill KS, et al. Human mesenchymal stem cells differentiate to a cardiomyocyte phenotype in the adult murine heart. *Circulation* 2002;105:93–8.
36. Sakai D, Mochida J, Iwashina T, et al. Regenerative effects of transplanting mesenchymal stem cells embedded in Atelocollagen to the degenerated intervertebral disc. *Biomaterials* in press.
37. Rajpurohit R, Risbud MV, Ducheyne P, et al. Phenotypic characteristics of the nucleus pulposus: expression of hypoxia inducing factor-1, glucose transporter-1 and MMP-2. *Cell Tissue Res* 2002;30:401–7.
38. Roughley PJ. Biology of intervertebral disc aging and degeneration: involvement of the extracellular matrix. *Spine* 2004;29:2691–9.

Effect of Reinsertion of Activated Nucleus Pulposus on Disc Degeneration: An Experimental Study on Various Types of Collagen in Degenerative Discs

Keita Watanabe,¹ Joji Mochida,¹ Takeshi Nomura,¹ Masahiko Okuma,¹
Kou Sakabe,² and Kanji Seiki²

¹Department of Orthopedic Surgery Functional Reconstruction, Tokai University School of Medicine, Isehara, Kanagawa, Japan

²Department of Morphology, Tokai University School of Medicine, Isehara, Kanagawa, Japan

We examined the emergence and sequential changes in type I, II, and VI collagen production in an experimental rabbit model of disc degeneration. Type I collagen was minimally present initially and did not change over 24 weeks. Type I collagen seemed to have no effect on the degenerative process in this model. Staining for type II collagen was positive circumferentially in chondrocytelike cells and was mild in the early phase of disc degeneration, when the chondrocytelike cells began to appear in the inner layers of the annulus fibrosus. The stain became stronger during the middle phase when the chondrocytelike cells arranged themselves in cluster. Compared with type II collagen, the staining for type VI collagen was relatively strong early in the degenerative process. These findings led us to speculate that these chondrocytelike cells play an active role in the degenerative process. The reinsertion of nucleus pulposus cells cocultured with annulus fibrosus delayed disc degeneration and the emergence of chondrocytelike cells. Considering that the emergence of chondrocytelike cells which produce type II and type VI collagen is delayed in discs with the injection of cocultured nucleus pulposus cells by annulus fibrosus cells, we conclude that chondrocytelike cells that produce type VI collagen also seems to accelerate degeneration. Type VI collagen is produced at an earlier phase than type II collagen and may be both active agent and a marker for disc degeneration.

Keywords Low Back Pain, Disc Herniation, Nucleus Pulposus, Annulus Fibrosus.

Lumbar disc herniation is one of the main causes of chronic low back pain and neurologic disorders. Using an animal model, Nishimura et al. [7] found that the reinsertion of autogenous

nucleus pulposus can delay further disc degeneration. Okuma et al. [8] reported that the reinsertion of nucleus pulposus cells, which had been activated by coculture with annulus fibrosus cells, delayed the progression of degenerative change in an animal model. The authors also observed the emergence of chondrocytelike cells, which stained positively for type II collagen. On the other hand, while the main types of collagen in normal discs are types I and II, other types of collagen also exist in abundance. The purpose of this article is to document the emergence and record the chronologic change in types I, II, and VI collagen production stimulated by reinsertion of cocultured nucleus pulposus cells.

MATERIALS AND METHODS

This study was conducted under a protocol approved by the Animal Experimentation Committee, Tokai University School of Medicine.

Experimental Study

Twenty Japanese white rabbits weighing 1.3 kg were anesthetized with intravenous pentobarbital sodium (30 mg/kg). The nucleus pulposus from four intervertebral discs (L2/L3, L3/L4, L4/L5, and L5/L6) were aspirated through an anterolateral approach using a 21-gauge needle on a 10-ml syringe to create a model of a degenerative disc under aseptic conditions. After the nucleus pulposus was aspirated, the ventral one-fourth of the annulus fibrosus of the L2/L3 and L3/L4 intervertebral discs also were excised.

Cell Culture

Cell culture was performed according to the method of Shinmei et al. as described previously [14].

Received 3 January 2002; accepted 15 July 2002.

Address correspondence to Joji Mochida, Department of Orthopedic Surgery Functional Reconstruction, Tokai University School of Medicine, Bouseidai, Isehara, Kanagawa, 259-1193 Japan. E-mail: jomo@is.icc.u-tokai.ac.jp

The aspirated nucleus pulposus and the extirpated annulus fibrosus were used for cell culture. The nucleus pulposus was digested in a mixture of 0.2% pronase E (actinase E; Kakenkagaku, Tokyo) and 0.0125% collagenase P (Boehringer Mannheim GmbH, Mannheim, Germany) for 1 h. The annulus fibrosus was digested in 0.025% collagenase P for 6 h after treatment for 1 h with 0.3% actinase. The digested nucleus pulposus and annulus fibrosus cells were washed with Dulbecco's modified Eagle's medium (DMEM) (Nissui, Tokyo) three times. Centrifugation was done at 1500 rpm to collect the cells [13, 14].

A specially designed culture plate (Iwaki Glass, Chiba, Japan) was used for coculture of the two cellular components. A membrane culture insert (Iwaki Glass, Chiba, Japan) was set in each well of the tissue culture plate, dividing the well into two chambers. This membrane was a mixed cellulose ester with 0.45- μ m pores at the bottom and prevented any exchange of cellular components. The cells collected by centrifugation were used for coculture of the nucleus pulposus and annulus fibrosus (Figure 1). Coculture was performed for 2 weeks in the medium containing 10% fetal bovine serum (FBS; lot SF10305, Bocknet, Ontario, Canada) and 50 mg/L ascorbic acid (Daiichi Chemical, Tokyo).

Preparation for Reinsertion

After 2 weeks, the cocultured nucleus pulposus cells had increased in number and adhered to the bottom of the culture plate. These cocultured cells were scraped and gathered for reinsertion. The cells were resuspended in 0.02% DMEM. A second procedure was performed on the contralateral side to the first operation under anesthesia induced by intravenous injection of pentobarbital sodium. Cocultured nucleus pulposus cells suspended in 0.02 ml DMEM were injected into the L5/L6 intervertebral disc using a 27-gauge needle with a microinjector.




Figure 1. A specially designed culture plate (Iwaki Glass, Chiba, Japan) was used for coculture of the two cellular components. A membrane culture insert (Iwaki Glass, Chiba, Japan) was set in each well of the tissue culture plate, dividing the well into two chambers. This membrane was a mixed cellulose ester with 0.45- μ m pores at the bottom and prevented any exchange of cellular components. The cells collected by centrifugation were used for coculture of the nucleus pulposus and annulus fibrosus.

Tissue Preparation

Five rabbits each were killed by intravenous injection of high-dose pentobarbital sodium (120 ml/kg) 2, 4, 8, and 24 weeks after the second procedure, and the L4/L5 disc (negative control group), the L5/L6 disc (injected cell group), and the L6/L7 disc (normal control group) were excised immediately. Frozen sections were made for histologic examination. Samples were fixed with 4% paraformaldehyde in phosphate-buffered saline immediately after their removal, decalcified in Plank-Rychlo solution, dehydrated in a graded alcohol series (70%, 80%, 85%, 90%, 95%, and 99%), and embedded in optimal cutting temperature (OCT) compound (Sakura Finetek, Torrance, CA). Seven-micrometer-thick sections were stained with hematoxylin and eosin.

Immunohistochemistry

Each section was stained with goat anti-type I, II, or VI collagen (Southern Biotechnology Associates, Birmingham). Data from injected cell (L5/L6), negative control (L4/L5), and normal control (L6/L7) discs were compared.

RESULTS

Histological Findings

The cultured cells were subconfluent at reinsertion. No evidence of infection was observed. In the normal control group, the structure of the annulus fibrosus and nucleus pulposus tissue had been morphologically preserved in all discs at each stage (Figure 2). Although the nucleus pulposus cells were still present at week 8 in the negative control group, any nucleus pulposus cells disappeared at week 24. In addition the proliferation of fibrous tissue and a serpentine or indistinct appearance of the annulus fibrosus architecture were observed. There were typical degenerative changes in disc structure. Chondrocytelike cells were seen mainly in the inner layer of the annulus fibrosus from week 4 to 8. At the week 24, chondrocytelike cells were present throughout the annulus fibrosus (Figure 3). In the injected cell group, the nucleus pulposus cells were still present at week 24. Chondrocytelike cells in the inner layer of annulus fibrosus first emerged at week 8 and continued to increase in number through week 24. However, the number of chondrocytelike





Figure 2. Normal control group. The structure of the annulus fibrosus and nucleus pulposus tissue had been morphologically preserved (week 24, H&E, $\times 6.6$).

cells in this group was smaller than in the negative control group. Degenerative changes in the disc structure were less extensive in the injected cell group than in the negative control group (Figure 4).



Immunohistochemistry

Type I Collagen

Staining for type I collagen tended to occur in the outer layer of the annulus fibrosus in the normal control group (Figure 5). However, the staining was not prominent and did not change over time. No significant differences between normal control group, negative control group, and injected cell group were found.

Type II Collagen

Staining for type II collagen was relatively more prominent in the nucleus pulposus and the inner layer of annulus fibrosus in the normal control group (Figure 5). In the negative control group, faint circumferential staining for type II collagen was detected inside the chondrocytelike cells in the inner layer at week 4, with stronger staining at week 8. By week 24, all layers stained more strongly as the number of chondrocytelike cells increased (Figure 3). In contrast, in the injected cell group, in which the emergence of chondrocytelike cells was delayed, staining for type II collagen inside the chondrocytelike cells did not appear until week 8, and all layers stained positive for type II collagen by week 24 (Figure 4).

Type VI Collagen

Staining for type VI collagen was stronger in the normal control group than staining for type I or II collagen. Staining for type VI collagen was observed throughout the disc, including the nucleus pulposus and the inner and outer layers of the annulus fibrosus (Figure 5). In the negative control group, strong circumferential staining for type VI collagen was detected inside the chondrocytelike cells in the inner layer at week 4. Staining for type VI collagen was stronger than for type II collagen at the week 4. Proliferating fibrous tissues were also stained. Strong circumferential staining inside the chondrocytelike cells also was observed at weeks 8 and 24 (Figure 3).

In the injected cell group, the staining patterns for type VI collagen were similar to that in the negative control group. However, the additional staining for type VI collagen appeared along with the emergence of chondrocytelike cells as it did with type II collagen. In short, the delayed emergence of chondrocytelike cells in the injected cell group correlated with a delay in type VI collagen production. But chondrocytelike cells were present in all layers at week 24, and all layers stained positively for type VI collagen (Figure 4).

Figure 3. Negative control group. Although the nucleus pulposus cells were still present at week 8 in the negative control group, any nucleus pulposus cells disappeared at week 24. In addition, proliferation of fibrous tissue and a serpentine or indistinct appearance of the annulus fibrosus architecture were observed. There were typical degenerative changes in disc structure (H&E, $\times 6.6$). Chondrocytelike cells were seen mainly in the inner layer of the annulus fibrosus from week 4 to 8. At week 24, chondrocytelike cells were present throughout the annulus fibrosus. The difference between the staining for type VI collagen and that for type II collagen was detected at week 4. Staining for type VI collagen was stronger than for type II collagen at the week 4 (H&E, type II collagen, type VI collagen, $\times 50$).




Figure 4. Injected cell group. The nucleus pulposus cells were still present at week 24. Degenerative changes in the disc structure were less extensive in the injected cell group than in the negative control group (H&E, $\times 6.6$). Chondrocytelike cells in the inner layer of annulus fibrosus first emerged at week 8 and continued to increase in number through week 24. However, the number of chondrocytelike cells in this group was smaller than in the negative control group. When the emergence of chondrocytelike cells was delayed, staining for type II and VI collagen inside the chondrocytelike cells did not appear until week 8 (H&E, type II collagen, type VI collagen, $\times 50$).




Figure 5. Staining for various types of collagen in normal disc. Staining for type I collagen was not prominent, but this staining tended to occur in the outer layer of the annulus fibrosus. Staining for type II collagen was relatively more prominent in the nucleus pulposus and the inner layer of the annulus fibrosus. Staining for type VI collagen was stronger than that for type I or II collagen. Staining for type VI collagen was observed throughout the disc, including the nucleus pulposus and the inner and outer layers of the annulus fibrosus. Type I collagen, type II collagen, type VI collagen, $\times 6.6$.

DISCUSSION

Structural degeneration and mechanical failure of the intervertebral disc is a major clinical problem, and some corrective surgical procedures, such as herniotomy via the posterior approach, percutaneous nucleotomy, and laser disc decompression, can actually accelerate disc degeneration under some circumstances [9].

To better understand the degenerative process of the disc components, it is helpful to have a better appreciation of the changes in the amount and distribution of the different types of collagen. The collagen framework of the intervertebral disc is composed primarily type I and II collagen. Takaishi [12] reported that the gene expression of type II collagen increases early in the process of disc degeneration. In addition, biochemical and immunohistologic analyses have demonstrated other types of collagen in the intervertebral disc [1–6, 10, 11]. Type VI collagen is surprisingly abundant in the bovine disc [10], and staining intensity for type VI collagen shows a tendency to be reduced in disc disease [1]. Andreas [6] reported immunolocalization of major interstitial collagen (types I–III, V, VI, and IX) in human lumbar intervertebral discs in patients of various ages, but the reason for this finding remains unclear.

In this experimental rabbit model of disc degeneration, staining for type I collagen tended to be limited to the outer layer of the annulus fibrosus in the normal control group. However, staining was rather mild. In addition, the staining pattern did not change over time, and there was no significant difference between normal control group, negative control group, and injected cell group. Therefore type I collagen does not seem to figure prominently in the degenerative process in this experimental model. The staining for type II collagen was observed in a circumferential pattern within the chondrocytelike cells, which signaled the beginning of the degenerative process, as reported by Nishimura and by Okuma [7, 8]. The intensity of staining within the chondrocytelike cells for type II collagen was mild at week 4, when the chondrocytelike cells first began to appear in the inner layers of the annulus fibrosus, and became stronger by the week 8, when the chondrocytelike cells began to arrange themselves in clusters. Staining remained strong through week 24, when the chondrocytelike cells were observed in all layers of the annulus fibrosus.

The staining pattern for type VI collagen resembled that of type II collagen. Strong circumferential staining for type VI collagen was detected inside the chondrocytelike cells. The difference between the staining for type VI collagen and that for type II collagen was detected at week 4. Staining for type VI collagen was stronger than for type II collagen at the week 4. Additionally, strong staining was observed in fibrous tissue that proliferated to fill the void in the nucleus pulposus. Many reports suggest that the deposition of type IV collagen is beneficial to healing. Although a similar increased deposition of type IV collagen was observed during the early phase of disc degeneration in our experimental model, the degeneration actually appeared to progress rather than regress as a result. Further, many chondrocytelike cells were observed at the week 24 in this experimental model. We hypothesize that these cells produce type VI collagen early in disc degeneration, but subsequently change to produce both type II and type VI collagen. It was not possible to determine whether the chondrocytelike cells that produced type VI collagen during the early phase of degeneration were the same cells that produced type II collagen subsequently.

Nishimura et al. [7] have developed an innovative procedure in which autogenous nucleus pulposus is reinserted into a degenerated disc. Using an experimental model, the authors showed that this intervention retards further degeneration of the disc. Furthermore, Okuma et al. [8] reported that reinsertion of nucleus pulposus cells activated by coculture with annulus fibrosus cells delays disc degeneration in an animal model similar to the one we used in the current study. Reinsertion of nucleus pulposus cells retarded disc degeneration and fewer chondrocytelike cells were observed. Okuma et al. [8] also have reported that the presence of type II collagen correlated with the appearance

of chondrocytelike cells in the inner layer of annulus fibrosus during progression of the degenerative process. The authors suggested that the presence of type II collagen appears to accelerate degeneration. We identified chondrocytelike cells that produced type VI collagen prior to the production of type II collagen. Considering that the emergence of chondrocytelike cells that produce type II and type VI collagen is delayed in the discs in which cocultured nucleus pulposus cells had been reinserted, we conclude that chondrocytelike cells that produce type VI collagen also seems to accelerate degeneration. Type VI collagen emerges at an earlier phase than type II collagen, and therefore may be an important marker for disc degeneration in addition to the role it plays in the actual pathophysiology.

REFERENCES

- [1] Roberts, S., Menage, J., Duance, V., Wotton, S., and Ayad, S. (1991). Collagen types around the cells of the intervertebral disc and cartilage end plate: An immunolocalization study. *Spine* 16:1030-1038.
- [2] Roberts, S., Menage, J., Duance, V., and Wotton, S. (1991). Type III collagen in the intervertebral disc. *Histochem. J.* 23:503-508.
- [3] Kaapa, E., Zhang, L.-Q., Muona, P., Holm, S., Vanharanta, H., and Peltonen, J. (1994). Expression of type I, III, and VI collagen mRNAs in experimentally injured porcine intervertebral disc. *Connect. Tissue Res.* 30:203-214.
- [4] Kaapa, E., Han, X., Holm, S., Peltonen, J., Takala, T., and Vanhanta, H. (1995). Collagen synthesis and types I, III, IV, and VI collagens in an animal model of disc degeneration. *Spine* 20:59-67.
- [5] Roberts, S., Ayad, S., and Menage, P.J. (1991). Immunolocalization of type VI collagen in the intervertebral disc. *Ann. Rheum. Dis.* 50:787-791.
- [6] Andreas, G.N., Boss, N., and Wiest, I. (1998). Immunolocalization of major interstitial collagen types in human lumbar intervertebral discs of various ages. *Virchows Arch.* 432:67-76.
- [7] Nishimura, K., and Mochida, J. (1998). Percutaneous reinsertion of the nucleus pulposus: An experimental study. *Spine* 23:1531-1538.
- [8] Okuma, M., Mochida, J., Nishimura, K., Sakabe, K., and Seiki, K. (2000). Reinsertion of stimulated nucleus pulposus cells retards intervertebral disc degeneration: An in vitro and in vivo experimental study. *J. Orthop. Res.* 18(6):988-997.
- [9] Mochida, J., Nishimura, K., Nomura, T., Toh, E., and Chiba, M. (1996). The importance of preserving disc structure in surgical approaches to lumbar disc herniation. *Spine* 21:1556-1563.
- [10] Jiann-Jiu, W., Eyre, D.R., and Slayter, H.S. (1987). Type VI collagen of the intervertebral disc. *Biochem. J.* 248:373-381.
- [11] Rufai, A., Benjamin, M., and Ralphs, J.R. (1995). The development of fibrocartilage in the rat intervertebral disc. *Anat. Embryol.* 192:53-62.
- [12] Takaishi, H., Nemoto, O., Shiota, M., Kikuchi, T., Yamada, H., Yamagishi, M., and Yabe, Y. (1997). Type-II collagen gene expression is transiently upregulated in experimentally induced degeneration of rabbit intervertebral disc. *J. Orthop. Res.* 15:528-538.
- [13] Shinmei, M., Kikuchi, T., Yamagishi, M., and Shimomura, Y. (1988). The role of interleukin-1 on proteoglycan metabolism of rabbit annulus fibrosus cells cultured in vitro. *Spine* 13:1284-1290.
- [14] Shinmei, M., Kikuchi, T., Yamagishi, M., and Shimomura, Y. (1988). Isolation and culture of the cells from annulus fibrosus and nucleus pulposus of rabbit intervertebral disc. *Trans. Orthop. Res. Soc.* 13:376.



Transplantation of mesenchymal stem cells embedded in Atelocollagen[®] gel to the intervertebral disc: a potential therapeutic model for disc degeneration

Daisuke Sakai^{a,b,*}, Joji Mochida^{a,b}, Yukihiro Yamamoto^{a,b}, Takeshi Nomura^a, Masahiko Okuma^a, Kazuhiro Nishimura^a, Tomoko Nakai^b, Kiyoshi Ando^b, Tomomitsu Hotta^b

^a Department of Orthopaedic Surgery, Functional Reconstruction, Tokai University School of Medicine, Bohseidai, Isehara, Kanagawa, 259-1193, Japan

^b Center for Regenerative Medicine, Tokai University School of Medicine, Bohseidai, Isehara, Kanagawa, 259-1193, Japan

Received 2 December 2002; accepted 24 March 2003

Abstract

Intervertebral disc degeneration is considered to be one of the major causes of low back pain. Despite this irreversible phenomenon, attempts to decelerate disc degeneration using various techniques have been reported. However, to date there has been no proven technique effective for broad clinical application. Based on previous studies, we hypothesize that maintenance of proteoglycan content in the disc is achieved by avoiding the depletion of nucleus pulposus and preserving the structure of the annulus is a primary factor in decelerating disc degeneration.

One novel approach to solve the dilemma of intervertebral disc degeneration is found at the stem cell level. Mesenchymal stem cells (MSCs) are known to possess the ability to differentiate into various kinds of cells from mesenchymal origin. Although the majority of cells that contribute to disc formation are known to obtain chondrocyte-like phenotypes, no reported study has emphasized the correlation with mesenchymal stem cells.

To evaluate the possible potential of MSCs in disc cell research and treatment of degenerative disc disease, autologous MSCs embedded in Atelocollagen[®] gel were transplanted into the discs of rabbits which had undergone a procedure proven to induce degeneration.

The results suggest that MSC transplantation is effective in decelerating disc degeneration in experimental models and provided new hopes for treatment of degenerative disc disease in humans. Atelocollagen[®] gel served as an important carrier of MSCs in transplantation, permitting proliferation, matrix synthesis and differentiation of MSCs. This study strengthens the viable efficacy of practical application of MSCs in treatment of intervertebral disc disease.

© 2003 Elsevier Science Ltd. All rights reserved.

Keywords: Intervertebral disc; Mesenchymal stem cells; Cell transplantation; Tissue engineering; Atelocollagen[®]

1. Introduction

Low back pain is a common medical and social problem in the adult population in the world today [1–3]. Intervertebral disc (IVD) degeneration is an irreversible phenomenon, which is often associated with low back pain [1–2]. Multiple elements contribute to disc

degeneration including genetic factors, environmental factors and aging [2]. Chronic weight loading, which occurs in the context of heavy work, intense physical activity, and obesity, are conditions, which are found to accelerate the process of disc degeneration [2,3]. Moreover, degeneration secondary to certain corrective surgical approaches such as long fusion, posterior herniotomy, and percutaneous nucleotomy or laser disc decompression sometimes elicit lumbago and neurological dysfunction [3–5]. Recent experimental studies focusing on decelerating disc degeneration have provided insight into the future management of degenerative disc diseases [3,6–12]. However, adequate methods

*Corresponding author. Department of Orthopaedic Surgery, Functional Reconstruction, Tokai University School of Medicine, Bohseidai, Isehara, Kanagawa, 259-1193, Japan. Tel.: +81-463-93-1121; fax: +81-463-96-4404.

E-mail address: daisakai@is.icc.u-tokai.ac.jp (D. Sakai).

for broad clinical application have not yet been available.

The etiology and pathophysiology of disc degeneration is still unknown [2,3]. However, disc degeneration can be described clinically as a loss of proper stability and mobility, which are the two major roles of the disc [13,14]. Morphologically and histopathologically, disc degeneration can be described as a decrease in water content associated with proteoglycan diminution of the nucleus pulposus and inner annulus. This effect results in destruction of annular structure and flattening of the disc [15–17]. Disc tissues have a limited ability to regenerate, since they are avascular and nutritionally supported only by passive diffusion at the end plates [3,18,19]. Consequently, once the degenerative process is activated, it is difficult to decelerate, and is ultimately considered to be an irreversible condition [3].

Nishimura and Mochida, coworkers in this study, developed an innovative procedure in which autogenic nucleus pulposus is reinserted into a degenerated disc, resulting in retardation of degeneration [6]. Okuma advanced the efficacy of this procedure by activating the biochemical properties of reinserted nucleus pulposus cells using a nucleus pulposus/annulus fibrosus coculture method, since cell cultures of nucleus pulposus cells alone expresses poor viability [7]. From these studies, we established the concept of utilizing the contained structure of the disc for cell transplantation in attempt to treat disc degeneration.

To date, *in vitro* studies of the IVD cells have shown that most IVD cells have the capacity to synthesize proteoglycan epitopes similar to chondrocytes and cellular morphology expressing chondrocyte-like behavior [20–25]. However, notochordal cells, which are only found in the nucleus pulposus of animals or in humans of young age, do not resemble chondrocyte-like behavior, since they are said to be the remnants of the notochord [26]. From these facts, it can be assumed that the majority of disc cells originate from the mesenchyme.

Mesenchymal stem cells (MSCs) are undifferentiated cells found in small numbers in the periosteum or in the bone marrow. They possess the unique ability to differentiate into varieties of cells found in connective tissues of mesenchymal origin [27]. Their ability to differentiate is governed mainly by the environment in which they are placed. Conditions favorable for inducing osteogenesis, chondrogenesis and adipogenesis have been well established [28,29]. Recent studies of MSCs have reported successful applications of MSC transplantation in various fields [30–33]. Several studies have reported repair of articular cartilage defects by transplanting MSCs embedded in collagen gels or gelatin sponge [31–33].

Despite the fact that IVD cells possess chondrocyte-like phenotypes as mentioned previously, the association

between disc cells and MSCs has never been addressed in the literature. In this study, we describe the first application of MSCs in IVD experimental model by transplantation of autologous MSCs to the degenerated disc.

Atelocollagen[®] gel was chosen as a cell carrier because of its immunogenic and safety advantages over other carriers. In Atelocollagen[®], the antigenic telopeptide region is removed by pepsin digestion and differential salt precipitation during purification. This immunogenic advantage has enabled Atelocollagen[®] to be used in various clinical settings. It is most widely used as a wrinkle eraser by injecting Atelocollagen[®] into the skin. Atelocollagen[®] is also used as a scaffold for keratinocytes and fibroblasts in artificial skin. Basic science research on culturing chondrocytes in Atelocollagen[®] gel has shown that cultures using Atelocollagen[®] result in greater matrix synthesis when compared to cells cultured in monolayer [34,35].

2. Materials and methods

2.1. Mesenchymal stem cell isolation and culture

After approval by the Animal Experimentation Committee at the author's institution, 15 Japanese white rabbits weighing an average of 1.3 kg were divided equally into five groups (normal controls; degeneration-induced models; MSC transplantation models evaluated at 2 weeks; 4 weeks; 8 weeks). Under inhalation anesthesia with 2.5% isoflurane (Abbott Laboratories, North Chicago, IL, USA), 5 ml of bone marrow blood was aspirated from the iliac crests of MSC transplantation models into 3000 units of heparin using 18-gauge needles and 10 ml syringes. Collected marrow was added to 0.7 ml Ficoll-Hypaque (Sigma, St Louis, MI, USA) and centrifuged at 150 times gravity for 15 min. The cells obtained, following buffy coat isolation and RBC destruction by RBC lysis buffer (Sigma), were washed three times with phosphate-buffered saline (PBS). Mono-nucleated cells were then cultured with 25 cm² culture flask in low-glucose Dulbecco's modified eagles medium (DMEM, Gibco, Green Island, NY, USA) containing 10% fetal bovine serum (Gibco) and antibiotics (penicillin G, 100 U/ml; streptomycin, 0.1 mg/ml; amphotericin B, 0.25 µg/ml) at 37°C, 5% CO₂. Medium was changed every 48 h with removal of non-adherent cells and cultured 12–15 days until reaching subconfluency.

2.2. Labeling of MSCs with recombinant adenovirus vector expressing *E. coli lacZ* Gene (*Ad-lacZ*)

To confirm the viability of MSCs after transplantation, the cultured MSCs were infected with Ad-lacZ

expressing *E. coli lacZ* gene under the control of CAG promoter consisting of cytomegalovirus IE enhancer, chicken beta-actin promoter, and rabbit beta-globulin polyadenylation signal (kindly provided by Dr. Izumu Saito, Tokyo University, Tokyo, Japan) [36,37]. The MSCs were infected overnight with 1×10^4 PFU/ml of Ad-lacZ at multiplicity of infection (MOI) 1, 5, 10 and 25. To examine the potential of Ad-lacZ infection, 2×10^4 infected MSCs were sampled and stained with 0.1% 5-bromo-4-chloro-3-indolyl-B-D-galactopyranoside (X-gal; Takara, Ohtsu, Shiga, Japan) following PBS wash. Nearly 100% of the cells were stained by infection with Ad-lacZ at $\text{MOI} \geq 5$.

2.3. Confirmation of multilineage differentiation in the MSCs

Osteogenic, chondrogenic and adipogenic induction were assessed to prove multilineage differentiation in MSCs used for this study. Osteogenic induction was performed in monolayered MSCs grown in medium containing 100 nM dexamethasone (Sigma), 10 mM beta-glycerol phosphate (Sigma), and 50 μM ascorbic acid 2-phosphate (Wako Pure Chemical Industries, Ltd., Osaka, Japan). Alkaline phosphatase activity and calcium deposition shown by cresolphthalein complexone (Sigma) staining were evaluated.

In chondrogenic induction, MSCs were pelleted by 1500 times gravity for 5 min centrifugation, and cultured in induction medium containing high-glucose DMEM, 1% ITS+ (Beckton Dickinson, Franklin Lakes, NJ, USA), 100 nM dexamethasone, 50 μM ascorbic acid 2-phosphate, 35 $\mu\text{g}/\text{ml}$ proline (Sigma) and 5 ng/ml TGF-beta1 (Sigma). The pellet was cultured for 21 days, then cut into frozen sections, and evaluated by histology for Safranin O and immunohistochemistry for type II collagen and proteoglycan.

Adipogenesis was induced by an induction using a medium containing high-glucose DMEM, 100 nM dexamethasone, 200 μM indomethacin (Wako), 10 $\mu\text{g}/\text{ml}$ insulin (Sigma), 500 μM 3-isobutyl-1-methylxanthin (Wako). Induction was confirmed by oil red staining.

2.4. Embedding of MSCs into Atelocollagen[®] gel

Since type II collagen is the main collagen component of IVD, Atelocollagen[®] gel solution (KOKENCELL-GEN; Koken Co., Ltd., Tokyo, Japan) composed of 0.3% type II collagen (pH 3.0) was used as a carrier. Atelocollagen[®] solution is in liquid form when cooled to 4°C but gelatinizes as the temperature rises to 37°C. Eight milliliter of cooled Atelocollagen[®] solution was quickly prepared for cell culture by adding 1 ml of $10 \times$ concentrated low-glucose DMEM, 0.1 ml HEPES (pH 7.4) (Sigma) and 0.1 ml 2.2% NaHCO_3 (Sigma). All MSCs were infected with Ad-lacZ and transplanted no

later than 21 days after isolation. Gene labeled MSCs were embedded in this solution with a final cell density of 1×10^6 cells/ml. Viability of MSCs checked after embedding showed that more than 95% were viable.

2.5. Disc degeneration induction

Disc degeneration in rabbits was performed under inhalation anesthesia to degeneration-induced and MSC transplantation models by aspirating the nucleus pulposus from L2-L3, L3-L4, L4-L5 IVDs through an anterolateral approach using a 21-gauge needle and a 10 ml syringe, as described by Okuma and Nomura [7,8]. Aspirated disc fragments were carefully examined under dissecting microscopy to confirm that only nucleus pulposus was aspirated. The weight of nucleus pulposus inside the needle ranged from 0.005 to 0.008 g.

2.6. Autologous transplantation of MSCs in Atelocollagen[®]

Autogenous MSCs embedded in Atelocollagen[®] were transplanted to degeneration-induced L2-L3, L3-L4, L4-L5 discs in anesthetized MSC transplantation models at 2 weeks after second operation in the following manner. Atelocollagen[®] gel-medium solution (0.04 ml), in which autogenous MSCs were embedded, was injected through a 27-gauge insulin injector to each discs with careful manipulation.

2.7. Evaluations

2.7.1. Histology

At 2, 4, and 8 weeks after the operation, five rabbits including one degeneration-induced model and one normal control were euthanized by intravenous injection of high-dose pentobarbital sodium (120 mg/kg) (Abbott). The L2-L3, L3-L4, L4-L5 discs were excised immediately in continuity with the cranial and caudal vertebral bodies. Femurs with thigh muscles were also harvested. The vertebral body–disc–vertebral body units were then fixed in 10% formalin neutral buffer solution (Wako), decalcified in Plant-Rychlo solution (Decalcifying Solution A; Wako), dehydrated in a graded series of ethanol (70%, 90%, 99%, Wako), and processed individually with paraffin wax embedding. The paraffin blocks were sectioned longitudinally using a microtome into 4- μm sections. The sections were stained with hematoxylin and eosin, and Safranin O for evaluation. Grading system for disc degeneration by Nishimura and Mochida [6] focusing on the morphological changes in the annular structure was used for evaluation: Grade 0, normal structure; Grade 1, mildly serpentine appearance of the annulus fibrosus with rupture; Grade 2, moderately serpentine appearance of the annulus fibrosus with rupture; Grade 3, severely serpentine appearance of the

annulus fibrosus with mildly reversed contour; Grade 4, severely reversed contour; and Grade 5, indistinct.

2.7.2. Immunohistochemistry

Proteoglycan accumulation changes in the discs were studied to evaluate disc degeneration. The sections were labeled overnight at 4°C with anti-rabbit proteoglycan monoclonal antibody (Chemicon, Temecula, CA, USA), prepared at a dilution of 1:200 in PBS. This antibody recognizes short peptides substituted with keratan sulfate side chains and core proteins of proteoglycan in the articular cartilage of mouse. The samples were washed with PBS three times and reacted with anti-mouse HRP (Dako A/S, Profuktionsvej, Denmark) at a dilution of 1:100 in PBS for 30 min at 4°C. Finally, the sections were counter-stained with hematoxylin for histologic examination. Cartilage of the femoral head served as a positive control and muscle as a negative control.

2.7.3. Confocal laser microscopy

In order to visualize changes in proteoglycan accumulation more clearly, the sections labeled with proteoglycan antibodies were stained with anti-goat Alexa fluor[®] 488 (Molecular Probes, Eugene, OR, USA) a fluorescent second antibody, and evaluated under confocal laser scanning microscopy. Statistical analyses were performed between the groups by Wilcoxon's paired signed rank test using Stat View[™] software. Significance in all cases were set at $p < 0.05$.

2.7.4. Viability of transplanted MSCs

The sections were labeled with monoclonal anti beta-galactosidase antibody (GAL-40, Sigma) and stained with Alexa fluor[®] 488 using the same technique. Viability of the transplanted cells in the discs was examined under fluorescent microscopy.

3. Results

3.1. Confirmation of multilineage differentiation in rabbit MSCs

All of the inductions performed on the obtained MSCs were successful, confirming multilineage differentiation of obtained MSCs (Fig. 1). MSCs grown in osteogenic induction medium demonstrated mineralized bone matrix after 10 days. Alkaline phosphatase was positively stained and cresolphthalein complexone staining identified calcium depositions. Immunohistochemistry demonstrated positive staining for type I collagen. In chondrogenic induction, MSCs cultured in pellet showed slight growth over a 21-day period. Frozen sections of the pellet demonstrated cartilage-like matrix, which were stained positive for Safranin O, type II

collagen and proteoglycan. MSCs grown in adipogenic induction medium started to show fat deposits after 7 days of culture. These deposits were stained positive by oil red staining.

3.2. Histological analysis

In normal control discs, nucleus pulposus and inner annulus with no major structural changes were seen after 2, 4, and 8 weeks; grade 0 (Fig. 2). The extra cellular matrix of nucleus pulposus was rich in gelatinous structure clearly distinguishable from the inner annulus with cells containing cytoplasm with large vacuoles.

In degeneration-induced models, progressive degenerative changes in both nucleus pulposus and inner annulus fibrosus were apparent over time. Two weeks after operation, the number of vacuolated cells in the nucleus pulposus had markedly decreased with mild rupture of the annulus; grade 1 (Fig. 3a). At 4 weeks, the curved structure of inner annulus fibrosus had collapsed; grade 2–3, with appearance of cells with large nucleus around the nucleus-annulus boarder (Fig. 3b). At 8 weeks post-operation, nucleus pulposus cell were rarely observed and lamellae collapse of the inner annulus had accelerated with invasion of proliferative connective tissue; grade 4–5 (Fig. 3c). Disc height flattening began to appear at this point and Safranin O staining became less intense.

In models receiving MSC transplantation, degenerative changes such as collapse of annular structure and invasion of proliferative connective tissue was normal to minimal through out all time periods; grade 0–1 (Fig. 4a–c). Instead, newly formed tissue had replaced nucleus pulposus cells in areas where it had disappeared (Fig. 4d). These tissues were formed mostly by spindle like cells with fairly loose extra cellular matrix in longitudinal layers with high intensity staining of Safranin O (Fig. 4e). These discs did not show apparent flattening. There was no obvious difference among disc levels in overall grading results in all groups. No apparent changes were noted in the adjacent non-operated sites in all groups.

3.3. Immunohistochemical analysis

Positive staining for proteoglycan was strongly identified in the extra cellular matrix of nucleus pulposus and inner annulus fibrosus in normal controls (Fig. 5a). In the degeneration-induced models, the staining intensity decreased or did not show significant intensity in areas where nucleus pulposus cells were depleted, or in areas where proliferative connective tissues had invaded according to the time course (Fig. 5b). Extra cellular matrix of tissue composed of transplanted cells showed




Fig. 1. Confirmation of multilineage differentiation of rabbit MSCs used in the study. (a) Rabbit MSCs cultured for 15 days after isolation. Magnification $4\times$. (b) MSCs cultured for 14 days in osteogenic induction medium. Mineralized bone matrix formation is observed with positive histological staining for alkaline phosphatase and calcium deposition shown by cresolphthalein complexone staining. Magnification $4\times$. (c) Immunohistochemistry of type II collagen showing positive staining in frozen section of MSCs after 21 days of pellet culture by chondrogenic induction. Magnification $10\times$. (d) MSCs cultured for 10 days in adipogenic induction medium showing oil red positive staining. Magnification $10\times$.




Fig. 2. Disc of a normal control group in week 8 showing no apparent abnormality; annulus degeneration grade 0. Magnification $4\times$.

a high staining intensity in the extra cellular matrix of the spindle like cells (Fig. 5c).

3.4. Proteoglycan accumulations evaluated by confocal laser scanning microscopy

In evaluation of sections using confocal laser scanning microscopy, results were compatible with that of the non-fluorescent staining but clearer (Fig. 5). Results were significantly different ($p < 0.005$) in average peak intensity levels of intensity measured in the center of the disc at 8 weeks post-operation were 3.3 ± 0.5 in normal controls, 2.8 ± 0.4 in MSC-transplanted models and 1.1 ± 0.3 in degeneration-induced models.

3.5. Viability of the transplanted cells

X-gal staining of the MSCs done prior to embedding into Atelocollagen[®] gel showed good positive staining




Fig. 3. (a) Week 2 degeneration-induced model. Annular structure is mildly affected although majority of the nucleus pulposus is depleted; annular degeneration grade 1. Magnification $4\times$. (b) Week 4 degeneration-induced model. Annular structure is severely serpentine with few nucleus pulposus left; grade 3. Magnification $4\times$. (c) Week 8 degeneration-induced model. Most of the annular structure is collapsed with severe invasion of proliferative connective tissue; grade 4. Magnification $4\times$.

in nearly all of the cells infected at each MOI. Fluorescent immunohistochemical staining of the sections harvested from MSC-transplanted models demonstrated positive staining of beta-galactosidase in sections at 2 and 4 weeks after operation (Fig. 6). However, the number of positive cells had decreased over time.

4. Discussion

In this study, partial aspiration of nucleus pulposus tissue was used in order to induce degeneration. Disc annular lesion models and partial nucleus pulposus aspiration models are an established method for creating disc degeneration experimentally [7–9,17,19]. It is obvious from the results of histology and immunohistochemistry in the degeneration-induced models of our experiment that degeneration was successfully induced. This evidence confirms the fact that preserving viability of the nucleus pulposus and annular structure is the primary factor in decelerating disc degeneration.

The approach to delivering cells into the degenerated discs of rabbits was established by Okuma et al. [7]. In that study, decelerating disc degeneration was successful by reinserting nucleus pulposus cells that were activated by coculture with annulus fibrosus cells. Activated nucleus pulposus was shown to have proliferated inside the disc and possibly helped to maintain proteoglycan synthesis and morphological structure. However, in Okuma's experimental model, reinserted cells were suspended in medium, in spite the fact that nucleus pulposus cells retain their character, shared prominent matrix production when cultured in a three-dimensional environment. In addition, no appropriate markers were used to check the viability of the cells after reinsertion.

Chondrocytes when cultured in monolayer are known to express a fibroblastic differentiated morphology and lose their ability to synthesize matrix predominantly of type I collagen distribution [35,38]. Human IVD cells are known to possess chondrocyte phenotypes [20–25]. Like chondrocytes, they also express fibroblastic morphology in monolayer but regain their features when cultured in a three-dimensional environment [40–43]. In




Fig. 4. Week 2 (a), 4 (b) and 8 (c) MSC transplantation models. Transplanted cells are fulfilling the disc space. Magnification $4\times$. The rupture of the annular structure is mild; grade 0–1. (d) Newly formed tissue in the disc 8 weeks after MSC transplantation with rich extra cellular matrix. Magnification $100\times$. (e) Safranin O staining of the tissue. Magnification $4\times$.

studies of MSC transplantation in cartilage defects, Atelocollagen[®] gels and gelatin sponges have been used as a successful carrier for cell delivery [39]. Based upon these elements, Type II collagen-based Atelocollagen[®] gel was used as a cell carrier, making it close to disc component. The type II Atelocollagen solution used in our study was prepared from bovine cartilage, which was decomposed by pepsin digestion with incubation and isolated by differential salt precipitation into acidic solution. This solution is in liquid form when cooled to temperatures near 4°C but gelatinizes when heated to about 37°C. In order to use in cell cultures, Atelocollagen[®] solution was neutralized by adding HEPES buffer (pH 7.4) and NaHCO₃.

The results of cell delivery in our transplant models shared successful construction of tissues without any

defects. Proliferation of cells after transplantation was found to be successful with minimal disorientation of annular structure. Using beta -glycosidase staining, it was confirmed that the transplanted MSCs were viable, and it can be assumed that cells in the tissue of transplanted models had differentiated from these MSCs. It can be speculated that expression of beta -glycosidase lasted only 4 weeks at the most, because delivery of genes by recombinant adenovirus vectors are not considered permanent. However, use of Ad-lacZ for marking was considered appropriate in this study due to success in confirming viability and ease of handling.

Histologically, MSC-transplanted models had preserved disc structure with minimal degeneration in all time periods compared to degeneration-induced models. We found that primary morphological features of disc




Fig. 5. (a) Proteoglycan staining of a normal disc tissue. Magnification 20 \times . Staining intensity by confocal laser scanning microscopy is shown below. (b) Week 8 degeneration-induced model. Staining intensity of proteoglycan is low. Magnification 20 \times . (c) Week 8 MSC transplantation model shows high intensity peak levels similar to normal control group. Magnification 20 \times .




Fig. 5 (continued).

degeneration, cell depletion in the nucleus pulposus and disorientation of oval annular structure were prevented by injection of Atelocollagen[®] gel embedded with MSCs. This unique growth-enhancing process served to maintain cell volume with high viability and matrix gel formation. Surprisingly, transplanted MSCs survived in the environment of the disc, differentiating into spindle-shaped cells arranged in longitudinal layers, which resembled original disc cells. Extra cellular matrix, which was suspected to originate from the MSCs, was observed in interlineating layers. Safranin O stained with markedly high intensity in this matrix and immunohistochemical staining for proteoglycan also showed high intensity, suggesting the potential for rich matrix production, including proteoglycan that these differentiated cells possessed. Proteoglycan staining intensity levels with confocal laser scanning microscopy demonstrated that the intensity patterns of the MSC-



Fig. 6. Cells in disc tissue of a week 4 MSC-transplanted model shows positive staining for beta-galactosidase, which confirms the survival of MSCs at harvest. Magnification $63\times$.

transplanted models were significantly higher than the degeneration-induced models ($p < 0.005$) and similar to controls at all time points. This proves that deceleration of degeneration and maintaining proteoglycan preservation was possible.

We conclude that in attempt to decelerate IVD degeneration, MSC transplantation was effective in preserving annular structure by the use of MSC/Atelocollagen[®] conjugates. These amalgamates served to fill depleted nucleus pulposus, and preventing proteoglycan decrease by increasing production from the differentiated cells of transplanted MSCs. Results from our study provide initial evidence for the potential of MSCs to differentiate into IVD cells, which provides new information in MSC research. However, further long-term studies with detailed analysis of the newly formed tissue by transplanted MSCs will be needed in the future.

Acknowledgements

Authors thank Dr. Adam Tucker for review of the manuscript, Joubu Itoh, Akihiko Serizawa and staffs at Laboratories for Structure and Function Research, Tokai University School of Medicine for expert technical assistance. This work was supported in part

by Grant-in-Aid for Scientific Research grant number 14571406 and a Grant of The Science Frontier Program from the Ministry of Education, Culture, Sports, Science and Technology of Japan, a grant from Japan Orthopaedic and Traumatology Foundation grant number 0139 and a grant from 2002 Tokai University School of Medicine Research Aid.

References

- [1] Kraemer J. Natural course and prognosis of intervertebral disc diseases. *Spine* 1995;20:635–9.
- [2] Buckwalter JA. Aging and degeneration of the human intervertebral disc. *Spine* 1995;20:1307–14.
- [3] Nishida K, Kang JD, Gilbertson LG, Moon S-H, Suh J-K, Vogt MT, Robbins PD, Evans CH. 1999 Volvo Award in basic science: modulation of the biologic activity of the rabbit intervertebral disc by gene therapy: an in vivo study of adenovirus-mediated transfer of the human transforming growth factor beta1 encoding gene. *Spine* 1999;24:2419–25.
- [4] Mochida J, Nishimura K, Nomura T, Toh E, Chiba M. The importance of preserving disc structure in surgical approaches to lumbar disc herniation. *Spine* 1996;21:1556–64.
- [5] Lee CK. Accelerated degeneration of the segment adjacent to a lumbar fusion. *Spine* 1988;13:375–7.
- [6] Nishimura K, Mochida J. Percutaneous reinsertion of the nucleus pulposus: an experimental study. *Spine* 1998;23:1531–638.
- [7] Okuma M, Mochida J, Nishimura K, Sakabe K, Seiki K. Reinsertion of stimulated nucleus pulposus cells retards intervertebral disc degeneration: an in vitro and in vivo experimental study. *J Orthop Res* 2000;18:988–97.
- [8] Nomura T, Mochida J, Okuma M, Nishimura K, Sakabe K. Nucleus pulposus allograft retards intervertebral disc degeneration. *Clin Orthop* 2001;389:94–101.
- [9] Gruber HE, Johnson TL, Leslie K, Ingram JA, Martin D, Hoelscher G, Banks D, Phieffer L, Coldham G, Hanley Jr EN. Autologous intervertebral disc cell implantation: a model using *Psamomys obesus*, the sand rat. *Spine* 2002;27:1626–33.
- [10] Nishida K, Kang JD, Suh JK, Robbins PD, Evans CH, Gilbertson LG. Adenovirus-mediated gene transfer to nucleus pulposus cells: implications for the treatment of intervertebral disc degeneration. *Spine* 1998;23:2437–43.
- [11] Nishida K, Gilbertson LG, Robbins PD, Evans CH, Kang JD. Potential applications of gene therapy to the treatment of intervertebral disc disorders. *Clin Orthop* 2000;379S:S234.
- [12] Takegami K, Thonar EJ, An HS, Kamada H, Masuda K. Osteogenic Protein-1 enhances matrix replenishment by intervertebral disc cells previously exposed to interleukin-1. *Spine* 2002;27:1318–25.
- [13] Frick SL, Hanley EN, Meyer RA, Ramp WK, Chapman TM. Lumbar intervertebral disc transfer. *Spine* 1994;19:1826–35.
- [14] Bao Q-B, McCullen GM, Higham PA, Dumbleton JH, Yuan HA. The artificial disc: theory, design and materials. *Biomaterials* 1996;17:1157–67.
- [15] Hirsch C, Schajowicz F. Studies on structural changes in the lumbar annulus fibrosus. *Acta Orthop Scand* 1953;22:184–231.
- [16] Osti OL, Vernon-Roberts B, Moore R, Fraser RD. Annular tears and disc degeneration in the lumbar spine. *J Bone Joint Surg* 1992;74-B:678–82.
- [17] Melrose J, Roberts S, Smith S, Menage J, Ghosh P. Increased nerve and blood vessel ingrowth associated with proteoglycan depletion in an ovine annular lesion model of experimental disc degeneration. *Spine* 2002;27:1278–85.
- [18] Bradford DS, Cooper KM, Oegema Jr TR. Chymopapain, chemonucleolysis, and nucleus pulposus regeneration. *J Bone Joint Surg* 1983;65:1220–31.
- [19] Lipson SJ, Muir H. 1980 Volvo Award in basic science: proteoglycans in experimental intervertebral disc degeneration. *Spine* 1981;6:194–210.
- [20] Horner HA, Urban JPG. 2000 Volvo Award in basic science: effect of nutrient supply on the viability of cells from the nucleus pulposus of the intervertebral disc. *Spine* 2001;26:2543–9.
- [21] Horner HA, Roberts S, Bielby RC, Menage J, Evans H, Urban JPG. Cells from different regions of the intervertebral disc. *Spine* 2002;27:1018–28.
- [22] Gruber HE, Hanley Jr EN. Human disc cells in monolayer vs 3D culture: cell shape, division and matrix formation. *BMC Musculoskeletal Disorders* 2000;1(1):1.
- [23] Antoniou J, Steffen T, Nelson F, Winterbottom N, Hollander AP, Poole RA, Aebi M, Alini M. Human lumbar intervertebral disc: evidence for changes in the biosynthesis and denaturation of the extracellular matrix with growth, maturation, ageing, and degeneration. *J Clin Invest* 1996;98(4):996–1003.
- [24] Desai BJ, Gruber HE, Hanley Jr EN. The influence of Matrigel or growth factor reduced Matrigel on human intervertebral disc cell growth and proliferation. *Histol Histopathol* 1999;14:359–68.
- [25] Gruber HE, Stasky AA, Hanley Jr EN. Characterization and phenotypic stability of human disc cells in vitro. *Matrix Biol* 1997;16:285–8.
- [26] Aguiar DJ, Johnson SL, Oegema TR. Notochordal cells interact with nucleus pulposus cells: regulation of proteoglycan synthesis. *Exp Cell Res* 1999;246:129–37.
- [27] Caplan AI. Mesenchymal stem cells. *J Orthop Res* 1991;9:641–50.
- [28] Caplan AI, Elyanderani M, Mochizuki Y, Wakitani S, Goldberg VM. Principles of cartilage repair and regeneration. *Clin Orthop Rel Res* 1997;342:254–69.
- [29] Haynesworth SE, Goshima J, Goldberg VM, Caplan AI. Characterization of cells with osteogenic potential from human marrow. *Bone* 1992;13:81–8.
- [30] Haynesworth SE, Reuben D, Caplan AI. Cell-based tissue engineering therapies: the influence of whole body physiology. *Adv Drug Del Rev* 1998;33:3–14.
- [31] Quintavalla J, Uziel-Fusi S, Yin J, Boehnlein E, Pastor G, Blancuzzi V, Singh HN, Kraus KH, O'Byrne E, Pellas TC. Fluorescently labeled mesenchymal stem cells (MSCs) maintain multilineage potential and can be detected following implantation into articular cartilage defects. *Biomaterials* 2002;23:109–19.
- [32] Wakitani S, Goto T, Pineda SJ, Young RG, Mansour JM, Caplan AI, Goldberg VM. Mesenchymal cell-based repair of large, full-thickness defects of articular cartilage. *J Bone Joint Surg* 1994;76-A:579–92.
- [33] Im G-I, Kim D-Y, Shin J-H, Hyun C-W, Cho W-H. Repair of cartilage defect in the rabbit with cultured mesenchymal stem cells from bone marrow. *J Bone Joint Surg* 2001;83-B:289–94.
- [34] Nishikiori T, Ochi M, Uchio Y, Maniwa S, Kataoka H, Kawasaki K, Katsube K, Kuriwaka M. Effects of low-intensity pulsed ultrasound on proliferation and chondroitin sulfate synthesis of cultured chondrocytes embedded in Atelocollagen gel. *J Biomed Mater Res* 2001;59:201–6.
- [35] Uchio Y, Ochi M, Matsusaki M, Kurioka H, Katsube K. Human chondrocyte proliferation and matrix synthesis cultured in Atelocollagen gel. *J Biomed Mater Res* 2000;50:138–43.
- [36] Kanegae Y, Makimura M, Saito I. A simple and efficient method for purification of infectious recombinant adenovirus. *Jpn J Med Sci Biol* 1995;47:157–66.
- [37] Tsubota K, Inoue H, Ando K, Ono M, Yoshino K, Saito I. Adenovirus-mediated gene transfer to the ocular surface epithelium. *Exp Eye Res* 1998;67:531–8.

- [38] Kimura T, Yasui N, Ohsawa S, Ono K. Chondrocytes embedded in collagen gels maintain cartilage phenotype during long-term cultures. *Clin Orthop* 1984;186:231–40.
- [39] Wakitani S, Kimura T, Hirooka A, Ochi T, Yoneda M, Yasui N, Owaki H, Ono K. Repair of rabbit articular surfaces with allograft chondrocytes embedded in collagen gel. *J Bone Joint Surg* 1989;71:74–80.
- [40] Gruber HE, Fisher Jr EC, Desai B, Stasky AA, Hoelscher G, Hanley Jr EN. Human intervertebral disc cells from the annulus: three-dimensional culture in agarose or alginate and responsiveness to TGF-beta 1. *Exp Cell Res* 1997;235:13–21.
- [41] Lee JY, Hall R, Pelinkovic D, Cassinelli E, Usas A, Gilbertson L, Huard J, Kang J. New use of a three-dimensional pellet culture system for human intervertebral disc cells. *Spine* 2001;26:2316–22.
- [42] Chiba K, Andersson GB, Masuda K, Momohara S, Williams JM, Thonar EJ. A new culture system to study the metabolism of the intervertebral disc in vitro. *Spine* 1998;23:1821–7.
- [43] Chiba K, Andersson GB, Masuda K, Thonar EJ. Metabolism of the extracellular matrix formed by intervertebral disc cells cultured in alginate. *Spine* 1997;15:2885–93.