

the differences between metaplastic and normal cells, even though the histological appearance is similar.

Acknowledgements

We thank Masayoshi Takatani and Katsunari Ogawa for excellent technical assistance and advice. We thank the Analysis Center of Life Science, Hiroshima University for the use of their facilities. This work was supported, in part, by Grants-in-Aid for Cancer Research from the Ministry of Education, Culture, Science, Sports, and Technology of Japan, and from the Ministry of Health, Labour, and Welfare of Japan.

References

- Buckhaults P, Rago C, St Croix B, Romans KE, Saha S, Zhang L, et al. Secreted and cell surface genes expressed in benign and malignant colorectal tumors. *Cancer Res* 2001;61:6996–7001.
- Yasui W, Oue N, Ito R, Kuraoka K, Nakayama H. Search for new biomarkers of gastric cancer through serial analysis of gene expression and its clinical implications. *Cancer Sci* 2004;95:385–392.
- Oue N, Hamai Y, Mitani Y, Matsumura S, Oshimo Y, Aung PP, et al. Gene expression profile of gastric carcinoma: identification of genes and tags potentially involved in invasion, metastasis, and carcinogenesis by serial analysis of gene expression. *Cancer Res* 2004;64:2397–2405.
- Hartupée JC, Zhang H, Bonaldo MF, Soares MB, Dieckgraefe BK. Isolation and characterization of a cDNA encoding a novel member of the human regenerating protein family: Reg IV. *Biochim Biophys Acta* 2001;1518:287–293.
- Violette S, Festor E, Pandrea-Vasile I, Mitchell V, Adida C, Dussaulx E, et al. Reg IV, a new member of the regenerating gene family, is overexpressed in colorectal carcinomas. *Int J Cancer* 2003;103:185–193.
- Zhang Y, Lai M, Lv B, Gu X, Wang H, Zhu Y, et al. Overexpression of Reg IV in colorectal adenoma. *Cancer Lett* 2003;200:69–76.
- Kamarainen M, Heiskala K, Knuutila S, Heiskala M, Winqvist O, Andersson LC. RELP, a novel human REG-like protein with up-regulated expression in inflammatory and metaplastic gastrointestinal mucosa. *Am J Pathol* 2003;163:11–20.
- Kuraoka K, Taniyama K, Fujitaka T, Nakatsuka H, Nakayama H, Yasui W. Small cell carcinoma of the extrahepatic bile duct: case report and immunohistochemical analysis. *Pathol Int* 2003;53:887–891.
- Sobin LH, Wittekind CH (eds). *TNM Classification of Malignant Tumors* (5th edn). Wiley-Liss: New York, 1997; 59–62.
- Ochiai A, Yasui W, Tahara E. Growth-promoting effect of gastrin on human gastric carcinoma cell line TMK-1. *Jpn J Cancer Res* 1985;76:1064–1071.
- Yanagihara K, Seyama T, Tsumuraya M, Kamada N, Yokoro K. Establishment and characterization of human signet ring cell gastric carcinoma cell lines with amplification of the c-myc oncogene. *Cancer Res* 1991;51:381–386.
- Kondo T, Oue N, Yoshida K, Mitani Y, Naka K, Nakayama H, et al. Expression of POT1 is associated with tumor stage and telomere length in gastric carcinoma. *Cancer Res* 2004;64:523–529.
- Yasui W, Ayhan A, Kitadai Y, Nishimura K, Yokozaki H, Ito H, et al. Increased expression of p34cdc2 and its kinase activity in human gastric and colonic carcinomas. *Int J Cancer* 1993;53:36–41.
- Yasui W, Ji ZQ, Kuniyasu H, Ayhan A, Yokozaki H, Ito H, et al. Expression of transforming growth factor alpha in human tissues: immunohistochemical study and northern blot analysis. *Virchows Arch A Pathol Anat Histopathol* 1992;421:513–519.
- Mizoshita T, Tsukamoto T, Nakanishi H, Inada K, Ogasawara N, Joh T, et al. Expression of Cdx2 and the phenotype of advanced gastric cancers: relationship with prognosis. *J Cancer Res Clin Oncol* 2003;129:727–734.
- Mantel N. Evaluation of survival data and two new rank order statistics arising in its consideration. *Cancer Chemother Rep* 1966;50:163–170.
- Li A, Crimmins DL, Luo Q, Hartupée J, Landt Y, Ladenson JH, et al. Expression of a novel regenerating gene product, Reg IV, by high density fermentation in *Pichia pastoris*: production, purification, and characterization. *Protein Expr Purif* 2003;31:197–206.
- Yamamoto H, Bai YQ, Yuasa Y. Homeodomain protein CDX2 regulates goblet-specific MUC2 gene expression. *Biochem Biophys Res Commun* 2003;300:813–818.
- Mallo GV, Rechreche H, Frigerio JM, Rocha D, Zweibaum A, Lacasa M, et al. Molecular cloning, sequencing and expression of the mRNA encoding human Cdx1 and Cdx2 homeobox. Down-regulation of Cdx1 and Cdx2 mRNA expression during colorectal carcinogenesis. *Int J Cancer* 1997;74:35–44.
- Silberg DG, Swain GP, Suh ER, Traber PG. Cdx1 and cdx2 expression during intestinal development. *Gastroenterology* 2000;119:961–971.
- Silberg DG, Sullivan J, Kang E, Swain GP, Moffett J, Sund NJ, et al. Cdx2 ectopic expression induces gastric intestinal metaplasia in transgenic mice. *Gastroenterology* 2002;122:689–696.
- Bai YQ, Yamamoto H, Akiyama Y, Tanaka H, Takizawa T, Koike M, et al. Ectopic expression of homeodomain protein CDX2 in intestinal metaplasia and carcinomas of the stomach. *Cancer Lett* 2002;176:47–55.
- Almeida R, Silva E, Santos-Silva F, Silberg DG, Wang J, De Bolos C, et al. Expression of intestine-specific transcription factors, CDX1 and CDX2, in intestinal metaplasia and gastric carcinomas. *J Pathol* 2003;199:36–40.
- Tatematsu M, Tsukamoto T, Inada K. Stem cells and gastric cancer: role of gastric and intestinal mixed intestinal metaplasia. *Cancer Sci* 2003;94:135–141.
- Wong WM, Stamp GW, Elia G, Poulosom R, Wright NA. Proliferative populations in intestinal metaplasia: evidence of deregulation in Paneth and goblet cells, but not endocrine cells. *J Pathol* 2000;190:107–113.
- Lee CS, Perreault N, Brestelli JE, Kaestner KH. Neurogenin 3 is essential for the proper specification of gastric enteroendocrine cells and the maintenance of gastric epithelial cell identity. *Genes Dev* 2002;16:1488–1497.
- Tajima Y, Shimoda T, Nakanishi Y, Yokoyama N, Tanaka T, Shimizu K, et al. Association of gastric and intestinal phenotypic marker expression of gastric carcinomas with tumor thymidylate synthase expression and response to postoperative chemotherapy with 5-fluorouracil. *J Cancer Res Clin Oncol* 2003;129:683–690.
- Dhar DK, Udagawa J, Ishihara S, Otani H, Kinoshita Y, Takasawa S, et al. Expression of regenerating gene I in gastric adenocarcinomas: correlation with tumor differentiation status and patient survival. *Cancer* 2004;100:1130–1136.
- Vogelstein B, Kinzler KW. The multistep nature of cancer. *Trends Genet* 1993;9:138–141.

DNA Methylation of Genes Linked with Retinoid Signaling in Gastric Carcinoma

Expression of the Retinoid Acid Receptor β , Cellular Retinol-Binding Protein 1, and Tazarotene-Induced Gene 1 Genes Is Associated with DNA Methylation

Mariko Shutoh, M.D.^{1,2}

Naohide Oue, M.D., Ph.D.¹

Phyu Phyu Aung, M.B.B.S.¹

Tsuyoshi Nōguchi, M.D., Ph.D.²

Kazuya Kuraoka, M.D., Ph.D.¹

Hirofumi Nakayama, M.D., Ph.D.¹

Katsunobu Kawahara, M.D., Ph.D.²

Wataru Yasui, M.D., Ph.D.¹

¹ Department of Molecular Pathology, Hiroshima University Graduate School of Biomedical Sciences, Hiroshima, Japan.

² Department of Oncological Science (Surgery II), Faculty of Medicine, Oita University, Oita, Japan.

Supported in part by Grants-in-Aid for Cancer Research from the Ministry of Education, Culture, Science, Sports, and Technology of Japan and from the Ministry of Health, Labor, and Welfare of Japan.

The authors thank Masayoshi Takatani for excellent technical assistance and advice. This work was performed with the kind cooperation of the Research Center for Molecular Medicine, Faculty of Medicine, Hiroshima University. The authors also thank the Analysis Center of Life Science, Hiroshima University, for the use of their facilities.

Address for reprints: Wataru Yasui, M.D., Ph.D., Department of Molecular Pathology, Hiroshima University Graduate School of Biomedical Sciences, 1-2-3 Kasumi, Minami-ku, Hiroshima, 734-8551, Japan; Fax: (011) 81 822575149; E-mail: wyasui@hiroshima-u.ac.jp

Received January 27, 2005; revision received May 7, 2005; accepted May 31, 2005.

BACKGROUND. Hypermethylation of CpG islands has been associated with silencing of various tumor suppressor genes, and the retinoid acid receptor β (*RAR β*), cellular retinol-binding protein 1 (*CRBPI*), and tazarotene-induced gene 1 (*TIG1*) genes have been associated with retinoic acid signaling. To the authors' knowledge, little is known regarding the involvement of these three genes in gastric carcinoma (GC). In this study, the authors investigated the methylation status of these genes and analyzed the role of their DNA methylation in GC.

METHODS. DNA methylation of 3 retinoic acid-associated genes was analyzed in 42 samples of GC from 42 patients and in 8 GC cell lines by methylation-specific polymerase chain reaction (PCR) analysis. The mRNA expression levels for these three genes were measured by quantitative reverse transcription-PCR.

RESULTS. In 7 of 8 GC cell lines, the *CRBPI* gene was hypermethylated, and *CRBPI* transcription was inactive. In 6 of 8 GC cell lines, the *TIG1* gene was hypermethylated, and *TIG1* transcription was inactive. Treatment with demethylating agent 5-aza-2'-deoxycytidine restored both *CRBPI* and *TIG1* transcription. DNA methylation of the *RAR β* , *CRBPI*, and *TIG1* genes was detected in 15 of 42 GC samples (36%), 14 of 42 GC samples (33%), and 4 of 42 GC samples (10%), respectively, and in 6 of 30 samples (20%), 0 of 30 samples (0%), and 1 of 30 samples (3%) of corresponding nonneoplastic mucosa. None of the 10 normal gastric mucosa samples from young, healthy individuals demonstrated hypermethylation of any of these genes. DNA methylation of each gene was associated significantly with low mRNA expression of the respective gene. Twenty-four of 42 GC samples (57%) demonstrated hypermethylation of at least 1 of the 3 genes. However, no significant, concordant hypermethylation of these genes was observed.

CONCLUSIONS. The results suggested that gastric carcinogenesis involves transcriptional inactivation by aberrant DNA methylation of genes related to retinoid signaling. *Cancer* 2005;104:1609–19. © 2005 American Cancer Society.

KEYWORDS: DNA methylation, gastric carcinoma, retinoic acid, cellular retinol-binding protein 1, tazarotene-induced gene 1.

Various genetic and epigenetic alterations are associated with gastric carcinoma (GC).^{1,2} Epigenetic alterations, such as DNA methylation of CpG islands, are common in human malignancies. Hypermethylation of CpG islands is associated with the silencing of several genes and has been proposed as a mechanism for inactivation of tumor suppressor genes in carcinoma.^{3,4} The identification of methylated genes may be useful in the diagnosis and treatment of carcinoma and may provide insight into gastric carcinogenesis. It has been demonstrated that DNA hypermethylation is a crucial mechanism for

transcriptional silencing of tumor suppressor genes and tumor-related genes in GC⁵⁻⁸ and that DNA methylation of tumor suppressor genes occurs in early-stage GC and accumulates along the multiple pathways of gastric carcinogenesis.⁹

A frequently hypermethylated gene, not only in GC but also in breast and lung carcinomas, is retinoic acid receptor β (*RAR β*).¹⁰⁻¹² Retinoids regulate the growth, differentiation, and apoptosis of normal cells during embryonic development and of premalignant and malignant cells during carcinogenesis. The effects of retinoids are mediated predominantly by the RARs (*RAR α* , *RAR β* , and *RAR γ*), which act as retinoic acid-dependent transcriptional activators in heterodimeric association with the retinoid X receptors (*RXR α* , *RXR β* , and *RXR γ*) and in conjunction with multiple corepressors.^{13,14} RARs can induce the expression of certain genes in a ligand-dependent manner by binding to retinoic acid-responsive elements in the promoter regions.

Other key components of retinoid signaling include cellular retinol-binding proteins (CRBPs), which belong to the family of fatty acid-binding proteins. Three CRBP members have been described: *CRBP1*, which is expressed widely,¹⁵ and *CRBP2* and *CRBP3*, which demonstrate tissue-specific expression.^{16,17} Retinoic acid is present in the circulation, and most tissues rely on the uptake and cytosolic metabolism of retinoic acid to activate RARs and RXRs. CRBPs possess high-affinity binding for retinoic acid and possibly function as chaperone-like proteins to regulate the prenuclear phase of retinoic acid signaling.¹⁶ A critical role of *CRBP1* was demonstrated in *CRBP1* knock-out mice, which are susceptible to hypovitaminosis A syndrome.¹⁸ The *CRBP1* gene is hypermethylated in various human malignancies.¹⁹ Possible involvement of *CRBP2* in N-methyl-N'-nitro-N-nitrosoguanidine-induced rat stomach carcinogenesis has been reported.²⁰ However, to our knowledge, little is known regarding the DNA methylation status of *CRBP1* in GC.

Tazarotene, which is a synthetic retinoid that binds *RAR β* and *RAR γ* , is used in the treatment of psoriasis, a hyperproliferative disorder. One gene induced by tazarotene is tazarotene-induced gene 1 (*TIG1*).²¹ The *TIG1* gene also is hypermethylated in various human malignancies.^{22,23} *TIG1* may function as a cell adhesion molecule, and its expression on the cell surface may increase cell-cell contact and reduce proliferation.²⁴ Therefore, it is important to consider the potential function of *TIG1* and its role in RAR-mediated biology. *TIG1* has been identified as a potential tumor suppressor gene in human prostate carcinoma.²⁴ However, to our knowledge, DNA

hypermethylation of the *TIG1* gene in GC has not been investigated.

In the current study, we examined the methylation status of the *RAR β* , *CRBP1*, and *TIG1* genes and the expression levels of these genes in 42 primary GC samples and in corresponding samples of nonneoplastic mucosa. To determine whether hypermethylation induced transcriptional inactivation, we compared methylation status with mRNA expression. We also examined whether correlations exist between the methylation status of each of these genes.

MATERIALS AND METHODS

Cell Lines

Eight cell lines derived from human GC were used. The TMK-1 cell line was established in our laboratory from a poorly differentiated adenocarcinoma.²⁵ Five GC cell lines of the MKN series (*MKN-1*, adenocarcinoma; *MKN-7*; *MKN-28*; *MKN-74*, well differentiated adenocarcinoma; and *MKN-45*, poorly differentiated adenocarcinoma) kindly were provided by Dr. T. Suzuki. The KATO-III and HSC-39 cell lines, which were established from signet ring cell carcinomas, kindly were provided by Dr. M. Sekiguchi and Dr. K. Yanagihara, respectively.²⁶ All cell lines were maintained in RPMI 1640 (Nissui Pharmaceutical Co., Ltd., Tokyo, Japan) containing 10% fetal bovine serum (Whittaker, Walkersville, MD) in a humidified atmosphere of 5% CO₂ and 95% air at 37 °C. To analyze restoration of *CRBP1* and *TIG1* gene expression, *MKN-28* cells were incubated for 5 days with 1 μ M Aza-2'-deoxycytidine (Aza-dC; Sigma Chemical Company, St. Louis, MO).

Tissue Samples

Frozen tissue samples were collected from 42 patients with GC who underwent surgery between 1998 and 2001 at the Department of Oncological Science, Oita University Hospital (Oita, Japan). All patients underwent curative resection. GC tissue specimens from 42 patients were analyzed for methylation of the *RAR β* , *CRBP1*, and *TIG1* genes. Thirty samples of corresponding nonneoplastic mucosa, including 22 samples with intestinal metaplasia, also were analyzed. Tumors and corresponding nonneoplastic mucosa were removed surgically, immediately frozen in liquid nitrogen, and stored at -80 °C until use. We confirmed microscopically that the tumor tissue specimens consisted mainly (> 50%, on a nuclear basis) of tumor tissue and that nonneoplastic mucosa did not exhibit any tumor cell invasion or show significant inflammatory involvement. *Helicobacter pylori* status was confirmed by histology and/or ¹³C-urea breath

test. Tumor staging was performed according to TNM stage grouping.²⁷ In addition, methylation status of these genes was determined in 10 samples of normal gastric mucosa obtained endoscopically from 10 healthy, young individuals (age range, 22–35 yrs; average age, 26.4 yrs) with no clinical symptoms and no microscopic mucosal changes. Because written informed consent was not obtained, for strict privacy protection, all samples were disidentified before analyzing DNA methylation status. This procedure is in accordance with Ethical Guidelines for Human Genome/Gene Research enacted by the Japanese Government.

Genomic DNA Extraction and Methylation Analysis

To examine DNA methylation patterns of the *RARβ*, *CRBP1*, and *TIG1* genes, we extracted genomic DNAs with a genomic DNA purification kit (Promega, Madison, WI) and performed methylation-specific polymerase chain reaction (PCR) (MSP) analysis.²⁸ In brief, 2 μg of genomic DNA were denatured by treatment with 2 M NaOH and modified with 3 M sodium bisulfate for 16 hours. DNA samples were purified with Wizard DNA purification resin (Promega), treated with 3 M NaOH, precipitated with ethanol, and resuspended in 25 μL water. Two-microliter aliquots were used as templates for PCR reactions. Primer sequences and annealing temperatures for MSP were as described previously (Table 1).^{19,22,29}

Reverse Transcriptase-PCR

mRNA expression of *RARβ*, *CRBP1*, and *TIG1* was analyzed by reverse transcriptase (RT)-PCR. Total RNA was extracted with an RNeasy Mini Kit (QIAGEN, Tokyo, Japan), and 1 μg of total RNA was converted to cyclic DNA (cDNA) with a first-strand cDNA synthesis kit (Amersham Pharmacia Biotech, Uppsala, Sweden). Primer sequences and annealing temperatures were as described previously (Table 1).^{19,23} RT-PCR products were subjected to 1.5% agarose gel electrophoresis, stained with ethidium bromide, and examined under ultraviolet light. *ACTB*-specific PCR products were amplified from the same RNA samples and served as internal controls.

Quantitative RT-PCR Analysis

Total RNA was extracted with an RNeasy Mini Kit, and 1 μg of total RNA was converted to cDNA with a first-strand cDNA synthesis kit (Amersham Pharmacia Biotech). PCR was performed with the SYBR Green PCR Core Reagents kit (Applied Biosystems, Foster City, CA). Real-time detection of the emission intensity of SYBR Green bound to double-stranded DNA was performed with an ABI PRISM 7700 Se-

TABLE 1
Primer Sequences for Methylation-Specific PCR, RT-PCR, and Quantitative RT-PCR

Primer sequence	Annealing temperature
MSP (<i>RARβ</i> , methylated) F: 5'-GGT TAG TAG TTC GGG TAG GGT TTA TC-3' R: 5'-CCG AAT CCT ACC CCG ACG-3'	64 °C
MSP (<i>RARβ</i> , unmethylated) F: 5'-TTA GTA GTT TGG GTA GGG TTT ATT-3' R: 5'-CCA AAT CCT ACC CCA ACA-3'	55 °C
MSP (<i>CRBP1</i> , methylated) F: 5'-TTG GGA ATT TAG TTG TCG TCG TTT C-3' R: 5'-AAA CAA CGA CTA CCG ATA CTA CGG G-3'	70 °C
MSP (<i>CRBP1</i> , unmethylated) F: 5'-GTG TTG GGA ATT TAG TTG TTG TTT-3' R: 5'-ACT ACC AAA ACA ACA ACT ACC AAT ACT ACA-3'	67 °C
MSP (<i>TIG1</i> , methylated) F: 5'-GCG GGG TTC GGG GAT TTC-3' R: 5'-GTA CGC GAA CAA ACA AAC G-3'	56 °C
MSP (<i>TIG1</i> , unmethylated) F: 5'-GTG GGG TTT GGG GAT TTT GAT-3' R: 5'-ATA CAC AAA CAA ACA AAC ACA-3'	55 °C
RT-PCR (<i>CRBP1</i>) F: 5'-TTG TTG CCA AAC TGG CTC CA-3' R: 5'-ACA CTG GAG CTT GTC TCC GT-3'	53 °C
RT-PCR (<i>TIG1</i>) F: 5'-GAA AAA CCC CTT GGA AAT AGT CAG C-3' R: 5'-AGT GTG ACA CCT GTG TTG TCA TTT CC-3'	68 °C
RT-PCR (<i>ACTB</i>) F: 5'-CTGTCTGGCGCACCACCAT-3' R: 5'-GCAACTAAGTCATAGTCCGC-3'	55 °C
Quantitative RT-PCR (<i>RARβ</i>) F: 5'-ACC ACT GGA CCA TGT AAC TCT AGT GT-3' R: 5'-GGC ATC AAG AAG GGC TGG A-3'	60 °C
Quantitative RT-PCR (<i>CRBP1</i>) F: 5'-CAA CAG TGA GCT GGG ACG G-3' R: 5'-GCC ACG CCC CTC CIT C-3'	60 °C
Quantitative RT-PCR (<i>TIG1</i>) F: 5'-GGC CGC GCG TGG AT-3' R: 5'-GGT TGT AGC GCT CTG TGC TG-3'	60 °C
Quantitative RT-PCR (<i>ACTB</i>) F: 5'-TCA CCG AGC GCG GCT-3' R: 5'-TAA TGT CAC GCA CGA TTT CCC-3'	60 °C

PCR: polymerase chain reaction; RT-PCR: reverse transcriptase-polymerase chain reaction; MSP: methylation-specific polymerase chain reaction; *RARβ*: retinoid acid receptor β; F: forward; R: reverse; *CRBP1*: cellular retinol-binding protein 1; *TIG1*: tazarotone-induced gene 1; RT-PCR: reverse transcriptase-polymerase chain reaction.

quence Detection System (Applied Biosystems) as described previously.³⁰ Primer sequences are listed in Table 1. We calculated the ratio of target gene mRNA expression levels between GC samples (T) and samples of corresponding nonneoplastic mucosa (N), with a T/N ratio < 0.5 representing reduced expression. *ACTB*-specific PCR products were amplified from the same RNA samples and served as internal controls.




FIGURE 1. mRNA expression and DNA methylation status in gastric carcinoma (GC) cell lines. (A) Reverse transcriptase-polymerase chain reaction (RT-PCR) analysis of GC cell lines. Cellular retinol-binding protein 1 (*CRBP1*) was expressed only in the MKN-1 cell line. Tazarotene-induced gene 1 (*TIG1*) expression was identified in the MKN-7 and KATO-III cell lines. A plus sign indicates cell lines with retinoid acid receptor β (*RAR* β) expression and a minus sign indicates cell lines without *RAR* β expression (see Hayashi et al.¹⁰). Note that at least one of the three genes was inactivated in all of the GC cell lines tested. (B) Methylation-specific PCR (MSP) of *CRBP1* and *TIG1* in GC cell lines. Primer sets were either unmethylated (U) or methylated (M). The methylated allele for *CRBP1* was detected in the MKN-7, MKN-28, MKN-45, MKN-74, TMK-1, HSC-39, and KATO-III cell lines. The methylated allele for *TIG1* was detected in the MKN-1, MKN-28, MKN-45, MKN-74, TMK-1, and HSC-39 cell lines. U* indicates cell lines with an unmethylated *RAR* β gene, M* indicates cell lines with a methylated *RAR* β gene, and U/M* indicates cell lines with an unmethylated and a methylated *RAR* β gene (see Hayashi et al.¹⁰). (C) MSP in untreated and Aza-2'-deoxycytidine (Aza-dC)-treated MKN-28 cells. An unmethylated allele of both *CRBP1* and *TIG1* genes was identified in Aza-dC-treated cells. (D) RT-PCR of untreated and Aza-dC-treated MKN-28 cells. mRNA expression of both *CRBP1* and *TIG1* genes was identified in Aza-dC-treated cells.

Statistical Methods

Differences were analyzed statistically with the Fisher exact test and the Mann-Whitney *U* test. *P* values < 0.05 were considered statistically significant.

RESULTS

DNA Methylation Status and Expression of CRBP1 and TIG1 in GC Cell Lines

To determine whether hypermethylation of the *CRBP1* and *TIG1* genes induces transcriptional inactivation, we performed in vitro analysis with 8 GC cell lines. RT-PCR analysis showed that seven of the cell lines (MKN-7, MKN-28, MKN-45, MKN-74, TMK-1, HSC-39, and KATO-III) did not express *CRBP1* mRNA, and six

of the cell lines (MKN-1, MKN-28, MKN-45, MKN-74, TMK-1, and HSC-39) did not express *TIG1* mRNA. Figure 1A,B shows that all GC cell lines that lacked *CRBP1* gene expression showed methylation of CpG sites within the *CRBP1* gene, and all GC cell lines that lacked *TIG1* gene expression showed methylation of CpG sites within the *TIG1* gene. Therefore, methylation status is associated with expression of both *CRBP1* and *TIG1*.

Next, we investigated whether methylation status of the *RAR* β , *CRBP1*, and *TIG1* genes was related (Fig. 1A,B). We previously reported on the *RAR* β methylation status in GC cell lines.¹⁰ No clear relation was noted for the methylation status among the three




FIGURE 2. Methylation-specific polymerase chain reaction (MSP) analysis of retinoid acid receptor β (*RAR\beta*), cellular retinol-binding protein 1 (*CRBP1*), and tazarotene-induced gene 1 (*TIG1*) genes in gastric carcinoma, in corresponding nonneoplastic gastric mucosa, and in normal gastric mucosa obtained from young, healthy individuals. U indicates unmethylated polymerase chain reaction (PCR) products and M indicates methylated PCR products. A methylated allele of the *RAR\beta* gene was detected in Samples 1T and 3T. A methylated allele of the *CRBP1* gene was detected in Samples 4T and 5T. A methylated allele of the *TIG1* gene was detected in Sample 5T. In corresponding nonneoplastic gastric mucosa, a methylated allele of the *RAR\beta* gene was detected in Sample 1N. Methylated alleles of *CRBP1* and *TIG1* were not detected. In normal gastric mucosa from young, healthy individuals, no methylation was detected.

genes; however, our results showed that at least one of these genes was hypermethylated and inactivated in all of GC cell lines tested.

To confirm whether transcriptional inactivation of *CRBP1* and *TIG1* causes DNA methylation, we treated MKN-28 cells with Aza-dC and then performed RT-PCR (Fig. 1C) and MSP (Fig. 1D) analyses. Treatment with Aza-dC restored *CRBP1* and *TIG1* expression, and an unmethylated allele was detected after Aza-dC treatment.

***RAR\beta*, *CRBP1*, and *TIG1* Methylation Status and mRNA Expression Levels in GC Tissues**

Representative MSP data for *RAR\beta*, *CRBP1*, and *TIG1* in GC tissues, in corresponding nonneoplastic mucosa, and in normal gastric mucosa from young, healthy individuals are shown in Figure 2. In the 42 GC samples, CpG island hypermethylation was detected at the following frequencies: 15 of 42 GC samples (36%) were hypermethylated for *RAR\beta*, 14 of 42 GC samples (33%) were hypermethylated for *CRBP1*, and 4 of 42 samples (10%) were hypermethylated for *TIG1*. The overall results are shown in Table 2. No clear association was found between the 3 genes in terms of methylation status (Table 3). Of the 42 GC samples, at least 1 of the 3 genes was hypermethylated in 24 samples (57%). We analyzed the correlation of methylation status of each gene with clinical characteristics (age, gender, *H. pylori* infection, T classification [depth of tumor invasion], N status [degree of lymph node metastasis], tumor stage, and histologic type) (Table 4). DNA methylation of the *CRBP1* gene was found more frequently in female patients than in male patients. No association was found between the GC samples with hypermethylation of at least one gene and clinical data (data not shown).

In 30 samples of corresponding nonneoplastic mucosa, DNA methylation of *RAR\beta* was found in 6 of

30 samples (20%), DNA methylation of *CRBP1* was found in 0 of 30 samples (0%), DNA methylation of *TIG1* was found in 1 of 30 samples (3%) (Fig. 2). Only *CRBP1* was methylated significantly more frequently in tumor tissue compared with nonneoplastic tissue (Table 5). Among 30 corresponding nonneoplastic mucosa samples, intestinal metaplasia was found in 22 samples. No correlation was observed between methylation of any of these genes and the presence of intestinal metaplasia (data not shown). There was no correlation of methylation status between tumor and nonneoplastic tissue from the same patient. Methylated alleles were not detected in any of 10 samples of normal gastric mucosa from young, healthy individuals (Fig. 2).

We also used quantitative RT-PCR analysis to determine whether methylation of the *RAR\beta*, *CRBP1*, and *TIG1* genes affects expression of the respective mRNA. Reduced expression of *RAR\beta*, *CRBP1*, and *TIG1* was found in 17 of 42 GC samples (40%), 16 of 42 GC samples (38%), and 8 of 42 GC samples (19%) (Table 2). The *RAR\beta* mRNA expression levels in tumor tissues with DNA methylation of *RAR\beta* (mean \pm standard error [in arbitrary units], 2.40 ± 0.58) were significantly lower compared in the levels in tumor tissues without DNA methylation (8.35 ± 1.56 ; $P = 0.0087$; Mann-Whitney *U* test) (Table 6). The mRNA expression levels of both *CRBP1* and *TIG1* also were correlated with hypermethylation of the respective genes ($P = 0.0109$ for *CRBP1*; $P = 0.0003$ for *TIG1*; Mann-Whitney *U* test) (Table 6).

DISCUSSION

It is believed that *RAR\beta* functions as a tumor suppressor. Previous studies have shown that overexpression of *RAR\beta* induces growth arrest and apoptosis in several carcinoma cell lines.^{10,31,32} In the current study, we investigated three genes associated with retinoic

TABLE 2
Summary of DNA Methylation and mRNA Expression of *RARβ*, *CRBP1*, and *TIG1* in Gastric Carcinoma

Patient no.	Gender	Age (yrs)	<i>H. pylori</i>	TNM classification ^a				Histology ^b	DNA methylation			RNA expression		
				T	N	M	Stage		<i>RARβ</i>	<i>CRBP1</i>	<i>TIG1</i>	<i>RARβ</i>	<i>CRBP1</i>	<i>TIG1</i>
8	Female	73	Negative	2	1	0	II	Intestinal	MET	MET	MET	R	R	R
4	Male	69	Positive	4	1	0	IV	Diffuse	MET	MET	MET	R	R	R
20	Male	77	Negative	2	0	0	IB	Intestinal	MET	MET	— ^c	R	R	— ^d
10	Male	81	Positive	2	2	0	IIIA	Diffuse	MET	MET	—	—	R	—
6	Female	61	Negative	3	2	0	IIIB	Intestinal	MET	MET	—	—	—	—
31	Female	88	Positive	4	3	0	IV	Diffuse	MET	MET	—	R	R	—
3	Female	62	Positive	3	3	0	IV	Diffuse	MET	—	MET	R	—	R
15	Male	76	Positive	1	0	0	IA	Intestinal	MET	—	—	R	—	—
41	Male	61	Positive	2	1	0	II	Diffuse	MET	—	—	—	—	R
2	Male	69	Positive	2	1	0	II	Intestinal	MET	—	—	—	—	—
25	Male	65	Negative	3	1	0	IIIA	Diffuse	MET	—	—	R	—	—
16	Male	69	Positive	3	1	0	IIIA	Intestinal	MET	—	—	R	—	—
36	Male	70	Positive	4	0	0	IV	Intestinal	MET	—	—	—	—	—
9	Female	74	Negative	4	2	0	IV	Diffuse	MET	—	—	R	—	—
5	Male	34	Negative	4	2	0	IV	Intestinal	MET	—	—	R	—	—
37	Female	74	Positive	2	0	0	IB	Intestinal	—	MET	—	—	R	—
34	Female	54	Positive	3	0	0	II	Diffuse	—	MET	—	—	R	—
38	Male	76	Positive	2	1	0	II	Diffuse	—	MET	—	—	R	—
33	Male	63	Positive	2	1	0	II	Intestinal	—	MET	—	—	—	—
13	Female	77	Positive	3	0	0	II	Intestinal	—	MET	—	—	R	—
30	Male	74	Positive	3	1	0	IIIA	Diffuse	—	MET	—	—	—	—
17	Female	53	Positive	2	2	0	IIIA	Diffuse	—	MET	—	—	R	R
32	Female	40	Positive	3	3	0	IV	Diffuse	—	MET	—	—	—	—
11	Male	66	Negative	3	1	0	IIIA	Intestinal	—	—	MET	—	—	R
1	Male	71	Positive	2	0	0	IB	Intestinal	—	—	—	—	—	—
24	Male	75	Negative	2	0	0	IB	Intestinal	—	—	—	—	—	R
40	Male	72	Positive	2	0	0	IB	Intestinal	—	—	—	—	—	—
42	Male	73	Positive	2	0	0	IB	Diffuse	—	—	—	R	R	—
23	Female	69	Positive	2	1	0	II	Intestinal	—	—	—	R	R	—
29	Male	79	Positive	2	1	0	II	Diffuse	—	—	—	—	—	—
28	Male	75	Positive	3	2	0	IIIB	Diffuse	—	—	—	—	—	—
12	Male	85	Positive	3	1	0	IIIA	Diffuse	—	—	—	—	—	—
21	Male	78	Negative	3	1	0	IIIA	Diffuse	—	—	—	—	—	R
26	Male	87	Positive	3	2	0	IIIB	Diffuse	—	—	—	—	—	—
35	Male	78	Positive	3	1	0	IIIA	Diffuse	—	—	—	R	R	—
18	Female	57	Positive	3	1	0	IIIA	Intestinal	—	—	—	R	R	—
22	Male	54	Negative	3	2	0	IIIB	Intestinal	—	—	—	R	R	—
27	Female	67	Positive	3	3	0	IV	Diffuse	—	—	—	IV	—	—
14	Male	72	Positive	3	3	0	IV	Diffuse	—	—	—	R	R	—
39	Male	71	Positive	3	3	0	IV	Diffuse	—	—	—	—	—	—
7	Male	72	Positive	3	3	0	IV	Intestinal	—	—	—	—	—	—
19	Male	58	Positive	3	3	0	IV	Intestinal	—	—	—	R	—	—

RARβ: retinoid acid receptor β; *CRBP1*: cellular retinol-binding protein 1; *TIG1*: tazarotone-induced gene 1; *H. pylori*, *Helicobacter pylori*; T: tumor; N: lymph node; M: metastasis; MET: methylated; R: reduced expression.

^a Stage was classified according to the criteria of the International Union Against Cancer TNM classification system, 6th edition, 2002.

^b Histology was classified according to the criteria of Lauren.

^c Unmethylated.

^d Not reduced.

acid signaling. DNA hypermethylation of *CRBP1* was found in 33% of GC samples and was associated with low levels of mRNA expression. DNA hypermethylation of *TIG1* was found in 10% of GC samples and also was associated with low levels of mRNA expres-

sion. In addition, Aza-dC treatment induced demethylation of both the *CRBP1* gene and the *TIG1* gene and restored expression of both mRNAs in MKN-28 cells, suggesting that hypermethylation of both *CRBP1* and *TIG1* plays an important role in transcriptional silenc-

TABLE 3
Correlation of DNA Methylation Status of Retinoic Acid Signaling-Associated Genes in Gastric Carcinoma

Methylation status	Methylation status				P value ^a
	<i>RARβ</i>		<i>CRBP1</i>		
	Methylated	Unmethylated	Methylated	Unmethylated	
<i>CRBP1</i>					
Methylated	6	8			0.5159
Unmethylated	9	19			
<i>TIG1</i>					
Methylated	3	1			0.1220
Unmethylated	12	26			
<i>TIG1</i>					
Methylated			2	2	0.5902
Unmethylated			12	26	

RARβ: retinoid acid receptor β; *CRBP1*: cellular retinol-binding protein 1; *TIG1*: tazarotone-induced gene 1.
^a Statistical significance determined using the Fisher exact test.

TABLE 4
Association between DNA Methylation of Retinoic Acid Signaling-Associated Genes and Clinicopathologic Parameters

Characteristic	No. of patients (%)								
	<i>RARβ</i>			<i>CRBP1</i>			<i>TIG1</i>		
	M	U	P value ^a	M	U	P value	M	U	P value
Age									
> 65 yrs	10 (33)	20	NS	9 (30)	21	NS	3 (10)	27	NS
≤ 65 yrs	5 (42)	7		5 (42)	7		1 (8)	11	
Gender									
Male	10 (34)	19	NS	6 (21)	23	0.0148	2 (7)	27	NS
Female	5 (38)	8		8 (62)	5		2 (15)	11	
<i>H. pylori</i> infection									
Positive	9 (28)	23	NS	11 (34)	21	NS	2 (6)	30	NS
Negative	6 (60)	4		3 (30)	7		2 (20)	8	
T classification ^b									
T1/T2	6 (38)	10	NS	7 (44)	9	NS	1 (6)	15	NS
T3/T4	9 (35)	17		7 (27)	19		3 (12)	23	
N status ^b									
N0	3 (30)	7	NS	4 (40)	6	NS	0 (0)	10	NS
N1/N2/N3	12 (38)	20		10 (31)	22		4 (13)	28	
Stage ^b									
Stage I/II	5 (31)	11	NS	7 (44)	9	NS	1 (6)	15	NS
Stage III/IV	10 (38)	16		7 (27)	19		3 (12)	23	
Histologic type ^c									
Intestinal	8 (40)	12	NS	6 (30)	14	NS	2 (10)	18	NS
Diffuse	7 (32)	15		8 (36)	14		2 (9)	20	

RARβ: retinoid acid receptor β; *CRBP1*: cellular retinol-binding protein 1; *TIG1*: tazarotone-induced gene 1; M: methylated; U: unmethylated; NS: not significant; *H. pylori*: *Helicobacter pylori*.

^a Statistical significance was determined using the Fisher exact test.

^b Stage was classified according to the criteria of the International Union Against Cancer TNM classification of malignant tumors, 6th edition, 2002.²⁷

^c Histology was classified according to the criteria of Lauren.

ing in GC. We also confirmed that DNA methylation of *RARβ* is associated with low levels of mRNA expression in this study. It is important to note that there were several tumor samples with DNA hypermethylation that did not show low gene expression. This may

be related to the extreme sensitivity of the MSP technique, which theoretically can detect as few as 0.1% of methylated cells.²⁸ In contrast, several samples showed low gene expression in the absence of DNA methylation. Alternative inactivating pathways, such

TABLE 5
Frequency of DNA Methylation of Retinoic Acid Signaling-Associated Genes in Gastric Carcinoma and Corresponding Nonneoplastic Mucosa

Measure	No. of patients (%)		P value ^a
	Methylated	Unmethylated	
<i>RARβ</i> methylation status			
Gastric carcinoma	15 (36)	27	0.1922
Corresponding nonneoplastic mucosa	6 (20)	24	
<i>CRBP1</i> methylation status			
Gastric carcinoma	14 (33)	28	0.0002
Corresponding nonneoplastic mucosa	0 (0)	30	
<i>TIG1</i> methylation status			
Gastric carcinoma	4 (10)	38	0.3932
Corresponding nonneoplastic mucosa	1 (3)	29	

RARβ: retinoid acid receptor β; *CRBP1*: cellular retinol-binding protein 1; *TIG1*: tazarotene-induced gene 1.

^a Statistical significance was determined using the Fisher exact test.

TABLE 6
Association between DNA Methylation of Retinoic Acid Signaling-Associated Genes and mRNA Expression

Sample	No. of patients	Expression level ^a	P value ^b
<i>RARβ</i>			
Gastric carcinoma	42	6.22 ± 1.11 ^c	0.0087
Methylated	15	2.40 ± 0.58	
Unmethylated	27	8.35 ± 1.56	
Corresponding nonneoplastic mucosa	42	10.24 ± 2.14	
<i>CRBP1</i>			
Gastric carcinoma	42	4.22 ± 1.83 ^c	0.0109
Methylated	14	1.40 ± 0.28	
Unmethylated	28	5.63 ± 2.72	
Corresponding nonneoplastic mucosa	42	6.07 ± 0.91	
<i>TIG1</i>			
Gastric carcinoma	42	17.68 ± 5.59 ^d	0.0003
Methylated	4	0.41 ± 0.05	
Unmethylated	38	19.50 ± 6.11	
Corresponding nonneoplastic mucosa	42	15.86 ± 5.18	

RARβ: retinoid acid receptor β; *CRBP1*: cellular retinol-binding protein 1; *TIG1*: tazarotene-induced gene 1.

^a Mean ± standard error.

^b Mann-Whitney U test.

^c The units were arbitrary, and the *RARβ* and *CRBP1* mRNA expression levels were calculated by standardization with 1 μg total RNA of the MKN-1 cells, taken as 1.0.

^d The units were arbitrary, and the *TIG1* mRNA expression levels were calculated by standardization with 1 μg total RNA of the MKN-7 cells, taken as 1.0.

as hemizygous deletion and alteration of transcription factors, may account for the low gene expression in these samples. The *RARβ* gene is located on chromosome 3p, and both the *CRBP1* gene and the *TIG1* gene are located on chromosome 3q. Loss of heterozygosity of chromosome 3p has been reported in 33% of GCs.³³

DNA methylation of *RARβ* and *TIG1* was detected in corresponding nonneoplastic mucosa. Although the frequency of methylation of all 3 genes in GC was greater than in nonneoplastic tissue, statistical analysis revealed that only *CRBP1* was methylated signifi-

cantly more frequently in tumor tissue compared with corresponding nonneoplastic mucosa. This suggests that DNA methylation of the *CRBP1* gene may contribute to gastric carcinogenesis. Conversely, there was no statistical difference between the *RARβ* and *TIG1* methylation status of tumor tissue and nonneoplastic tissue. Because the mRNA expression levels of these two genes in tumor tissues were correlated with DNA methylation, it is possible that these two genes may demonstrate monoallelic methylation in nonneoplastic tissue and biallelic methylation in tumors. Epi-

TABLE 7
Association between DNA Methylation of Retinoic Acid Signaling-Associated Genes and mRNA Expression

	No. cases	<i>RAR-beta</i> expression levels ^a	<i>P</i> value ^b
Gastric cancer	42	6.22 ± 1.11 ^c	0.0087
Methylated	15	2.40 ± 0.58	
Unmethylated	27	8.35 ± 1.56	
Corresponding non-neoplastic mucosa	42	10.24 ± 2.14	
		<i>CRBP1</i> expression levels	
Gastric cancer	42	4.22 ± 1.83 ^c	0.0109
Methylated	14	1.40 ± 0.28	
Unmethylated	28	5.63 ± 2.72	
Corresponding non-neoplastic mucosa	42	6.07 ± 0.91	
		<i>TIG1</i> expression levels	
Gastric cancer	42	17.68 ± 5.59 ^d	0.0003
Methylated	4	0.41 ± 0.05	
Unmethylated	38	19.50 ± 6.11	
Corresponding non-neoplastic mucosa	42	15.86 ± 5.18	

^a Mean ± SE

^b Mann-Whitney *U*-test.

^c The units are arbitrary, and we calculated the *RAR-beta* and *CRBP1* mRNA expression level by standardization with 1 µg total RNA of the MKN-1 cells, taken as 1.0.

^d The units are arbitrary, and we calculated the *TIG1* mRNA expression level by standardization with 1 µg total RNA of the MKN-7 cells, taken as 1.0.

genetic changes, including DNA methylation, occur in premalignant and histologically normal gastric epithelium.³⁴ Recent evidence suggests that methylation of certain genes is associated with aging.³⁴ We confirmed that normal gastric mucosa from young, healthy individuals did not show hypermethylation of the three genes. Thus, aging may be involved in the methylation status of nonneoplastic mucosa. Age-related methylation of the *RARβ* and *TIG1* genes also may explain the possible link between aging and increased risk for GC. Because *CRBP1* methylation was not detected in corresponding nonneoplastic mucosa, *CRBP1* methylation may serve as a good biomarker for GC. It is known that *CRBP1* regulates intracellular retinoic acid transport, metabolism, and transcriptional activity. Thus, *CRBP1* affects the transcription of retinoic acid-responsive genes that regulate cell proliferation. It is possible that, even if *RARβ* is expressed, the loss of *CRBP1* may compromise retinoic acid metabolism by diminishing retinoic acid transport, resulting in blockage of the transcription of retinoic acid-responsive genes that regulate cell proliferation. In the current study, there was no clear association between DNA methylation of *RARβ* and *CRBP1*, indicating that inactivation of the two genes may not create a synergistic effect but may be a random event in gastric carcinogenesis. *TIG1* hypermethylation was detected in 10% of GC samples. In other malignancies, *TIG1* gene hypermethylation has been reported in 33–90% of sam-

ples.^{22,23,35} Although *TIG1* has been identified as a potential tumor suppressor gene for prostate carcinoma,²⁴ the significance of *TIG1* inactivation in GC may be weak compared with that in other malignancies. In the current study, there was a significant difference between the methylation of *TIG1* in cell lines (6 of 8 cell lines; 75%) and in GC tissue samples (4 of 42 samples; 10%). The reason for this discrepancy is unclear. Because DNA methylation of *TIG1* reportedly has been correlated with tumor invasiveness in prostate carcinoma,²² DNA methylation of *TIG1* may occur frequently in GC cell lines that were established from metastatic foci.³⁶

DNA methylation occurs early in the multistep process of gastric carcinogenesis^{9,37}; thus, it is believed that DNA methylation contributes to carcinogenesis but not to tumor progression. In the current study, although DNA methylation of the *CRBP1* gene was observed more frequently in female patients than in male patients, the other clinical characteristics, such as tumor stage, were not correlated with DNA methylation. In addition, the frequencies of methylation of three genes in GC were greater than those in corresponding nonneoplastic mucosa, indicating that DNA methylation of these genes may participate in gastric carcinogenesis but not in GC progression. In addition, although concordant hypermethylation of *RARβ* and *CRBP1*¹⁹ and of *RARβ* and *TIG1*²² has been reported, we found no statistically significant concordant hypermethylation among these

genes, and greater than 50% of the GC samples we studied showed DNA methylation of at least 1 of the 3 genes. These results suggest that alterations of retinoic acid signaling are involved widely in GC.

In conclusion, the results of the current study demonstrate that inactivation of retinoic acid signaling-associated genes, *RAR β* , *CRBP1*, and *TIG1*, due to DNA methylation occurs frequently in GC. Because DNA methylation can be reversed by demethylating agents, such as Aza-dC, it may be better to use a combination of retinoids and demethylating agents for therapeutic strategies in patients with GC.

REFERENCES

1. Yasui W, Oue N, Ono S, Mitani Y, Ito R, Nakayama H. Histone acetylation and gastrointestinal carcinogenesis. *Ann NY Acad Sci*. 2003;983:220–231.
2. Oue N, Hamai Y, Mitani Y, et al. Gene expression profile of gastric carcinoma: identification of genes and tags potentially involved in invasion, metastasis, and carcinogenesis by serial analysis of gene expression. *Cancer Res*. 2004;64:2397–2405.
3. Jones PA, Baylin SB. The fundamental role of epigenetic events in cancer. *Nat Rev Genet*. 2002;3:415–428.
4. Kaneda A, Wakazono K, Tsukamoto T, et al. Lysyl oxidase is a tumor suppressor gene inactivated by methylation and loss of heterozygosity in human gastric cancers. *Cancer Res*. 2004;64:6410–6415.
5. Rashid A, Issa JP. CpG island methylation in gastroenterologic neoplasia: a maturing field. *Gastroenterology*. 2004;127:1578–1588.
6. Toyota M, Ahuja N, Suzuki H, et al. Aberrant methylation in gastric cancer associated with the CpG island methylator phenotype. *Cancer Res*. 1999;59:5438–5442.
7. Kaneda A, Kaminishi M, Yanagihara K, Sugimura T, Ushijima T. Identification of silencing of nine genes in human gastric cancers. *Cancer Res*. 2002;62:6645–6650.
8. Oue N, Motoshita J, Yokozaki H, et al. Distinct promoter hypermethylation of p16INK4a, CDH1, and RAR-beta in intestinal, diffuse-adherent, and diffuse-scattered type gastric carcinomas. *J Pathol*. 2002;198:55–59.
9. Kang GH, Lee S, Kim JS, Jung HY. Profile of aberrant CpG island methylation along the multistep pathway of gastric carcinogenesis. *Lab Invest*. 2003;83:635–641.
10. Hayashi K, Yokozaki H, Goodison S, et al. Inactivation of retinoic acid receptor beta by promoter CpG hypermethylation in gastric cancer. *Differentiation*. 2001;68:13–21.
11. Widschwendter M, Berger J, Hermann M, et al. Methylation and silencing of the retinoic acid receptor-beta2 gene in breast cancer. *J Natl Cancer Inst*. 2000;92:826–832.
12. Virmani AK, Rathi A, Zochbauer-Muller S, et al. Promoter methylation and silencing of the retinoic acid receptor-beta gene in lung carcinomas. *J Natl Cancer Inst*. 2000;92:1303–1307.
13. Chambon P. The retinoid signaling pathway: molecular and genetic analyses. *Semin Cell Biol*. 1994;5:115–125.
14. Chambon P. A decade of molecular biology of retinoic acid receptors. *FASEB J*. 1996;10:940–954.
15. Folli C, Calderone V, Ottonello S, et al. Identification, retinoid binding, and X-ray analysis of a human retinol-binding protein. *Proc Natl Acad Sci USA*. 2001;98:3710–3715.
16. Ong DE, Newcomer ME, Chytil F. Cellular retinoid-binding proteins. In: Sporn MB, Roberts AB, Goodman DS, editors. *The retinoids: biology, chemistry and medicine*, 2nd ed. New York: Raven Press, 1994:283–317.
17. Vogel S, Mendelsohn CL, Mertz JR, et al. Characterization of a new member of the fatty acid-binding protein family that binds all-trans-retinol. *J Biol Chem*. 2001;276:1353–1360.
18. Ghyselinck NB, Bavik C, Sapin V, et al. Cellular retinol-binding protein I is essential for vitamin A homeostasis. *EMBO J*. 1999;18:4903–4914.
19. Esteller M, Guo M, Moreno V, et al. Hypermethylation-associated inactivation of the cellular retinol-binding-protein 1 gene in human cancer. *Cancer Res*. 2002;62:5902–5905.
20. Yamashita S, Nomoto T, Ohta T, Ohki M, Sugimura T, Ushijima T. Differential expression of genes related to levels of mucosal cell proliferation among multiple rat strains by using oligonucleotide microarrays. *Mamm Genome*. 2003;14:845–852.
21. Nagpal S, Patel S, Asano AT, Johnson AT, Duvic M, Chandraratna RA. Tazarotene-induced gene 1 (TIG1), a novel retinoic acid receptor-responsive gene in skin. *J Invest Dermatol*. 1996;106:269–274.
22. Zhang J, Liu L, Pfeifer GP. Methylation of the retinoid response gene TIG1 in prostate cancer correlates with methylation of the retinoic acid receptor beta gene. *Oncogene*. 2004;23:2241–2249.
23. Youssef EM, Chen XQ, Higuchi E, et al. Hypermethylation and silencing of the putative tumor suppressor Tazarotene-induced gene 1 in human cancers. *Cancer Res*. 2004;64:2411–2417.
24. Jing C, El-Ghany MA, Beesley C, et al. Tazarotene-induced gene 1 (TIG1) expression in prostate carcinomas and its relationship to tumorigenicity. *J Natl Cancer Inst*. 2002;94:482–490.
25. Ochiai A, Yasui W, Tahara E. Growth-promoting effect of gastrin on human gastric carcinoma cell line TMK-1. *Jpn J Cancer Res*. 1985;76:1064–1071.
26. Yanagihara K, Seyama T, Tsumuraya M, Kamada N, Yokoro K. Establishment and characterization of human signet ring cell gastric carcinoma cell lines with amplification of the c-myc oncogene. *Cancer Res*. 1991;51:381–386.
27. Sobin LH, Wittekind CH. Stomach. In: Sobin LH, Wittekind CH, editors. *TNM classification of malignant tumors*, 6th ed. New York: John Wiley & Sons, 2002:65–68.
28. Herman JG, Graff JR, Myohanen S, Nelkin BD, Baylin SB. Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci USA*. 1996;93:9821–9826.
29. Sirchia SM, Ferguson AT, Sironi E, et al. Evidence of epigenetic changes affecting the chromatin state of the retinoic acid receptor beta2 promoter in breast cancer cells. *Oncogene*. 2000;19:1556–1563.
30. Kondo T, Oue N, Yoshida K, et al. Expression of POT1 is associated with tumor stage and telomere length in gastric carcinoma. *Cancer Res*. 2004;64:523–529.
31. Seewaldt VL, Johnson BS, Parker MB, Collins SJ, Swisshelm K. Expression of retinoic acid receptor beta mediates retinoic acid-induced growth arrest and apoptosis in breast cancer cells. *Cell Growth Differ*. 1995;6:1077–1088.
32. Wan H, Oridate N, Lotan D, Hong WK, Lotan R. Overexpression of retinoic acid receptor beta in head and neck squamous cell carcinoma cells increases their sensitivity to retinoid-induced suppression of squamous differentiation by retinoids. *Cancer Res*. 1999;59:3518–3526.

33. Kim KM, Kwon MS, Hong SJ, et al. Genetic classification of intestinal-type and diffuse-type gastric cancers based on chromosomal loss and microsatellite instability. *Virchows Arch.* 2003;443:491–500. Epub 2003 Aug 15.
34. Waki T, Tamura G, Tsuchiya T, Sato K, Nishizuka S, Motoyama T. Promoter methylation status of E-cadherin, hMLH1, and p16 genes in nonneoplastic gastric epithelia. *Am J Pathol.* 2002;161:399–403.
35. Kwong J, Lo KW, Chow LS, Chan FL, To KF, Huang DP. Silencing of the retinoid response gene TIG1 by promoter hypermethylation in nasopharyngeal carcinoma. *Int J Cancer.* 2005;113:386–392.
36. Yokozaki H. Molecular characteristics of eight gastric cancer cell lines established in Japan. *Pathol Int.* 2000;50:767–777.
37. Oue N, Sentani K, Yokozaki H, Kitadai Y, Ito R, Yasui W. Promoter methylation status of the DNA repair genes hMLH1 and MGMT in gastric carcinoma and metaplastic mucosa. *Pathobiology.* 2001;69:143–149.

Shunji Matsumura · Naohide Oue · Hirofumi Nakayama
Yasuhiko Kitadai · Kazuhiro Yoshida
Yoshiyuki Yamaguchi · Kazue Imai
Kei Nakachi · Keisuke Matsusaki · Kazuaki Chayama
Wataru Yasui

A single nucleotide polymorphism in the *MMP-9* promoter affects tumor progression and invasive phenotype of gastric cancer

Received: 18 June 2004 / Accepted: 16 August 2004 / Published online: 30 September 2004
© Springer-Verlag 2004

Abstract Purpose: Matrix metalloproteinase-9 (MMP-9, gelatinase B) plays a key role in cancer invasion and metastasis by degrading the extracellular matrix (ECM) and basement membrane barriers. A cytosine (C)-thymidine (T) single nucleotide polymorphism (SNP) at position -1562 in the *MMP-9* promoter is reported to affect expression of this gene. The purpose of this study was to investigate the relation between the -1562 C/T polymorphism and the development and progression of gastric cancer. **Methods:** The study population included 177 gastric cancer patients and 224 healthy control subjects. The SNP in the *MMP-9* promoter was analyzed by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) and sequencing. Genotype frequencies were compared between patients and controls, and the association of

genotypes with clinicopathological features was studied. **Results:** Genotype frequencies in gastric cancer patients were similar to those in control subjects ($P = 0.223$). However, significant association was found between degree of tumor invasion, clinical stage, and lymphatic invasion and the *MMP-9* polymorphism in gastric cancer patients ($P < 0.05$, for each). **Conclusions:** Our results indicate that the T allele in the *MMP-9* promoter is associated with the invasive phenotype of gastric cancer.

Keywords MMP-9 · SNP · Gastric cancer · Depth of tumor invasion · Lymphatic invasion

Abbreviations MMP-9: Matrix metalloproteinase-9 · SNP: Single nucleotide polymorphism · C: Cytosine · T: Thymidine · RFLP: Restriction fragment length polymorphism · ECM: Extracellular matrix · PCR: Polymerase chain reaction · OR: Odds ratio · CI: Confidence interval

S. Matsumura · N. Oue · H. Nakayama · W. Yasui (✉)
Department of Molecular Pathology,
Hiroshima University Graduate School
of Biomedical Sciences, 1-2-3 Kasumi, Minami-ku,
734-8551 Hiroshima,
Japan
E-mail: wyasui@hiroshima-u.ac.jp
Tel.: + 81-82-2575145
Fax: + 81-82-2575149

Y. Kitadai · K. Chayama
Department of Medicine and Molecular Science,
Hiroshima University Graduate School of Biomedical Sciences,
Hiroshima, Japan

K. Yoshida · Y. Yamaguchi
Department of Surgical Oncology,
Research Institute for Radiation Biology and Medicine,
Hiroshima University, Hiroshima, Japan

K. Imai · K. Nakachi
Department of Radiobiology/Molecular Epidemiology,
Radiation Effects Research Foundation, Hiroshima,
Japan

K. Matsusaki
Department of Surgery, Hofu Institute of Gastroenterology,
Yamaguchi, Japan

Introduction

Gastric cancer is one of the most common malignancies in the world and is the leading cause of death in Japan. Poor prognosis reflects the invasive and metastatic capabilities of cancer cells. Degradation of the extracellular matrix (ECM) and basement membrane barriers by matrix metalloproteinases (MMPs) plays an important role in tumor invasion and metastasis (Forget et al. 1999; Kohn and Liotta 1995; Liotta et al. 1991). MMP-9 (92-kDa gelatinase, type IV collagenase) is a member of the family of MMP genes, which encode zinc-dependent enzymes that break down ECM through the degradation of type IV collagen (Nagase and Woessner 1999), and promote tumor cell invasion. The prognostic value of MMP-9 expression by tumor tissues has been reported in relation to a variety of cancers (McDonnell and Matri-

sian 1990; Kallakury et al. 2001; Baker and Leaper 2003; Tanioka et al. 2003; Ozalp et al. 2003; Sakata et al. 2004).

Zhang et al. (Zhang et al. 1999) reported that a cytosine (C)-to-thymidine (T) transition at nucleotide -1562 in the *MMP-9* gene promoter generates low-activity (C/C) and high-activity (C/T, T/T) promoter genotypes, which influence gene transcription. This polymorphism is associated with the severity of coronary atherosclerosis in patients with coronary artery disease. However, there have been no studies of the relation between this polymorphism and malignancies. Various genetic and epigenetic alterations are associated with gastric carcinoma (Yasui et al. 2000; Oue et al. 2002; Oue et al. 2003). We previously reported that several polymorphisms are significantly associated with gastric cancer (Kuraoka et al. 2003; Matsumura et al. 2004). With respect to the role of the C/T polymorphism in transcriptional activity and degradation of ECM, we hypothesized that this polymorphism might also act as a genetic modifier in the development and progression of gastric cancer. Therefore, we conducted a case-control study to investigate the association between the different *MMP-9* promoter alleles and gastric cancer. Moreover, we examined the relation between the C/T polymorphism and the clinicopathological features of gastric cancer patients.

Materials and methods

Samples

A total of 401 peripheral blood samples from 224 healthy control subjects and 177 gastric cancer patients were used in the present study. Control subjects were randomly selected from among individuals visiting hospitals for regular health checks or because of symptoms such as appetite loss or epigastralgia. Control subjects were confirmed to be free of malignancy by gastrointestinal endoscopy and biopsy. Gastric cancer patients underwent surgical or endoscopic mucosal resection (EMR) at Hiroshima University Hospital or Hofu Institute of Gastroenterology. Gastric cancer patients were 129 men and 48 women; the median age was 65.1 ± 11.7 years. Gastric cancer was classified histologically according to the criteria of Lauren (Lauren 1965); 103 patients had intestinal type gastric cancer, and 74 patients had diffuse type. Intestinal type and diffuse type correspond to well-differentiated type and poorly differentiated type, respectively, of the histological classification system of the Japanese Gastric Cancer Association (Japanese Gastric Cancer Association 1998). Gastric cancer patients were grouped according to TNM classification (Sobin and Wittekind 2002) on the basis of the post-operative histopathological evaluation. Moreover, patients were assigned to two subgroups on the basis of whether they were positive or negative for lymphatic invasion or venous invasion at the time of diagnosis (Japanese Gastric Cancer Association 1998). All patients

and control subjects gave written informed consent prior to enrollment in the study. The Human Genome Research Ethics Screening Committee of Hiroshima University School of Medicine approved the study.

DNA extraction and polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) analysis

A genomic DNA purification kit (Promega, Madison, Wisc., USA) and a QIAamp 96 DNA Blood kit (QIAGEN, Valencia, Calif., USA) were used for DNA extraction. PCR-RFLP assay was used to determine *MMP-9* genotypes as previously described (Zhang et al. 1999). To analyze the -1562 C/T polymorphism, we amplified a region of the *MMP-9* promoter with forward primer 5'-GCC TGG CAC ATA GTA GGC CC-3' and reverse primer 5'-CTT CCT AGC CAG CCG GCA TC -3' (Zhang et al. 1999). The target sequence was amplified in a 25- μ l reaction volume containing 10–20 ng of genomic DNA, 0.2 μ M dNTP, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 2 mM MgCl₂, 0.3 μ M of each primer, and 0.75 units of AmpliTaq Gold (Perkin-Elmer, Norwalk, Conn., USA). Amplification conditions were an initial activation step of 94 °C for 10 min followed by 35 cycles of denaturation at 94 °C for 30 s, annealing at 59 °C for 30 s, and extension at 72 °C for 30 s. The 435 bp fragment was then digested with *SphI* (New England BioLabs, Beverly, Mass., USA) overnight at 37 °C. *SphI* does not digest the C allele (435 bp) but generates 188 bp and 247 bp fragments for the T allele. Digests were separated by electrophoresis on 2.5% NuSieve 3:1 agarose (FMC BioProducts, Rockland, Me., USA) gels for 60 min at 100 V. Heterozygotes had a combination of both alleles (435 bp, 247 bp, and 188 bp bands).

Sequencing analysis of PCR products

The PCR products were purified and cloned into pCR2.1 (Invitrogen, Carlsbad, Calif., USA). Plasmid DNA was extracted from individual clones by alkaline lysis plasmid miniprep. The cloned PCR fragments obtained from each sample were sequenced with both M13 reverse and M13 forward primers and the PRISM AmpliTaq DNA Polymerase FS Ready Reaction Dye Terminator Sequencing Kit (PerkinElmer ABI, Foster City, Calif., USA). Reamplified DNA fragments were purified with Centri-Sep Columns (Princeton Separations, Adelphia, N.J., USA) and sequenced with an ABI PRISM 310 Genetic Analyzer (PerkinElmer ABI).

Helicobacter pylori status

Confirmation of *H. pylori* status was based on combinations of histology, ¹³C-urea breath test, and serum anti-*H. pylori* IgG antibody. One hundred and seventeen





Fig. 1 PCR-RFLP analysis of the -1562 C/T polymorphism in ten patients with gastric cancer. The ethidium bromide-stained 2.5% NuSieve 3:1 agarose gel used for genotyping is shown. The target region (435 bp) of the *MMP-9* gene promoter was PCR amplified and digested with *SphI*, which cleaved the T allele to generate two fragments (247 bp and 188 bp) are indicated, but did not cut the C allele (435 bp). Numbers above the panel are case numbers. Genotypes are indicated below each case

of 224 healthy control subjects and 73 of 177 gastric cancer patients underwent these examinations. Four biopsy specimens were taken, two from the greater curvature of the antrum and two from the upper body of the stomach. When lesions suspected to be cancerous were noted, additional biopsies were performed. Of these four specimens were fixed in formalin and assessed for *H. pylori* by Giemsa staining. The cutoff value of ^{13}C -urea breath test was 3.5‰. If one or more test were positive, patient was judged to be infected with *H. pylori*. If all *H. pylori* tests were negative, the patient was considered uninfected.

Statistical analysis

Statistical analysis was performed with the use of Fisher's exact test. $P < 0.05$ was considered statistically significant. Odds ratios (ORs) and 95% confidence intervals (95% CIs) were used to estimate the risk of association with genotypes. ORs for the genotypes were

Fig. 2 Representative electropherogram for each genotype for the -1562 C/T polymorphism of *MMP-9*. Genotype of the SNP was C/C in case 2 and T/T in case 5



calculated by the logistic regression model, with adjustments for age and sex; logistic regression analysis was performed for the association between genotypes and clinicopathological factors (SPSS 11.0, SPSS, Chicago, Ill., USA).

Results

Genotyping and genotype distributions in gastric cancer patients and control subjects

We examined the -1562 C/T polymorphism in the *MMP-9* gene promoter by PCR-RFLP in gastric cancer patients and control subjects. Three PCR-RFLP patterns were detected (Fig. 1). Patients 2, 4, 6, 7, 8, and 9 had a single 435 bp band (C/C genotype), and patient 5 had 247 bp and 188 bp bands (T/T genotype). Patients 1, 3, and 10 had all three bands, indicating the heterozygous genotype (C/T). Subsequent sequencing of representative cases confirmed the genotypes of patients 2 and 5 (Fig. 2). Distribution of the -1562 C/T polymorphism genotypes in gastric cancer patients and control subjects is shown in Table 1. Distribution of genotypes in controls was in good agreement with Hardy-Weinberg equilibrium; distribution in cases also agreed with Hardy-Weinberg equilibrium. Frequency of the T allele (C/T and T/T) was similar between gastric cancer patients (24.9%) and control subjects (30.3%; $P = 0.22$, OR=0.75, 95% CI=0.48–1.18). In addition, genotype frequencies (C/C vs C/T + T/T) did not differ by sex, age, or *H. pylori* status.

Table 1 *MMP-9* Genotype distributions in study subjects (OR odds ratio, CI confidence interval)

	Controls ^a (%) (n = 224)	Patients (%) (n = 177)	P value ^b	Crude OR ^c (95% CI)
Overall				
C/C	156 (69.7)	133 (75.1)	0.223	0.758 (0.486–1.182)
C/T	63 (28.1)	38 (21.5)		
T/T	5 (2.2)	6 (3.4)		
Sex				
Male				
C/C	73 (68.2)	97 (75.2)	0.235	0.708 (0.400–1.252)
C/T	32 (29.9)	28 (21.7)		
T/T	2 (1.9)	4 (3.1)		
Female				
C/C	83 (70.9)	36 (75.2)	0.597	0.813 (0.378–1.748)
C/T	31 (26.5)	10 (20.8)		
T/T	3 (2.6)	2 (4.2)		
Age (years)				
<65				
C/C	78 (67.8)	54 (71.1)	0.636	0.858 (0.456–1.613)
C/T	35 (30.4)	20 (26.3)		
T/T	2 (1.8)	2 (2.6)		
≥65				
C/C	78 (71.6)	79 (78.2)	0.267	0.701 (0.373–1.316)
C/T	28 (35.7)	18 (17.8)		
T/T	3 (2.7)	4 (4.0)		
<i>H. pylori</i> infection				
Negative	n = 42	n = 29		
C/C	32 (76.2)	25 (86.2)	0.297	0.512 (0.143–1.827)
C/T	9 (21.4)	3 (21.7)		
T/T	1 (2.4)	1 (3.5)		
Positive	n = 75	n = 44		
C/C	55 (73.3)	32 (72.7)	0.942	1.031 (0.446–2.383)
C/T	19 (25.3)	12 (27.3)		
T/T	1 (1.4)	0 (0.0)		

^aObserved genotype distribution of controls was in agreement with Hardy-Weinberg equilibrium

^bAssociation was analyzed by Fisher's exact test. P are values for C/T+T/T genotype relative to C/C genotype

^cOdds ratios are for C/T+T/T genotypes relative to C/C genotype

Association between genotypes and clinicopathological features

We next analyzed the association between the -1562 C/T genotypes and clinicopathological features among gastric cancer patients (Table 2). Gastric cancer patients with the T allele (C/T and T/T) showed deeper invasion over the submucosal (sm) layer than did patients with the C/C genotype (crude OR=2.61, 95% CI=1.07–6.33). Advanced-stage gastric cancers were significantly more frequent in patients with the T allele (C/T and T/T) than in patients with the C/C genotype (stage I vs stages II, III, and IV, crude OR=2.25, 95% CI=1.12–4.50). Furthermore, there was a significant difference in genotype distribution (C/T+T/T vs C/C) with respect to lymphatic invasion (lymphatic invasion positive vs negative, crude OR=2.27, 95% CI=1.09–4.73), and risk of venous invasion of gastric cancer was increased (enhanced risk=1.98) in patients with the T allele (venous invasion positive vs negative, crude OR=1.98, 95% CI=0.99–3.96). Histological classification (diffuse type vs intestinal type) and lymph node metastasis was not significantly associated with genotype. Logistic regression analysis revealed that genotype associations with depth of tumor invasion (adjusted OR=2.61, 95% CI=1.07–6.34), TNM classification (adjusted OR=2.26, 95% CI=1.12–4.54), and lymphatic invasion (adjusted OR=2.27, 95%

CI=1.09–4.73) remained significant even after adjustments for age and sex.

Discussion

In the present study, we examined whether risk of gastric cancer is associated with the -1562 C/T polymorphism in the promoter of *MMP-9* in a Japanese population. It has been suggested that carcinogenesis is a multicellular and multistage process in which breakdown of the microenvironment is required for conversion of normal tissue to tumor (Park et al. 2000). Although we hypothesized that the *MMP-9* polymorphism alters the microenvironment and may be involved in the process of carcinogenesis, the allele frequencies in the gastric cancer patients were similar to those in controls. Our findings suggest that the T allele does not enhance susceptibility to the development of gastric cancer. In contrast, we found a significant association between this polymorphism and clinicopathological features, specifically depth of tumor invasion, TNM classification, and lymphatic invasion. The T allele was detected more frequently in patients with advanced-stage gastric cancers than in those with the C/C genotype (OR=2.26). Moreover, lymphatic invasion was significantly enhanced in gastric cancer patients with the T allele. However, lymph node metastasis showed no signifi-

Table 2 Association between -1562 C/T polymorphism in *MMP-9* promoter and clinicopathological features of gastric cancer

	Genotype (%)			OR ^a (95%CI)	
	C/C	C/T	T/T	Crude	Adjusted ^b
Age (year)					
<65 (<i>n</i> = 76)	54 (71.1)	20 (26.3)	2 (2.6)	0.683 (0.345–1.356)	
≥65 (<i>n</i> = 101)	79 (78.2)	18 (17.8)	4 (4.0)		
Sex					
Male (<i>n</i> = 129)	97 (75.2)	28 (21.7)	4 (3.1)	0.990 (0.460–2.129)	
Female (<i>n</i> = 48)	36 (75.0)	10 (20.8)	2 (4.2)		
Histological classification ^c					
Intestinal (<i>n</i> = 103)	81 (78.6)	18 (17.5)	4 (3.9)	1.558 (0.784–3.093)	1.601 (0.778–3.297) <i>P</i> = 0.201
Diffuse (<i>n</i> = 74)	52 (70.3)	20 (27.0)	2 (2.7)		
Depth ^d					
m (<i>n</i> = 51)	44 (86.3)	6 (11.8)	1 (1.9)	2.613 (1.079–6.331)	2.610 (1.074–6.340) <i>P</i> = 0.034
sm~ (<i>n</i> = 126)	89 (70.6)	32 (25.4)	5 (4.0)		
Lymphatic invasion ^e					
Negative (<i>n</i> = 78)	65 (83.3)	12 (15.4)	1 (1.3)	2.279 (1.097–4.736)	2.274 (1.092–4.736) <i>P</i> = 0.028
Positive (<i>n</i> = 99)	68 (68.7)	26 (26.3)	5 (5.0)		
Venous invasion ^e					
Negative (<i>n</i> = 95)	77 (81.1)	16 (16.8)	2 (2.1)	1.986 (0.994–3.969)	1.984 (0.993–3.967) <i>P</i> = 0.053
Positive (<i>n</i> = 82)	56 (68.3)	22 (26.8)	4 (4.9)		
Lymph node metastasis					
N (-) (<i>n</i> = 114)	89 (78.1)	22 (19.3)	3 (2.6)	1.537 (0.765–3.088)	1.537 (0.761–3.102) <i>P</i> = 0.230
N (+) (<i>n</i> = 63)	44 (69.8)	16 (25.4)	3 (4.8)		
TNM classification ^f					
Stage I (<i>n</i> = 99)	81 (81.8)	16 (16.2)	2 (2.0)	2.250 (1.123–4.507)	2.260 (1.124–4.547) <i>P</i> = 0.022
Stage II, III, IV (<i>n</i> = 78)	52 (66.7)	22 (28.2)	4 (5.1)		
<i>H. pylori</i> infection ^g					
Negative (<i>n</i> = 29)	25 (86.2)	3 (10.3)	1 (3.5)	0.427 (0.123–1.484)	0.416 (0.117–1.473) <i>P</i> = 0.174
Positive (<i>n</i> = 44)	32 (72.7)	12 (27.3)	0 (0.0)		

^aOdds ratios (ORs) and 95% confidence intervals (CIs) for clinicopathological features with reference to the *MMP-9* promoter polymorphism (C/T+T/T to C/C genotypes)

^bAdjusted for age and sex, with a logistic regression model

^cGastric cancer classified histologically according to the criteria of Lauren

^dDepth of tumor invasion. (m tumour without invasion of the submucosa. sm~ tumour invades over the submucosal layer)

^eLymphatic invasion and venous invasion classified according to the criteria of Japanese Classification of Gastric Carcinoma, 2nd English edn

^fTNM grades were according to the criteria of the TNM Classification of Malignant Tumors, 6th edn

^g*H. pylori* status was based on histology, ¹³C-urea breath test, and serum anti-*H. pylori* IgG antibody

cantly associated with genotype. Establishment of metastasis requires the serial processes such as invasion, migration, implantation, and regrowth of cancer cells at the metastatic site. The *MMP-9* polymorphism may affect the initial invasion step of lymph node metastasis. The present findings support our hypothesis that the -1562 C/T polymorphism may have a profound impact on progression and invasion of gastric cancer. To our knowledge, this is the first study to investigate the association of the -1562 C/T polymorphism of *MMP-9* with development and malignant phenotypes of gastric cancer.

The association between the *MMP-9* polymorphism and invasive phenotype of gastric cancer is consistent with the biological function of *MMP-9*. Overexpression of *MMP-9* has been observed in a variety of cancers including gastric cancer, breast cancer, prostate cancer, brain cancer, melanoma, and lymphoma (Murray et al.

1998; Hujanen et al. 1994; Jones et al. 1999; Rao et al. 1993; Sehgal et al. 1996; Chicoine et al. 2002). In gastric cancer, expression of *MMP-9* is associated with pathological features such as TNM stage, lymphatic invasion, and tumor depth (Zhang et al. 2003). Furthermore, expression of *MMP-9* is associated with poor prognosis in renal cell carcinoma patients (Kallakury et al. 2001). In breast cancer patients, plasma *MMP-9* activity is also associated with prognosis (Baker and Leaper 2003). These findings clearly indicate that *MMP-9* plays an important role in tumor progression and invasion. Moreover, the C/T polymorphism in *MMP-9* has functional significance. The *MMP-9* promoter with the T allele shows significantly higher transcriptional activity than the C allele in cultured macrophages, and this polymorphism influences the atherosclerotic phenotype (Zhang et al. 1999). The *MMP-9* gene promoter contains binding sites for AP-1, NF-κB, Sp-1, and Ets

transcription factors (Gum et al. 1996). In particular, Ets-1 expression is upregulated together with MMP-9 (Behrens et al. 2001), and this upregulation correlates with tumor invasion in gastric cancer (Nakayama et al. 1996). This polymorphism is located within a transcription factor binding site, and that changing the allele may affect transcription factor binding and therefore expression of MMP-9 (Zhang et al. 1999). We also found the *MMP-9* polymorphism to be associated with depth of tumor invasion, lymphatic invasion, and TNM classification of gastric cancer. Taken together, these results suggest that the T allele may affect expression of MMP-9 in gastric cancer, increasing ECM degradation and leading to subsequent invasion of cancer cells.

Conclusion

In conclusion, our findings suggest that the T allele of the *MMP-9* promoter may affect expression of MMP-9 and is closely related to the invasive phenotype of gastric cancer. Recently, *in vitro* studies showed that MMP-9 production is increased more than two times by tumor necrosis factor- α (TNF- α) (Ferrand et al. 2002) and more than seven times by gastrin (Wroblewski et al. 2002). Moreover, Chicoine et al. (Chicoine et al. 2002) reported that MMP-9 expression is controlled by the methylation status of the *MMP-9* promoter. Further studies of the association between the levels of MMP-9 expression and the -1562 C/T polymorphism in gastric cancer patients are needed with increased numbers of cases to confirm our findings.

Acknowledgements We thank Masayoshi Takatani and Mutsumi Ueda, Department of Molecular Pathology, Hiroshima University Graduate School of Biomedical Sciences, for their excellent technical assistance and advice.

References

- Baker EA, Leaper DJ (2003) The plasminogen activator and matrix metalloproteinase systems in colorectal cancer: relationship to tumour pathology. *Eur J Cancer* 39:981–988
- Behrens P, Rothe M, Wellmann A, Krischler J, Wernert N (2001) The Ets-1 transcription factor is up-regulated together with MMP-1 and MMP-9 in the stroma of pre-invasive breast cancer. *J Pathol* 194:43–50
- Chicoine E, Esteve PO, Robledo O, Van Themsche C, Potworowski EF, St-Pierre Y (2002) Evidence for the role of promoter methylation in the regulation of MMP-9 gene expression. *Biochem Biophys Res Commun* 297:765–772
- Ferrand PE, Parry S, Sammel M, Macones GA, Kuivaniemi H, Romero R, Strauss JF 3rd (2002) A polymorphism in the matrix metalloproteinase-9 promoter is associated with increased risk of preterm premature rupture of membranes in African Americans. *Mol Hum Reprod* 8:494–501
- Forget MA, Desrosiers RR, Beliveau R (1999) Physiological roles of matrix metalloproteinases; implications for tumor growth and metastasis. *Can J Physiol Pharmacol* 77:465–480
- Gum R, Lengyel E, Juarez J, Chen JH, Sato H, Seiki M, Boyd D (1996) Stimulation of 92-kDa gelatinase B promoter activity by ras is mitogen-activated protein kinase kinase 1-independent and requires multiple transcription factor binding sites including closely spaced PEA3/ets and AP-1 sequences. *J Biol Chem* 271:10672–10680
- Hujanen ES, Vaisanen A, Zheng A, Tryggvason K, Turpeenniemi-Hujanen T (1994) Modulation of M(r) 72,000 and M(r) 92,000 type-IV collagenase (gelatinase A and B) gene expression by interferons alpha and gamma in human melanoma. *Int J Cancer* 58:582–586
- Japanese Gastric Cancer Association (1998) Japanese classification of gastric carcinoma. 2nd English Edition. *Gastric Cancer* 1:10–24
- Jones JL, Glynn P, Walker RA (1999) Expression of MMP-2 and MMP-9, their inhibitors, and the activator MT1-MMP in primary breast carcinomas. *J Pathol* 189:161–168
- Kallakury BV, Karikehalli S, Haholu A, Sheehan CE, Azumi N, Ross JS (2001) Increased expression of matrix metalloproteinases 2 and 9 and tissue inhibitors of metalloproteinases 1 and 2 correlate with poor prognostic variables in renal cell carcinoma. *Clin Cancer Res* 7:3113–3119
- Kohn EC, Liotta LA (1995) Molecular insights into cancer invasion: strategies for prevention and intervention. *Cancer Res* 55:1856–1862
- Kuraoka K, Matsumura S, Hamai Y, Nakachi K, Imai K, Matsusaki K, Oue N, Ito R, Nakayama H, Yasui W (2003) A single nucleotide polymorphism in the transmembrane domain coding region of HER-2 is associated with development and malignant phenotype of gastric cancer. *Int J Cancer* 107:593–596
- Lauren P (1965) The two histological main types of gastric carcinoma. Diffuse and so-called intestinal type carcinoma: an attempt at histological classification. *Acta Pathol Microbiol Scand* 64:31–49
- Liotta LA, Steeg PS, Stetler-Stevenson WG (1991) Cancer metastasis and angiogenesis: an imbalance of positive and negative regulation. *Cell* 64:327–336
- Matsumura S, Oue N, Kitadai Y, Chayama K, Yoshida K, Yamaguchi Y, Toge T, Imai K, Nakachi K, Yasui W (2004) A single nucleotide polymorphism in the MMP-1 promoter is correlated with histological differentiation of gastric cancer. *J Cancer Res Clin Oncol* 130:259–265
- McDonnell S, Matrisian LM (1990) Stromelysin in tumor progression and metastasis. *Cancer Metastasis Rev* 9:305–319
- Murray GI, Duncan ME, Arbuckle E, Melvin WT, Fothergill JE (1998) Matrix metalloproteinases and their inhibitors in gastric cancer. *Gut* 43:791–797
- Nakayama T, Ito M, Ohtsuru A, Naito S, Nakashima M, Fagin JA, Yamashita S, Sekine I (1996) Expression of the Ets-1 proto-oncogene in human gastric carcinoma: correlation with tumor invasion. *Am J Pathol* 149:1931–1939
- Nagase H, Woessner JF Jr (1999) Matrix metalloproteinases. *J Biol Chem* 274:21491–494
- Oue N, Motoshita J, Yokozaki H, Hayashi K, Tahara E, Taniyama K, Matsusaki K, Yasui W (2002) Distinct promoter hypermethylation of p16INK4a, CDH1, and RAR- β in intestinal, diffuse-adherent, and diffuse-scattered type gastric carcinomas. *J Pathol* 198:55–59
- Oue N, Oshimo Y, Nakayama H, Ito R, Yoshida K, Matsusaki K, Yasui W (2003) DNA methylation of multiple genes in gastric carcinoma: association with histological type and CpG island methylator phenotype. *Cancer Sci* 94:901–905
- Ozalp S, Tanir HM, Yalcin OT, Kabukcuoglu S, Oner U, Uray M (2003) Prognostic value of matrix metalloproteinase-9 (gelatinase-B) expression in epithelial ovarian tumors. *Eur J Gynaecol Oncol* 24:417–420
- Park CC, Bissell MJ, Barcellos-Hoff MH (2000) The influence of the microenvironment on the malignant phenotype. *Mol Med Today* 6:324–329
- Rao JS, Steck PA, Mohanam S, Stetler-Stevenson WG, Liotta LA, Sawaya R (1993) Elevated levels of M(r) 92,000 type IV collagenase in human brain tumors. *Cancer Res* 53:2208–2211
- Sakata K, Satoh M, Someya M, Asanuma H, Nagakura H, Oouchi A, Nakata K, Kogawa K, Koito K, Hareyama M, Himi T

- (2004) Expression of matrix metalloproteinase 9 is a prognostic factor in patients with non-Hodgkin lymphoma. *Cancer* 100:356–365
- Sehgal I, Baley PA, Thompson TC (1996) Transforming growth factor beta1 stimulates contrasting responses in metastatic versus primary mouse prostate cancer-derived cell lines in vitro. *Cancer Res* 56:3359–3365
- Sobin LH, Wittekind CH (eds) (2002) TNM classification of malignant tumors, 6th edn. Wiley-Liss, New York, pp 65–68
- Tanioka Y, Yoshida T, Yagawa T, Saiki Y, Takeo S, Harada T, Okazawa T, Yanai H, Okita K (2003) Matrix metalloproteinase-7 and matrix metalloproteinase-9 are associated with unfavourable prognosis in superficial oesophageal cancer. *Br J Cancer* 89:2116–2121
- Wroblewski LE, Pritchard DM, Carter S, Varro A (2002) Gastrin-stimulated gastric epithelial cell invasion: the role and mechanism of increased matrix metalloproteinase 9 expression. *Biochem J* 365:873–879
- Yasui W, Yokozaki H, Fujimoto J, Naka K, Kuniyasu H, Tahara E (2000) Genetic and epigenetic alterations in multistep carcinogenesis of the stomach. *J Gastroenterol* 35:111–115
- Zhang B, Ye S, Hermann SM, Eriksson P, de Maat M, Evans A, Arveiler D, Luc G, Cambien F, Hamsten A, Watkins H, Henney AM (1999) Functional polymorphism in the regulatory region of gelatinase B gene in relation to severity of coronary atherosclerosis. *Circulation* 99:1788–794
- Zhang S, Li L, Lin JY, Lin H (2003) Imbalance between expression of matrix metalloproteinase-9 and tissue inhibitor of metalloproteinase-1 in invasiveness and metastasis of human gastric carcinoma. *World J Gastroenterol* 9:899–904

Eiji Tahara Jr · Hidetoshi Tahara · Masamoto Kanno
Kazuhito Naka · Yayoi Takeda · Takeshi Matsuzaki
Ryuta Yamazaki · Hiroto Ishihara · Wataru Yasui
J Carl Barrett · Toshinori Ide · Eiichi Tahara

G1P3, an interferon inducible gene 6-16, is expressed in gastric cancers and inhibits mitochondrial-mediated apoptosis in gastric cancer cell line TMK-1 cell

Received: 13 August 2004 / Accepted: 17 November 2004 / Published online: 1 February 2005
© Springer-Verlag 2005

Abstract Expression of an interferon inducible gene 6-16, G1P3, increases not only in type I interferon-treated cells but also in human senescent fibroblasts. However, the function of 6-16 protein is unknown. Here we report that 6-16 is 34 kDa glycosylated protein and localized at mitochondria. Interestingly, 6-16 is expressed at high levels in gastric cancer cell lines and tissues. One of exceptional gastric cancer cell line, TMK-1, which do not express detectable 6-16, is sensitive to apoptosis induced by cycloheximide (CHX), 5-fluorouracil (5-FU) and serum-deprivation. Ectopic expression of 6-16 gene

restored the induction of apoptosis and inhibited caspase-3 activity in TMK-1 cells. Thus 6-16 protein has anti-apoptotic function through inhibiting caspas-3. This anti-apoptotic function is expressed through inhibition of the depolarization of mitochondrial membrane potential and release of cytochrome c. By two-hybrid screening, we found that 6-16 protein interacts with calcium and integrin binding protein, CIB/KIP/Calmyrin (CIB), which interacts with presenilin 2, a protein involved in Alzheimer's disease. These protein interactions possibly play a pivotal role in the regulation of apoptosis, for which further detailed analyses are need. These results overall indicate that 6-16 protein may have function as a cell survival protein by inhibiting mitochondrial-mediated apoptosis.

E. Tahara Jr · H. Tahara (✉) · Y. Takeda · T. Ide
Department of Cellular and Molecular Biology,
Division of Integrated Medical Science,
Graduate School of Biomedical Science,
Hiroshima University, 1-2-3 Kasumi, Minamiku-ku,
Hiroshima 734-8551, Japan
E-mail: toshi@hiroshima-u.ac.jp
Tel.: +81-82-2575293
Fax: +81-82-2575293

Keywords Interferon inducible protein 6-16 · G1P3 · Mitochondria · Apoptosis · CIB · Survival gene · Bcl-2

M. Kanno · H. Ishihara · E. Tahara Jr
Department of Immunology and Parasitology,
Graduate School of Biomedical Science,
Hiroshima University, 1-2-3 Kasumi, Minamiku-ku,
Hiroshima 734-8551, Japan

K. Naka · W. Yasui
Department of Molecular Pathology,
Graduate School of Biomedical Science,
Hiroshima University, 1-2-3 Kasumi, Minamiku-ku,
Hiroshima 734-8551, Japan

T. Matsuzaki · R. Yamazaki · J C. Barrett
Yakult Central Institute for Microbiological Research,
1796 Yaho, Kunitachi 186-8650, Japan

E. Tahara
Center for Cancer Research,
National Cancer Institute, Bldg. 31,
Room 3A11, 31 Center Drive, MSC-2440,
Bethesda, MD 20892-2440, USA

Radiation Effects Research Foundation, 5-2 Hijiyama Park,
Minamiku-ku, Hiroshima 732-0815, Japan

Introduction

Interferon inducible gene 6-16, G1P3, was first identified as one of the genes that are induced by interferon α and β [1]. It locates on chromosome 1p35 [2] and produces three types of mRNA by differential splicing between exon2 and exon3 [3]. The most abundant mRNA (B type) encodes a 14 kDa hydrophobic protein of 130 amino acids. Although extensive studies on the upstream regulatory region of 6-16 gene are well reported [4], little is known that the function or role of 6-16 protein in interferon treated cells. We isolated several cDNA clones that are expressed at higher level in senescent human fibroblasts than in young counterparts and found that one of these cDNA clones is 6-16 [5]. One of the characteristics of senescent human fibroblasts is that the cells do not proliferate but are viable for relatively long period of time, usually for more than a year in a