

to these two interaction models are easily derived and equivalent to those above, except that the $a(x)$ terms include the full joint-effect risk model with interaction:

$$AF_{\text{add,int}}^{\text{O}}(x|z) = 1 - \frac{\frac{1+ERO_z}{1+ERO_x+ERO_z+\Delta xz} + f_0(1 + ERO_z)}{1 + f_0(1 + ERO_z)},$$

and

$$AF_{\text{mult,int}}^{\text{O}}(x|z) = 1 - \frac{\frac{1}{(1+ERO_x) \times (1+ERO_z) \times e^{\delta xz}} + f_0(1 + ERO_z)}{1 + f_0(1 + ERO_z)}.$$

References

- [1] Greenland S 1999 Relation of probability of causation to relative risk and doubling dose: a methodologic error that has become a social problem *Am. J. Public Health* **89** 1166–9
- [2] Robins J M and Greenland S 1989 Estimability and estimation of excess and etiologic fractions *Stat. Med.* **8** 845–59
- [3] Greenland S and Robins J M 1988 Conceptual problems in the definition and interpretation of attributable fractions *Am. J. Epidemiol.* **128** 1185–97
- [4] Mosteller F 1987 Compensating for radiation-related cancers by probability of causation or assigned shares *Bull. Int. Stat. Inst.* **52** 571–7
- [5] NIH/NCI 2003 Report of the NCI-CDC Working Group to Revise the 1985 NIH Radioepidemiological Tables *NIH Publication No. 03-5387* (Bethesda, MD: National Institutes of Health/National Cancer Institute) p 118
- [6] Whittemore A S and McMillan A 1983 Lung cancer mortality among US uranium miners: a reappraisal *JNCI* **71** 489–99
- [7] Thomas D C, McNeill K G and Dougherty C 1985 Estimates of lifetime lung cancer risks resulting from Rn progeny exposure *Health Phys.* **49** 825–46
- [8] Lubin J H and Steindorf K 1995 Cigarette use and the estimation of lung cancer attributable to radon in the United States *Radiat. Res.* **141** 79–85
- [9] Little M P 2002 Comparisons of lung tumour mortality risk in the Japanese A-bomb survivors and in the Colorado Plateau uranium miners: support for the ICRP lung model *Int. J. Radiat. Biol.* **78** 145–63
- [10] Seiler F A and Scott B R 1987 Mixtures of toxic agents and attributable risk calculations *Risk Anal.* **7** 81–90
- [11] Luebeck E G, Heidenreich W F, Hazelton W D, Paretzke H G and Moolgavkar S H 1999 Biologically based analysis of the data for the Colorado uranium miners cohort: age, dose and dose-rate effects *Radiat. Res.* **152** 339–51
- [12] Hazelton W D, Luebeck E G, Heidenreich W F and Moolgavkar S H 2001 Analysis of a historical cohort of Chinese tin miners with arsenic, radon, cigarette smoke, and pipe smoke exposures using the biologically based two-stage clonal expansion model *Radiat. Res.* **156** 78–94
- [13] Land C E, Hayakawa N, Machado S G, Yamada Y, Pike M C, Akiba S and Tokunaga M 1994 A case-control interview study of breast cancer among Japanese A-bomb survivors. II. Interactions with radiation dose *Cancer Causes Control* **5** 167–76
- [14] Beyea J and Greenland S 1999 The importance of specifying the underlying biologic model in estimating the probability of causation *Health Phys.* **76** 269–74
- [15] Fujiwara S, Kusumi S, Cologne J, Akahoshi M, Kodama K and Yoshizawa H 2000 Prevalence of anti-hepatitis C virus antibody and chronic liver disease among atomic bomb survivors *Radiat. Res.* **154** 12–9
- [16] Wong F L, Yamada M, Sasaki H, Kodama K, Akiba S, Shimaoka K and Hosoda Y 1993 Noncancer disease incidence in the atomic bomb survivors: 1958–1986 *Radiat. Res.* **135** 418–30
- [17] Cologne J B, Tokuoka S, Beebe G W, Fukuhara T and Mabuchi K 1999 Effects of radiation on incidence of primary liver cancer among atomic bomb survivors *Radiat. Res.* **152** 364–73
- [18] Neriishi K, Akiba S, Amano T, Ogino T and Kodama K 1995 Prevalence of hepatitis B surface antigen, hepatitis B e antigen and antibody, and antigen subtypes in atomic bomb survivors *Radiat. Res.* **144** 215–21
- [19] Sharp G B, Mizuno T, Cologne J B, Fukuhara T, Fujiwara S, Tokuoka S and Mabuchi K 2003 Hepatocellular carcinoma among atomic bomb survivors: significant interaction of radiation with hepatitis C virus infections *Int. J. Cancer* **103** 531–7

- [20] Sharp G B, Cologne J B, Fukuhara T, Itakura H, Yamamoto M and Tokuoka S 2001 Temporal changes in liver cancer incidence rates in Japan: accounting for death certificate inaccuracies and improving diagnostic techniques *Int. J. Cancer* **93** 751–8
- [21] Yoshizawa H 2002 Hepatocellular carcinoma associated with hepatitis C virus infection in Japan: projection to other countries in the foreseeable future *Oncology* **62** (Suppl. 1) 8–17
- [22] Greenland S and Rothman K J 1998 Concepts of interaction *Modern Epidemiology* 2nd edn, ed K J Rothman and S Greenland (Philadelphia, PA: Lippincott Williams&Wilkins) pp 329–42
- [23] Donato F *et al* 2002 Alcohol and hepatocellular carcinoma: the effect of lifetime intake and hepatitis virus infections in men and women *Am. J. Epidemiol.* **155** 323–31
- [24] McCullagh P and Nelder J A 1989 *Generalized Linear Models* 2nd edn (London: Chapman and Hall)
- [25] Thomas D C 1981 General relative-risk models for survival time and matched case-control analysis *Biometrics* **37** 673–86
- [26] Preston D L, Shimizu Y, Pierce D A, Suyama A and Mabuchi K 2003 Studies of mortality of atomic bomb survivors. Report 13: solid cancer and noncancer disease mortality: 1950–1997 *Radiat. Res.* **160** 381–7
- [27] Committee on Assessment of Centers for Disease Control and Prevention Radiation Studies from DOE Contractor Sites: Subcommittee to Review Radioepidemiological Tables, Board on Radiation Effects Research, Commission on Life Sciences, National Academy of Sciences 2000 *A Review of the Draft Report of the NCI-CDC Working Group to Revise the 1985 Radioepidemiological Tables* (Washington, DC: National Academy Press)
- [28] Yang Q, Khoury M J, Friedman J M and Flanders W D 2003 On the use of population attributable fraction to determine sample size for case-control studies of gene-environment interaction *Epidemiology* **14** 161–7
- [29] Seiler F A 1986 Assigned shares and combined insults *Risk Anal.* **6** 371–2

A meta-analysis of previous fracture and subsequent fracture risk

J.A. Kanis,^{a,*} O. Johnell,^b C. De Laet,^c H. Johansson,^d A. Oden,^d P. Delmas,^e J. Eisman,^f
S. Fujiwara,^g P. Garnero,^c H. Kroger,^h E.V. McCloskey,^a D. Mellstrom,ⁱ L.J. Melton,^j
H. Pols,^c J. Reeve,^k A. Silman,^l and A. Tenenhouse^m

^aWHO Collaborating Centre for Metabolic Bone Diseases, WHO Collaborating Centre, University of Sheffield Medical School, Sheffield, UK

^bDepartment of Orthopaedics, Malmö General Hospital, Sweden

^cDepartment of Public Health, Erasmus Medical Center, Rotterdam, The Netherlands

^dConsulting Statistician, Gothenburg, Sweden

^eINSERM Unite 403, Hopital Edouard Herriot, Lyon, France

^fBone and Mineral Research, Garvan Institute of Medical Research, NSW, Australia

^gRadiation Effects Research Foundation, Hiroshima, Japan

^hDepartment of Surgery, Kuopio University Hospital, Kuopio, Finland

ⁱDepartment of Geriatric Medicine, University of Goteborg, Sweden

^jDivision of Epidemiology, Mayo Clinic, Rochester, MN, USA

^kStrangeways Research Laboratories, Cambridge, UK

^lARC Epidemiology Research Unit, University of Manchester, Manchester, UK

^mDivision of Bone Metabolism, The Montreal General Hospital, Montreal, Canada

Received 15 January 2004; revised 17 March 2004; accepted 24 March 2004

Abstract

Previous fracture is a well-documented risk factor for future fracture. The aim of this study was to quantify this risk on an international basis and to explore the relationship of this risk with age, sex, and bone mineral density (BMD). We studied 15 259 men and 44 902 women from 11 cohorts comprising EVOS/EPOS, OFELY, CaMos, Rochester, Sheffield, Rotterdam, DOES, Hiroshima, and two cohorts from Gothenburg. Cohorts were followed for a total of 250 000 person-years. The effect of a prior history of fracture on the risk of any fracture, any osteoporotic fracture, and hip fracture alone was examined using a Poisson model for each sex from each cohort. Covariates examined were age, sex, and BMD. The results of the different studies were merged by using the weighted β -coefficients.

A previous fracture history was associated with a significantly increased risk of any fracture compared with individuals without a prior fracture (RR = 1.86; 95% CI = 1.75–1.98). The risk ratio was similar for the outcome of osteoporotic fracture or for hip fracture. There was no significant difference in risk ratio between men and women. Risk ratio (RR) was marginally downward adjusted when account was taken of BMD. Low BMD explained a minority of the risk for any fracture (8%) and for hip fracture (22%). The risk ratio was stable with age except in the case of hip fracture outcome where the risk ratio decreased significantly with age.

We conclude that previous history of fracture confers an increased risk of fracture of substantial importance beyond that explained by measurement of BMD. Its validation on an international basis permits the use of this risk factor in case finding strategies.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Prior fracture; Meta-analysis; Hip fracture; Osteoporotic fracture

Introduction

It is well established from many cohort, case-control, and cross-sectional studies that a prior osteoporotic fracture

increases the risk of future fractures [1–8]. A prior forearm fracture is associated with about a twofold increase in the subsequent risk of fracture [9–13]. More recently, significant increases in risk have been described for prior fractures at other sites characteristic of osteoporosis [6,7,14–21]. The risk of another vertebral fracture is particularly high after a spine fracture [7,22–24]. Similar observations are found in the setting of randomized clinical trials. In the placebo arm, the risk of vertebral deformities is approximately fivefold

* Corresponding author. WHO Collaborating Centre for Metabolic Bone Diseases, University of Sheffield Medical School, Beech Hill Road, Sheffield S10 2RX, UK. Fax: +44-114-285-1813.

E-mail address: w.j.Pontefract@shef.ac.uk (J.A. Kanis).

higher in patients with a prior vertebral deformity than in those without [4,25,26]. The interrelationships between the site of prior fracture and site of subsequent fracture have been summarized by meta-analysis [27] and a large case-control study, published more recently found broadly similar relationships [8].

Increased fracture risk may be in part due to the fact that patients with fracture have low bone mineral density (BMD). Studies that have adjusted for BMD suggest that the relative risk is only modestly downward adjusted [3,20,24,28–31].

The consistent association between a prior fracture and subsequent fracture risk has led to the inclusion of prior fracture as a risk factor to be used in assessment guidelines [32–35]. For example, in Europe, it is recommended that patients be identified on the basis of risk factors for subsequent assessment by BMD [33–35]. Patients are then considered for intervention on the finding of osteoporosis (i.e., a *T* score of ≤ -2.5 SD). This approach is conservative since it does not recognize the independent contribution of the risk factor from BMD. This has been recognized in some guidelines where the intervention threshold is less conservative in the presence of a risk factor such as a prior fragility fracture [32,36]. The consideration of several independent risk factors permits the more accurate categorization of risk [37], and attention has focussed recently on the assessment of fracture risk using multiple risk factors, rather than the use of BMD alone, to define intervention thresholds [38,39]. This demands knowledge of the interrelationships between these risk factors.

The aim of the present study was to quantify the risk associated with a history of prior fracture for future fracture in an international setting and to explore the dependence of this risk with age, sex, and BMD.

Methods

We studied 60 161 men and women of whom 26% had a prior fracture history taken from 11 prospectively studied

cohorts. Brief details of the cohorts studied are given below and summarized in Table 1.

OFELY

The OFELY cohort comprises an age-stratified cohort of 1039 women aged 31–89 years randomly selected from the regional section of a large health insurance company (Mutuelle Generale d'Education Nationale, Lyon) [40]. Eighteen percent of women contacted participated in the study. Baseline characteristics were obtained using a standardized questionnaire, including the documentation of prior wrist, humeral, vertebral, and hip fracture that occurred after the age of 45 years. Only low trauma fractures (falls from a standing height or less) were recorded. BMD was measured at the lumbar spine, at the proximal femur, distal radius, and whole body by DXA using a Hologic QDR 2000. Women were reviewed annually and fractures registered. Peripheral fractures were confirmed by radiography. Vertebral fractures were identified from sequential X-rays of the thoracic and lumbar spine by morphometry in 80% of patients, but only clinical fractures were used for this analysis.

EVOS

The European Vertebral Osteoporosis Study (EVOS) comprised age- and sex-stratified random samples from 36 centers in 19 European countries [41–43]. Equal numbers of men and women were drawn in each center within six 5-year age bands (50–54 to 75–79 years). A baseline radiograph for vertebral fracture prevalence was undertaken in 15570 men and women (response rate, 29%). BMD was measured in 3461 men and women from 13 centers by DXA at the femoral neck using pencil beam machines that were cross-calibrated using the European spine phantom. The sample provided the framework for the European Prospective Osteoporosis study (EPOS) where repeated assessment was undertaken in 29 of the centers. Information on clinical fractures were used for this report.

Table 1

Details of cohorts studied including individuals with information on follow-up time, prior fracture, and subsequent fracture

Cohort	Sample size	Person-years	Mean age (years)	Age range (years)	% Female	Fracture history (%)	Any fracture	Osteoporotic fracture	Hip fracture
EPOS	13366	40160	63.8	41–91	52	36	715	715	44
OFELY	426	2124	64.2	50–89	100	16	53	–	–
CaMos	9400	26653	62.1	25–103	69	44	586	316	42
Rochester	1001	6228	56.8	21–94	65	18	289	244	42
Sheffield	2147	6826	80.0	74–96	100	51	284	236	62
Rotterdam	7774	43606	70.3	55–106	61	14	992	768	284
Kuopio	11798	56602	52.3	47–57	100	17	1053	–	–
Gothenburg I	2375	16439	78.8	69–86	61	9	431	431	336
Gothenburg II	7098	29750	58.9	21–89	100	18	441	312	29
Dubbo	2163	16333	70.7	57–96	61	15	532	418	107
Hiroshima	2613	9861	65.1	47–95	70	26	187	90	32
Totals	60161	254582	62.9	21–106	75	26	5563	3530	978

CaMos

The Canadian Multicentre Osteoporosis Study (CaMos) is an ongoing prospective age-stratified cohort. The study is documenting the incidence of fractures and risk factors in a random sample of 9424 men and women aged 25 years or more selected by telephone listings. The sampling frame is from nine study centers in nine provinces [44]. Characterization of individuals was by interview. BMD was measured by DXA at the hip (Hologic QDR 1000) and lumbar spine, and an ultrasound scan taken at the heel in individuals aged 50 years or more.

Rochester

The Rochester cohort was recruited from two random population samples stratified by decade of age, one comprising women who were subsequently followed for up to 20 years [45] and another sample of women and men followed for 8 years [46]. The response rates were 49.8% in the women and 38.7% in the men. BMD of the right femoral neck was measured by dual photon absorptiometry in the first cohort (cross calibrated to DXA) and by DXA (Hologic QDR 2000) in the second group. Fractures were ascertained by periodic interview combined with review of the inpatient and outpatient medical records of all local care providers.

Sheffield

The Sheffield cohort comprised women aged 75 years or more selected randomly from the population of Sheffield, UK, and surrounding districts between 1993 and 1999. Approximately 35 000 women, identified from general practitioner listings, were contacted by letter and invited to attend for assessment of their skeletal status. Five thousand eight hundred and seventy-three women were willing to attend for the screening visit. Of these, 281 were excluded, and the remainder randomly allocated after informed consent to treatment with the bisphosphonate clodronate, or to an identical placebo. This study is still in progress, and the material used for the present paper comprised 2148 women allocated to treatment with placebo [47]. All women had baseline assessment of BMD undertaken at the femoral neck using the Hologic 4500. Outcomes were assessed by 6 monthly home visits.

Rotterdam

The Rotterdam study, begun in 1990, is a prospective cohort study that aimed to examine and follow-up all residents aged 55 years and older living in Ommoord, a district of Rotterdam [48]. By 1993, 7983 residents had been included (response rate, 78%). Bone mineral density was assessed at the femoral neck by DXA using a Lunar DPX-L [49]. Fracture follow-up was undertaken using an automated link with general practitioner computer systems

and hospital admission data. Fracture data were collected and validated by two independent research physicians. For this analysis, validated fracture follow up was available for 7774 participants (3065 men) with an average follow up time of 6 years. Femoral neck BMD was measured in 5776 individuals (2432 men).

Kuopio

The Kuopio osteoporosis risk factor and prevention (OSTPRE) study in Finland comprised a postal inquiry sent to all 14220 women aged 47–56 who were residents of Kuopio province in 1989. Thirteen thousand and one hundred women responded to the inquiry, of whom 1214 were excluded for incomplete information. This left a study population of 11 886 women. A random stratified sample of 3222 women underwent bone mineral densitometry by DXA using the Lunar DPX [50].

Gothenburg I

This study comprised four birth cohorts of 2375 randomly sampled men and women aged 70 years or more followed for up to 20 years at Gothenburg [51,52] after a baseline BMD measurement. The participation rate was 73%. The participants were drawn randomly from the population register in Gothenburg by the date of birth to provide cohorts aged 70, 76, 79, and 85 years at the time of investigation. Bone mineral density was measured at the right heel using dual photon absorptiometry.

Gothenburg II

The Gothenburg study comprised a randomly drawn population cohort of women aged 21–89 years followed up to 7.9 years (mean 4.2 years) [53]. Seventy percent of those invited (approximately 7000 women) participated in the study that examined risk factors for osteoporosis by use of a standardized questionnaire. BMD was assessed at baseline at the distal forearm using the Osteometer DTX 200.

DOES

The Dubbo Osteoporosis Epidemiology Study (DOES) is a population-based study with multiple assessments of skeletal status in men and women aged 60 years or more from Dubbo, Australia [54]. Participation in the study was 56% of the population. Baseline measurements included BMD at the femoral neck assessed using DXA (GE-Lunar, DPX). Fractures are identified through radiologists reports from the two centers servicing the region.

Hiroshima

The Adult Health Study in Hiroshima (AHS) documents the late health effects of radiation exposure among atomic

Table 2
Details of the construct of the questionnaire on fracture history in the cohorts studied

Cohort	Question
EVOS/EPOS	Have you ever suffered a fracture?
CaMos	Have you ever fractured any bones?
Rochester	Prior fracture with moderate trauma
Rotterdam	History of any fracture in the past 5 years
Gothenburg I	Evidence of any fracture (in hospital records)
OFELY	History of a fragility fracture from low trauma since age of 30 years
Sheffield	History of any fracture
Kuopio	History of any fracture from the age of 15 years
Gothenburg II	History of fracture after the age of 25 years
Dubbo	Self-reported
Hiroshima	Self-reported

bomb survivors in Hiroshima and Nagasaki. The original AHS cohort consisted of about 15 000 atomic bomb survivors and 5000 controls selected from residents in Hiroshima and Nagasaki using the 1950 national census supplementary schedules and the Atomic Bomb Survivors Survey. AHS subjects have been followed through biennial medical examinations since 1958 with a participation rate of approximately 80% throughout this period. BMD at the lumbar spine and proximal femur has been measured at each biennial health examination using DXA (Hologic QDR-2000) since December 1993. Trained nurses interviewed the subjects about baseline risk factors and measured height and weight at each biennial visit [55,56].

Baseline and outcome variables

The construct of the question to determine a prior fracture history differed between the cohorts studied (Table 2). Prospective fracture ascertainment was undertaken by self-report (Sheffield, Kuopio, EVOS/EPOS; Hiroshima) and/or verified from hospital central data bases (CaMos, Sheffield, EVOS/EPOS, Rochester, Rotterdam, Kuopio, Gothenburg I and II, CaMos, DOES). The EPOS and OFELY study also included sequential systematic radiography to define incident vertebral deformities, but were not used in this analysis. Information on all clinical fractures was used for this report. In addition, fractures considered to be due to osteoporosis were analyzed, and finally, hip fracture alone was considered separately. For Kuopio and OFELY, all fractures were recorded and no distinction made between fracture sites. In the case of the EPOS study and Gothenburg I, osteoporotic fractures only were recorded. In the other cohorts, an osteoporotic fracture was one considered to be due to osteoporosis either by the investigator or by the Coordinating Centre. For the EVOS study, osteoporotic fractures comprised hip, forearm, humeral, or limb fractures. For the CaMos study, they comprised fractures of the spine, pelvis, ribs, distal forearm, forearm, and hip. In the other cohorts (Sheffield, Rotterdam, Rochester, Gothenburg I, DOES,

Hiroshima), fractures at sites considered to be characteristic for osteoporosis were [38] extracted from the data.

Statistical methods

The risk of fracture was estimated by Poisson regression applied to each cohort and each sex separately. Covariates included time since start of follow up, current age, prior history of fracture, and BMD. We additionally excluded BMD from the model. A further model included the interaction term prior fracture \times time to determine whether the strength of the association of prior fracture and fracture risk waned with time. The beta value for each sex in each cohort is age-dependent, $\beta_{k+1} \cdot \text{age}$. The estimated value of $\beta_{k+1} \cdot \text{age}$ was determined for each age from 50 to 85 years, together with the variance. Results of each cohort and the two sexes were weighted according to the variance and merged to determine the weighted means and standard deviations. The risk ratio (RR) of those with a prior fracture history versus those without a prior fracture history was equal to e^{mean} .

The component of the risk ratio explained by BMD was computed from a meta-analysis of BMD and fracture risk [57]. The risk of any fracture was assumed to increase 1.6-fold for each SD decrease in BMD. For hip fracture, the gradient of risk was assumed to be 2.6 per SD. The proportion of risk attributed to a low BMD was computed as

$$\frac{[\log RR_a / \log GR] - [\log RR_b / \log GR]}{[\log RR_a / \log GR]}$$

where RR_a is the unadjusted risk ratio, RR_b is the risk ratio adjusted for BMD, and GR is the gradient of risk.

Results

Of 60 161 men and women studied, 877 men and 4686 women were identified as having a subsequent fracture of any kind, of which 680 and 2850 were characterized as osteoporotic in men and women, respectively. Two hundred and eleven men and 767 women sustained a hip fracture. The total follow-up was 61 938 person years in men and 192 644 in women. BMD measurements were available in 62% of individuals.

Table 3
Prevalence of a prior fracture history in men and women by age

Age (years)	Probability of fracture history (%)		
	Men	Women	Combined
30	44	15	24
40	43	18	27
50	42	23	30
60	41	29	34
70	40	35	37
80	39	41	41
90	38	48	45

Table 4
Risk ratio (RR) and 95% confidence interval (CI) of fracture associated with a history of prior fracture in men and women, without and with adjustment for BMD

Outcome fracture	Men		Women		Combined	
	RR	95% CI	RR	95% CI	RR	95% CI
A. Without BMD						
Any	2.02	1.73–2.38	1.84	1.72–1.96	1.86	1.75–1.98
Osteoporotic	1.93	1.61–2.33	1.85	1.70–2.01	1.86	1.72–2.01
Hip	2.30	1.56–3.41	1.77	1.49–2.11	1.85	1.58–2.17
B. With BMD						
Any	2.04	1.67–2.48	1.73	1.59–1.88	1.77	1.64–1.91
Osteoporotic	1.91	1.50–2.43	1.74	1.57–1.92	1.76	1.60–1.93
Hip	1.97	1.12–3.48	1.56	1.23–1.98	1.62	1.30–2.01

Probability of fracture history rose almost linearly with age (Table 3). The probability of recording a history of a prior fracture was higher in men than in women (OR = 1.19; 95% CI = 1.14–1.25).

Risk of any fracture

Previous fracture was associated with a significantly increased risk of any subsequent fracture (Table 4). There was no difference in the risk ratio between men and women. In men and women combined, the risk ratio ranged from 1.83 to 2.03 depending upon age (Table 5). The risk ratio was marginally lower by approximately 10% when account was taken of BMD. If it is assumed that the risk of any fracture increases 1.60-fold for each SD deviation decrease in hip BMD, then the difference in risk between those with and without a prior fracture is equal to an expected difference in BMD of 1.32 SD [$\log(1.86)/\log(1.60)$]. In reality, the difference in BMD at all ages in men and women combined was approximately 0.11 SD ($[\log(1.86)/\log(1.60)] - [\log(1.77)/\log(1.60)]$). Thus, low BMD accounts for the minority (8%; 0.11/1.32) of the difference in risk between those with or without a prior fracture.

Table 5
Risk ratio (RR) for any fracture and 95% confidence intervals (CI) comparing men and women with and without a previous fracture by age, with or without adjustment for BMI

Age (years)	RR without BMD ^a		RR with BMD ^a	
	Mean	95% CI	Mean	95% CI
50	1.92	1.63–2.20	1.91	1.59–2.29
55	1.90	1.73–2.09	1.83	1.60–2.10
60	1.98	1.80–2.18	1.94	1.73–2.17
65	2.02	1.86–2.20	1.99	1.81–2.20
70	2.03	1.87–2.21	1.98	1.79–2.18
75	1.96	1.80–2.13	1.82	1.65–2.02
80	1.88	1.72–2.06	1.72	1.54–1.91
85	1.83	1.65–2.04	1.72	1.51–1.96
All ages	1.86	1.75–1.98	1.77	1.64–1.91

^a Prior fracture versus no fracture.

History of previous fracture and the risk of osteoporotic fracture




Fig. 1. Risk ratio for an osteoporotic fracture in men and women with a prior history of fracture with and without adjustment for BMD.

Risk of osteoporotic fracture

Previous fracture was also associated with a significantly increased risk of an osteoporotic fracture at all ages with and without adjustment for BMD (Table 4). The unadjusted risk ratios for an osteoporotic fracture were almost identical with the risks of a prior fracture for any fracture. For example, at the age of 80 years, the risk of any fracture was 1.88 (95% CI = 1.72–2.06) and for an osteoporotic fracture was 1.89 (95% CI = 1.72–2.09). There was no difference in risk ratio between men and women. Fracture risk decreased somewhat with age by about 10% per decade of age (Fig. 1), but the trend was short of conventional significance ($P = 0.089$).

Risk of hip fracture

A prior fracture history was a significant risk factor for hip fracture at all ages (Table 6). The risk ratio was highest at younger ages and decreased progressively with age ($P < 0.002$ for the interaction term). The risk decreased by 3%

Table 6
Risk ratio (RR) for hip fracture and 95% confidence intervals (CI) comparing men and women with and without a prior fracture by age, with and without BMD

Age (years)	RR without BMD ^a		RR with BMD ^a	
	Mean	95% CI	Mean	95% CI
50	5.04	2.66–9.56	3.88	1.79–8.43
55	4.20	2.46–7.15	3.98	2.08–7.62
60	3.40	2.21–5.24	3.16	1.88–5.32
65	2.60	1.85–3.64	2.28	1.52–3.41
70	2.31	1.76–3.02	1.90	1.37–2.65
75	2.14	1.71–2.68	1.64	1.24–2.17
80	1.90	1.58–2.28	1.41	1.12–1.78
85	1.66	1.39–1.98	1.32	1.04–1.68
All ages	1.85	1.58–2.17	1.62	1.30–2.01

^a Prior fracture versus no fracture.

(95% CI = 1–5%) for each year of age. The RR was significantly increased at all ages, but at ages less than 60 years, the confidence estimates were wide (very few hip fractures). There was no difference in RR between men and women. Adjustment for BMD had an effect on the risk estimate for hip fracture that was quantitatively greater than for all fractures. The RR adjusted for BMD fell by approximately 30%. As in the case of all fractures, differences in BMD explained a minority of the increased risk ratio for hip fracture. In men and women combined, low BMD explained 22% of the increase in risk ratio and was constant by age (assuming a gradient of risk for hip fracture of 2.6/SD decrease in BMD).

The exclusion of data from Gothenburg (where BMD was assessed at the forearm or heel) had no material effect on these results (data not shown). There was no significant interaction of fracture history with time since baseline assessment.

Discussion

The present study confirms that a history of prior fracture is a significant risk factor for future fractures. In addition, the effect is over and above that which can be explained by variations in BMD. The risk of subsequent fractures is not as great as that identified in some studies [8], but as expected, falls within the confidence estimates of most estimates [27]. Discrepancies may be related to the duration of follow-up since the risk of subsequent fracture may not be linear over time [4,5]. Other possible reasons may relate to differences in the populations studied and the questionnaire used to identify prior fractures. A particular strength of the present study is that the estimate of risk is made in an international setting from randomly selected population cohorts. Calculations were based on the primary data, decreasing the risk of publication biases. The consistency of the association between cohorts (data not shown) additionally indicates the international validity of the importance of this risk factor. The risk of any subsequent fracture was comparable to the risk of a new osteoporotic fracture or a hip fracture. The large sample size permitted the quantification of risk by age. For all fractures and for osteoporotic fractures, the risk ratios were relatively constant with age. In the case of hip fracture, risk ratios decreased with age.

The present study also quantifies the independent contributions of low BMD and prior fracture. At all ages, low BMD explained a minority of the total risk, a proportion that decreased with age. The mechanism for the BMD-independent increase in risk could not be determined from this study but is likely due, in part, to coexisting morbidity that might increase the risk of falls or impair the protective responses to injury [6,21,29,58]. In addition, changes in the microarchitecture of cancellous and cortical bone with rapid bone loss after fracture or immobilization [59–61] may weaken the

resistance to mechanical force out of proportion to any effect on BMD.

Irrespective of the underlying mechanism, these data indicate that the risk of fractures is substantially greater in individuals with a prior fragility fracture than in individuals of the same age, sex, and BMD without such a fracture. This has important implications for intervention thresholds. Health economic analysis suggests that intervention is cost-effective when treatment is targeted to women without a prior fracture with a *T* score of -2.5 SD at the femoral neck [39]. Since a prior fracture confers a risk over and above that provided by BMD, intervention thresholds for BMD can be less stringent (say at a *T* score of -1.5 SD) for those with a prior fracture, and still yield the same cost effectiveness. This approach has been incorporated into health economic analyses [32,62]. However, a large number of additional independent risk factors for fracture have been identified. These include smoking, corticosteroid exposure, a family history of fracture, secondary osteoporosis [39], and possibly the biochemical indices of bone turnover [63]. The interrelationships of all these risk factors will need to be determined before they can be easily used for assessing fracture risk in the general population.

The majority of reports have provided risk ratios for fracture in those with a prior fracture history compared to those without. For practical use, it is appropriate to express risk relative to the general population since risks expressed in this way can be more readily adjusted with other risk factors, such as the risk provided by BMD measurements [64]. For this purpose, knowledge of the prevalence of the risk factor is required. The adjustment decreases the risk ratios by a factor proportional to the prevalence of the risk factor. The relative risk is computed as $RR/[p \cdot RR + (1 - p)]$ where RR is the risk ratio and *p* the prevalence of the risk factor [64]. Since the prevalence of fracture history is high (26%), the quantitative effect of adjustment is substantial.

The present study has some limitations that should be mentioned. As with nearly all randomly drawn populations, nonresponse biases may have occurred, which we were unable to document for all cohorts. The effect is likely to exclude sicker members of society, and may underestimate the absolute risk of fracture. Thus, the probability of a prior fracture may be underestimated from a societal perspective (Table 3), but this is unlikely to affect risk ratios. The greatest problem is the construct of the question concerning prior fractures and the methods of documenting and characterizing subsequent fracture events. These differed substantially between cohorts. The effect of this heterogeneity is likely, however, to weaken rather than to strengthen the associations that we found, that is, the association in reality may be stronger.

We conclude that prior fracture confers a substantial risk for future fractures and that this risk is largely independent of BMD. The consistency of the association in an international setting provides the rationale for the use of this risk factor in case of finding strategies. Moreover, patients

identified can be targeted for treatment at a higher BMD than individuals of the same age without a fracture history.

Acknowledgments

We are grateful to the International Society for Clinical Densitometry, the National Osteoporosis Foundation, the International Osteoporosis Foundation, and the European Community (EU FP 3/5). We are also grateful to Lilly, Hologic, GE Lunar, Roche, Pfizer, IGEA, the Alliance for Better Bone Health, Novartis, Wyeth, for their unrestricted support of this work.

References

- [1] Cummings SR, Melton LJ. Epidemiology and outcomes of osteoporotic fractures. *Lancet* 2002;359:1761–7.
- [2] Robinson CM, Royds M, Abraham A, McQueen MM, Court-Brown CM, Christie J. Refractures in patients at least forty-five years old. A prospective analysis of twenty-two thousand and sixty patients. *J Bone Joint Surg Am* 2002;84:1528–33.
- [3] Nevitt MC, Ross PD, Palermo L, Musliner T, Genant HK, Thompson DE. Association of prevalent vertebral fractures, bone density and alendronate treatment with incident vertebral fractures: effect of number and spinal location of fractures. *Bone* 1999;25:613–9.
- [4] Lindsay R, Silverman SL, Cooper C, Hanley DA, Barton I, Broy SB, et al. Risk of new vertebral fracture in the year following a fracture. *JAMA* 2001;285:320–3.
- [5] Johnell O, Oden A, Caullin F, Kanis JA. Acute and long-term increase in fracture risk after hospitalisation for vertebral fracture. *Osteoporosis Int* 2001;12:207–14.
- [6] Gunnes M, Mellstrom D, Johnell O. How well can a previous fracture indicate a new fracture? A questionnaire study of 29,802 postmenopausal women. *Acta Orthop Scand* 1998;69:508–12.
- [7] Ross PD, Davis JW, Epstein RS, Wasnich RD. Pre-existing fractures and bone mass predict vertebral fracture incidence in women. *Ann Intern Med* 1991;114:919–23.
- [8] Van Staa TP, Leufkens HGM, Cooper C. Does a fracture at one site predict later fractures at other sites? A British cohort study. *Osteoporosis Int* 2002;13:624–9.
- [9] Gay JD. Radial fracture as an indicator of osteoporosis: a 10-year follow-up study. *Can Med J Assoc* 1974;111:156–7.
- [10] Mallmin H, Ljunghall S, Persson I, et al. Fracture of the distal forearm as a forecaster of subsequent hip fracture: a population based cohort study with 24 years of follow-up. *Calcif Tissue Int* 1993;52:269–72.
- [11] Lauritzen JB, Schwarz P, McNair P, et al. Radial and humeral fractures as predictors of subsequent hip, radial or humeral fractures in women, and their seasonal variation. *Osteoporosis Int* 1993;3:133–7.
- [12] Cuddihy MT, Gabriel SE, Crowson CS, O'Fallon WM, Melton LJ. Forearm fractures as predictors of subsequent osteoporotic fractures. *Osteoporosis Int* 1999;9:469–75.
- [13] Finsen V, Benum P. Colles fracture as an indicator of increased risk of hip fracture. An epidemiological study. *Ann Chir Gynaecol* 1987;76:114–8.
- [14] Kotowicz MA, Melton LJ, Cooper C, Atkinson EJ, O'Fallon WM, Riggs BL. Risk of hip fracture in women with vertebral fracture. *J Bone Miner Res* 1994;9:599–605.
- [15] Lauritzen JB, Lund B. Risk of hip fracture after osteoporosis fractures: 451 women with fracture of the lumbar spine, olecranon, knee or ankle. *Acta Orthop Scand* 1993;64:297–300.
- [16] Melton LJ, Atkinson EJ, Cooper C, O'Fallon WM, Riggs BL. Vertebral fractures predict subsequent fractures. *Osteoporosis Int* 1999;10:214–21.
- [17] Tuppurainen M, Kroger H, Honkanen R, Puntilla E, Huopio J, Saarikoski S, et al. Risks of perimenopausal fractures—A prospective population-based study. *Acta Obstet Gynecol Scand* 1995;74:624–8.
- [18] Black DM, Arden NK, Palermo L, Pearson J, Cummings SR, For the Study of Osteoporotic Fractures Research Group. Prevalent vertebral deformities predict hip fractures and new vertebral deformities but not wrist fractures. *J Bone Miner Res* 1999;14:821–8.
- [19] Honkanen R, Tuppurainen M, Kroger H, Alhava E, Puntilla E. Associations of early premenopausal fractures with subsequent fractures vary by sites and mechanisms of fractures. *Calcif Tissue Int* 1997;60:327–31.
- [20] Burger H, van Dade PLA, Algra D, Hofman A, Grobbee DE, Schutte HA, et al. Vertebral deformities as predictors of non-vertebral fractures. *Br Med J* 1994;309:991–2.
- [21] Karlsson MK, Hasserijs R, Obrant KJ. Individuals who sustain non-osteoporotic fractures continue to also sustain fragility fractures. *Calcif Tissue Int* 1993;53:229–31.
- [22] Davis JW, Grove JS, Wasnich RD, et al. Spatial relationships between prevalent and incident spine fractures. *Bone* 1999;24:261–4.
- [23] Ross PD, Genant HK, Davis JW, Miller PD, Wasnich RD. Predicting vertebral fracture incidence from prevalent fractures and bone density among non-black, osteoporotic women. *Osteoporosis Int* 1993;3:120–6.
- [24] Wasnich RD, Davis JW, Ross PD. Spine fracture risk is predicted by non-spine fractures. *Osteoporosis Int* 1994;4:1–5.
- [25] Black DM, Cummings SR, Karpf DB, Cauley JA, Thompson DE, Nevitt MC. Randomised trial of effect of alendronate on risk of fracture in women with existing vertebral fractures. Fracture intervention trial research group. *Lancet* 1996;348:1535–41.
- [26] Cummings S, Black D, Thompson DE, Applegate WB, Barrett C, Musliner TA. Effect of alendronate on risk of fracture in women with low bone density but without vertebral fractures. *JAMA* 1998;280:2077–82.
- [27] Klotzbuecher CM, Ross PD, Landsman PB, Abbot TA, Berger M. Patients with prior fractures have an increased risk of future fractures: a summary of the literature and statistical synthesis. *JBM* 2000;15:721–39.
- [28] Nguyen TV, Eisman JA, Kelly PJ, Sambrook PN. Risk factors for osteoporotic fractures in elderly men. *Am J Epidemiol* 1996;144:255–63.
- [29] Ensrud KG, Nevitt MC, Unis C, Cauley JA, Seeley DG, Fox KM, et al. Correlates of impaired function in older women. *J Am Geriatr Soc* 1994;42:481–9.
- [30] Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE, et al., for the Study of Osteoporotic Fractures Research Group. Risk factors for hip fracture in white women. *N Engl J Med* 1995;332:767–73.
- [31] Torgerson DJ, Campbell MK, Thomas RE, Reid DM. Predictor of perimenopausal fractures by bone mineral density and other factors. *J Bone Miner Res* 1996;11:293–7.
- [32] National Osteoporosis Foundation DM. Osteoporosis: review of the evidence for prevention, diagnosis and treatment and cost-effectiveness analysis. *Osteoporosis Int* 1998;8(Suppl. 4):1–88.
- [33] Kanis JA, Delmas P, Burckhardt P, Cooper C, Torgerson D, on behalf of the EFFE. Guidelines for the diagnosis and management of osteoporosis. *Osteoporosis Int* 1997;7:390–406.
- [34] Royal College of Physicians. Osteoporosis: clinical guidelines for prevention and treatment. London: The Royal College of Physicians of London; 1999.
- [35] Royal College of Physicians. Osteoporosis: clinical guidelines for prevention and treatment. Update on pharmacological interventions and an algorithm for management. London: RCP; 2000.
- [36] Guidelines Writing Group. Glucocorticoid-induced osteoporosis. Guidelines for prevention and treatment. Bone and Tooth Society

- of Great Britain, National Osteoporosis Society and Royal College of Physicians. London: RCP; 2002.
- [37] Kanis JA, Johnell O, Oden A, De Laet C, Jonsson B, Dawson A. Ten year risk of osteoporotic fracture and the effect of risk factors on screening strategies. *Bone* 2002;30:251–8.
- [38] Kanis JA, Oden A, Johnell O, Jonsson B, De Laet C, Dawson A. The burden of osteoporotic fractures: a method for setting intervention thresholds. *Osteoporosis Int* 2001;12:417–27.
- [39] Kanis JA. Diagnosis of osteoporosis and assessment of fracture risk. *Lancet* 2002;359:1929–36.
- [40] Garnero P, Sornay-Rendu E, Chapuy M-C, Delmas PD. Increased bone turnover in late postmenopausal women is a major determinant of osteoporosis. *J Bone Miner Res* 1996;11:337–49.
- [41] Felsenberg D, Silman AJ, Lunt M, Ambrecht G, Ismail AA, Finn JD, et al. Incidence of vertebral fracture in Europe: results from the European Prospective Osteoporosis Study (EPOS). *J Bone Miner Res* 2002;17:716–24.
- [42] Ismail AA, Pye SR, Cockerill WC, Lunt M, Silman AJ, Reeve J, et al. Incidence of limb fracture across Europe: results from the European prospective osteoporosis study (EPOS). *Osteoporosis Int* 2002;13:565–71.
- [43] O'Neill TW, Felsenberg D, Varlow J, Cooper C, Kanis JA, Silman AJ. The prevalence of vertebral deformity in European men and women: European vertebral osteoporosis study. *J Bone Miner Res* 1996;11:1010–7.
- [44] Kreiger N, Tenenhouse A, Joseph L, et al. The Canadian Multicenter Osteoporosis Study (CaMos): background, rationale, methods. *Can J Aging* 1999;18:376–87.
- [45] Melton III LJ, Crowson CS, O'Fallon WM, Wahner HW, Riggs BL. Relative contributions of bone density, bone turnover and clinical risk factors to long-term fracture prediction. *J Bone Miner Res* 2003;18:312–8.
- [46] Melton III LJ, Atkinson EJ, O'Connor MK, O'Fallon WM, Riggs BL. Bone density and fracture risk in men. *J Bone Miner Res* 1998;13:1915–23.
- [47] Johansson H, Oden A, Johnell O, Jonsson B, De Laet C, Oglesby A, et al. Optimisation of BMD measurements to identify high risk groups for treatment—A test analysis. *J Bone Miner Res* (in press).
- [48] Hofman A, Grobbee DE, De Jong PT, van den Ouweland FA. Determinants of disease and disability in the elderly: the Rotterdam study. *Eur J Epidemiol* 1991;7:403–22.
- [49] De Laet CEDH, Van Hout BA, Burger H, Hofman A, Weel AEAM, Pols HAP. Hip fracture prediction in elderly men and women: validation of the Rotterdam study. *J Bone Miner Res* 1998;13:1587–93.
- [50] Honkanen R, Kroger H, Tuppurainen M, Alhava E, Saarikoski S. Fractures and low axial bone density in perimenopausal women. *J Clin Epidemiol* 1995;48:881–8.
- [51] Svanborg A. 70 year old people in Gothenburg. A population study in an industrialised Swedish city ii. Journal presentation of social and medical conditions. *Acta Med Scand* 1977;611(Suppl.):5.
- [52] Johansson C, Black D, Johnell O, Oden A, Mellstrom D. Bone mineral density is a predictor of survival. *Calcif Tissue Int* 1998;63:190–6.
- [53] Stenstrom M, Olsson J-O, Mellstrom D. Thyroid hormone replacement is not related to increased risk of osteoporosis. *Osteoporosis Int* 2000;11(Suppl. 2):S144.
- [54] Jones G, Nguyen TV, Sambrook PN, Kelly PJ, Gilbert C, Eisman JA. Symptomatic fracture incidence in elderly men and women. The Dubbo osteoporosis epidemiology study DOES. *Osteoporosis Int* 1994;4:277–82.
- [55] Fujiwara S, Kasagi F, Yamada M, Kodama K. Risk factors for hip fracture in Japanese cohort. *J Bone Miner Res* 1997;12:998–1004.
- [56] Fujiwara S, Fumiyoshi K, Masunari N, Naito K, Suzuki G, Fukunage M. Fracture prediction from bone mineral density in Japanese men and women. *J Bone Miner Res* 2003;18:1533–47.
- [57] Marshall D, Johnell O, Wedel H. Meta-analysis of how well measures of bone mineral density predict occurrence of osteoporotic fractures. *Lancet* 1996;312:1254–9.
- [58] Nelson HD, Nevitt ME, Scott JC, Stone KL, Cummings SR. Smoking, alcohol and neuromuscular and physical function of older women. *JAMA* 1994;272:1825–31.
- [59] Kanis JA. Textbook of osteoporosis. Oxford, UK: Blackwell; 1996.
- [60] Mayhew P, Loveridge N, Power J, Kroger HPJ, Parker M, Reeve J. Discrimination between cases of hip fracture and controls is improved by hip structural analysis. An ex vivo study of the femoral neck. *Bone* (in press).
- [61] Crabtree N, Loveridge N, Parker M, Rushton N, Power J, Bell KL, et al. Intracapsular hip fracture and the region-specific loss of cortical bone: analysis by quantitative computed tomography (pQCT). *J Bone Miner Res* 2001;16:1318–28.
- [62] Kanis JA, Johnell O, Oden A, De Laet C, Jonsson B. Intervention thresholds for osteoporosis. *Bone* 2001;31:26–31.
- [63] Delmas PD, Eastell R, Garnero P, Seibel MJ, Stepan J. The use of biochemical markers of bone turnover in osteoporosis. *Osteoporosis Int* 2000;11(Suppl. 6):S2–17.
- [64] Kanis JA, Johnell O, Oden A, Jonsson B, Dawson A, Dere W. Risk of hip fracture derived from relative risks: an analysis applied to the population of Sweden. *Osteoporosis Int* 2000;11:120–7.

Case-control study of risk factors for fractures of the distal radius and proximal humerus among the Japanese population

H. Hagino · S. Fujiwara · E. Nakashima · Y. Nanjo
R. Teshima

Received: 25 March 2003 / Accepted: 13 October 2003 / Published online: 17 January 2004
© International Osteoporosis Foundation and National Osteoporosis Foundation 2004

Abstract We conducted a case-control study to identify risk factors for fractures of the distal radius and proximal humerus. Subjects were selected from women aged 45 and over with distal radius and proximal humerus fractures, resulting from minor trauma. Two age- and gender-matched controls for each case were selected from patients who subsequently visited the same clinic for treatment of conditions other than fractures. Questionnaires including anthropometric data, past and current physical activity, and lifestyle were sent by mail to both subjects and controls. A total of 140 women with distal radius fractures (mean age 67.4 years) and 242 controls were analyzed. Falls during the previous year were a significant risk factor, while futon use (instead of bed use) before fracture was a protective factor for distal radius fractures. A total of 37 women with proximal humerus fractures (mean age 76.3 years) and 67 controls were analyzed. Weight loss was a significant risk factor, while greater frequency of going outside significantly decreased the risk of proximal humerus fracture. There was no significant correlation with eating habits, milk and alcohol consumption, or smoking to the risk of either fracture.

Keywords Fractures · Humerus · Japanese · Radius · Risk factors

Introduction

With the rapid increase in the elderly population, osteoporosis and related fractures are major health and socioeconomic issues. Osteoporosis increases the risk of vertebral fractures as well as fractures of the hip, distal radius, and proximal humerus [1]. Although recent anti-osteoporosis pharmaceuticals could reduce the risk of fragile fractures up to 50% [2], osteoporosis is not curable and the number of patients with these fractures is increasing rapidly in both Western and Asian countries.

The incidence of fractures of the hip, distal radius, and proximal humerus is lower in Asians, including Japanese, than in Caucasians in Northern Europe and North America [3]. As bone mass in Asians is known to be lower than or similar to that in Caucasians, bone mass difference does not account for the difference in the incidence of hip, distal radius, and proximal humerus fractures between these groups. Elucidation of the factors underlying the racial difference in the incidence of these fractures will suggest preventive measures that may protect against osteoporosis-related fractures.

Since osteoporosis-related fractures result from the coincidence of bone fragility and falls, falls are important in the pathogenesis of osteoporotic fracture. Aoyagi et al. found that the incidence of falls in Japanese was about half that of Caucasians, and concluded that the difference in hip fracture incidence was closely related to the incidence of falls [4]. Recent surveys found that age- and gender-specific rates of hip as well as distal radius and proximal humerus fractures are increasing among the Japanese population [3, 5]. This trend may be due to the fairly rapid change from a traditional Japanese lifestyle to a Western one. The traditional Japanese lifestyle, including squatting to toilet, use of the Japanese straw mat room, and sleeping on the floor, may

H. Hagino (✉)
Rehabilitation Division,
Tottori University Hospital,
Yonago, Tottori 683-8504, Japan
E-mail: hagino@grape.med.tottori-u.ac.jp
Tel.: +81-859-348155
Fax: +81-859-348093

S. Fujiwara
Department of Clinical Studies,
Radiation Effects Research Foundation, Hiroshima, Japan

E. Nakashima
Department of Statistics,
Radiation Effects Research Foundation, Hiroshima, Japan

Y. Nanjo · R. Teshima
Department of Orthopedic Surgery,
Tottori University, Tottori, Japan

maintain muscle strength, resulting in fewer falls and lower incidence of hip, distal radius, and proximal humerus fractures.

Therefore, to understand the basis underlying reduced osteoporosis-related fracture risk, it is necessary to evaluate risk factors in Japanese subjects. Although two studies of risk factors for hip fractures among Japanese have been described [6, 7], no study has examined risk factors for fractures of the distal radius and proximal humerus. Therefore, we conducted a case-control study to identify risk factors for these fractures among the Japanese population.

Materials and methods

Cases

Subjects were selected from women aged 45 and over with distal radius and proximal humerus fractures, who were treated at six orthopedic clinics in Tottori Prefecture from January 1999 to December 2000. In Tottori Prefecture, 27 hospitals had a department of orthopedic surgery and orthopedic clinics for outpatients. The six clinics selected were among those that dealt with the most fracture patients and agreed to collect data for this study. General orthopedic disease such as osteoarthritis of the joints and spine, other forms of arthritis, traumas, and fractures, are treated at these clinics.

Patients were selected by reviewing hospital records and diagnoses were confirmed by radiograms read by doctors at each clinic. Patients were excluded if 1) they died before this study was completed, 2) the fracture was caused by a high-energy impact, such as a traffic accident or a fall from greater than standing height, 3) the fracture was caused by neoplastic disease, or 4) the patient was severely demented and unable to satisfy the requirement of informed consent. Patients were subsequently contacted by mail and asked to respond to an enclosed questionnaire.

A total of 144 women aged 46–95 years (mean 67.5 years) with distal radius fractures and a total of 38 women aged 47–90 years (mean 76.4 years) with proximal humerus fractures were registered. Thirty-eight patients with distal radius fractures and 21 patients with proximal humerus fractures were admitted, while all others were managed as outpatients.

Controls

For each eligible patient registered, two age- (± 3 years) and gender-matched controls were chosen from patients who subsequently (within 3 days) visited the same orthopedic clinic for the first time for treatment of a condition other than fractures. Most controls visited because of joint pain, low back pain, and neck pain. Patients who were demented or needed assistance to walk were excluded.

Data collection

A questionnaire consisting of four parts with 43 items was mailed. The first part concerned sociodemographic data, anthropometric data, and daily physical activity such as abiding place; marital status; occupation; body height; body weight; dominant hand; distance able to walk without rest; frequency of going outdoors; vision; and ability to dress, eat, perform household tasks, stand up from a chair, ascend stairs, walk outdoors, and bath. Part two evaluated likes and dislikes and consumption of coffee, tea, Japanese tea, milk, yogurt, cheese, meat, fish, alcohol, and cigarettes. Part three concerned past history of medications and diseases. Part

four dealt with previous or recent lifestyle data including sports participation in youth, room style (western or tatami), type of bed (bed or futon) prior to fracture, toilet style (western or Japanese), and frequency of falls. Missing observations were completed by telephone, when necessary and possible, for both cases and controls.

Statistical analysis

Comparisons of background characteristics such as age, body height, body weight, and body mass index (BMI) between the two groups were performed with *t*-tests, while comparison of number of children was performed with a Mann-Whitney test.

Odds ratios for the risk variables were calculated using a conditional logistic regression model. Multivariate analysis was performed for the variables selected by univariate analysis. First, variables were screened by univariate analysis with a significance level of $P=0.10$. Next, multivariate adjusted odds ratios were calculated for variables selected in the first step, where all *P*-values were calculated using the chi-square likelihood ratio test with appropriate degrees of freedom. When considering the effect of a variable with more than two levels, we constructed several indicator variables that represent each level of the original variable. The trend test was performed when the variable had more than two levels. Multivariate analysis made it possible to estimate the odds ratio of a variable after adjusting for the effects of other variables. The Epicure software package for conditional logistic regression, specifically the PECAN program, (Hirosoft International Corporation, Seattle, USA) was used for analysis.

Results

Distal radius fractures

Among 144 patients with distal radius fractures, both controls replied in 102 cases, one control replied in 38 cases, and no controls replied in four cases. Therefore, 140 cases with at least one control and 242 controls in total were selected and analyzed. There was no significant difference in age, body height, body weight, BMI, or number of children between patients with fractures and controls (Table 1).

Table 1 Selected characteristics of cases with distal radius fracture and controls. A significance level of $P < 0.05$ was using for this analysis

Characteristics	Case (<i>n</i> = 140)	Control (<i>n</i> = 242)	<i>P</i>
<i>Age group</i>			
< 49	4	6	–
50–59	27	58	–
60–69	51	80	–
70–79	43	79	–
80–89	12	18	–
90+	3	1	–
Mean age \pm SD (years)	67.4 \pm 9.6	66.8 \pm 9.2	NS
Body height (cm)	153.7 \pm 34.3	151.6 \pm 28.2	NS
Body weight (kg)	51.5 \pm 8.6	52.2 \pm 8.4	NS
BMI (kg/m ²)	22.3 \pm 4.4	22.9 \pm 4.3	NS
Number of children (mean)	0–6 (2.1)	0–6 (2.3)	NS
<i>Dominant hand</i>			
Right	131	223	–
Left	4	15	–
Unknown	5	4	–

Table 2 Selected risk factors for distal radius fracture by univariate analysis. Factors with significance level $P < 0.10$ are shown

	Odds ratio	95% CI	P	P trend
<i>Ascend and descend stairs</i>				
Easy	1.00	Reference	–	0.017
Difficult	0.43	(0.21–0.89)	0.017	–
Unable	0.45	(0.13–1.56)	0.19	–
<i>Walk outside</i>				
Easy	1.00	Reference	–	0.050
Difficult	0.31	(0.14–0.67)	0.0012	–
Unable	1.24	(0.34–4.48)	0.75	–
<i>Walkable distance without rest</i>				
< 100 m	1.00	Reference	–	0.26
100 m to 1 km	2.28	(1.10–4.73)	0.023	–
> 1 km	1.94	(0.97–3.89)	0.054	–
<i>Frequency of going outside</i>				
Rarely or not at all	1.00	Reference	–	0.045
Once in 2 or 3 days	0.69	(0.22–2.15)	0.52	–
Once a day	1.37	(0.55–3.41)	0.49	–
> Once a day	1.94	(0.78–4.78)	0.14	–
<i>Bed or futon use when sleeping</i>				
Bed use	1.00	Reference	–	–
Futon use	0.62	(0.37–1.03)	0.064	–
<i>Falls during the previous year</i>				
Never	1.00	Reference	–	–
> Once a year	2.26	(1.45–3.54)	0.0002	–
<i>Number of children</i>				
Each one	0.86	(0.71–1.03)	0.090	–

In the first step by univariate analysis, ability to ascend and descend stairs, ability to walk outside, distance capable of walking without rest, frequency of going outside, bed or futon use before the fracture, falls during previous years, and number of children were selected ($P < 0.1$) (Table 2). Physical activity such as greater ability to ascend and descend stairs, increased walkable distance without rest, and greater frequency of going outside increased the risk of distal radius fracture. However, other variables of daily physical activity such as ability to dress, walking inside, household tasks, and bathing showed no association with fracture risk. Recent futon use (as opposed to bed use) before the fracture and increased number of children both decreased the risk of fracture. No variables related to dietary habits (coffee, tea, Japanese tea, milk, cheese, fish), past alcohol use, and medication (antihypertensive drugs and diuretics) showed any association with the risk of distal radius fracture.

In the second step by multivariate analysis, falls during the previous year was a significant risk factor, while futon use (instead of bed use) before fracture was a protective factor (Table 3).

Proximal humerus fractures

For proximal humerus fractures, both controls replied in 30 cases, one control replied in seven cases, and no controls replied in one of 38 cases. Therefore, 37 cases with at least one control and 67 controls in total were selected and analyzed. There were no significant differences in age, body height, body weight, BMI, or number

Table 3 Selected risk factors for distal radius fracture by multivariate analysis. Factors with significance level $P < 0.10$ by univariate analysis are presented

	Odds ratio	95% CI	P	P trend
<i>Ascend and descend stairs</i>				
Easy	1.00	Reference	–	0.23
Difficult	0.74	(0.23–2.26)	0.59	–
Unable	0.06	(0.0014–1.25)	0.072	–
<i>Walk outside</i>				
Easy	1.00	Reference	–	0.84
Difficult	0.42	(0.12–1.43)	0.17	–
Unable	28.0	(1.22–641.9)	0.031	–
<i>Walkable distance without rest</i>				
< 100 m	1.00	Reference	–	0.97
100 m to 1 km	2.07	(0.79–5.70)	0.14	–
> 1 km	1.53	(0.58–4.28)	0.40	–
<i>Frequency of going outside</i>				
Rarely or not at all	1.00	Reference	–	0.051
Once in 2 or 3 days	1.69	(0.41–7.13)	0.47	–
Once a day	2.33	(0.73–8.53)	0.16	–
> Once a day	3.20	(1.00–11.9)	0.049	–
<i>Bed or futon use when sleeping</i>				
Bed use	1.00	Reference	–	–
Futon use	0.55	(0.31–0.97)	0.039	–
<i>Falls during the previous year</i>				
Never	1.00	Reference	–	–
> Once a year	2.52	(1.54–4.24)	0.0002	–
<i>Number of children</i>				
Each one	0.84	(0.68–1.04)	0.10	–

Table 4 Selected characteristics of cases with proximal humerus fracture and controls. Significance level of $P < 0.05$ was using for this analysis

Characteristics	Case (n = 37)	Control (n = 67)	P
<i>Age group</i>			
–49	1	1	–
50–59	3	6	–
60–69	7	15	–
70–79	10	17	–
80–89	15	24	–
90+	1	4	–
Mean age \pm SD (years)	76.3 \pm 11.1	74.9 \pm 11.2	NS
Body height (cm)	148.7 \pm 9.9	150.7 \pm 6.7	NS
Body weight (kg)	46.5 \pm 10.4	49.0 \pm 7.4	NS
BMI (kg/m ²)	21.0 \pm 4.2	21.8 \pm 2.8	NS
Number of children (mean)	0–6 (2.0)	0–6 (2.4)	NS
<i>Dominant hand</i>			
Right	* 37	66	–
Left	0	1	–
Unknown	0	0	–

of children, between patients with fractures and controls (Table 4).

In the univariate analysis, abiding place, body weight, weight loss, frequency of going outside, alcohol consumption, and falls during the past year were selected ($P < 0.1$) (Table 5). Increased physical activity, such as greater frequency of going outside, decreased the risk of proximal humerus fracture. No variables related to dietary habits (coffee, tea, Japanese tea, milk, cheese, fish), and medication (antihypertensive drugs and diuretics) showed any association with the risk of proximal humerus fracture.

Table 5 Selected risk factors for proximal humerus fracture by univariate analysis. Factors with significance level $P < 0.10$ are presenting

	Odds ratio	95% CI	P
<i>Residence</i>			
Home	1.00	Reference	–
Nursing home	5.88	(1.20–28.8)	0.015
<i>Body weight</i>			
Each 10 kg	0.57	(0.30–1.10)	0.087
<i>Weight loss</i>			
No	1.00	Reference	–
Yes	2.15	(0.90–5.10)	0.077
<i>Frequency of going outside</i>			
Rarely or not at all	1.00	Reference	–
More than once every 2 or 3 days	0.22	(0.07–0.68)	0.004
<i>Alcohol consumption</i>			
Non-drinker	1.00	Reference	–
Drinker	0.19	(0.055–0.68)	0.003
<i>Falls during the previous year</i>			
Never	1.00	Reference	–
> Once a year	2.87	(1.03–8.00)	0.038

Table 6 Selected risk factors for proximal humerus fracture by multivariate analysis. Factors with significance level $P < 0.10$ by univariate analysis are presenting

	Odds ratio	95% CI	P
<i>Residence</i>			
Home	1.00	Reference	–
Nursing home	1.76	(0.071–111.2)	0.75
<i>Body weight</i>			
each 10 kg	0.58	(0.23–1.31)	0.19
<i>Weight loss</i>			
No	1.00	Reference	–
Yes	5.00	(1.37–24.9)	0.013
<i>Frequency of going outside</i>			
Rarely or not at all	1.00	Reference	–
More than once every 2 or 3 days	0.14	(0.015–0.88)	0.035
<i>Alcohol drinking</i>			
Non-drinker	1.00	Reference	–
Drinker	0.29	(0.052–1.16)	0.083
<i>Falls during the previous year</i>			
Never	1.00	Reference	–
> Once a year	2.60	(0.55–14.9)	0.23

Multivariate analysis showed that weight loss was a significant risk factor, while greater frequency of going outside significantly decreased the risk of proximal humerus fracture (Table 6).

Discussion

The incidence of distal radius fractures increases with age, from the late 50s. However, the increase correlated with aging is not observed in subjects older than 60 years [3]. In contrast, the incidence of proximal humerus fractures, similar to that for hip fractures, increases markedly in subjects older than 80 years [3, 8]. Therefore, risk factors for distal radius fractures are different from those for proximal humerus fractures. Mean ages of the subjects in the present study were 67.4 years for distal radius fractures and 76.3 years for proximal humerus fractures, ages in which fractures are caused by bone fragility.

Epidemiological studies have indicated that risk factors for distal forearm fracture are low bone mass, estrogen deficiency [9], falls [10], drinking alcohol [11, 12], and consumption of animal protein [13]. Poor visual acuity [14, 15], frequent walking [14, 16], and walking at a brisk pace [17] are also risk factors for distal forearm fractures. A recent cohort study showed a 60% reduction in the risk of wrist fractures in women with no vigorous exercise in the past 2 weeks [15]. The authors suggested that strategies to increase exercise level among older people might have the adverse effect of increasing the number of wrist fractures.

This study demonstrated that increased physical activity, in particular increased walking ability, is a risk factor for distal radius fractures. This is in agreement with factors identified in previous studies, which concluded that increased physical activity, increased walking ability, and frequent outdoor walking all increase the

risk of falls. Since nearly all distal radius fractures are caused by falls [18], these fractures are most likely to occur in highly active patients with low bone mass. This study also confirmed that tendency to fall is the most significant risk factor for distal radius fractures after adjusting for other factors by multivariate analysis.

Risk factors for proximal humerus fractures include low level of physical activity [19] and infrequent walking [14]. This contrasts with risk factors for distal radius fractures, indicating that distal radius fractures are most likely in patients with fragile bones and increased physical activity, while proximal humerus fractures are most likely in patients with fragile bones and decreased physical activity. Our results agree with those obtained in past cohort and case-control studies. Patients with reduced physical activity tend to fall before supporting themselves with their hands, which could result in injury of the proximal humerus. Greater weight loss was also a significant risk factor for proximal humerus fractures. Low body weight is a significant risk factor for bone loss [20], and therefore, becoming thinner increases the risk of fracture.

The finding that greater physical activity increases the risk of distal radius fractures and decreases the risk of proximal humerus fractures corresponds with changes in the incidence of fractures with aging. Distal radius fractures are most likely in 50- to 79-year-old patients with normal or high physical activity, while proximal humerus and hip fractures are most likely in those older than 80 years with decreased physical activity.

Although risk factors for hip fractures have been well defined in Caucasian subjects, only two epidemiological studies have examined the risk factors for hip fractures in Japanese subjects [6, 7]. Suzuki et al. performed a case-control study on hip fractures and found that sleeping on a bed (as opposed to a futon) was a significant risk factor (OR = 1.95), as were known risk factors

such as reduced physical activity [6]. They also reported that Japanese lifestyle, such as drinking Japanese tea, was effective in preventing fractures.

One significant preventive factor for distal radius fractures among Japanese was the use of a futon. This was also a preventive factor for hip fractures, as reported by Suzuki et al. [6]. Therefore, futon use (as opposed to bed use) may be important in reducing the risk of falls. One reason may be that spreading futons and putting them away in a closet everyday contributes to the maintenance of muscular strength in the lower limbs, and thereby reduces the risk of falls.

This study had several limitations. Since controls were selected from outpatients at orthopedic clinic, bias from other orthopedic disease cannot be dismissed. All control subjects were outpatients, and all patients with reduced activity of daily living, such as inability to walk outdoors without assistance, were excluded. However, control subjects may have lower walking ability than healthy subjects, which might result in the finding of an apparent risk of physical activity among subjects. It should be noted that tendency to fall was a significant risk factor for distal radius fractures, independent of the ability to walk outside. Second, this study was a cross-sectional study and information was obtained after the fracture event, which could bias the findings. It is necessary to perform case-control studies with a larger number of subjects, including population-based controls, as well as cohort studies in Asian countries where the incidence of osteoporosis related fractures is lower than in Caucasians.

In conclusion, risk factors for fractures of the upper limbs were elucidated here for the first time in Asian subjects. Falling is a significant risk factor for both fractures and futon use (instead of bed use) is a protective factor for distal radius fractures. Efforts to maintain physical activity, while being cautious to avoid falls, should be considered to reduce fracture risk among the elderly.

Acknowledgements The authors thank Akira Fukushima, Yu Oh-sako, Masaaki Shimizu, and Noboru Yamagata, for their help and support. This study was supported by a Grant-in-Aid for Scientific Research C (14571378) from the ministry of Education, Culture, Sports, Science, and Technology of Japan.

References

- Seeley DG, Browner WS, Nevitt MC, Genant HK, Scott JC, Cummings SR (1991) Which fractures are associated with low appendicular bone mass in elderly women? The Study of Osteoporotic Fractures Research Group. *Ann Int Med* 115:837-842
- Marcus R, Wong M, Heath H 3rd, Stock JL (2002) Antiresorptive treatment of postmenopausal osteoporosis: comparison of study designs and outcomes in large clinical trials with fracture as an endpoint. *Endocrine Rev* 23:16-37
- Hagino H, Yamamoto K, Ohshiro H, Nakamura T, Kishimoto H, Nose T (1999) Changing incidence of hip, distal radius, and proximal humerus fractures in Tottori Prefecture, Japan. *Bone* 24:265-270
- Aoyagi K, Ross PD, Davis JW, Wasnich RD, Hayashi T, Takemoto T (1998) Falls among community-dwelling elderly in Japan. *J Bone Miner Res* 13:1468-1474
- Iga T, Dohmae Y, Endo N, Takahashi HE (1999) Increase in the incidence of cervical and trochanteric fractures of the proximal femur in Niigata Prefecture, Japan. *J Bone Miner Metab* 17:224-231
- Suzuki T, Yoshida H, Hashimoto T et al. (1997) Case-control study of risk factors for hip fractures in the Japanese elderly by a Mediterranean Osteoporosis Study (MEDOS) questionnaire. *Bone* 21:461-467
- Fujiwara S, Kasagi F, Yamada M, Kodama K (1997) Risk factors for hip fracture in a Japanese cohort. *J Bone Miner Res* 12:998-1004
- Melton LJ 3rd (1995) Epidemiology of fractures. In: Riggs BL, Melton LJ 3rd (eds) *Osteoporosis. Etiology, diagnosis, and management*. Lippincott-Raven, Philadelphia, pp 225-247
- Mallmin H, Ljunghall S, Persson I, Bergstrom R (1994) Risk factors for fractures of the distal forearm: a population-based case-control study. *Osteoporos Int* 4:298-304
- Winner SJ, Morgan CA, Evans JG (1989) Perimenopausal risk of falling and incidence of distal forearm fracture. *BMJ* 298:1486-1488
- Hemenway D, Colditz GA, Willett WC, Stampfer MJ, Speizer FE (1988) Fractures and lifestyle: effect of cigarette smoking, alcohol intake, and relative weight on the risk of hip and forearm fractures in middle-aged women. *Am J Public Health* 78:1554-1558
- Hernandez-Avila M, Colditz GA, Stampfer MJ, Rosner B, Speizer FE, Willett WC (1991) Caffeine, moderate alcohol intake, and risk of fractures of the hip and forearm in middle-aged women. *Am J Clin Nutr* 54:157-163
- Feskanich D, Willett WC, Stampfer MJ, Colditz GA (1996) Protein consumption and bone fractures in women. *Am J Epidemiol* 143:472-479
- Kelsey JL, Browner WS, Seeley DG, Nevitt MC, Cummings SR (1992) Risk factors for fractures of the distal forearm and proximal humerus. The Study of Osteoporotic Fractures Research Group. *Am J Epidemiol* 135:477-489
- Ivers RQ, Cumming RG, Mitchell P, Peduto AJ (2002) Risk factors for fractures of the wrist, shoulder and ankle: the Blue Mountains Eye Study. *Osteoporos Int* 13:513-518
- Silman AJ (2003) Risk factors for Colles' fracture in men and women: results from the European Prospective Osteoporosis Study. *Osteoporos Int* 14:213-218
- O'Neill TW, Marsden D, Adams JE, Silman AJ (1996) Risk factors, falls, and fracture of the distal forearm in Manchester, UK. *J Epidemiol Commun Health* 50:288-292
- Nevitt MC, Cummings SR (1993) Type of fall and risk of osteoporotic fractures. The Study of Osteoporotic Fractures Research Group. *J Am Geriatr Soc* 41:1226-1234
- Lee SH, Dargent-Molina P, Breart G (2002) Risk factors for fractures of the proximal humerus: results from the EPIDOS prospective study. *J Bone Miner Res* 17:817-825
- Woolf AD, Akesson K (2003) Preventing fractures in elderly people. *BMJ* 327:89-95

Self-reported number of remaining teeth is associated with bone mineral density of the femoral neck, but not of the spine, in Japanese men and women

Akira Taguchi · Saeko Fujiwara
Naomi Masunari · Gen Suzuki

Received: 20 October 2003 / Accepted: 2 February 2004 / Published online: 26 February 2004
© International Osteoporosis Foundation and National Osteoporosis Foundation 2004

Abstract Recent studies suggest that a small number of remaining teeth may be associated with low skeletal bone mineral density (BMD) in postmenopausal women. Estrogen deficiency after menopause is considered potential cause relating to tooth loss accompanied by low skeletal BMD in women. Since estrogen plays a dominant role in regulating the male skeleton, it is likely that a small number of remaining teeth also may be associated with low skeletal BMD in men. However, it remains uncertain whether tooth loss is associated with low skeletal BMD in both men and women. We investigated the association between self-reported number of remaining teeth and BMD of the spine and the femoral neck in a cohort of 1914 Japanese subjects aged 48–95 years who were recruited from the Adult Health Study conducted by the Radiation Effects Research Foundation (RERF). BMD of the spine and the femoral neck was measured by dual energy X-ray absorptiometry (DXA). Tooth count was self-reported in response to a simple question to subjects about the number of remaining teeth they had at the time of the survey. Multiple regression analysis adjusted for age, weight, height, smoking, estrogen use, and years since menopause revealed a significant association between number of remaining teeth and BMD of the femoral neck in both men and women; however, no association was found between number of remaining teeth and BMD of the spine in both sexes. Retention of four teeth was significantly associated with a 0.004 g/cm^2 increase in femoral neck BMD in men ($P < 0.05$), which was similar to that

observed in women ($P < 0.01$). Our results suggest the presence of common causes, except age and body weight, relating to tooth loss accompanied by low BMD of the femoral neck in both men and women.

Keywords Bone mineral density · Femoral neck · Osteoporosis · Self-reported · Tooth loss

Introduction

Osteoporotic fractures are associated with substantial morbidity, increased medical costs and high mortality risk in the elderly [1]. Tooth loss is also associated with deterioration in the systemic health of the elderly through alteration of dietary intake [2,3]. Causes of tooth loss include dental caries, periodontal disease, eruption problems, trauma, and orthodontics, among others [4]. It is generally known that women lose significant amounts of bone after menopause; however, women also lose more teeth after 50 years of age, which is mean menopausal age, than do men of the same age, in spite of a higher frequency of tooth brushing and a smaller number of untreated teeth [5].

Since Daniell [6] first reported a significant association between postmenopausal tooth loss and metacarpal bone mass, some investigators have linked tooth loss with low general skeletal bone mineral density (BMD) and high bone loss rates in postmenopausal women [7,8,9,10,11,12,13]. However, others failed to find an association between tooth loss and skeletal BMD in postmenopausal women [14,15,16,17,18]. It is likely that variance in the size and the age range of study populations may have contributed to this controversy.

Three epidemiological studies in the United States suggested the protective effect of estrogen use on tooth retention in postmenopausal women [19,20,21]. Men lose significant amounts of bone with age, although they do not have the equivalent of menopause. Recent reports suggested that estrogen plays a dominant role in

A. Taguchi (✉)
Department of Oral and Maxillofacial Radiology,
Hiroshima University Hospital, 1-2-3 Kasumi,
Minami-ku, 734-8553 Hiroshima, Japan
E-mail: akiro@hiroshima-u.ac.jp
Tel.: +81-82-2575691
Fax: +81-82-2575692

A. Taguchi · S. Fujiwara · N. Masunari · G. Suzuki
Department of Clinical Studies,
Radiation Effects Research Foundation, Hiroshima, Japan

regulating the skeleton in both sexes [22,23,24,25,26]. Riggs et al. proposed a new unitary model for the pathophysiology of involutional osteoporosis that identifies estrogen deficiency as the cause of both the early, accelerated and the late, slow phases of bone loss in postmenopausal women and as a contributing cause of the continuous phase of bone loss in aging men [27].

The potential role of estrogen deficiency linking tooth loss and low skeletal BMD in postmenopausal women suggests that tooth loss also may be associated with low skeletal BMD in men. Only one study reported that a small self-reported number of remaining teeth was associated with low BMD of the spine and the hip in Caucasian men [16]. However, since the age range of the subjects in this study was limited to only 10 years (65–76 years of age), it remains uncertain whether tooth loss is associated with low BMD in both men and women.

The purpose of this study was therefore to investigate whether self-reported number of remaining teeth is associated with BMD of the spine and the femoral neck in a large population, including both men and women, with a wide age range.

Materials and methods

Subjects

The Adult Health Study (AHS) was established by the Radiation Effects Research Foundation (RERF) in 1958 to document the late health effects of radiation exposure among atomic bomb survivors in Hiroshima and Nagasaki. The original AHS cohort consisted of about 15,000 atomic bomb survivors and about 5000 controls selected from residents in Hiroshima and Nagasaki using Japan's 1950 national census supplementary schedules and the Atomic Bomb Survivors Survey. AHS subjects have been followed through biennial medical examinations since 1 July 1958. The participation rate has been around 80% throughout this period. Details concerning recruitment and examination of participants have been reported [28].

Of AHS subjects who underwent physical examinations in Hiroshima between 2000 and 2002, 1914 subjects (616 men and 1298 women) aged 48–95 years were recruited for this study. Exclusion criteria were: impaired bone metabolism (such as hyperparathyroidism, renal osteodystrophy, and bilateral oophorectomy), and use of medications that affect bone metabolism (such as corticosteroids, calcitonin, vitamin D, bisphosphonate, and vitamin K) except estrogen.

BMD measurement and self-reported tooth count

BMD of the spine (L2–L4) and the femoral neck was measured using dual X-ray absorptiometry (DXA, QDR-2000, Hologic Inc., Waltham, Mass., USA). An anthropomorphic spine phantom was scanned daily to calibrate the instrument. There was no drift in machine performance during the study period. The instrument's precision was also carefully monitored over the study period using the anthropomorphic phantom, and was found to be less than 1%. No subjects had undergone spinal X-ray radiographs prior to DXA assessment between 2000 and 2002, because of concerns about exposure to high radiation doses. In medical examinations between 1998 and 2000 in this cohort, 3.7% of men and 10.2% of women had spinal fractures determined by X-ray radiographs. Vertebral fracture was diagnosed by lateral and posterior-anterior chest and spine X-ray examination. The vertebral fracture was assessed using

semi-quantitative assessment [29,30]. All subjects gave written informed consent for BMD measurements and all other health examinations. Tooth count was self-reported in response to a simple question that asked subjects about the number of remaining teeth they had at the time of the survey.

Statistical analysis

The linear regression model was used to analyze the relationship between potential risk factors and BMD of the spine and the femoral neck. Since we previously demonstrated no association between exposure to atomic bomb radiation and skeletal BMD in this AHS cohort [28], exposure to atomic bomb radiation was not included in potential risk factors. Multivariate analysis was performed in addition to univariate analysis. First, variables were selected by univariate analysis adjusting only for age, using the level of $P=0.05$ as an indicator of significance. Next, a search for the best model was conducted in stages. Initially, all factors selected in the univariate analysis were included in the model. Non-significant variables were successively eliminated until all remaining variables except age and number of teeth were significant, with $P<0.05$. All computations were carried out using the Statistical Analysis System (SAS) package of programs. There were numerous unknown responses to each item on the questionnaire, as is common in mail surveys. For example, information about smoking was missing in 28% of the responses. In the linear regression analysis, the missing values were grouped together as one category, without excluding them from the study sample, to avoid a decrease in statistical power.

Results

Characteristics of the study population are shown in Table 1. BMD of the spine in men was significantly associated with age ($P<0.001$). In univariate analysis after adjusting for age, BMD of the spine in men was significantly related to weight ($P<0.001$), and height ($P<0.001$), but not to the number of remaining teeth ($P=0.30$). In women, BMD of the spine was significantly

Table 1 Characteristics of the study population (mean \pm SD or %)

	Men	Women
No. of subjects	616	1298
Age (years)	66.9 \pm 9.0	70.8 \pm 9.0
Height (cm) ^a	162.9 \pm 6.4	149.4 \pm 6.1
Weight (kg) ^a	60.6 \pm 9.2	51.3 \pm 9.0
Age at menopause (years) ^b	–	48.4 \pm 4.9
<i>Bone mineral density (g/cm²)</i>		
Lumbar spine ^c	1.00 \pm 0.18	0.80 \pm 0.16
Femoral neck ^d	0.74 \pm 0.12	0.59 \pm 0.11
No. of remaining teeth	15.7 \pm 10.6	13.4 \pm 10.2
<i>Smoking habit^e</i>		
No. (%) who never smoked	88 (20.3%)	811 (86.2%)
No. (%) who used to smoke	155 (35.8%)	43 (4.6%)
No. (%) who smoke currently	190 (43.9%)	87 (9.2%)
No. (%) who use estrogen	–	48 (3.7%)

^aFour women had no data on height and weight

^bNine hundred and fifty-one women were postmenopausal, 839 of whom had information about age at menopause

^cTwo men and one woman had no BMD data of the spine

^dOne man and six women had no BMD data of the femoral neck

^eOne hundred and eighty-three men and 357 women had no information

Table 2 Correlation with bone mineral density of the spine and the femoral neck by univariate analysis, after adjusting for age. The results are presented as the parameter estimate \pm SE

Items	Bone mineral density of the spine				Bone mineral density of the femoral neck			
	Men	r-square	Women	r-square	Men	r-square	Women	r-square
Age (10 years)	0.011 \pm 0.008 ^a	0.003	-0.037 \pm 0.005 ^a	0.050	-0.039 \pm 0.005 ^a	0.075	-0.055 \pm 0.003 ^a	0.220
Weight (10 kg)	0.070 \pm 0.007 ^a	0.128	0.069 \pm 0.005 ^a	0.196	0.053 \pm 0.005 ^a	0.230	0.041 \pm 0.003 ^a	0.329
Height (10 cm)	0.048 \pm 0.012 ^a	0.031	0.044 \pm 0.008 ^a	0.076	0.046 \pm 0.007 ^a	0.128	0.033 \pm 0.047 ^a	0.248
Smoking (vs non-smoker)	-	0.016	-	0.053	-	0.080	-	0.225
Past	0.042 \pm 0.024	-	-0.001 \pm 0.023	-	0.017 \pm 0.016	-	-0.027 \pm 0.015 ^d	-
Current	0.010 \pm 0.023	-	-0.023 \pm 0.017	-	-0.006 \pm 0.015	-	-0.021 \pm 0.011 ^c	-
No information	0.049 \pm 0.023 ^c	-	0.009 \pm 0.010	-	0.005 \pm 0.015	-	0.050 \pm 0.006	-
Tooth (4 teeth)	0.003 \pm 0.003	0.005	0.002 \pm 0.002	0.052	0.006 \pm 0.002 ^b	0.089	0.004 \pm 0.001 ^b	0.226
Estrogen use (vs no)	-	-	0.008 \pm 0.023	0.051	-	-	0.005 \pm 0.014	0.220
Pre-menopause (vs post)	-	-	0.061 \pm 0.019 ^b	0.052	-	-	0.019 \pm 0.012	0.211
Years since menopause (10 years)	-	-	-0.038 \pm 0.009 ^a	0.056	-	-	-0.026 \pm 0.006 ^a	0.221

^a $P < 0.001$, ^b $P < 0.01$, ^c $P < 0.05$, ^d $0.05 < P < 0.10$

associated with age ($P < 0.001$), weight ($P < 0.001$), height ($P < 0.001$), menopausal status ($P < 0.01$), and years since menopause ($P < 0.001$), but not with number of remaining teeth ($P = 0.25$). BMD of the femoral neck was significantly associated with age ($P < 0.001$), weight ($P < 0.001$), height ($P < 0.001$), and number of remaining teeth ($P < 0.01$) in men, and with age ($P < 0.001$), weight ($P < 0.001$), height ($P < 0.001$), current smoking ($P < 0.05$), years since menopause ($P < 0.001$), and number of remaining teeth ($P < 0.01$) in women (Table 2).

Multiple regression analysis adjusted for all potential confounding factors that were selected by univariate revealed a significant association between number of remaining teeth and BMD of the femoral neck in both men and women; however, no association was found between number of remaining teeth and BMD of the spine in either men ($P = 0.67$) or women ($P = 0.37$) (Table 3). Retention of four teeth was significantly associated with 0.004 g/cm² ($P < 0.05$) increase of femoral neck BMD in men and 0.004 g/cm² ($P < 0.01$) increase of femoral neck BMD in women, although the variance of tooth loss related to femoral neck BMD was small (0.75% in men and 0.58% in women). Advancing age was significantly associated with increased BMD of the spine ($P < 0.001$) and decreased BMD of the femoral neck ($P < 0.001$) in men. Similarly, BMD tended to be

higher at the spine ($P = 0.07$) and lower at the femoral neck ($P = 0.08$) with advancing age in women. However, a great number of years since menopause was significantly associated with decreased BMD of the spine ($P < 0.05$) and the femoral neck ($P < 0.001$) in postmenopausal women. Premenopausal women had significantly higher BMD of the spine than did postmenopausal women ($P < 0.01$). Higher body weight was significantly associated with increased BMD of the spine ($P < 0.001$) and the femoral neck ($P < 0.001$) in both sexes.

Discussion

BMD of the femoral neck was significantly associated with number of remaining teeth in both men and women in our study. Krall et al. found a significant association between number of remaining teeth and BMD of the femoral neck in 329 healthy Caucasian postmenopausal women aged 41–71 years [8]. May et al. observed this association in women aged 65–76 years [16], although it did not reach statistical significance. Earnshaw et al. failed to find an association, but their subjects were limited to early postmenopausal women [18]. They concluded that the lack of association in their study might have been because dental status in younger women is a

Table 3 Relationship between number of remaining teeth and bone mineral density of the spine and the femoral neck by multiple regression analysis. The results are presented as the parameter estimate \pm SE

Items	Bone mineral density of spine		Bone mineral density of femoral neck	
	Men	Women	Men	Women
Age (10 years)	0.029 \pm 0.001 ^a	0.002 \pm 0.001 ^d	-0.020 \pm 0.005 ^a	-0.012 \pm 0.007 ^d
Weight (10 kg)	0.070 \pm 0.007 ^a	0.068 \pm 0.005 ^a	0.053 \pm 0.005 ^a	0.041 \pm 0.003 ^a
Tooth (4 teeth)	0.001 \pm 0.003	0.002 \pm 0.002	0.004 \pm 0.002 ^c	0.004 \pm 0.001 ^b
Pre-menopause (vs post)	-	0.051 \pm 0.019 ^b	-	-
Years since menopause (10 years)	-	-0.023 \pm 0.003 ^c	-	-0.023 \pm 0.005 ^a
r-square	0.129	0.200	0.238	0.339

^a $P < 0.001$, ^b $P < 0.01$, ^c $P < 0.05$, ^d $0.05 < P < 0.10$

reflection more of dietary habits and previous dental surgery than of age-related bone loss. Gur et al. also failed to find an association between the two factors in 1171 postmenopausal women aged 40–86 years recruited from multiple locations in Turkey [12], but their results were not adjusted for age, weight and smoking habit, which may have had some kind of effect on both BMD of the femoral neck and number of remaining teeth.

Only one previous study found a significant association between number of remaining teeth and BMD of the femoral neck in 608 men aged 65–76 years [16]. Our results in 605 men agreed with these results. Furthermore, the extent (0.004 g/cm^2) of BMD increase of the femoral neck associated with retention of four teeth in women was similar to that (0.004 g/cm^2) in men. This implies that a common cause, or a combination of factors, may play a role in the link between number of remaining teeth and BMD of the femoral neck in both sexes. Loss of oral bone surrounding the teeth as a consequent of accelerated skeletal bone loss may be considered one of potential causes linking number of remaining teeth and BMD of the femoral neck in women; however, it is doubtful that slow continuous phase of skeletal bone loss in men causes oral bone loss, resulting in tooth loss.

Other potential causes linking number of remaining teeth and BMD of the femoral neck may include an increased inflammation of periodontal tissue surrounding the teeth. Previous studies in Japan [10] and in Finland [15] suggested that women with higher skeletal BMD tended to have more teeth than did those with lower skeletal BMD even when both had the same degree of oral bone surrounding the teeth. Ronderos et al. demonstrated that women with osteoporosis based on femoral neck BMD were at increased risk of periodontal disease, and that this risk may be attenuated by the use of estrogen replacement therapy [31]. Similar association between increased risk of periodontal disease and low femoral BMD was also observed in men in their study. These facts indicate that increased risk of periodontal disease may be an additional cause linking number of remaining teeth and BMD of the femoral neck in both sexes. Morishita et al. recently indicated that estradiol and progesterone inhibited the production of interleukin-1 from human peripheral monocytes, although testosterone did not show any significant effect on interleukin-1 production [32]. Increased level of interleukin-1 in both men and women with low concentration of estrogen may contribute to increased inflammation of periodontal tissue surrounding the teeth.

BMD of the spine was not associated with number of remaining teeth in women in this study. We previously found a significant association between number of remaining teeth and BMD of the spine in 90 Japanese postmenopausal women without spinal fracture aged 40–68 years [10]. Krall et al. also found an association in postmenopausal women without spinal fracture [8]. Gur et al. reported an association in postmenopausal women without the fractures after 25 years of age [12], although their results were not adjusted for confounding variables

related to both tooth loss and BMD of the spine. The large rate of spinal fracture incidence in medical examinations between 1998 and 2000 in our cohort suggests the possibility that spinal fracture or deformity may have influenced BMD of the spine in women in our study.

We found no association between number of remaining teeth and BMD of the spine in men, which did not agree with the finding reported by May et al. [16]. However, a significant association between advancing age and increased BMD of the spine in men strongly suggests that other factors related to advancing age such as spinal fracture, spinal deformity, aortic calcification and/or osteophyte formation may have influenced BMD of the spine in our study.

This study has limitations. Self-reported tooth count might be inaccurate in comparison with the number of remaining teeth that dentists or trained professionals can determine clinically or radiographically. However, Douglass et al. demonstrated that self-reported number of remaining teeth was highly correlated with the actual number of teeth found in clinical examinations in a general population ($r=0.97$) [33]. Pitiphat et al. also reported that the self-reported numbers of remaining teeth, fillings, root canal therapy, and prosthesis were strongly correlated with clinical records ($r=0.74-1.0$), although self-reporting was less accurate for measuring periodontal disease ($r=0.56$) [34]. Self-reported number of remaining teeth is used to investigate the association between tooth loss and systemic diseases such as stroke [35] or peripheral arterial disease [36] in large population studies. Although there have been no previous studies demonstrating the validity of self-reported tooth count in Japan, it is likely that the self-reported number of remaining teeth may accurately reflect the actual number of remaining teeth in our subjects, because all subjects expressed a strong interest in their general health, including oral health.

In conclusion, self-reported number of remaining teeth was significantly associated with BMD of the femoral neck in both men and women, but not with BMD of the spine. Our results suggest that there may be common causes relating to tooth loss accompanied by low BMD of the femoral neck in both men and women, although it is unknown whether these causes include the effect of estrogen.

Acknowledgements The Radiation Effects Research Foundation (RERF), Hiroshima and Nagasaki, Japan is a private, non-profit foundation founded by the Japanese Ministry of Health, Labour and Welfare (MHLW) and the US Department of Energy (DOE) the latter through the National Academy of Sciences. This publication was supported by RERF Research Protocol RP #3-98 and Grant of "Determination of risk factors for vertebral deformity and its prevention among the elderly" (H12-Longivity-032) by MHLW.

References

1. Melton LJ 3rd (2003) Adverse outcomes of osteoporotic fractures in the general population. *J Bone Miner Res* 18:1139–1141

2. Shimazaki Y, Soh I, Saito T, Yamashita Y, Koga T, Miyazaki H et al. (2001) Influence of dentition status on physical disability, mental impairment, and mortality in institutionalized elderly people. *J Dent Res* 80:340-345
3. Ritchie CS, Joshipura K, Hung HC, Douglass CW (2002) Nutrition as a mediator in the relation between oral and systemic disease: associations between specific measures of adult oral health and nutrition outcomes. *Crit Rev Oral Biol Med* 13:291-300
4. Morita M, Kimura T, Kanegae M, Ishikawa A, Watanabe T (1994) Reasons for extraction of permanent teeth in Japan. *Commun Dent Oral Epidemiol* 22:303-306
5. Dental Health Division of Health Policy Bureau Ministry of Health, Labour and Welfare Japan (1999) Report on the survey of dental disease. Oral Health Association, Tokyo
6. Daniell HW (1982) Postmenopausal tooth loss. Contributions to edentulism by osteoporosis and cigarette smoking. *Arch Int Med* 143:1678-1682
7. Kribbs PJ (1990) Comparison of mandibular bone in normal and osteoporotic women. *J Prosthet Dent* 63:218-222
8. Krall EA, Dawson-Hughes B, Papas A, Garcia RI (1994) Tooth loss and skeletal bone density in healthy postmenopausal women. *Osteoporos Int* 4:104-109
9. Krall EA, Garcia RI, Dawson-Hughes B (1996) Increased risk of tooth loss is related to bone loss at the whole body, hip, and spine. *Calcif Tissue Int* 59:433-437
10. Taguchi A, Sueti Y, Ohtsuka M, Otani K, Tanimoto K, Hollender LG (1999) Relationship between bone mineral density and tooth loss in elderly Japanese women. *Dentomaxillofac Radiol* 28:219-223
11. Inagaki K, Kurosu Y, Kamiya T et al. (2001) Low metacarpal bone density, tooth loss, and periodontal disease in Japanese women. *J Dent Res* 80:1818-1822
12. Gur A, Nas K, Kayhan O, Atay MB, Akyuz G, Sindal D, et al (2003) The relation between tooth loss and bone mass in postmenopausal osteoporotic women in Turkey: a multicenter study. *J Bone Miner Metab* 21:43-47
13. Mohammad AR, Hooper DA, Vermilyea SG, Mariotti A, Preshaw PM (2003) An investigation of the relationship between systemic bone density and clinical periodontal status in post-menopausal Asian-American women. *Int Dent J* 53:121-125
14. Elders PJ, Habets LL, Netelenbos JC, van der Linden LW, van der Stelt PF (1992) The relation between periodontitis and systemic bone mass in women between 46 and 55 years of age. *J Clin Periodontol* 19:492-496
15. Klemetti E, Collin HL, Forss H, Markkanen H, Lassila V (1994) Mineral status of skeleton and advanced periodontal disease. *J Clin Periodontol* 21:184-188
16. May H, Reader R, Murphy S, Khaw KT (1995) Self-reported tooth loss and bone mineral density in older men and women. *Age Ageing* 24:217-221
17. Mohammad AR, Bauer RL, Yeh CK (1997) Spinal bone density and tooth loss in a cohort of postmenopausal women. *Int J Prosthodont* 10:381-385
18. Earnshaw SA, Keating N, Hosking DJ, Chilvers CE, Ravn P, McClung M et al. (1998) Tooth counts do not predict bone mineral density in early postmenopausal Caucasian women. EPIC study group. *Int J Epidemiol* 27:479-483
19. Paganini-Hill A (1995) The benefits of estrogen replacement therapy on oral health. The Leisure World cohort. *Arch Int Med* 155:2325-2329
20. Grodstein F, Colditz GA, Stampfer MJ (1996) Post-menopausal hormone use and tooth loss: a prospective study. *J Am Dent Assoc* 127:370-377
21. Krall EA, Dawson-Hughes B, Hannan MT, Wilson PW, Kiel DP (1997) Postmenopausal estrogen replacement and tooth retention. *Am J Med* 102:536-542
22. Smith EP, Boyd J, Frank GR, Takahashi H, Cohen RM, Specker B et al. (1994) Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. *N Engl J Med* 331:1056-1061
23. Morishima A, Grumbach MM, Simpson ER, Fisher C, Qin K (1995) Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. *J Clin Endocrinol Metab* 80:3689-3698
24. Bilezikian JP, Morishima A, Bell J, Grumbach MM (1998) Increased bone mass as a result of estrogen therapy in a man with aromatase deficiency. *N Engl J Med* 339:599-603
25. Slemenda CW, Longcope C, Zhou L, Hui SL, Peacock M, Johnston CC (1997) Sex steroids and bone mass in older men. Positive associations with serum estrogens and negative associations with androgens. *J Clin Invest* 100:1755-1759
26. Falahati-Nini A, Riggs BL, Atkinson EJ, O'Fallon WM, Eastell R, Khosla S (2000) Relative contributions of testosterone and estrogen in regulating bone resorption and formation in normal elderly men. *J Clin Invest* 106:1553-1560
27. Riggs BL, Khosla S, Melton LJ 3rd (1998) A unitary model for involutional osteoporosis: estrogen deficiency causes both type I and type II osteoporosis in postmenopausal women and contributes to bone loss in aging men. *J Bone Miner Res* 13:763-773
28. Fujiwara S, Kasagi F, Masunari N, Naito K, Suzuki G, Fukunaga M (2003) Fracture prediction from bone mineral density in Japanese men and women. *J Bone Miner Res* 18:1547-1553
29. Genant HK, Wu CY, van Kuijk C, Nevitt MC (1993) Vertebral fracture assessment using a semiquantitative technique. *J Bone Miner Res* 8:1137-1148
30. National Osteoporosis Foundation Working Group on Vertebral Fractures (1995) Report: assessing vertebral fractures. *J Bone Miner Res* 10:518-523
31. Ronderos M, Jacobs DR, Himes JH, Pihlstrom BL (2000) Associations of periodontal disease with femoral bone mineral density and estrogen replacement therapy: cross-sectional evaluation of US adults from NHANES III. *J Clin Periodontol* 27:778-786
32. Morishita M, Miyagi M, Iwamoto Y (1999) Effects of sex hormones on production of interleukin-1 by human peripheral monocytes. *J Periodontol* 70:757-760
33. Douglass CW, Berlin J, Tennstedt S (1991) The validity of self-reported oral health status in the elderly. *J Public Health Dent* 51:220-222
34. Pitiphat W, Garcia RI, Douglass CW, Joshipura KJ (2002) Validation of self-reported oral health measures. *J Public Health Dent* 62:122-128
35. Joshipura KJ, Hung HC, Rimm EB, Willett WC, Ascherio A (2003) Periodontal disease, tooth loss, and incidence of ischemic stroke. *Stroke* 34:47-52
36. Hung HC, Willett W, Merchant A, Rosner BA, Ascherio A, Joshipura KJ (2003) Oral health and peripheral arterial disease. *Circulation* 107:1152-1157