

cant even after been adjusted for age, BMI, and contraceptive use. The risk reduction of endometrial cancer was associated not only with breastfeeding itself but also with time since the last breastfeeding. From 1982 to 2000, seven case-control studies conducted in six countries, including four developing countries, examined the relationship between breastfeeding and the risk of endometrial cancer. Four early studies, two of which were Japanese, failed to support an association (Kelsey et al. 1982; Brinton et al. 1992; Hirose et al. 1996, 1999). Three recent Western studies, however, suggested a protective effect of breastfeeding (Rosenblatt and Thomas 1995; Salazar-Martinez et al. 1999; Newcomb and Trentham-Dietz 2000). This effect was more pronounced with recent breastfeeding, diminishing as the history of breastfeeding became more remote (Rosenblatt and Thomas 1995; Newcomb and Trentham-Dietz 2000). Our findings were consistent with those of the latter studies, making this the first report that notes an inverse association between breastfeeding and the risk of endometrial cancer among Japanese women.

Exposure of the endometrium to estrogen in the absence of progesterone is thought to increase the risk of endometrial cancer (Key and Pike 1988- see comment for citation). In lactating women, the ovarian cycle is suppressed and blood estrogen levels are reduced (Baird et al. 1979). In the case of oral contraceptives, progesterone continually opposes estrogen, minimizing the duration of time the endometrium is exposed to unopposed estrogen. Thus, suppression of circulating estrogen levels, or opposition of estrogen by progesterone, may represent a biological mechanism accounting for the protective effects of pregnancy, oral contraceptives, and breastfeeding against carcinogenesis of endometrial tissue.

Among Japanese women, the birth rate decreased 28.1 to 9.3 per 1,000 during 1950-2000, and the proportion of women who exclusively breastfed decreased from 70.5% to 44.8% during the same period (Kaneda 2003). In our study, the proportion of women who breastfed for 13 months was 52.4%. The observed lower risk associated with breastfeeding in this study sug-

gests that the recent increase in incidence of endometrial cancer in Japan may be in part attributed to a decrease in both the number of pregnancies and the prevalence of breastfeeding.

A limitation of this study was its lack of age matching. This resulted in a mean age of cases that was 6 years older than that of controls. It is unlikely that the lack of age matching resulted in serious distortion of our observations because all analyses were adjusted for continuous age. Furthermore, the findings in these studies were consistent with data obtained in several previous studies. Another limitation of the study was the small number of control. To overcome these limitations, in progress is our new case control study which matched ages of cases and controls and included two times more subjects of control. These data will confirm the present observations.

Acknowledgments

This study was supported by a grant for Scientific Research in a priority area from the Ministry of Education, Science, Sports and Culture, Japan, a grant from the Ministry of Education, Science, Sports and Culture, Japan, a grant from the Ministry of Health, Labor and Welfare, Japan, and the 21st Century COE Program Special Research Grant (Tohoku University) from the Ministry of Education Science, Sports and Culture, Japan.

References

- Austin, H., Austin, J.M., Partridge, E.E., Hatch, K.D. & Shingleton, H.M. (1991) Endometrial cancer, obesity, and body fat distribution. *Cancer Res.*, **51**, 568-572.
- Baird, D.T., McNeilly, A.S., Sawers, R.S. & Sharpe, R.M. (1979) Failure of estrogen-induced discharge of luteinizing hormone in lactating women. *J. Clin. Endocrinol. Metab.*, **49**, 500-506.
- Brinton, L.A., Berman, M.L., Mortel, R., Twiggs, L.B., Barrett, R.J., Wilbanks, G.D., Lannom, L. & Hoover, R.N. (1992) Reproductive, menstrual, and medical risk factors for endometrial cancer: results from a case-control study. *Am. J. Obstet. Gynecol.*, **167**, 1317-1325.
- Herrinton, L.J., Voigt, L.F., Weiss, N.S., Beresford, S.A. & Wingo, P.A. (2001) Risk factors for synchronous primary endometrial and ovarian cancers. *Ann. Epidemiol.*, **11**, 529-533.
- Hirose, K., Tajima, K., Hamajima, N., Takezaki, T., Inoue, M., Kuroishi, T., Kuzuya, K., Nakamura, S. & Tokudome, S. (1996) Subsite (cervix/endometrium)-specific risk and protective factors in uterus cancer. *Jpn. J. Cancer Res.*, **87**, 1001-1009.
- Hirose, K., Tajima, K., Hamajima, N., Kuroishi, T., Kuzuya, K., Miura, S. & Tokudome, S. (1999) Comparative case-

- referent study of risk factors among hormone-related female cancers in Japan. *Jpn. J. Cancer Res.*, **90**, 255-261.
- Inoue, M., Okayama, A., Fujita, M., Enomoto, T., Tanizawa, O. & Ueshima, H. (1994) A case-control study on risk factors for uterine endometrial cancer in Japan. *Jpn. J. Cancer Res.*, **85**, 346-350.
- Iemura, A., Douchi, T., Yamamoto, S. & Nagata, Y. (2000) Body fat distribution as a risk factor of endometrial cancer. *J. Obstet. Gynaecol. Res.*, **26**, 421-425.
- Kalandidi, A., Tzonou, A., Lipworth, L., Gamatsi, I., Filippa, D. & Trichopoulos, D. (1996) A case-control study of endometrial cancer in relation to reproductive, somatometric, and life-style variables. *Oncology*, **53**, 354-359.
- Kaneda, I. (2003) Maternal and child health statistics of Japan, edited by Tokyo: Mothers' and Children's Health Organization.
- Kelsey, J.L., LiVolsi, V.A., Holford, T.R., Fischer, D.B., Mostow, E.D., Schwartz, P.E., O'Connor, T. & White, C. (1982) A case-control study of cancer of the endometrium. *Am. J. Epidemiol.*, **116**, 333-342.
- Kirschner, M.A., Ertel, N. & Schneider, G. (1981) Obesity, hormones, and cancer. *Cancer Res.*, **41**, 3711-3717.
- McPherson, C.P., Sellers, T.A. & Potter, J.D. (1996) Reproductive factors and risk of endometrial cancer. The Iowa Women's Health Study. *Am. J. Epidemiol.*, **143**, 1195-1202.
- Newcomb, P.A. & Trentham-Dietz, A. (2000) Breastfeeding practices in relation to endometrial cancer risk, USA. *Cancer Causes Control.*, **11**, 663-667.
- Rosenblatt, K.A. & Thomas, D.B. (1995) Prolonged lactation and endometrial cancer. WHO Collaborative Study of Neoplasia and Steroid Contraceptives. *Int. J. Epidemiol.*, **24**, 499-503.
- Salazar-Martinez, E., Lazcano-Ponce, E.C., Gonzalez Lira-Lira, G., Escudero De los Rios, P., Salmeron-Castro, J. & Hernandez-Avila, M. (1999) Reproductive factor of ovarian and endometrial cancer risk in a high fertility population in Mexico. *Cancer Res.*, **59**, 3658-3662.
- Schapira, D.V., Kumar, N.B., Lyman, G.H., Cavanagh, D., Roberts, W.S. & LaPolla, J. (1991) Upper-body fat distribution and endometrial cancer risk. *JAMA*, **266**, 1808-1811.
- Shu, X.O., Brinton, L.A., Zheng, W., Swanson, C.A., Hatch, M.C., Gao, Y.T. & Fraumeni, J.F., Jr. (1992) Relation of obesity and body fat distribution to endometrial cancer in Shanghai, China. *Cancer Res.*, **52**, 3865-3870.
- Zumoff, B. (1982) Relationship of obesity to blood estrogens. *Cancer Res.*, **42**, Suppl. 8, 3289s-3294s.

Lymph-vascular space invasion and number of positive para-aortic node groups predict survival in node-positive patients with endometrial cancer[☆]

Hidemichi Watari^{*}, Yukiharu Todo, Mahito Takeda, Yasuhiko Ebina,
Ritsu Yamamoto, Noriaki Sakuragi

Department of Obstetrics and Gynecology, Hokkaido University School of Medicine, North 15, West 7, Kita-Ku,
Sapporo, 060-8638, Japan

Received 19 May 2004
Available online 5 January 2005

Abstract

Objective. The aim of this study was to determine pathologic variables associated with disease-specific survival of node-positive patients with endometrial carcinoma treated with combination of surgery including pelvic and para-aortic lymphadenectomy and adjuvant chemotherapy.

Methods. Survival of 55 node-positive endometrial carcinoma patients prospectively treated with surgery and adjuvant chemotherapy between 1982 and 2002 at Hokkaido University Hospital was compared to various histopathologic variables. All patients underwent primary surgical treatment including pelvic and para-aortic lymphadenectomy followed by adjuvant chemotherapy consisting of intravenous cisplatin, doxorubicin, and cyclophosphamide. Survival analyses were performed by the Kaplan–Meier curves and the log-rank test. Independent prognostic factors were determined by multivariate Cox regression analysis using a forward stepwise selection.

Results. Among 303 consecutive endometrial cancer patients treated during the period of this study, 55 patients (18.2%), including 44 without peritoneal metastasis (FIGO stage IIIc) and 11 with peritoneal metastasis (FIGO stage IV), were found to have retroperitoneal lymph node metastasis. Multivariate Cox regression analysis revealed that peritoneal metastasis and lymph-vascular space invasion (LVSI) were independently related to poor survival in node-positive endometrial carcinoma. The estimated 5-year survival rate of stage IIIc patients with or without moderate/prominent LVSI was 50.9% and 93.3%, respectively with statistically significant difference ($P = 0.0024$). The estimated 5-year survival rate of stage IV patients was 20.0%. Prognosis of stage IIIc patients could be stratified into three groups by the number of positive para-aortic node (PAN) with an estimated 5-year survival rate of 86.4% for no positive PAN ($n = 23$), 60.4% for one positive PAN ($n = 13$), and 20.0% for ≥ 2 positive PAN ($n = 8$). The difference of survival rate between no or one positive PAN and ≥ 2 positive PAN was statistically significant ($P = 0.0007$ for no positive PAN vs. ≥ 2 positive PAN, $P = 0.0319$ for one positive PAN vs. ≥ 2 positive PAN). Multivariate analysis including number of positive PAN groups showed that LVSI, number of positive PAN groups were independent prognostic factors for survival. Survival of patients with stage IIIc disease could be stratified into three groups by combination of LVSI and number of positive PAN groups with an estimated 5-year survival rate of 93.3% for no or one positive PAN group with nil or minimal LVSI, 62.6% for no or one positive PAN group with intermediate or prominent LVSI, and 20.0% for ≥ 2 positive PAN groups irrespective of LVSI ($P = 0.0002$ for no or one positive PAN group with nil or minimal LVSI vs. ≥ 2 positive PAN groups, $P = 0.0223$ for no or one positive PAN group with nil or minimal LVSI vs. no or one positive PAN group with intermediate or prominent LVSI, $P = 0.0388$ for no or one positive PAN group with intermediate or prominent LVSI vs. ≥ 2 positive PAN groups).

Conclusions. LVSI and number of positive PAN groups were independent prognostic factors for stage IIIc endometrial cancer patients. Postoperative therapy and follow-up modality need to be individualized according to LVSI and the number of positive PAN for stage IIIc patients. New molecular markers to predict the prognosis of endometrial cancer patients preoperatively should be found for individualization

[☆] This work was presented at the 40th annual meeting of American Society of Clinical Oncology (ASCO), New Orleans, LA, June 2004.

^{*} Corresponding author. Fax: +81 11 706 7711.

E-mail address: watarih@med.hokudai.ac.jp (H. Watari).

of treatment. New chemotherapy regimen including taxane needs to be considered as an adjuvant therapy for patients with node-positive endometrial cancer.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Endometrial carcinoma; Prognostic factor; Stage IIIc; Para-aortic lymph node metastasis; Lymph-vascular space invasion

Introduction

The incidence of endometrial carcinoma is rapidly increasing in Japan, with an estimated 4800 new cases in the year 2000; this number is still much lower than that in the United States, which was 40,100 in the year 2003 [1].

A majority of patients are diagnosed as being without clinical evidence of extrauterine spread, the International Federation of Gynecology and Obstetrics (FIGO) stages I and II, and have a 5-year survival of approximately 90%. Involvement of pelvic and para-aortic lymph nodes has been recognized as a poor prognostic factor in endometrial carcinoma. The new FIGO surgical staging system classifies endometrial carcinoma with metastasis to the pelvic and/or para-aortic lymph nodes as stage IIIc. The overall 5-year survival for patients with stage IIIc disease varies considerably, depending on the presence of various risk factors, extent of lymphadenectomy, and postoperative adjuvant therapy. Data on surgically staged endometrial cancer patients with nodal involvement have been limited [2–5]. The distribution of nodal disease and its importance in predicting survival are largely unknown, since various procedures have been used to assess para-aortic and pelvic nodes in endometrial cancer, that is, biopsies from enlarged nodes only, selective nodal sampling from multiple sites, pelvic lymphadenectomy, and para-aortic and pelvic lymphadenectomy. It is obvious that para-aortic and pelvic lymphadenectomy is most accurate of these methods. However, para-aortic and pelvic lymphadenectomy is not regarded as the standard surgical procedure for endometrial cancer because the therapeutic significance of the procedure has not yet been sufficiently demonstrated [6,7]. We have routinely performed complete lymphadenectomy in all patients with endometrial cancer because (i) nodal status is the most important prognosticator [2], (ii) results of lymphadenectomy allow tailoring of postoperative adjuvant treatment, (iii) there is an apparent small survival advantage after lymphadenectomy [8], and (iv) there is no increased morbidity with lymphadenectomy [9].

We have previously reported that para-aortic lymph node metastasis is an independent prognostic factor for endometrial cancer patients as well as cell type, grade and LVSI [10,11]. However, the fact that patients with para-aortic lymph node metastasis sometimes have shown long survival suggest that specific prognostic factors may exist for node-positive endometrial carcinoma.

In this study, we tried to identify the independent histopathologic prognostic factors for endometrial carcinoma patients with lymph node metastasis, who were

uniformly treated with extensive surgical staging that includes modified radical hysterectomy and systematic pelvic and para-aortic lymphadenectomy followed by adjuvant chemotherapy. We also described the distribution of nodal disease in FIGO stage IIIc endometrial cancer and evaluated whether nodal distribution is related to survival.

Materials and methods

Patients

A total of 303 patients with endometrial carcinoma underwent primary radical surgical treatment from 1982 to 2002 at the Department of Obstetrics and Gynecology, Hokkaido University Hospital. Among 303 patients, 55 patients showed positive retroperitoneal lymph nodes. All subjects underwent modified radical hysterectomy, bilateral salpingo-oophorectomy, and systematic retroperitoneal lymphadenectomy which consisted of complete dissection of pelvic and para-aortic lymph nodes from the femoral ring to the level of the renal vein. All lymphatic tissues that surrounded the arteries and veins were completely removed. The FIGO (1988) stage of the patients was as follows: 44 (80%) stage IIIc and 11 (20%) stage IV. Stage IV disease with distant metastasis (liver or lung metastasis) was excluded from this analysis. We, therefore, defined stage IV patients as node-positive patients with peritoneal metastasis. Median follow-up period was 43 months (1–200 months). The patients' characteristics are shown in Table 1. All patients were treated with an adjuvant chemotherapy of CAP (cyclophosphamide: 350 mg/m², adriamycin: 40 mg/m² and cisplatin: 50–70 mg/m²) every 3 weeks.

The following histopathologic prognostic factors were included in the survival analyses: FIGO (1988) stage, histologic subtype, depth of myometrial invasion, architectural grade (AG), nuclear grade (NG), LVSI, ovarian metastasis, PAN metastasis, and number of positive PAN group. All risk factors except number of positive PAN were determined as previously described [10,11]. We defined a node group by its laterality and location. Pelvic node groups include common iliac nodes, external iliac nodes, internal iliac nodes, obturator nodes, median deep inguinal nodes, deep inguinal nodes, parametrial nodes, and sacral nodes. PAN metastasis was investigated by dividing the metastatic group according to the sites above and below the inferior mesenteric artery and left and right sections bordering on the midline of the aorta as previously described [3]. In this analysis, caval (pre-, peri-, retro-) and aortocaval nodes were

Table 1
Clinicopathologic characteristics of 55 patients with lymph node metastasis

	No.	%
FIGO stage (1988)		
IIIc	44	80.0
IV	11	20.0
Histologic subtype		
Endometrioid	43	78.2
Serous/Clear	12	21.8
Architectural grade		
1	16	29.1
2	27	49.1
3	12	21.8
Nuclear grade		
1	14	25.4
2	25	45.5
3	16	29.1
Depth of myometrial invasion		
≤ 1/2	15	27.3
> 1/2	40	72.7
Lymph-vascular space invasion		
Nil/minimal	20	36.4
Moderate/prominent	35	63.6
Cervical invasion		
Negative	28	50.9
Positive	27	49.1
Ovarian metastasis		
Negative	37	67.3
Positive	18	32.7
Para-aortic lymph node metastasis		
Negative	26	47.3
Positive	29	52.7

included in right para-aortic nodes. PAN groups, therefore, consists of four groups.

Statistics

Correlation between the variables was analyzed using χ^2 test. Patients survival was calculated using Kaplan–Meier method. The significance of the survival difference was examined by the log-rank test. Univariate and multivariate survival analyses were performed using the Cox regression model with disease-specific overall survival as the outcome measure. Forward stepwise procedure was used to select the independent variable in multivariate analysis. $P < 0.05$ was considered statistically significant. Statistical analyses were performed with the Statview software package (SAS Institute, Inc, Cary, NC).

Results

Univariate and multivariate survival analysis for all node-positive patients

Age of patients ranged from 40 to 75 (median 58) years. The univariate analysis revealed that the FIGO (1988) stage (IIIc vs. IV, $P < 0.0001$), the histologic subtype (endometrioid vs. serous/clear, $P = 0.0216$), architectural grade (G1/2

vs. G3, $P = 0.0024$), nuclear grade (G1 vs. G2/3, $P = 0.0201$), depth of myometrial invasion (absence or presence of serosal invasion, $P < 0.0001$), LVSI (none/minimal vs. moderate/prominent, $P = 0.0026$), cervical invasion ($P = 0.0159$), and PAN metastasis ($P = 0.0078$) were shown to be related to poor survival. Ovarian metastasis was not related to survival ($P = 0.3306$) (Table 2).

Multivariate analysis, which included the prognostic factors determined by univariate analysis to have statistical significance, was performed using a forward stepwise procedure (Table 3). It was shown that FIGO (1988) stage ($P < 0.0001$) and LVSI ($P = 0.0002$) were independent prognostic factors. We could stratify the patients into three prognostic risk-groups by integrating those two histopathologic risk factors, that is, low risk group (group A: stage IIIc with nil/minimal LVSI, $n = 19$), intermediate risk group (group B: stage IIIc with moderate/prominent LVSI, $n = 25$) and high risk group (group C: stage IV with any LVSI, $n = 11$) with an estimated 5-year survival rate of 93.3%, 50.9%, and 20.0%, respectively (Fig. 1). There was statistically significant difference of survival rate between each group (A vs. B: $P = 0.0024$, B vs. C: $P < 0.0001$, A vs. C: $P < 0.0001$). Because prognostic impact of FIGO (1988) stage was extremely strong for the survival of node-positive patients, we performed further analysis on stage IIIc patients alone ($n = 44$).

Lymph node metastasis in stage IIIc patients

Incidences of pelvic lymph node (PLN) metastasis alone, PAN metastasis alone, and both PLN and PAN metastasis were 52.3% (23/44), 4.5% (2/44), and 43.2% (19/44),

Table 2
Univariate and multivariate Cox regression analysis of prognostic factors of node positive endometrial carcinoma

Prognostic factor	Univariate		Multivariate	
	<i>P</i> value	<i>P</i> value	Risk ratio	95% CI
FIGO (1988) stage	< 0.0001	11.2	4.0–31.3	< 0.0001
Histologic subtype	0.0216	–	–	NS
Architectural grade	0.0024	–	–	NS
Nuclear grade	0.0201	–	–	NS
Lymph-vascular space invasion	0.0026	9.3	2.1–41.7	0.0033
Myometrial invasion	< 0.0001	–	–	NS
Cervical invasion	0.0159	–	–	NS
Ovarian metastasis	0.3306	–	–	S
Para-aortic node metastasis	0.0078	–	–	NS

NS: not significant.

FIGO (1988) stage: stage IIIc vs stage IV determined by the presence of peritoneal metastasis.

Tumor cell type: endometrioid vs serous/clear cell.

Architectural grade: G1/2 vs G3.

Nuclear grade: G1 vs G2/3.

Lymph-vascular space invasion: (–)/(+) vs (++)/(+++).

Myometrial invasion: serosal invasion (–) vs (+).

Cervical invasion: (–) vs (+).

Ovarian metastasis: (–) vs (+).

Para-aortic lymph node metastasis: (–) vs (+).

Table 3
Univariate and multivariate Cox regression analysis of prognostic factors of stage IIIc endometrial carcinoma

Prognostic factor	Univariate <i>P</i> value	Multivariate		
		Risk ratio	95% CI	<i>P</i> value
Histologic subtype	0.7226	–	–	NS
Architectural grade	0.3911	–	–	NS
Nuclear grade	0.0605	–	–	NS
Lymph-vascular space invasion	0.0173	8.8	1.1–71.4	0.0413
Myometrial invasion	0.6036	–	–	NS
Cervical invasion	0.4577	–	–	NS
Ovarian metastasis	0.1815	–	–	NS
Number of positive PAN	0.0016	3.9	1.2–13.0	0.0260

NS: not significant.

Tumor cell type: endometrioid vs serous/clear cell.

Architectural grade: G1/2 vs G3.

Nuclear grade: G1 vs G2/3.

Lymph-vascular space invasion: (-)/(+) vs (++)/(+++).

Myometrial invasion: serosal invasion (-) vs (+).

Cervical invasion: (-) vs (+).

Ovarian metastasis: (-) vs (+).

Number of positive PAN: 0, 1 vs ≥ 2 .

respectively. The estimated 5-year survival rate of patients without or with PAN metastasis was 86.4%, 48.1%, respectively (Fig. 2). The difference was statistically significant ($P = 0.0108$).

Prognostic impact of the number of positive PLN groups in stage IIIc patients

The estimated 5-year survival rate for patients with one positive PLN group was 79.3% and that for patients with ≥ 2 positive PLN groups was 60.8%. The difference of survival was not statistically significant. However, 21 of 22 patients (95.5%) with no or one positive PLN group had no or one positive PAN group and only one of 22 patients (4.5%) had two positive PAN groups, while 15 of 22 patients (68.2%) with ≥ 2 positive PLN groups had no or one positive PAN group and 7 of 22 patients (31.8%) had ≥ 2




Fig. 1. Survival of node-positive patients with endometrial carcinoma by combination of FIGO (1988) stage and LVSI.




Fig. 2. Survival of patients with stage IIIc endometrial carcinoma by PAN metastasis.

2 positive PAN groups. There was a statistically significant difference in incidence of ≥ 2 positive PAN groups between patients who had no or one positive PLN group and those who had ≥ 2 positive PLN groups ($P = 0.023$). (Table 4).

Prognostic impact of the number of positive PAN groups in stage IIIc patients

Fig. 3 shows the survival of stage IIIc patients according to the number of positive PAN groups. The estimated 5-year survival rate was 86.4% for patients without positive PAN group ($n = 23$), 60.4% for those with one positive PAN group ($n = 13$), and 20.0% for those with ≥ 2 positive PAN groups ($n = 8$). There was statistically significant difference between no positive PAN group and ≥ 2 positive PAN groups ($P < 0.0007$), between one positive PAN group and ≥ 2 positive PAN groups ($P = 0.0319$). There was no statistically significant difference between no positive PAN group and one positive PAN group ($P = 0.1354$).

Univariate and multivariate survival analysis for stage IIIc patients

Since the number of positive PAN group was shown to have significant impact on the survival of stage IIIc patients,

Table 4

Incidence of para-aortic lymph node metastasis according to number of positive pelvic lymph node groups in stage IIIc endometrial cancer patients

	Positive PAN group		Total
	0, 1	≥ 2	
positive PLN group			
0, 1	21	1	22
> 2	15	7	22
Total	36	8	44

$P = 0.0023$

PLN: pelvic lymph node, PAN: para-aortic lymph node.

There was a statistically significant difference in incidence of ≥ 2 positive PAN groups between patients who had no or one positive PLN group and those who had ≥ 2 positive PLN groups ($P = 0.023$).




Fig. 3. Survival of patients with stage IIIc endometrial carcinoma by the number of positive PAN group.

we included number of positive PAN groups in the univariate analysis instead of presence or absence of PAN metastasis. The univariate analysis revealed that the LVSI ($P = 0.0173$), number of positive PAN groups ($P = 0.0016$) were shown to be related to poor survival. Histologic subtype ($P = 0.7226$), architectural grade ($P = 0.3911$), nuclear grade ($P = 0.0605$), depth of myometrial invasion ($P = 0.6036$), cervical invasion ($P = 0.4577$), ovarian metastasis ($P = 0.1815$) were not related to poor survival (Table 3, Fig. 4).

Multivariate analysis revealed that both LVSI ($P = 0.0413$) and number of positive PAN groups ($P = 0.026$) were independent prognostic factors. Survival of patients with stage IIIc disease could be stratified into three groups by combination of LVSI and number of positive PAN group with an estimated 5-year survival rate of 93.3% for no or one positive PAN group with nil or minimal LVSI (group D), 62.6% for no or one positive PAN group with intermediate or prominent LVSI (group E), and 20.0% for ≥ 2 positive PAN groups irrespective of LVSI (group F). The difference of survival rate between each group was statistically significant ($P = 0.0002$ for group D vs. group F, $P = 0.023$ for group D vs. group E, $P = 0.0388$ for group E vs. group F).

Discussion

Stages I and II endometrial carcinomas have shown a favorable prognosis by combination of surgery, radiotherapy, and/or chemotherapy. Some histopathologic factors have been found to be related to prognosis of endometrial carcinoma. Lymph node metastasis is one of the most important prognostic factors of endometrial carcinoma and advanced endometrial carcinoma with lymph node metastasis (IIIc/IV) has been shown to have poorer prognosis. In our series of patients treated in the same manner, the prognosis for patients with stage IIIa endometrial carcinoma was excellent with an estimated 5-year survival of over 90%

(patients with stage IIIb disease were not found in our series). However, the prognosis for patients with stage IIIc endometrial carcinoma with an estimated 5-year survival rate of 79.6% was poorer than that of stage IIIa in spite of intensive treatment consisting of extended surgery including pelvic and para-aortic lymphadenectomy and systemic adjuvant chemotherapy. We, therefore, performed retrospective analysis on the prognostic factors for node-positive patients to determine appropriate therapeutic and follow-up modality to achieve their favorable prognosis.

Concerning the distribution of lymph node metastases, 95.5% (42/44) of patients with nodal disease had pelvic node metastases and 45.2% (19/42) of patients with PLN metastases had concomitant PAN metastases. McMeekin et al. [4] analyzed nodal distribution in 47 cases of stage IIIc endometrial cancer and found that an increasing number of positive PLN was associated with PAN metastasis. Our result on nodal distribution in stage IIIc patients is similar to McMeekin et al. [4], i.e., patients with single positive PLN group rarely have multiple positive PAN groups. We also found that positive aortic nodes were associated with poorer prognosis than were positive pelvic nodes alone (estimated 5-year survival 48.1% for positive PAN vs. 86.4% for negative PAN, $P = 0.0108$), suggesting that involvement of pelvic lymph nodes alone does not necessarily carry a poor prognosis as previously reported by Onda et al [12] who performed the same operative procedure as ours. This can be explained in part due to the therapeutic significance of our operative procedure including systematic pelvic and para-aortic lymphadenectomy. Para-aortic lymphadenectomy until just below the renal vein may contribute to the favorable survival of patients with multiple positive PLN groups.

In this study, we firstly reported that the survival of patients with stage IIIc endometrial carcinoma can be stratified by the number of positive PAN groups. The survival of patients without positive PAN (PLN metastasis




Fig. 4. Survival of patients with stage IIIc endometrial carcinoma by combination of LVSI and number of positive PAN group.

alone), with one positive PAN group, and with ≥ 2 positive PAN groups was 86.4%, 60.4%, 20.0%, respectively. The survival of patients with ≥ 2 positive PAN groups was much worse than others. Notably, there was no statistically significant difference between the patients without positive PAN group and those with one positive PAN group ($P = 0.14$), suggesting that single PAN metastasis is still a local disease that can be cured by complete lymphadenectomy and subsequent chemotherapy. When patients have multiple PAN metastasis, lymphadenectomy does not prolong survival.

Radiotherapy and chemotherapy have been employed as adjuvant therapies for endometrial cancer. Radiotherapy has been considered as a standard adjuvant therapy in Western countries. The result of GOG 122, however, clearly demonstrated that adjuvant chemotherapy (adriamycin and CDDP) significantly improved progression free survival and overall survival than adjuvant radiotherapy (whole abdominal radiotherapy) for stage III/IV patients [13], indicating that chemotherapy should be considered as a standard adjuvant therapy for endometrial cancer. Systemic chemotherapy has been widely accepted as a standard adjuvant therapy for endometrial cancer in Japan. Adriamycin has been used as a key drug for endometrial cancer. We have used CAP regimen for endometrial carcinoma with risk factors for recurrence. However, the poorer survival of stage IIIc patients with multiple positive PAN groups than single positive PAN group, who received adjuvant chemotherapy (CAP), clearly indicates that we should consider a new chemotherapeutic regimen to improve prognosis of node-positive patients. The most promising drug for endometrial cancer is taxane [14].

In this study, we found that LVSI and number of positive PAN group are independent prognostic factors by multivariate analysis, indicating that para-aortic lymphadenectomy should be routinely included in the surgical procedure for endometrial cancer to predict the survival of node-positive patients. We also conclude that we should investigate LVSI with more careful attention for node-positive patients. Careful investigation of LVSI, however, is time-consuming and it is impossible to evaluate LVSI preoperatively and during operation by frozen section. To individualize the therapeutic modality for each patient, we need to search for new molecular markers which can be easily assessed and reflect disease status in regard to LVSI preoperatively. There have been few reports on the useful molecular markers for survival of endometrial carcinoma. We reported that p53 overexpression by immunohistochemical staining was found to be an independent prognostic factor and the estimated 5-year survival rate of patients with stage III/IV disease without p53 overexpression was significantly better than that with p53 overexpression, indicating that p53 missense mutation, which is closely related to immunohistochemical p53 overexpression, have a significant prognostic impact on the survival of advanced endometrial carcinoma [15]. Kanamori et al. [16] reported

that PTEN expression was found to be associated with prognosis for patients with advanced endometrial carcinoma undergoing postoperative chemotherapy. Yokoyama et al. [17] reported that high levels of immunoreactivity for vascular endothelial growth factor (VEGF)-D in stromal cells and its receptor, VEGF-R-3 in carcinoma cells were independent prognostic factors in endometrial carcinoma. We need to further investigate more useful prognostic factors for endometrial carcinoma using molecular biological techniques.

References

- [1] Jemal A, Murray T, Samuels A, Ghafoor A, Ward E, Thun MJ. Cancer statistics, 2003. *CA Cancer J Clin* 2003;53:5–26.
- [2] Morrow CP, Bundy BN, Kurman RJ, Creasman WT, Heller P, Homesley HD, et al. Relationship between surgical–pathological risk factors and outcome in clinical stage I and II carcinoma of the endometrium: a Gynecologic Oncology Group study. *Gynecol Oncol* 1991;40:55–65.
- [3] Hirahatake K, Hareyama H, Sakuragi N, Nishiya M, Makinoda S, Fujimoto S. A clinical and pathologic study on para-aortic lymph node metastasis in endometrial carcinoma. *J Surg Oncol* 1997;65:82–7.
- [4] McMeekin D, Iashbrook D, Gold M, Scribner DR, Kamelle S, Tillmanns TD, et al. Nodal distribution and its significance in FIGO stage IIIc endometrial cancer. *Gynecol Oncol* 2001;82:375–9.
- [5] Katz LA, Andrews SJ, Fanning J. Survival after multimodality treatment for stage IIIc endometrial cancer. *Am J Obstet Gynecol* 2001;184:1071–3.
- [6] Calais G, Descamps P, Vitu L, Body G, Lansac J, Bougnoux P, et al. Is lymphadenectomy useful in the treatment of endometrial carcinoma? *Gynecol Oncol* 1990;38:71–5.
- [7] Faught W, Krepart GV, Lotocki R, Heywood M. Should selective para-aortic lymphadenectomy be part of surgical staging for endometrial cancer? *Gynecol Oncol* 1994;55:51–5.
- [8] Kilgore LC, Partridge EE, Alvarez RD, Austin JM, Shingleton HM, Noojin F, et al. Adenocarcinoma of the endometrium: survival comparisons of patients with and without pelvic node sampling. *Gynecol Oncol* 1995;56:29–33.
- [9] Fanning J, Firestein S. Prospective evaluation of the morbidity of complete lymphadenectomy in endometrial cancer. *Int J Gynecol Cancer* 1998;8:270–3.
- [10] Nishiya M, Sakuragi N, Hareyama H, Ebina Y, Furuya M, Oikawa M, et al. Cox multivariate regression models for estimating prognosis of patients with endometrioid adenocarcinoma of the uterine corpus who underwent through surgical staging. *Int J Cancer* 1998;79:521–5.
- [11] Sakuragi N, Hareyama H, Todo Y, Yamada H, Yamamoto R, Fujino T, et al. Prognostic significance of serous and clear cell adenocarcinoma in surgically staged endometrial carcinoma. *Acta Obstet Gynecol Scand* 2000;79:311–6.
- [12] Onda T, Yoshikawa H, Mizutani K, Mishima M, Yokota H, Nagano H, et al. Treatment of node-positive endometrial cancer with complete node dissection, chemotherapy and radiation therapy. *Br J Cancer* 1997;75:1836–41.
- [13] Randall ME, Brunetto G, Mussetal RS, et al. Whole abdominal radiotherapy versus combination chemotherapy with doxorubicin and cisplatin in advanced endometrial carcinoma (phase III): Gynecologic Oncology Group study no. 122. *Proc Am Soc Clin Oncol* 2003;21:2.
- [14] Ball HG, Blessing JA, Lentz SS, et al. A phase II trial of paclitaxel in patients with advanced or recurrent adenocarcinoma of the endometrium: a Gynecologic Oncology Group study. *Gynecol Oncol* 1996;62:278–81.

- [15] Ohkouchi T, Sakuragi N, Watari H, Nomura E, Todo Y, Yamada H, et al. Prognostic significance of Bcl-2, p53 overexpression and lymph node metastasis in surgically staged endometrial carcinoma. *Am J Obstet Gynecol* 2002;187:353–9.
- [16] Kanamori Y, Kigawa J, Itamochi H, Sultana H, Suzuki M, Ohwada M, et al. PTEN expression is associated with prognosis for patients with advanced endometrial carcinoma undergoing postoperative chemotherapy. *Int J Cancer* 2002;100:686–9.
- [17] Yokoyama Y, Charnock-Jones DS, Licence D, Yanaihara A, Hastings JM, Holland CM, et al. Expression of vascular endothelial growth factor (VEGF)-D and its receptor, VEGF receptor 3, as a prognostic factor in endometrial carcinoma. *Clin Cancer Res* 2003;9:1361–9.

Functional analysis of *p53* gene and the prognostic impact of dominant-negative *p53* mutation in endometrial cancer

Noriaki Sakuragi^{1*}, Hidemichi Watari¹, Yasuhiko Ebina¹, Ritsu Yamamoto¹, Eric Steiner², Heinz Koelbl², Masahiro Yano³, Mitsuhiro Tada³ and Tetsuya Moriuchi³

¹Department of Gynecology, Hokkaido University Graduate School of Medicine and School of Medicine, Sapporo, Japan

²Department of Obstetrics and Gynecology, Mainz University, Mainz, Germany

³Division of Cancer-Related Genes, Institute for Genetic Medicine, Hokkaido University, Sapporo, Japan

In addition to the loss of function, mutant *p53* can possess a dominant-negative effect on wild-type *p53* and may also exert gain-of-function activity. It is not clear whether the functional status of *p53* mutation contributes to differences in outcome in endometrial cancer. We collected a total of 92 RNA samples of high quality from endometrial cancer tissues, and the samples were subjected to yeast functional assay and sequencing for *p53* mutations. The detected mutant *p53* genes were further investigated for their dominant-negative activity using a yeast-based transdominance assay. *p53* mutation was found in 24 out of 92 (26.1%) tumors, of which 10 exhibited no dominant-negative activity (recessive mutation) and 14 showed dominant-negative activity. Dominant-negative *p53* mutation was related to advanced stages ($p = 0.01$), non-endometrioid type tumors ($p = 0.01$) and grade 3 tumors ($p = 0.04$). The patients with dominant-negative mutation had significantly shorter survival than patients with no mutation ($p < 0.0001$) and those with a recessive mutation ($p = 0.01$) in the *p53* gene. No difference in survival was found between the patients with tumors harboring a recessive *p53* mutation and those with tumors harboring a wild-type *p53*. Multivariate analysis revealed that dominant-negative *p53* mutation ($p = 0.019$), FIGO stage ($p = 0.0037$) and histologic subtype ($p = 0.014$) were independently related to patient survival. Dominant-negative *p53* mutation was the most important prognostic factor for stage III/IV endometrial cancer ($p = 0.0023$). In conclusion, dominant-negative *p53* mutation is often found in advanced stages and aggressive histologic subtypes of endometrial cancer and it is a strong predictor of survival of patients with advanced endometrial cancer. To elucidate further the role of *p53* mutation in endometrial cancer, it is necessary to investigate gain-of-function activity involving dominant-negative *p53* mutant proteins.

© 2005 Wiley-Liss, Inc.

Key words: endometrial cancer; *p53*; mutation; dominant negative; survival; serous adenocarcinoma

The *p53* tumor suppressor gene is mutated in about 50% of all tumors,^{1–3} and more than 19,000 different somatic mutations have been identified.⁴ Mutation of the *p53* gene plays a key role in the carcinogenesis and progression of many different malignancies, including endometrial cancer. *p53* overexpression has been shown to predict patient survival in endometrial cancer.^{5–7} *p53* overexpression, as determined by immunohistochemistry (IHC), is a surrogate marker of missense mutation of *p53* protein. Because MDM2 is a transcriptional target of *p53*, loss of *p53* function reduces the production of MDM2. MDM2 degrades *p53* protein through ubiquitination of the protein, and reduced MDM2 production will lead to an accumulation of *p53* protein in the nucleus, which is detected as overexpression by IHC.⁸ IHC is a convenient method for the investigation of *p53* status. However, it can be affected by many factors, such as antibody used, antigen retrieval technique and subjectivity of criteria for *p53* overexpression. *p53* overexpression does not necessarily correspond to *p53* gene mutation. One report showed that only 32% of tumors with exclusively nuclear staining were found to contain a *p53* gene mutation.⁶ Dominant-negative *p53* mutation will lead to decreased MDM2 production, irrespective of the status of the second allele of *p53*. *p53* mutations abolishing the production of *p53* protein will lead to loss of *p53* protein when it is associated with loss of heterozygosity (LOH) of the second allele. Therefore, *p53* overexpres-

sion can be related to both dominant-negative mutation and recessive missense mutation with LOH. It is therefore reasonable to expect that *p53* overexpression does not necessarily correspond to a dominant-negative mutation of *p53*.

At least 2 distinct pathways have been proposed that contribute to cisplatin-induced apoptosis *in vitro*. One involves *p53* tumor suppressor protein, and the other is mediated by the *p53*-related protein p73. Inhibition of p73 function by dominant-negative p73 proteins or by mutant *p53* abrogates apoptosis and cytotoxicity induced by these agents.⁹ Therefore, investigation into the status of p73 in endometrial cancer is expected to provide further information regarding the response to adjuvant chemotherapy in cases involving this type of cancer.

An understanding of the role played by *p53* mutation in endometrial cancer may lead to tailored treatment planning and more rational targeted approaches for treating this disease. *p53* gene mutation and LOH result in the loss of *p53* function. In addition to the loss of function, mutant *p53* can possess a dominant-negative effect that suppresses wild-type *p53*, and which may also exert gain-of-function activity.¹⁰ There have been many studies of the prognostic significance of and/or therapeutic outcome related to the type of *p53* mutations. However, the results of such studies have been inconsistent,¹¹ and the significance of the dominant-negative function in terms of both prognosis and therapeutic success remains unclear at present.¹²

In order to elucidate the prognostic importance of dominant-negative activity in endometrial cancer, we surveyed the functional status of *p53* protein in endometrial carcinoma using a yeast *p53* functional assay and a transdominance assay. The yeast *p53* functional assay tests the ability of *p53* to activate transcription *in vivo* in yeast.^{13,14} A modification of this method (transdominance assay) can identify the dominant-negative/recessive properties of the mutant *p53*.¹⁵ This is the first report to compare directly the functional status of *p53* (*i.e.*, the presence or absence of dominant-negative activity) to the survival of patients with endometrial cancer.

Material and methods

Tissue specimens

A total of 92 endometrial carcinoma tissue samples, which were obtained from the resected uterus of patients with endometrial carcinoma treated surgically at the Department of Obstetrics and Gynecology, Mainz University in Mainz, Germany, and at the

Presented in part at the 16th EORTC-NCI-AACR Symposium on Molecular Targets and Cancer Therapeutics, Geneva, Switzerland, on 1 October 2004.

*Correspondence to: Department of Obstetrics and Gynecology, Hokkaido University School of Medicine, N-15, W-7, Kitaku, Sapporo 060-8638, Japan. Fax: +81-11-706-7711.

E-mail: sakuragi@med.hokudai.ac.jp

Received 26 August 2004; Accepted after revision 18 January 2005

DOI 10.1002/ijc.21097

Published online 11 April 2005 in Wiley InterScience (www.interscience.wiley.com).

TABLE I - CLINICOPATHOLOGIC CHARACTERISTICS OF PATIENTS WITH ENDOMETRIAL CARCINOMA TREATED IN MAINZ UNIVERSITY AND HOKKAIDO UNIVERSITY

	Mainz, Germany (n = 49)	Sapporo, Japan (n = 43)	p
Age			
< 60	13	26	
> 60	36	17	0.001
FIGO stage			
I/II	37	28	
III/IV	12	15	0.27
Histologic subtype			
Endometrioid	44	35	
Nonendometrioid	5	8	0.25
Grade			
1	17	14	
2	19	17	
3	13	12	0.98
Follow-up period (months)	1-130 (median, 58.5)	5-125 (median, 41.0)	0.11

Department of Gynecology, Hokkaido University Graduate School of Medicine and School of Medicine in Sapporo, Japan, were used for this study. Informed consent was obtained from all study participants. From among the 92 samples, 49 were obtained from German patients and 43 from Japanese patients. Surgical treatment was initiated during the period between April 1990 and January 2000 for the German cohort and between February 1990 and December 2002 for the Japanese cohort. The data from 23 patients of the 43 Japanese cohort participants were reported in a previous study.¹⁶ The clinicopathologic variables, namely, FIGO stage, histologic subtype, grade of tumor and follow-up period, did not differ between the 2 cohorts, with the exception of patient age (Table I). The tissue samples included 79 cases of endometrioid-type adenocarcinoma and 13 cases of nonendometrioid adenocarcinoma. The group of nonendometrioid tumors included 11 serous adenocarcinomas, 1 clear cell adenocarcinoma and 1 squamous cell carcinoma. The treatment strategies employed in the 2 institutes were different. At Mainz University, the surgical procedure involved selective lymph node dissection rather than routine systematic lymphadenectomy. At Hokkaido University, routine systematic pelvic and paraaortic lymphadenectomy was carried out. The modality of adjuvant therapy employed at Mainz University was radiotherapy, whereas that used at Hokkaido University was chemotherapy.

RNA extraction and reverse transcription (RT)-PCR

Total RNA was extracted from 100-200 mg of each frozen tissue sample by the guanidinium/phenol/chloroform method (TRIzol reagent; Gibco-BRL, Gaithersburg, MD). RNA integrity was verified by electrophoresis on 1% agarose gel. p53 cDNA was synthesized at 37°C for 1 hr with 200 units of Moloney murine leukemia virus (MMLV) reverse transcriptase (Gibco-BRL) from 1-3 µg of total RNA in 20 µl of RT buffer containing 25 pmol p53-specific primer RT-1 (5'-CGGGAGGTAGAC-3'), 7.5 mM dithiothreitol (DTT), 0.5 mM MgCl₂ and 0.5 mM of each dNTP. The p53 cDNA was PCR-amplified in 20 µl of reaction mixture containing 2 µl of RT reaction product, 1.25 units of *Pfu* DNA polymerase (Stratagene, La Jolla, CA), 10% DMSO, 50 µM of each dNTP and 10 pmol of primers P3 [5'-ATTTGATGCTGTCCCGGACGATATTGAA(s)C-3', where (s) represents a phosphorothioate linkage] and P4 [5'-ACCCTTTTGGACTTCAGGTGGCTGGAGT(s)G-3']. PCR was run on a Thermal Cycler Model 2400 (Perkin-Elmer, Chiba, Japan) at 96°C for 1 min, then for 35 cycles of 95°C for 40 sec, 65°C for 70 sec and 78°C for 90 sec, followed by 78°C for 2 min. Satisfactory amplification was confirmed by examining the PCR product in a 1% agarose gel. Each crude PCR product was used for the transformation of yeast.




FIGURE 1 - An example of positive yeast p53 functional assay (a) and confirmation of p53 mutation by DNA sequencing of the reverse strand: (b) wild-type sequence; (c) missense mutation of CGT to CAT at codon 273.

Plasmids

The yeast expression vector pSS16¹⁷ was digested with excess amounts of *Hind*III and *Stu*I, dephosphorylated with calf intestinal alkaline phosphatase (Takara, Otsu, Japan) and electrophoresed on 1% low-melting-temperature agarose (Sea Plaque agarose; FMC, Rockland, ME). The linearized plasmids were recovered from the gel and purified with Wizard PCR prep kit (Promega, Madison, WI). A gap was created between codons 67 and 347.

Yeast p53 functional assay

The yeast functional assay was performed according to a method described previously.^{14,18} The yeast reporter strain yIG397¹⁵ was used throughout the study. The strain yIG397 contains an integrated plasmid with the ADE2 (phosphoribosylaminoimidazole carboxylase, EC 4.1.1.21) open reading frame under the control of a p53-responsive promoter. The genotype is *MATa ade2-1 leu2-3 112trp1-1 his3-11 15can1-100 ura3-1::[URA3 3xRGC-pCYC1-ADE2]*. When a yeast cell is transformed with a plasmid encoding mutant p53, the cell fails to express ADE2 and forms a red colony due to the accumulation of an oxidized polymerized derivative of phosphoribosyl-aminoimidazole.¹⁹ The yeast was cultured in 100 ml of YPD medium supplemented with 200 µg/ml of adenine until the OD₆₀₀ value reached 0.8. The cells

TABLE II - DISCOVERED *p53* MUTATIONS AND ITS TRANSDOMINANCE IN 92 ENDOMETRIAL CARCINOMAS

Institute and case number	FIGO stage	Histologic subtype	Grade	Mutation in <i>p53</i> gene	Base change	Type of mutation	Transdominance
Hokkaido University							
1	I	Endometrioid adenocarcinoma	2	213 Arg→Stop	CGA→TGA	Nonsense	
2	I	Endometrioid adenocarcinoma	2	321bp deletion (nt 673-993)		Deletion	
3	I	Endometrioid adenocarcinoma	1	133 Met→Arg	ATG→AGG	Missense	
4	I	Endometrioid adenocarcinoma	1	244 Gly→Asp	GGC→GAC	Missense	DN
5	I	Endometrioid adenocarcinoma	2	363bp deletion (nt 385-747)		Deletion	
6	I	Endometrioid adenocarcinoma	1	108 Gly→Ser	GGT→AGT	Missense	
7	I	Endometrioid adenocarcinoma	1	241 Ser→Ala	TCC→GCC	Missense	DN
8	I	Serous adenocarcinoma	3	273 Arg→His	CGT→CAT	Missense	DN
9	II	Serous adenocarcinoma	1	240-243 in frame deletion	AGTTCCTGC ATG→GTG	Deletion	DN
10	III	Endometrioid adenocarcinoma	3	264 Leu→Arg	CTA→CGA	Missense	
11	III	Endometrioid adenocarcinoma	2	280 Arg→Ile	AGA→ATA	Missense	DN
12	III	Endometrioid adenocarcinoma	2	173 Val→Leu	GTG→TTG	Missense	
13	III	Serous adenocarcinoma	2	273 Arg→His	CGT→CAT	Missense	DN
14	IV	Endometrioid adenocarcinoma	3	280 Arg→Gly	AGA→GGA	Missense	DN
15	IV	Serous adenocarcinoma	3	248 Arg→Trp	CGG→TGG	Missense	DN
Mainz University							
1	I	Endometrioid adenocarcinoma	3	306 Arg→Stop	CGA→TGA	Nonsense	
2	I	Endometrioid adenocarcinoma	2	280 Arg→Ser	AGA→AGT	Missense	DN
3	I	Endometrioid adenocarcinoma	3	273 Arg→Cys	CGT→TGT	Missense	DN
4	I	Clear cell adenocarcinoma	3	245 Gly→Val	GGC→GTC	Missense	
5	II	Endometrioid adenocarcinoma	2	257 Leu→Pro	CTG→CCG	Missense	
6	III	Endometrioid adenocarcinoma	3	248 Arg→Gln	CGG→CAG	Missense	DN
7	III	Endometrioid adenocarcinoma	1	273 Arg→His	CGT→CAT	Missense	DN
8	IV	Endometrioid adenocarcinoma	3	175 Arg→His	CGC→CAC	Missense	DN
9	IV	Serous adenocarcinoma	3	273 Arg→His	CGT→CAT	Missense	DN

DN, dominant negative.

were pelleted, washed with LiOAc solution containing 0.1 M lithium acetate, 10 mM Tris-HCl, pH 8.0, and 1 mM EDTANa₂; the cells were then pelleted again and resuspended in 500 µl of LiOAc solution. For each transformation, 50 µl of yeast suspension were mixed with 1-5 µl of unpurified *p53* cDNA PCR product, 50-100 ng of linearized plasmid, 5 µl of sonicated single-stranded salmon sperm DNA (10 mg/ml) and 300 µl of LiOAc containing 40% polyethylene glycol 4000 (Kanto, Tokyo, Japan). The mixture was incubated at 30°C for 30 min and heat-shocked at 42°C for 15 min. The yeast was then plated on a synthetic dropout (SD) medium minus leucine plus adenine (5 µg/ml) and was incubated for 48 hr in a 30°C humidified chamber. More than 200 colonies were examined on each culture plate. In this assay system, 16% was the cutoff value for *p53* mutation.²⁰

Recovery of *p53* plasmids from yeast and DNA sequencing

The yeast was digested with Zymolase-100T (Seikagaku-Kogyo, Tokyo, Japan), and *p53* expression plasmids were extracted by the alkaline lysis method (QIAprep plasmid kit; Qiagen, Hilden, Germany) and transfected into XL-1 blue *E. coli* by electroporation. The plasmids were recovered, purified and sequenced with a Dye-Deoxy Terminator Kit (Perkin-Elmer, Urayasu, Japan) on an ABI 377 automated sequencer (Applied Biosystems, Urayasu, Japan) as specified by the manufacturer's protocol and using the following primers: P3seq, 5'-ATTTGATGCTGTCCCCGGACGATATTGAAC-3'; P11seq, 5'-TACTCCCCCTGCCCTCAACAAGATG-3'; P12seq, 5'-TTGCGTGTGGAGTATTTGGATGAC-3'; and P13seq, 5'-GCC-CATCCTCACCATCATCACACT-3'.

Transdominance assay

The dominant-negative potential of *p53* mutation was tested using a yeast-based transdominance assay as described previously.¹⁵ Briefly, the yeast functional assay was performed using both a plasmid with wild-type *p53* and a plasmid with mutant *p53* that had been sequence-verified. For each transformation, 50 µl of yeast suspension were mixed with 100 ng of pTSHP53, 100 ng of

mutant *p53*-containing pSS16, 50 µg of sonicated single-stranded salmon sperm DNA and 300 µl of LiOAc containing 40% polyethylene glycol 4000. The mixture was incubated at 30°C for 30 min and heat-shocked at 42°C for 15 min. Yeast were then plated on SD medium minus leucine and tryptophan, but which contained a limited amount of adenine (5 µg/ml). The samples were then incubated for 48 hr in a 30°C humidified atmosphere. Double-transformant clones (Leu⁺, Trp⁺) giving rise to white (Ade⁺) or pink/red (Ade⁻) colonies were interpreted as expressing recessive and dominant-negative mutations, respectively.

Statistical analysis

The statistical significance of differences between the categorical variables was examined by the chi-square test. Disease-specific survival curves were obtained by the Kaplan-Meier method and the differences between curves were examined by the log-rank test. Independence of prognostic significance was examined using a Cox regression analysis with forward stepwise selection of the variables. $p < 0.05$ was considered to be statistically significant. Statistical analyses were performed using the Statview 5.0 software package (SAS Institute, Cary, NC).

Results

p53 status

The wild-type *p53* gene was observed in 68 tumors. A *p53* mutation was found in 24 (26.1%) tumors. An example of a tumor in which the yeast functional assay gave a positive result (*i.e.*, the number of red colonies vs. white colonies = 46:54), and in which DNA sequencing analysis revealed a mutation in codon 273, is shown in Figure 1. The *p53* mutations and their respective transdominance property observed here are summarized in Table II. The mutations included 19 missense mutations, 2 nonsense mutations and 3 deletion mutations. Missense mutation accounted for 79% of all the mutations observed in this study. Codon 273 was most frequently mutated (4 273Arg→His and 1 273Arg→Cys),

TABLE III - RELATIONSHIP BETWEEN p53 MUTATION AND CLINICAL FEATURES OF ENDOMETRIAL CANCER PATIENTS

	All mutations		Dominant-negative mutation	
	Number/total (%)	p	Number/total (%)	p
Age				
< 60	9/39 (23.1)	0.64	5/39 (12.8)	0.58
≥ 60	15/53 (28.3)		9/53 (17.0)	
FIGO Stage				
I/II	14/65 (21.5)	0.19	6/65 (9.2)	0.01
III/IV	10/27 (37.0)		8/27 (29.6)	
Histologic subtype				
Endometrioid	18/79 (22.8)	0.09	9/79 (11.4)	0.01
Nonendometrioid	6/13 (46.2)		5/13 (38.5)	
Grade				
1, 2	14/67 (20.9)	0.06	7/67 (10.4)	0.04
3	10/25 (40.0)		7/25 (28.0)	
Institute				
Mainz	9/49 (18.4)	0.07	6/49 (12.2)	0.39
Sapporo	15/43 (34.9)		8/43 (18.6)	




FIGURE 2 - Kaplan-Meier analysis and log-rank test for survival of patients according to FIGO stage.




FIGURE 3 - Kaplan-Meier analysis and log-rank test for survival of patients according to histologic subtype of tumors.

followed by codon 280 (280Arg→Ile, 280Arg→Gly, 280Arg→Ser and then codon 248 (248Arg→Trp and 248Arg→Gln). Regarding the transdominance of p53 mutation, 10 mutant p53 proteins (10.9%) exhibited recessive activity and 14 mutants




FIGURE 4 - Kaplan-Meier analysis and log-rank test for survival of patients according to grade of tumors.




FIGURE 5 - Kaplan-Meier analysis and log-rank test for survival of patients with wild-type, recessive mutant and dominant-negative mutant p53.

(15.2%) showed dominant-negative activity. Not all of the missense mutations had a dominant-negative effect. Only 13 of 19 (68%) missense mutations of p53 exhibited dominant-negative activity.

p53 mutation was compared to the clinical features (Table III). Total p53 mutation tended to be related to the following features: nonendometrioid subtype ($p = 0.09$), grade 3 tumors ($p = 0.06$) and Japanese cohort ($p = 0.07$). Dominant-negative mutation was significantly related to advanced FIGO stage ($p = 0.01$), nonendometrioid subtype ($p = 0.01$) and grade 3 tumors ($p = 0.04$).

Survival analysis

The Kaplan-Meier analysis and log-rank test revealed that the survival of patients in this study was significantly related to conventional prognostic factors: FIGO stage ($p < 0.0001$; Fig. 2), histologic subtype ($p < 0.0001$; Fig. 3) and grade of tumor ($p = 0.037$; Fig. 4). As regards p53 mutation, dominant-negative p53 mutation was found to be related to poor patient survival ($p < 0.0001$; Fig. 5). Missense p53 mutation was also significantly related to patient survival ($p = 0.0001$). The age and institute were not related to survival ($p = 0.17$ and 0.52 , respectively). When we further examined the functional status of p53 mutation in relation to patients' survival, the estimated 5-year

TABLE IV - COX REGRESSION ANALYSES FOR PATIENTS WITH ENDOMETRIAL CARCINOMA

Variable	Univariate Cox analysis <i>p</i>	Multivariate Cox analysis			
		coefficient	Standard error	Hazard ratio	<i>p</i>
Dominant negative <i>p53</i> mutation	< 0.0001	1.37	0.58	3.9	0.019
FIGO stage	0.0002	1.81	0.62	6.1	0.0037
Histologic subtype	0.0004	1.69	0.58	5.4	0.014
Grade	0.048				NS
Age	0.19				NS
Institute	0.48				NS

survival rate for patients with wild-type *p53* ($n = 68$), recessive *p53* mutation ($n = 10$) and dominant-negative *p53* mutation ($n = 14$) was 84.9%, 85.7% and 35.1%, respectively. There was a statistically significant difference in survival between the patients with recessive *p53* mutation and those with dominant-negative *p53* mutation ($p = 0.01$), as well as between the patients with wild-type *p53* and those with dominant-negative *p53* mutation ($p < 0.0001$). No difference in survival was found between the patients with recessive *p53* mutation and those with wild-type *p53* (Fig. 5). Furthermore, the survival of patients with missense *p53* mutation tended to be related to the transdominance property of mutant *p53*. The 5-year survival rate was 75.0% for patients with recessive missense mutations ($n = 10$) and 34.6% for patients with dominant-negative missense mutations ($n = 14$; $p = 0.09$).

Using multivariate Cox regression analysis, we found that dominant-negative *p53* mutation ($p = 0.019$), FIGO stage ($p = 0.0037$) and histologic subtype ($p = 0.014$) were independent prognostic factor for the patients in this study (Table IV). When only advanced-stage tumors were taken into consideration, dominant-negative *p53* mutation ($p = 0.0023$) was the most significant predictor of patient survival.

Discussion

The functional activities of mutant *p53* proteins have been grouped into 5 categories: retained wild-type activity, loss of function, gain of function, dominant-negative effect and temperature sensitivity.⁴ The dominant-negative activity of *p53* mutation corresponds to the capacity of the mutant protein to complex with the product of the remaining wild-type allele to inactivate its function. Thus, dominant-negative *p53* mutation results in the total abrogation of *p53* protein function, even if there is still wild-type protein expressed in the cell. Although the importance of analysis of the functional types of *p53* mutation, *i.e.*, the dominant-negative effect and gain of function, in terms of researching carcinogenesis and searching for novel human cancer therapies has been repeatedly emphasized,^{10,12,21} it remains unclear whether or not the dominant-negative activity of mutant *p53* proteins has a detrimental effect on the survival of cancer patients. Recessive *p53* mutation accompanied by loss of the second allele may be equal to dominant-negative *p53* mutation in terms of loss of function of the gene. Although we did not investigate LOH in this study, the present results did suggest that dominant-negative *p53* mutation is closely related to poor survival of patients with endometrial cancer, even after adjusting for established prognostic factors, that is, tumor stage, grade and histologic type. Future studies including the LOH status of the second allele will be of interest in this context. It is important to investigate whether or not the dominant-negative *p53* mutation exerts an influence on patient survival, not only through a loss of function, but also by other mechanisms attributable to mutant *p53* protein such as certain gain-of-function activities.

Also of interest in this context is that dominant-negative *p53* mutation was found to be closely related to the survival of patients with advanced-stage endometrial cancer. Because the prognosis of patients with early-stage endometrial cancer is generally excellent, gynecologic oncologists need to focus increasingly on the survival

of patients with advanced endometrial cancer. The histopathologic prognostic factors for endometrial cancer include depth of myometrial invasion, cervical involvement, serosal invasion, adnexal metastasis, positive peritoneal cytology, vaginal metastasis, lymph node metastasis, peritoneal metastasis and bladder/rectal involvement, which are incorporated in the FIGO surgical staging system.²² The grade of tumor, histologic subtype and lymph-vascular space invasion, which represent the aggressiveness of a tumor, are also important histopathologic prognostic factors that should be taken into consideration in planning treatment for patients with endometrial cancer.²³ The histologic subtypes of serous adenocarcinoma and clear cell adenocarcinoma exhibit more aggressive biologic behavior than common endometrioid adenocarcinoma and has been shown to lead to disproportionate mortality.²⁴⁻²⁶ Serous adenocarcinoma is frequently associated with *p53* overexpression or *p53* mutation. Our current study has shown that dominant-negative *p53* mutation is an important prognostic factor, in addition to the established predictors of survival, namely, FIGO stage and histologic subtype, in cases of endometrial cancer. This suggests that dominant-negative *p53* mutation may be a reasonable target for a novel therapy for cases of endometrial cancer with a poor prognosis.

In the present study, only 68% of the missense mutations in the cases of endometrial cancer studied here exhibited dominant-negative activity. This finding suggests that determining the dominant-negative activity of a *p53* mutation is more important than merely determining the presence or absence of a *p53* mutation as part of a tailored treatment or rational targeted treatment for endometrial cancer. Because of the high frequency of *p53* mutations in human cancers, and due to the pivotal role of *p53* in regulating growth, apoptosis and DNA repair, the introduction of the wild-type *p53* gene has been regarded as a reasonable strategy for a gene therapy designed to restore the lost activity of *p53*.²⁷ However, this approach has achieved substantial effectiveness to date.²⁸ A possible reason for the unsatisfactory results may be the accumulation of dominant-negative mutant *p53*, which results in a high amount of mutant protein, which would override the effects of the wild-type protein introduced by gene therapy. This explanation may in part account for the recently reported failure of gene therapy in the study of ovarian cancer.²⁹ In addition to overriding wild-type *p53*, some dominant-negative mutants are known to cause a gain of function related to tumor progression.^{10,12,30} Such mutants include 175 Arg→His, 273 Arg→His and 248 Arg→Trp, which accounted for 6 of the 14 dominant-negative mutants identified in the present study. The gain-of-function property of the *p53* mutants is considered to lend further malignant phenotypes to the tumor cells, such as enhancement of tumorigenicity, metastatic potential and therapy resistance. These properties may account for the extremely poor survival of endometrial cancer patients with dominant-negative *p53* mutations. The 6 cases with gain-of-function mutations in this study included 1 case of stage I serous adenocarcinoma (case 8 in the Japanese cohort), 1 case of stage III endometrioid adenocarcinoma (cases 7 in the German cohort), 1 case of stage III serous adenocarcinoma (case 13 in the Japanese cohort), 1 case of stage IV endometrioid adenocarcinoma (case 9 in the German cohort) and 2 cases of stage IV serous adenocarcinoma (case 15 in the Japanese cohort and case 9 in the German cohort). The gain-of-function mutation appears to be related to

advanced-stage tumors, although the number of cases studied was not large enough to draw any conclusions.

In summary, this study indicates that dominant-negative mutation of *p53* gene is often found in the advanced stages and aggressive histologic subtypes of endometrial cancer. Moreover, such

mutation is a strong predictor of the survival of patients with advanced endometrial cancer. Further investigation will be needed in order to clarify whether or not identification of the dominant-negative property of *p53* mutation may be useful for tailoring the treatment of endometrial cancer.

References

- Hainaut P, Hollstein M. *p53* and human cancer: the first ten thousand mutations. *Adv Cancer Res* 2000;77:81-137.
- Beroud C, Soussi T. *p53* gene mutation: software and database. *Nucleic Acids Res* 1998;26:200-4.
- Hollstein M, Moeckel G, Hergenbahn M, Spiegelhalter B, Keil M, Werle-Schneider G, Bartsch H, Brickmann J. On the origins of tumor mutations in cancer genes: insights from the *p53* gene. *Mutat Res* 1998;405:145-54.
- Olivier M, Eeles R, Hollstein M, Khan MA, Harris CC, Hainaut P. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum Mutat* 2002;19:607-14.
- Kohler MF, Carney P, Dodge R, Soper JT, Clarke-Pearson DL, Marks JR, Berchuck A. *p53* overexpression in advanced-stage endometrial adenocarcinoma. *Am J Obstet Gynecol* 1996;175:1246-52.
- Soong R, Knowles S, Williams KE, Hammond IG, Wysocki SJ, Iacopetta BJ. Overexpression of *p53* protein is an independent prognostic indicator in human endometrial carcinoma. *Br J Cancer* 1996;74:562-7.
- Ohkouchi T, Sakuragi N, Watari H, Nomura E, Todo Y, Yamada H, Fujimoto S. Prognostic significance of *Bcl-2*, *p53* overexpression, and lymph node metastasis in surgically staged endometrial carcinoma. *Am J Obstet Gynecol* 2002;187:353-9.
- Nylander K, Dabelsteen E, Hall PA. The *p53* molecule and its prognostic role in squamous cell carcinomas of the head and neck. *J Oral Pathol Med* 2000;29:413-25.
- Bergamaschi D, Gasco M, Hiller L, Sullivan A, Syed N, Trigiani G, Yulug I, Merlano M, Numico G, Comino A, Attard M, Reelfs O, Gusterson B, Bell AK, Heath V, Tavassoli M, Farrell PJ, Smith P, Lu X, Crook T. *p53* polymorphism influences response in cancer chemotherapy via modulation of *p73*-dependent apoptosis. *Cancer Cell* 2003;3:387-402.
- van Oijen MG, Slootweg PJ. Gain-of-function mutations in the tumor suppressor gene *p53*. *Clin Cancer Res* 2000;6:2138-45.
- Kirsch DG, Kastan MB. Tumor-suppressor *p53*: implications for tumor development and prognosis. *J Clin Oncol* 1998;16:3158-68.
- Roemer K. Mutant *p53*: gain-of-function oncoproteins and wild-type *p53* inactivators. *Biol Chem* 1999;380:879-87.
- Flaman JM, Frebourg T, Moreau V, Charbonnier F, Martin C, Chappuis P, Sappino AP, Limacher IM, Bron L, Benhattar J, Tada M, Van Mei EG, Estreicher A, Iggo RD. A simple *p53* functional assay for screening cell lines, blood, and tumors. *Proc Natl Acad Sci USA* 1995;92:3963-7.
- Kashiwazaki H, Tonoki H, Tada M, Chiba I, Shindoh M, Totsuka Y, Iggo R, Moriuchi T. High frequency of *p53* mutations in human oral epithelial dysplasia and primary squamous cell carcinoma detected by yeast functional assay. *Oncogene* 1997;15:2667-74.
- Marutani M, Tonoki H, Tada M, Takahashi M, Kashiwazaki H, Hida Y, Hamada J, Asaka M, Moriuchi T. Dominant-negative mutations of the tumor suppressor *p53* relating to early onset of glioblastoma multiforme. *Cancer Res* 1999;59:4765-9.
- Sakuragi N, Hirai A, Tada M, Yamada H, Yamamoto R, Fujimoto S, Moriuchi T. Dominant-negative mutation of *p53* tumor suppressor gene in endometrial carcinoma. *Gynecol Oncol* 2001;83:485-90.
- Ishioka C, Frebourg T, Yan YX, Vidal M, Friend SH, Schmidt S, Iggo R. Screening patients for heterozygous *p53* mutations using a functional assay in yeast. *Nat Genet* 1993;5:124-9.
- Tada M, Iggo RD, Waridel F, Nozaki M, Matsumoto R, Sawamura Y, Shinohe Y, Ikeda J, Abe H. Reappraisal of *p53* mutations in human malignant astrocytic neoplasms by *p53* functional assay: comparison with conventional structural analyses. *Mol Carcinogenesis* 1997;18:171-6.
- Stotz A, Linder P. The *ADE2* gene from *Saccharomyces cerevisiae*: sequence and new vectors. *Gene* 1990;95:91-8.
- Takahashi M, Tonoki H, Tada M, Kashiwazaki H, Furuuchi K, Hamada J, Fujioka Y, Sato Y, Takahashi H, Todo S, Sakuragi N, Moriuchi T. Distinct prognostic values of *p53* mutations and loss of estrogen receptor and their cumulative effect in primary breast cancers. *Int J Cancer* 2000;89:92-9.
- Brachmann RK, Vidal M, Boeke JD. Dominant-negative *p53* mutations selected in yeast hit cancer hot spots. *Proc Natl Acad Sci USA* 1996;93:4091-5.
- FIGO News. Corpus cancer staging. *Int J Gynecol Obstet* 1989;28:189-93.
- Nishiya M, Sakuragi N, Hareyama H, Ebina Y, Oikawa M, Yamamoto R, Fujino T, Fujimoto S. Cox multivariate regression models for estimating prognosis of patients with endometrioid adenocarcinoma of uterine corpus who underwent thorough surgical staging. *Int J Cancer* 1998;79:521-5.
- Wilson TO, Podratz KC, Gaffey TA, Malkasian GD, O'Brien PC, Naessens JM. Evaluation of unfavorable histologic subtypes in endometrial adenocarcinoma. *Am J Obstet Gynecol* 1990;162:418-26.
- Greven KM, Lanciano RM, Corn B, Case D, Randall ME. Pathologic stage III endometrial carcinoma: prognostic factors and patterns of recurrence. *Cancer* 1993;71:3697-702.
- Sakuragi N, Hareyama H, Todo Y, Yamada H, Yamamoto R, Fujino T, Sagawa T, Fujimoto S. Prognostic significance of serous and clear cell adenocarcinoma in surgically staged endometrial carcinoma. *Acta Obstet Gynecol Scand* 2000;79:311-6.
- Lebedeva IV, Su ZZ, Sarkar D, Fisher PB. Restoring apoptosis as a strategy for cancer gene therapy: focus on *p53* and *mda-7*. *Semin Cancer Biol* 2003;13:169-78.
- Westphal EM, Melchner HVH. Gene therapy approaches for the selective killing of cancer cells. *Curr Pharm Des* 2002;8:1683-94.
- Zeimet AG, Marth C. Why did *p53* gene therapy fail in ovarian cancer? *Lancet Oncol* 2003;4:415-22.
- Dittmer D, Pati S, Zambetti G, Chu S, Teresky AK, Moore M, Finlay C, Levine AJ. Gain of function mutations in *p53*. *Nat Genet* 1993;4:42-6.

HER-2 codon 655 polymorphism in cervical carcinogenesis

M. UEDA*, Y.-C. HUNG†, Y. TERAJ*, J. SAITO‡, O. NUNOBIKI‡, S. NODA‡ & M. UEKI*

*Department of Obstetrics and Gynecology, Osaka Medical College, Osaka, Japan; †Department of Obstetrics and Gynecology, China Medical College, Taichung, Taiwan; and ‡Osaka Cancer Prevention Center, Osaka, Japan

Abstract. Ueda M, Hung Y-C, Terai Y, Saito J, Nunobiki O, Noda S, Ueki M. HER-2 codon 655 polymorphism in cervical carcinogenesis. *Int J Gynecol Cancer* 2006;16:325–328.

HER-2 codon 655 polymorphism together with human papillomavirus (HPV) types were examined in a total of 279 cervical smear samples. Forty-nine patients with high-grade squamous intraepithelial lesion had higher frequency of high-risk HPV than 167 patients with low-grade squamous intraepithelial lesion and 63 controls. There was no statistical difference in the frequencies of HER-2 Ile/Ile, Ile/Val, and Val/Val genotypes between squamous intraepithelial lesions (SILs) and controls. When the Ile/Ile genotype was compared to the Ile/Val + Val/Val genotypes, there was also no statistical difference in the genotype prevalence between SILs and controls either in 91 or 188 patients with or without high-risk HPV, respectively. These results suggest that the HER-2 polymorphism at codon 655 in cervical cell samples is unlikely to be associated with HPV status and the onset of cervical cancer in a Japanese population.

KEYWORDS: cervical carcinogenesis, HER-2, polymorphism, SIL.

Cervical cancer is the second most common cancer in women worldwide and is both a preventable and a curable disease especially if identified at an early stage. It is widely accepted that specific human papillomavirus (HPV) types are the central etiologic agent of cervical carcinogenesis. Recently, several candidate markers for cervical cancer risk, such as glutathione-S-transferase and p53, have been described^(1–5). Despite extensive studies on germ line polymorphisms of these cancer susceptibility genes in the patients with pre-malignant and malignant cervical lesions, no correlation has been reported between their genotype prevalence and increased risk of cervical cancer^(1–3,5–7).

HER-2 (also known as c-erbB-2 or neu), the second member of the epidermal growth factor receptor family, encodes a transmembrane glycoprotein (p185) with tyrosine kinase activity^(8–10). Altered expression of the HER-2 gene has been implicated in the carcinogenesis and progression of breast, ovarian, gastric, prostate, and bladder cancer^(11–15). Previous studies have also

demonstrated that repression of the HER-2 function suppresses the malignant phenotype of HER-2 over-expressing cancer cells^(16,17). However, there have been only a few reports on the possible correlation between HER-2 expression and the development of cervical cancer^(18–21).

Sequence analysis of the human HER-2 gene identified a single nucleotide polymorphism at codon 655, resulting in a G to A transition (Ile 655 Val) in the transmembrane domain-coding region of this gene⁽²²⁾. Recently, Xie *et al.*⁽²³⁾ reported that the variant Val allele was associated with an elevated risk of breast cancer in Chinese women, whereas another study in a Caucasian population failed to find such a significant association⁽²⁴⁾. Although the biologic significance of the amino acid substitution at HER-2 codon 655 is still unclear, it would be of interest to investigate the relationship between the HER-2 polymorphism and pathogenesis in various kinds of human malignant tumors.

In this study, we investigated the HER-2 codon 655 polymorphism together with HPV status in exfoliated cervical cell samples from the patients with squamous intraepithelial lesion (SIL) of the cervix and evaluated the genotype prevalence of this gene in cervical carcinogenesis.

Address correspondence and reprint requests to: Masatsugu Ueda, MD, Department of Obstetrics and Gynecology, Osaka Medical College, 2-7 Daigakumachi, Takatsuki, Osaka 569-8686, Japan. Email: gyn017@poh.osaka-med.ac.jp

Materials and methods

Cell sample

We conducted HER-2 genotype analysis together with HPV typing in a total of 279 cervical smear samples from the patients who received cervical cancer screening. They consist of 63 normal, 167 low-grade SIL (LSIL), and 49 high-grade SIL (HSIL). All the 279 patients were Japanese women. Final histologic diagnosis was confirmed by colposcopy-directed biopsy.

DNA preparation

The exfoliated cervical cells were disrupted with lysis buffer (20 mM NaCl, 10 mM Tris-HCl [pH 8.0], 10 mM ethylenediaminetetraacetic acid [pH 8.0], 0.5% sodium dodecyl sulfate (SDS), 50 µg/mL proteinase K), and genomic DNA was extracted with phenol-chloroform and precipitated with ethanol using standard techniques. Purified DNA samples from the cells were stored at -20°C until use.

Genotyping of HER-2 codon 655

Polymerase chain reaction–restriction fragment length polymorphism analysis of codon 655 of the HER-2 gene, modified from a technique described by Wang *et al.*⁽²⁵⁾, was conducted to identify HER-2 genotypes using specific primers: sense 5'-AGAGAGCCAGCC-CTCTGACGTCCAT-3' and antisense 5'-TCCGTTT-CCTGCAGCAGTCTCCGCA-3'. One microliter of the DNA template from each cell sample was amplified by polymerase chain reaction (PCR) in a final volume of 50 µL reaction mixture containing 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 2 mM MgCl₂, 0.01% (w/v) gelatin, 200 µM deoxynucleoside triphosphate, 0.5 µM of each primer, and 1.25 units *Taq* polymerase (Applied Biosystems, Branchburg, NJ) as previously described⁽²⁶⁾. After an initial denaturation at 96°C for 3 min, 40 cycles of denaturation (94°C for 1 min), annealing (55°C for 1 min), and extension (72°C for 2 min) were carried out on a Perkin-Elmer GeneAmp PCR System 9700. The final extension was performed at 72°C for 10 min. After confirmation of an amplified fragment of the expected size (148 bp) by 1.5% agarose gel electrophoresis with ethidium bromide staining, 13 µL of each PCR product was digested with 25 units of restriction enzyme BsmAI (New England Biolabs Inc., Beverly, MA) at 55°C for 3 h, and DNA fragments were visualized on a 3.0% agarose gel. Restriction fragments were 130 and 26 bp for the Ile allele and 94, 36, and 26 bp for the Val allele. The heterozygote contains the following four bands: 130, 94, 36, and 26 bp.

HPV typing

The presence of various HPV types was examined using L1-PCR according to the method reported by Nagano *et al.*⁽²⁷⁾ Briefly, 100 ng of cellular DNA was subjected to PCR in the presence of published consensus primers (L1C1 and L1C2)⁽²⁸⁾. Amplified HPV fragments were typed on the basis of the restriction fragment length polymorphism among HPVs⁽²⁸⁾. L1-PCR can detect 22 registered low-risk (6, 11, 34, 40, 42, 43, 44) and high-risk (16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, 68, 69) HPV types.

Statistical analysis

The HPV status and polymorphic features of HER-2 gene in 279 cases were compared between normal, LSIL, and HSIL and checked by the Mann–Whitney and Chi-square tests. A level of $P < 0.05$ was accepted as statistically significant.

Results

The polymorphic site in codon 655 of the HER-2 gene was achieved by polymerase chain reaction–restriction fragment length polymorphism. As shown in Figure 1, fragments of 130 and 26 bp resulted from BsmAI digestion of the Ile allele. The fragments of 94, 36, and 26 bp indicated the digested PCR products from the Val allele. The heterozygote contains the following four bands: 130, 94, 36, and 26 bp. Although the 36 and 26 bp fragments were not clearly visualized, each genotype was judged by the presence or absence of the 130 and 94 bp fragments.

Table 1 shows HPV status and polymorphic frequency of HER-2 codon 655 in 279 samples examined.




Figure 1. Polymerase chain reaction–restriction fragment length polymorphism analysis on HER-2 codon 655 in DNA samples purified from exfoliated cervical cells. Restriction fragments treated by BsmAI are 130 and 26 bp for the Ile homozygote (lane 1) and 94, 36, and 26 bp for the Val homozygote (lane 3). The heterozygote is in lane 2. An undigested PCR fragment (148 bp) is shown in lane 4. Although the 36 and 26 bp fragments are not clearly visualized, each genotype is judged by the presence or absence of the 130 and 94 bp fragments. Lane 5 is negative control.

Table 1. HPV status and HER-2 codon 655 polymorphism

Lesions	n	Number with high-risk HPV	Amino acid at HER2 codon 655		
			Ile/Ile	Ile/Val	Val/Val
Normal	63	10 (15.9%)***	46 (73.0%)	16 (25.4%)	1 (1.6%)
LSIL	167	46 (27.5%)***	123 (73.7%)	38 (22.8%)	6 (3.6%)
HSIL	49	35 (71.4%)****	35 (71.4%)	14 (28.6%)	0 (0%)
All SIL	216	81 (37.5%)**	158 (73.1%)	52 (24.1%)	6 (2.8%)

* $P < 0.0001 \chi^2$ vs normal.** $P = 0.0013 \chi^2$ vs normal.*** $P < 0.0001 \chi^2$ vs LSIL.

The patients with HSIL had significantly higher frequency of high-risk HPV than those with LSIL and controls. There was also statistical difference in the frequency of high-risk HPV between all SIL and controls. However, the differences in the polymorphic frequency of HER-2 Ile/Ile, Ile/Val, and Val/Val genotypes between SILs and controls were statistically not significant.

When the Ile/Ile genotype was compared to the Ile/Val + Val/Val genotypes, there was no statistical difference in the genotype prevalence between SILs and controls in 188 patients without high-risk HPV, as shown in Table 2. The relative risks of the combined Ile/Val + Val/Val genotypes for SILs in 91 patients with high-risk HPV showed slightly increased odds ratios (2.345 for LSIL, 1.833 for HSIL, and 2.113 for all SIL); however, there was again no statistical difference between SILs and controls.

Discussion

The genes that encode growth factor receptors can be activated through mutations in their coding sequences. A point mutation in the transmembrane region of the neu gene, a rat homologue of human HER-2 gene, increased the transforming capacity of this gene⁽²⁹⁾.

Transgenic mice that carry a mutationally activated HER-2 gene or an overexpressed normal HER-2 gene frequently develop mammary adenocarcinoma⁽³⁰⁾. Although mutations in the human HER-2 gene have not been identified, sequence analysis revealed a single nucleotide polymorphism at codon 655⁽²²⁾. Since the HER-2 polymorphism at codon 655 is located in the transmembrane coding region, it may be also associated with the structural and functional alteration of the HER-2 protein. Several previous studies have reported the germ line polymorphism and ethnic variation of the HER-2 gene in breast, bladder, and gastric cancer^(23-25,31,32). However, there has been no report on the correlation between the HER-2 polymorphism and development of malignant tumors of gynecological origin.

In our investigation using exfoliated cervical cell samples from a Japanese population, the differences in the polymorphic frequency of HER-2 Ile/Ile, Ile/Val, and Val/Val genotypes between 216 SILs and 63 controls were statistically not significant. There was also no statistical difference in the genotype prevalence between SILs and controls in 188 patients without high-risk HPV. Although Val allele slightly increased the relative risk of LSIL or HSIL compared to controls in 91 patients with high-risk HPVs, the differences were

Table 2. Relationships between HER-2 codon 655 polymorphism and SIL grade

Study group	n	Amino acid at HER2 codon 655		OR	95% CI	P value
		Ile/Ile	Ile/Val + Val/Val			
High-risk HPV-						
Normal	53	38 (71.7%)	15 (28.3%)	1		
LSIL	121	94 (77.7%)	27 (22.3%)	0.728	0.350-1.514	0.396
HSIL	14	11 (78.6%)	3 (21.4%)	0.691	0.170-2.814	0.606
All SIL	135	105 (77.8%)	30 (22.2%)	0.724	0.352-1.487	0.379
High-risk HPV+						
Normal	10	8 (80.0%)	2 (20.0%)	1		
LSIL	46	29 (63.0%)	17 (37.0%)	2.345	0.461-11.930	0.305
HSIL	35	24 (68.6%)	11 (31.4%)	1.833	0.339-9.911	0.482
All SIL	81	53 (65.4%)	28 (34.6%)	2.113	0.433-10.315	0.355

again statistically not significant. To the best of our knowledge, this is the first study to examine the role of HER-2 codon 655 polymorphism in cytologic materials from women with premalignant cervical disease. Our present results suggest that Val655Ile of the HER-2 gene in cervical cell samples is unlikely to be associated with HPV status and the onset of cervical cancer in a Japanese population.

Acknowledgments

We are grateful to Dr Ken Ueki, Department of Obstetrics and Gynecology, Osaka Medical College, for collecting clinical materials. We also thank Kumiko Sato for her technical assistance. This work was supported in part by High-Tech Research Program of Osaka Medical College.

References

- Warwick A, Sarhanis P, Redman C *et al.* Theta class glutathione S-transferase GSTT1 genotypes and susceptibility to cervical neoplasia: interactions with GSTM1, CYP2D6 and smoking. *Carcinogenesis* 1994;15:2841-5.
- Chen C, Madeleine MM, Weiss NS, Daling JR. Glutathione S-transferase M1 genotypes and the risk of squamous carcinoma of the cervix: a population-based case-control study. *Am J Epidemiol* 1999;150:568-72.
- Goodman MT, McDuffie K, Hernandez B *et al.* CYP1A1, GSTM1, and GSTT1 polymorphisms and the risk of cervical squamous intraepithelial lesions in a multiethnic population. *Gynecol Oncol* 2001;81:263-9.
- Storey A, Thomas M, Kalita A *et al.* Role of a p53 polymorphism in the development of human papillomavirus-associated cancer. *Nature* 1998;393:229-34.
- Rosenthal AN, Ryan A, Al-Jehani RM, Storey A, Harwood CA, Jacobs IJ. p53 codon 72 polymorphism and the risk of cervical cancer in UK. *Lancet* 1998;352:871-2.
- Lanham S, Campbell I, Watt P, Gornall R. p53 polymorphism and risk of cervical cancer. *Lancet* 1998;352:1631.
- Hayes VM, Hofstra RMW, Buys CHCM, Hollema H, van der Zee AGJ. Homozygous arginine-72 in wild type p53 and risk of cervical cancer. *Lancet* 1998;352:1756.
- Semba K, Kamata N, Toyoshima K, Yamamoto T. A v-erbB-related protooncogene, c-erbB-2, is distinct from the c-erbB-1/epidermal growth factor-receptor gene and is amplified in a human salivary gland adenocarcinoma. *Proc Natl Acad Sci U S A* 1985;82:6497-501.
- Yamamoto T, Ikawa S, Akiyama T *et al.* Similarity of protein encoded by the human c-erbB-2 gene to epidermal growth factor receptor. *Nature* 1986;319:230-4.
- Di Fiore PP, Pierce JH, Kraus MH, Segatto O, King CR, Aaronson SA. erbB-2 is a potent oncogene when overexpressed in NIH/3T3 cells. *Science* 1987;237:178-82.
- Dowsett M, Cooke T, Ellis I *et al.* Assessment of HER2 status in breast cancer: why, when and how? *Eur J Cancer* 2000;36:170-6.
- Hengstler JG, Lange J, Kett A *et al.* Contribution of c-erbB-2 and topoisomerase IIalpha to chemoresistance in ovarian cancer. *Cancer Res* 1999;59:3206-14.
- Nakajima M, Sawada H, Yamada Y *et al.* The prognostic significance of amplification and overexpression of c-met and c-erbB-2 in human gastric carcinomas. *Cancer* 1999;85:1894-902.
- Morote J, de Torres I, Caceres C, Vallejo C, Schwartz S Jr, Reventos J. Prognostic value of immunohistochemical expression of the c-erbB-2 oncoprotein in metastatic prostate cancer. *Int J Cancer* 1999;84:421-5.
- Vollmer RT, Humphrey PA, Swanson PE, Wick MR, Hudson ML. Invasion of the bladder by transitional cell carcinoma: its relation to histologic grade and expression of p53, MIB-1, c-erbB-2, epidermal growth factor receptor, and bcl-2. *Cancer* 1998;82:715-23.
- Hung MC, Matin A, Zhang Y *et al.* HER-2/neu-targeting gene therapy—a review. *Gene* 1995;159:65-71.
- Kao MC, Liu GY, Chuang TC, Lin YS, Wu JA, Law SL. The N-terminal 178-amino-acid domain only of the SV40 large T antigen acts as a transforming suppressor of the HER-2/neu oncogene. *Oncogene* 1998;16:547-54.
- Bar JK, Harlozinska A, Sedlaczek P, Kasiak J, Markowska J. Relations between the expression of p53, c-erbB-2, Ki-67 and HPV infection in cervical carcinomas and cervical dysplasias. *Anticancer Res* 2001;21:1001-6.
- Kohlberger P, Edwards L, Hacker NF. Microinvasive squamous cell carcinoma of the cervix: immunohistochemically detected prognostic factors in a case with poor clinical outcome. *Gynecol Oncol* 2003;90:443-5.
- Bellone S, Palmieri M, Gokden M *et al.* Selection of HER-2/neu-positive tumor cells in early stage cervical cancer: implications for Herceptin-mediated therapy. *Gynecol Oncol* 2003;91:231-40.
- Niibe Y, Nakano T, Ohno T, Suzuki Y, Oka K, Tsujii H. Prognostic significance of c-erbB-2/HER2 expression in advanced uterine cervical carcinoma with para-aortic lymph node metastasis treated with radiation therapy. *Int J Gynecol Cancer* 2003;13:849-55.
- Papewalis J, Nikitin AY, Rajewsky MF. G to A polymorphism at amino acid codon 655 of the human erbB-2/HER2 gene. *Nucleic Acids Res* 1991;19:5452.
- Xie D, Shu XO, Deng Z *et al.* Population-based, case-control study of HER2 genetic polymorphism and breast cancer risk. *J Natl Cancer Inst* 2000;92:412-7.
- Baxter SW, Campbell IG. Re: population-based, case-control study of HER2 genetic polymorphism and breast cancer risk. *J Natl Cancer Inst* 2001;93:557-9.
- Wang L, Habuchi T, Takahashi T *et al.* No association between HER-2 gene polymorphism at codon 655 and a risk of bladder cancer. *Int J Cancer* 2002;97:787-90.
- Ueda M, Gemmill RM, West J *et al.* Mutations of the β - and γ -catenin genes are uncommon in human lung, breast, kidney, cervical and ovarian carcinomas. *Br J Cancer* 2001;85:64-8.
- Nagano H, Yoshikawa H, Kawana T *et al.* Association of multiple human papillomavirus types with vulvar neoplasias. *J Obstet Gynaecol Res* 1996;22:1-8.
- Yoshikawa H, Kawana T, Kitagawa K, Mizuno M, Yoshikura H, Iwamoto A. Detection and typing of multiple genital human papillomaviruses by DNA amplification with consensus primers. *Jpn J Cancer Res* 1991;82:524-31.
- Bargmann CI, Hung MC, Weinberg RA. Multiple independent activations of the neu oncogene by a point mutation altering the transmembrane domain of p185. *Cell* 1986;45:649-57.
- Siegel PM, Dankort DL, Hardy WR, Muller WJ. Novel activating mutations in the neu proto-oncogene involved in induction of mammary tumors. *Mol Cell Biol* 1994;14:7068-77.
- Ameyaw MM, Tayeb M, Thornton N *et al.* Ethnic variation in the HER-2 codon 655 genetic polymorphism previously associated with breast cancer. *J Hum Genet* 2002;47:172-5.
- Kuraoka K, Matsumura S, Hamai Y *et al.* A single nucleotide polymorphism in the transmembrane domain coding region of HER-2 is associated with development and malignant phenotype of gastric cancer. *Int J Cancer* 2003;107:593-6.

Accepted for publication February 9, 2005

Fas gene promoter –670 polymorphism in gynecological cancer

M. UEDA, Y. TERAJ, K. KANDA, M. KANEMURA, M. TAKEHARA, H. YAMAGUCHI, K. NISHIYAMA, M. YASUDA & M. UEKI

Department of Obstetrics and Gynecology, Osaka Medical College, Takatsuki Osaka, Japan

Abstract. Ueda M, Terai Y, Kanda K, Kanemura M, Takehara M, Yamaguchi H, Nishiyama K, Yasuda M, Ueki M. Fas gene promoter –670 polymorphism in gynecological cancer. *Int J Gynecol Cancer* 2006;16(Suppl. 1): 179–182.

Single-nucleotide polymorphism at –670 of Fas gene promoter (A/G) was examined in a total of 354 blood samples from normal healthy women and gynecological cancer patients. They consisted of 95 normal, 83 cervical, 108 endometrial, and 68 ovarian cancer cases. Eighty-three patients with cervical cancer had statistically higher frequency of GG genotype and G allele than 95 controls ($P = 0.0353$ and 0.0278 , respectively). There was no significant difference in the genotype or allele prevalence between control subjects and endometrial or ovarian cancer patients. The Fas –670 GG genotype was associated with an increased risk for the development of cervical cancer (OR = 2.56, 95% CI = 1.08–6.10) compared with the AA genotype. The G allele also increased the risk of cervical cancer (OR = 1.60, 95% CI = 1.05–2.43) compared with the A allele. Germ-line polymorphism of Fas gene promoter –670 may be associated with the risk of cervical cancer in a Japanese population.

KEYWORDS: gynecological cancer, Fas, polymorphism.

Apoptosis is a physiologic process for the elimination of specific types of cells, occurring extensively in embryonic development, metamorphosis, and differentiation. This process also plays an important role in eliminating unwanted or potentially dangerous cells throughout life. Abnormal regulation of apoptosis is likely to contribute to the pathogenesis of autoimmune diseases and malignant tumors. The acquired ability to resist apoptotic stimuli is shared by many types of malignant diseases, and genetic alteration in the components of apoptotic pathways is a pivotal mechanism in the development of cancer^(1,2).

Fas, also known as CD95 or APO-1, is a transmembrane receptor that plays a central role in apoptotic signaling in many cell types⁽³⁾. This receptor interacts with its natural ligand FASL, a member of the tumor necrosis factor superfamily, to initiate the death signal cascade, which results in apoptotic cell death^(3,4). Down-regulation of Fas with resultant resistance to death signals has been reported in many cancers. The transcriptional expression of Fas gene is regulated by a

number of genetic elements located in the 5'-upstream region of the gene. Single-nucleotide polymorphism (SNP) at –670 of Fas gene promoter (A/G) has recently been identified^(5,6). This SNP has been found with potentially different transcriptional efficiency⁽⁵⁻⁷⁾. Recent studies have demonstrated that the A/G SNP at Fas promoter –670 is closely associated with the pathogenesis of autoimmune diseases, such as rheumatoid arthritis and systemic lupus erythematosus^(7,8). However, the correlation between this SNP and cancer susceptibility including the risk of gynecological malignancies has not been extensively studied, and previous experimental results are controversial.

Recently, Lai *et al.*⁽⁹⁾ reported that SNP at position –670 in the Fas gene promoter is associated with the development of cervical cancer in a Chinese population. Engelmark *et al.*⁽¹⁰⁾ also conducted Fas promoter –670 polymorphism analysis and suggested that this SNP does not have a major impact on the susceptibility to cervical cancer in Swedish patients. In this study, we examined germ-line polymorphism of Fas gene promoter –670 in a total of 354 blood samples from normal healthy women and gynecological cancer patients to reevaluate the possible association between this SNP and the risk of cervical cancer in a Japanese population.

Address correspondence and reprint requests to: Masatsugu Ueda, MD, Department of Obstetrics and Gynecology, Osaka Medical College, 2-7 Daigakumachi, Takatsuki, Osaka 569-8686, Japan. Email: gyn017@poh.osaka-med.ac.jp

© 2006, Copyright the Authors
Journal compilation © 2006, IGCS