

radiation oncologists. However, it could not be determined in our study, because we did not encounter any severe toxicity.

Verification is a very important process, especially in hypofractionated stereotactic radiotherapy. Negoro *et al.* previously reported the details of our verification method and the results of setup error (10). We used A-P and lateral verification films obtained by a X-ray simulator after CT scan, to compare linacography (A-P and lateral port films) immediately before irradiation. X-ray simulation films have a higher resolution than digital reconstructed radiography, especially in the C-C direction. X-ray simulation films can be easily taken using our integrated system, in which the CT simulator and the X-ray simulator are employed on the same couch. Therefore, we used X-ray simulation films to verify patient setup.

Dose correction for lung inhomogeneity is still a controversial issue. The application of the Monte Carlo calculation method to routine clinics in the future is one of the solutions. In the present situation, we consider that dose correction using a method such as the generalized Batho method should be performed to deliver the true prescribed dose.

Target delineation and definition are other important issues in SRT for lung tumors. Interobserver variation in target delineation is not negligible in some cases. There has been no universal target definition for small solitary lung tumor in SRT. Although the only concept is proposed by International Commission on Radiation Units and Measurements Report 62, details of target definition depend on the treatment methods, such as the way to scan the CT of the tumor and the verification method. Further discussions on these issues are necessary.

In conclusion, the use of multiple noncoplanar static ports achieved homogeneous target dose distribution and avoided high dose to normal tissues, despite the limitation of the beam arrangement from the use of the body frame and couch structure. Tolerance doses to the normal tissues are yet unknown when using single high-dose irradiation. Therefore, we should continue to make treatment plans carefully. In addition, further follow-ups of clinical cases are required to know the tolerance dose to the normal tissues in stereotactic radiotherapy.

## REFERENCES

1. Uematsu M, Shioda A, Tahara K, *et al.* Focal, high dose, and fractionated modified stereotactic radiation therapy for lung carcinoma patients: a preliminary experience. *Cancer* 1998; 82:1062–1070.
2. Nakagawa K, Aoki Y, Tago M, *et al.* Megavoltage CT-assisted stereotactic radiosurgery for thoracic tumors: original research in the treatment of thoracic neoplasms. *Int J Radiat Oncol Biol Phys* 2000;48:449–457.
3. Wulf J, Hadinger U, Oppitz U, *et al.* Stereotactic radiotherapy of targets in the lung and liver. *Strahlenther Onkol* 2001;177: 645–655.
4. Hara R, Itami J, Kondo T, *et al.* Stereotactic single high dose irradiation of lung tumors under respiratory gating. *Radiother Oncol* 2002;63:159–163.
5. Nagata Y, Negoro Y, Aoki T, *et al.* Clinical outcomes of 3D conformal hypofractionated single high-dose radiotherapy for one or two lung tumors using a stereotactic body frame. *Int J Radiat Oncol Biol Phys* 2002;52:1041–1046.
6. Onimaru R, Shirato H, Shimizu S, *et al.* Tolerance of organs at risk in small-volume, hypofractionated, image-guided radiotherapy for primary and metastatic lung cancers. *Int J Radiat Oncol Biol Phys* 2003;56:126–135.
7. Hof H, Herfarth KK, Munter M, *et al.* Stereotactic single-dose radiotherapy of stage I non-small-cell lung cancer (NSCLC). *Int J Radiat Oncol Biol Phys* 2003;56:335–341.
8. Whyte RI, Crownover R, Murphy MJ, *et al.* Stereotactic radiosurgery for lung tumors: preliminary report of a phase I trial. *Ann Thorac Surg* 2003;75:1097–1101.
9. Takai Y, Mituya M, Nemoto K, *et al.* [Simple method of stereotactic radiotherapy without stereotactic body frame for extracranial tumors]. *Nippon Igaku Hoshasen Gakkai Zasshi* 2001;61:403–407.
10. Negoro Y, Nagata Y, Aoki T, *et al.* The effectiveness of an immobilization device in conformal radiotherapy for lung tumor: reduction of respiratory tumor movement and evaluation of the daily setup accuracy. *Int J Radiat Oncol Biol Phys* 2001;50: 889–898.
11. Koga Y, Yano S, Okada T, *et al.* Stereotactic radiotherapy using a stereotactic body frame: research on effective irradiation angle and correcting dose. *Nippon Hoshasen Gijutsu Gakkai Zasshi* 2001;57:1395–1405.
12. Lax I, Blomgren H, Naslund I, *et al.* Stereotactic radiotherapy of malignancies in the abdomen. Methodological aspects. *Acta Oncol* 1994;33:677–683.
13. Blomgren H, Lax I, Naslund I, *et al.* Stereotactic high dose fraction radiation therapy of extracranial tumors using an accelerator. Clinical experience of the first thirty-one patients. *Acta Oncol* 1995;34:861–870.
14. Herfarth KK, Debus J, Lohr F, *et al.* Extracranial stereotactic radiation therapy: set-up accuracy of patients treated for liver metastases. *Int J Radiat Oncol Biol Phys* 2000;46:329–335.
15. Uematsu M, Shioda A, Suda A, *et al.* Computed tomography-guided frameless stereotactic radiotherapy for stage I non-small cell lung cancer: a 5-year experience. *Int J Radiat Oncol Biol Phys* 2001;51:666–670.
16. Emami B, Lyman J, Brown A, *et al.* Tolerance of normal tissue to therapeutic irradiation. *Int J Radiat Oncol Biol Phys* 1991;21:109–122.
17. Gomi K, Koichi M, Oguchi M, *et al.* Clinical experience of stereotactic radiation therapy for stage Ia non-small cell lung cancer [abstract]. 6th International Stereotactic Radiosurgery Society Congress, Kyoto, Japan, June 22–26, 2003; 146.
18. Graham MV, Purdy JA, Emami B, *et al.* Clinical dose-volume histogram analysis for pneumonitis after 3D treatment for non-small cell lung cancer (NSCLC). *Int J Radiat Oncol Biol Phys* 1999;45:323–329.

---

## Original Article

---

# Bilateral Breast-Conserving Therapy for Bilateral Breast Cancer: Results and Consideration of Radiation Technique

Chikako Yamauchi<sup>\*1</sup>, Michihide Mitsumori<sup>\*1</sup>, Yasushi Nagata<sup>\*1</sup>, Masaki Kokubo<sup>\*2</sup>, Takashi Inamoto<sup>\*3</sup>, Keiichi Mise<sup>\*4</sup>, Hiroshi Kodama<sup>\*4</sup>, and Masahiro Hiraoka<sup>\*1</sup>

<sup>\*1</sup>Department of Therapeutic Radiology and Oncology, Graduate School of Medicine, Kyoto University, <sup>\*2</sup>Department of Image-based Medicine, Institute of Biomedical Research and Innovation, Kobe, Japan, <sup>\*3</sup>Gastroenterological Surgery, Graduate School of Medicine, Kyoto University, <sup>\*4</sup>Kodama Breast Clinic, Kyoto, Japan.

---

**Background:** Although breast-conserving surgery followed by definitive irradiation is an established treatment for patients with early breast cancer, the role of breast-conserving therapy (BCT) for patients with bilateral breast cancer has not been well studied and the radiation therapy technique is still under investigation. We examined the feasibility of breast-conserving therapy for bilateral breast cancer and present here our radiation therapy technique with CT simulator.

**Methods:** Between July 1990 and December 1998, we treated 17 patients with bilateral breast cancer who underwent bilateral breast-conserving surgery followed by definitive irradiation. Seven patients had synchronous bilateral breast cancer and ten had metachronous bilateral breast cancer. Radiation therapy consisted of 50 Gy to the bilateral whole breast in all patients but one. A CT simulator was used to plan a tangential radiation field to the breast in all patients. Boost irradiation of 10 Gy was administered to 8 tumors with close or positive margins.

**Results:** With a median follow-up periods of 95 months from each operation, no patients showed loco-regional recurrence on either side, and none suffered distant metastasis. Furthermore no serious late adverse effects were observed.

**Conclusion:** This study demonstrated that BCT is feasible for bilateral breast cancer and the CT simulator is useful for determining the radiation field, especially when lesions are metachronous.

*Breast Cancer 12:135-139, 2005.*

Key words: Bilateral breast cancer, Breast-conserving therapy, BCT, Breast-conserving surgery, Radiation therapy

---

The incidence of clinically observed bilateral breast cancer is reported to range from 1.4 to 11.8%<sup>1-3)</sup>, small but significant. Although breast-conserving surgery followed by irradiation is an established treatment for patients with early breast cancer, the frequency of patients receiving bilateral breast irradiation ranges from 0.4% to 5.5%<sup>4-6)</sup>. The role of breast-conserving therapy for patients with bilateral breast cancer has not been well studied and scant attention has been devoted to

the techniques for radiation therapy. We herein present our technique, which utilizes a CT simulator, and analyze the outcome of treatment for patients with bilateral breast cancer treated with breast-conserving therapy (BCT).

## Materials and Methods

Between July 1990 and December 1998, a total of 1036 patients with breast cancer were treated with BCT, defined as breast-conserving surgery and axillary lymph node dissection followed by definitive radiation therapy at the Department of Radiology at Kyoto University Hospital. Among them, 35 patients (3.4%) had bilateral breast cancer, and 17 of them were treated with bilateral BCT (Fig 1). Therefore, 17 patients treated with bilateral BCT were analyzed in the present study.

---

Reprint requests to Chikako Yamauchi, Department of Therapeutic Radiology and Oncology Graduate School of Medicine, Kyoto University 54 Kawahara-cho, Shogoin, Sakyo, Kyoto, 606-8507, Japan.  
E-mail: chikay@kuhp.kyoto-u.ac.jp

Abbreviations:  
BCT, Breast-conserving therapy

Received June 1, 2004; accepted November 24, 2004




Fig 1. Total patients treated between July 1990 and December 1998.

Table 1. Patient Characteristics

	Synchronous (7 Pt.)	Metachronous (10 Pt.)
Age at diagnosis	Median 53 (43-68)	Median 45 (28-54)*
Family history		
1st degree	0	3
2nd degree	1	0
Menstrual status		
Premenopausal	2	8
Perimenopausal	0	0
Postmenopausal	2	2
Unknown	1	

\* age at the diagnosis of the 1st tumor

Seven patients had synchronous bilateral breast cancer and 10 patients had metachronous bilateral breast cancer. They developed the newly diagnosed contralateral breast cancer 4 to 70 months after the first BCT with a median interval of 29 months. Synchronous breast cancer was defined as the diagnosis of both tumors within 1 month. The patients' characteristics and the characteristics of the 34 breast cancers are summarized in Tables 1 and 2.

As regards conservative surgery, 14 tumors were treated by quadrantectomy, while 20 tumors were treated by wide excision. All patients underwent axillary dissection bilaterally. Twenty-six tumors had negative margins of resection, 6 had close margins of resection, that is, within 5 mm

Table 2. Tumor Characteristics of the 34 Treated Breasts

	Number	%
Pathology		
DCIS	1	
Invasive ductal	32	94
Invasive lobular	1	
Clinical T Stage		
T0	1	3
T1	22	65
T2	11	32
Clinical UICC Stage		
I	22	65
IIA	9	26
IIB	3	9
Pathologic N stage		
N0	31	91
N1	3	9
Estrogen receptor status		
Negative	11	32
Positive	13	38
Not done/unknown	10	30

from the resected margin, and 1 had positive margins of resection, defined as microscopic involvement at the resected margin on the histological examination.

Following breast conserving surgery, a total dose of 50 Gy in daily fractions of 2 Gy was delivered over 5 weeks to the whole breast via opposing tangential fields. We used a CT simulator (Shimadzu Corp. CT-S, Kyoto) to plan the tangential

fields. We selected the beam energy for the tangential fields according to the breast size: twenty-seven unilateral breasts were treated with cobalt-60 gamma rays, 1 with 4-MV photons, and 5 with 6-MV photons for the tangential fields. One breast was irradiated with an en-face electron beam. Seven patients with simultaneous breast cancer were treated by matched midline technique with bilateral tangential fields using the CT simulator (Fig 2). On the other hand, we referred to the CT simulation images of the first tumors to avoid field overlapping when we determined the tangential fields for the second tumors in the patients with metachronous breast cancers (Fig 3). The primary site was boosted in the 7 patients with close or positive surgical margins. This boost irradiation comprised to a total dose of 10 Gy in 5 fractions of electron beams through a field 6 to 8 cm in diameter, including the tumor bed. The ipsilateral supraclavicular and ipsilateral internal mammary nodal areas were not included in the target volume.

All patients received oral 5-fluorouracil (5-FU)

or its derivatives, and also received tamoxifen for 2 years after the operation, regardless of the axillary node status or estrogen receptor (ER) status.




Fig 2. A case of simultaneous breast cancer: It is confirmed that there is no overlap by the skin markings.





Fig 3. A case of metachronous breast cancer. (a) Radiation field for the first treatment. (b) Radiation field for the second treatment. Identifying the first field by the skin reaction is impossible. (c) We could recognize the first field accurately with the use of images from the previous CT simulation.



**Fig 4.** Another case of metachronous breast cancer. (a) The first field crossed the midline. (b) The second tumor existed in the inner quadrant. Surgical clip shows the location of the tumor bed. If we avoided overlapping with the previous field using a tangential field, the tumor bed would not be included in the field. (c) In this case, only the tumor bed was irradiated with 9 MeV electron beams.

The patients were periodically followed-up at our clinic. They were examined every 3 to 6 months in the first 2 years, and every 6 to 12 months thereafter according to their pathological status. Locoregional recurrence, distant metastasis, complications and cosmetic outcomes were evaluated.

### Results

Of 17 patients, 15 patients were irradiated with matched tangential fields without overlapping, 1 patient was irradiated with matched tangential fields with overlapping of 1.2 cm, and 1 patient with a medially located metachronous tumor received en-face electron beam alone because overlapping with the previous field could not be avoided with tangential field (Fig 4).

No patients were lost to follow-up. The median follow-up period after each operation was 95 months. No patients showed loco-regional recurrence on either side or distant metastasis.

Regarding complications associated with treatment, severe arm edema was observed in one patient whose upper arms showed a 4 cm differ-

ence in circumference. One patient developed moderate fibrosis at the site of overlapping, but this did not affect cosmetic outcome. The patient who was irradiated with overlapping of 1.2 cm did not develop any skin or soft tissue complications. In other cases, complications were none or slight.

We also evaluated cosmetic outcome using the cosmetic score<sup>7</sup>. Six patients (35%) were scored as excellent and 10 (59%) were scored as good. Only one patient (6%) was graded fair because of unilateral breast contracture.

### Discussion

Although as many as 10% of the patients with breast cancer may develop bilateral cancer<sup>1,3)</sup> and radiation therapy is essential to breast conserving therapy, there is scant information on the technical aspects of such irradiation<sup>8,9)</sup>. To minimize late damage to skin and soft tissue, overlapping of bilateral tangential fields should be avoided. On the other hand, maintaining good coverage of breast tissue is important to minimize the risk of intra-breast recurrence. In the patients with meta-

chronous breast cancer patients, which account for 2/3 of all bilateral cases, it is necessary to reproduce the previous tangential field before planning the contralateral tangential beam. In a conventional X-ray simulator, it is almost impossible to reproduce the medial margin accurately. Tattooing, which is commonly used in Western countries and might be useful in such situations, is seldom used in Japan. CT-simulation is quite useful because the overlapping of bilateral tangential fields can be evaluated much more accurately than conventional simulation, although there are some limitations derived from the change of the patient's figure and the difference in positioning. In patients with thick subcutaneous tissue at the midline, or those with tumors located very near to the midline, overlapping may be unavoidable despite the use of a CT simulator. However, it is still possible to explore the use of a CT for planning tangential fields for irradiation of metachronous breast cancer patients.

### Conclusion

This study demonstrated that BCT is feasible for bilateral breast cancer and the CT simulator is useful for determining the radiation field, especially when they are metachronous. It is helpful in minimizing overlap of the radiation fields and pro-

vides the best possible treatment plan.

### References

- 1) Donovan AJ: Bilateral breast cancer. *Surg Clin North Am* 70:1141-1149, 1990.
- 2) Gogas J, Markopoulos C, Skandalakis P, *et al*: Bilateral breast cancer. *Am Surg* 59:733-735, 1993.
- 3) Michowitz M, Noy S, Lazebnik N, *et al*: Bilateral breast cancer. *J Surg Oncol* 30:109-112, 1985.
- 4) van Limbergen E, van den Bogaert W, van der Schueren E, *et al*: Tumor excision and radiotherapy as primary treatment of breast cancer. Analysis of patient and treatment parameters and local control. *Radiother Oncol* 8:1-9, 1987.
- 5) Chu AM, Cope O, Russo R, *et al*: Patterns of local-regional recurrence and results in Stages I and II breast cancer treated by irradiation following limited surgery. An update. *Am J Clin Oncol* 7:221-229, 1984.
- 6) Bedwinek JM, Brady L, Perez CA, *et al*: Irradiation as the primary management of stage I and II adenocarcinoma of the breast: analysis of the RTOG breast registry. *Cancer Clin Trials* 3:11-18, 1980.
- 7) Harris JR, Levene MB, Svensson G, *et al*: Analysis of cosmetic results following primary radiation therapy for stages I and II carcinoma of the breast. *Int J Radiat Oncol Biol Phys* 5:257-261, 1979.
- 8) Kopelson G, Munzenrider JE, Doppke K, *et al*: Bilateral breast cancer: radiation therapy results and technical considerations. *Int J Radiat Oncol Biol Phys* 7:335-341, 1981.
- 9) Fung MC, Schultz DJ, Solin LJ: Early-stage bilateral breast cancer treated with breast-conserving surgery and definitive irradiation: the University of Pennsylvania experience. *Int J Radiat Oncol Biol Phys* 38:959-967, 1997.

# Increased DNA Methyltransferase 1 (DNMT1) Protein Expression Correlates Significantly with Poorer Tumor Differentiation and Frequent DNA Hypermethylation of Multiple CpG Islands in Gastric Cancers

Tsuyoshi Etoh,<sup>\*,†</sup> Yae Kanai,<sup>\*</sup> Saori Ushijima,<sup>\*</sup>  
Tohru Nakagawa,<sup>\*</sup> Yukihiko Nakanishi,<sup>\*</sup>  
Mitsuru Sasako,<sup>‡</sup> Seigo Kitano,<sup>†</sup> and  
Setsuo Hirohashi<sup>\*</sup>

From the Pathology Division,<sup>\*</sup> National Cancer Center Research Institute, Tokyo, Japan; the Gastric Surgery Division,<sup>‡</sup> National Cancer Center Hospital, Tokyo, Japan; and the Department of Surgery I,<sup>†</sup> Oita Medical University, Oita, Japan

We evaluated the significance of aberrant DNA methyltransferase 1 (DNMT1) protein expression during gastric carcinogenesis. The protein expression of DNMT1, Muc2, human gastric mucin, E-cadherin, and proliferating cell nuclear antigen was examined immunohistochemically in gastric cancers and corresponding noncancerous mucosae from 134 patients. The DNA methylation status of the CpG islands of the *p16*, human MutL homologue 1 (*bMLH1*), *E-cadherin*, and *thrombospondin-1* (*THBS-1*) genes and the methylated in tumor (MINT)-1, -2, -12, and -31 clones was examined by methylation-specific polymerase chain reaction and combined bisulfite restriction enzyme analysis. Epstein-Barr virus (EBV) infection was detected by *in situ* hybridization. Nuclear immunoreactivity for DNMT1 was not detected in any of the noncancerous epithelia, except in proliferative zones (positive internal control), but was found in 97 (72%) of the gastric cancers. DNMT1 overexpression correlated significantly with poorer tumor differentiation ( $P < 0.001$ ), but not with the phenotype (gastric type versus intestinal type) of the cancer cells. It also correlated significantly with DNA hypermethylation of the CpG islands of the *bMLH1* ( $P = 0.024$ ) and *THBS-1* genes ( $P = 0.043$ ), and with the CpG island methylator phenotype in the gastric cancers ( $P = 0.007$ ). Reduced E-cadherin expression correlated significantly with poorer tumor differentiation ( $P = 0.002$ ), DNA hypermethylation of the *E-cadherin* gene ( $P < 0.001$ ) and DNMT1 overexpression ( $P = 0.014$ ). DNMT1 overexpression was also associated with EBV infection (a potential etiological factor in gastric car-

cinogenesis) but not with the proliferative activity of the cancer cells as indicated by the proliferating cell nuclear antigen-labeling index. These results suggest that DNMT1 overexpression may not be just a secondary effect of increased cancer cell proliferative activity, but may be associated with EBV infection and other etiological factors during gastric carcinogenesis. Furthermore, DNMT1 may play a significant role in the development of poorly differentiated gastric cancers by inducing frequent DNA hypermethylation of multiple CpG islands. (*Am J Pathol* 2004; 164:689–699)

DNA methylation plays an important role in transcriptional regulation and chromatin remodeling in mammalian cells.<sup>1</sup> Both overall DNA hypomethylation and more regional DNA hypermethylation have been well documented in various cancers.<sup>1–8</sup> Aberrant DNA methylation may be involved in carcinogenesis as a result of 1) increased gene mutagenicity because of deamination of 5-methylcytosine to thymine; 2) a possible association of aberrant DNA methylation with allelic loss; and 3) repression of gene transcription through methylation of CpG islands in regulatory regions of specific genes, including tumor-suppressor genes.<sup>1</sup>

To date, three enzymes, DNA methyltransferase 1 (DNMT1),<sup>9</sup> DNMT3a, and DNMT3b,<sup>10</sup> have been confirmed to possess DNMT activity. Of these, DNMT1 is the major and best known. As DNMT1 shows a preference for hemimethylated rather than unmethylated substrates *in vitro*,<sup>11</sup> and targets replication foci by binding to proliferating cell nuclear antigen (PCNA),<sup>12</sup> it seems to be a

Supported by a Grant-in-Aid for the Second Term Comprehensive 10-Year Strategy for Cancer Control; a Grant-in-Aid for Cancer Research from the Ministry of Health, Labor, and Welfare of Japan; and a research resident fellowship from the Foundation for Promotion of Cancer Research in Japan (to T.N.).

Accepted for publication October 8, 2003.

Address reprint requests to Yae Kanai, Pathology Division, National Cancer Center Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan. E-mail: ykanai@ncc.go.jp.

maintenance form of DNMT that copies methylation patterns after DNA replication. However, some workers have proposed that DNMT1 possesses both maintenance and *de novo* DNA methylation activity *in vivo*, regardless of its *in vitro* substrate preference.<sup>13</sup>

Overexpression of DNMT1 has been detected in several human cancers.<sup>14–16</sup> With regard to gastric cancer, we have reported that DNMT1 mRNA expression levels were significantly higher in cancer tissues than in normal gastric mucosae.<sup>16</sup> Moreover, during this previous study, we found that increased DNMT1 mRNA expression correlated significantly with the CpG island methylator phenotype (defined as frequent DNA hypermethylation of C-type CpG islands that are methylated in a cancer-specific but not an age-dependent manner<sup>17</sup>) in gastric and colorectal cancers.<sup>16</sup> However, most previous analyses concerning DNMT1 expression in human cancers have been performed at the mRNA level. To our knowledge, DNMT1 protein expression in gastric cancers has never been reported. The aim of this study was therefore to evaluate the significance of aberrant DNMT1 protein expression during gastric carcinogenesis. Firstly, we searched for correlations between DNMT1 protein expression and the clinicopathological features of gastric cancers. Secondly, to determine the targets of aberrantly expressed DNMT1 during gastric carcinogenesis, we examined the correlations between DNMT1 protein expression on the one hand and the DNA methylation status of multiple C-type CpG islands and E-cadherin expression on the other. Thirdly, to clarify the background behind aberrant DNMT1 protein expression, we investigated correlations with the proliferative activity of cancer cells (as indicated by the PCNA-labeling index) and with etiological factors that are believed to be involved in gastric carcinogenesis, such as *Helicobacter pylori*<sup>18</sup> and Epstein-Barr virus (EBV) infection.<sup>19</sup>

## Materials and Methods

### Patients and Tissue Samples

Cancerous tissues and corresponding noncancerous mucosae were obtained from 134 patients with primary gastric cancer. These patients underwent surgery at the National Cancer Center Hospital, Tokyo, Japan, between 1998 and 2002. They included 92 men and 42 women with a mean ( $\pm$  SD) age of  $61 \pm 7$  years (range, 45 to 83 years). None of the patients received any preoperative treatment, such as radiation or chemotherapy. Based on histological examinations, the 134 tumors were classified as 23 well differentiated, 31 moderately differentiated, and 80 poorly differentiated (including signet ring cell and mucinous carcinomas) adenocarcinomas.

### Immunohistochemistry

Five- $\mu$ m-thick sections of formalin-fixed, paraffin-embedded tissue specimens from all 134 patients

were deparaffinized and dehydrated. For antigen retrieval, the sections were heated for 10 minutes at 120°C in an autoclave. Nonspecific reactions were blocked with 2% normal swine serum. All sections were incubated with specific primary antibodies that recognized DNMT1 (goat polyclonal antibody, sc-10219, dilution 1:1000; Santa Cruz Biotechnology, Santa Cruz, CA), Muc2 (mouse monoclonal antibody, Ccp58, dilution 1:100; Novocastra, Newcastle-on-Tyne, UK), human gastric mucin (mouse monoclonal antibody, 45M1, dilution 1:50; Novocastra), E-cadherin (mouse monoclonal antibody, HECD-1,<sup>20</sup> dilution 1:500), PCNA (mouse monoclonal antibody, p56720, dilution 1:200; Transduction Laboratories, Lexington, KY) and *H. pylori* (rabbit polyclonal antibody, B0471, dilution 1:20; DAKO, Glostrup, Denmark), respectively. We previously confirmed the specificity of the goat anti-human DNMT1 polyclonal antibody by Western blotting analysis: an immunoreactive band of  $\sim$ 193.5 kd, corresponding to the molecular mass of DNMT1, was detected in human cancer cells, but no nonspecific bands were detected with this antibody.<sup>21</sup> All primary antibody incubations were conducted at 4°C overnight and were followed by incubation with biotinylated secondary antibodies (anti-goat IgG, anti-mouse IgG, or anti-rabbit IgG, dilution 1:200; Vector Laboratories, Burlingame, CA) at room temperature for 30 minutes. The sections were then treated with Vectastain Elite ABC reagent (Vector Laboratories). 3,3'-Diaminobenzidine tetrahydrochloride was used as the chromogen. All sections were counterstained with hematoxylin.

The gastric cancers were classified into three phenotypes according to previously described criteria:<sup>22</sup> the gastric type (positive for human gastric mucin), the intestinal type (positive for Muc2), and the mixed type (positive for both human gastric mucin and Muc2). For the evaluations of DNMT1 and PCNA expression, nuclear immunoreactivity in the proliferative zones of noncancerous foveolar epithelia was used as a positive internal control for all sections. Similarly, immunoreactivity in the cell membranes of noncancerous foveolar epithelia was used as a positive internal control for all sections during the evaluation of E-cadherin expression. As a negative control, the primary antibodies were omitted from the reaction sequence.

### Methylation-Specific Polymerase Chain Reaction (MSP) and Combined Bisulfite Restriction Enzyme Analysis (COBRA)

High-molecular-weight DNA was extracted from 105 fresh paired samples of cancerous tissues and their corresponding noncancerous mucosae by phenol-chloroform extraction and dialysis. Bisulfite conversion was performed using 1  $\mu$ g of genomic DNA and the reagents provided in the CpGenome DNA modification kit (Intergen, Purchase, NY). This process converts unmethylated cytosine residues to uracil, whereas methylated cytosine residues remain unchanged. The DNA methylation status of the CpG islands of the *p16*,

*MutL* homologue 1 (*hMLH1*), and *E-cadherin* genes was determined by MSP. This technique is based on the principle that the DNA sequences of methylated and unmethylated genomic regions differ after bisulfite conversion and can thus be distinguished by sequence-specific polymerase chain reaction (PCR) primers.<sup>23</sup> The bisulfite-modified DNA of the *p16* gene was amplified using the primer sets provided in the CpG WIZ amplification kit (Intergen) and that of the *hMLH1*<sup>24</sup> and *E-cadherin*<sup>23</sup> genes was amplified using the previously described primers. The DNA methylation status of the *thrombospondin-1* (*THBS-1*) gene and the methylated in tumor (MINT)-1, -2, -12, and -31 clones was determined by COBRA.<sup>25</sup> Bisulfite-modified DNA was amplified by PCR using previously described primers that were designed to amplify methylated and unmethylated genomic regions equally.<sup>17</sup> The amplified fragments were digested with restriction enzymes that digest DNA only if the CpG sites in their recognition sequences are methylated: *TaqI* for the *THBS-1* gene and the MINT-1 and -2 clones, *MaeII* for the MINT-12 clone and *BstUI* for the MINT-31 clone, respectively. The reaction products were separated electrophoretically on a 3% agarose gel and stained with ethidium bromide. Signal intensities were measured using an image analyzer (model FMBIO-2; Takara, Ohtsu, Japan).

### In Situ Hybridization

Five- $\mu$ m-thick sections of formalin-fixed, paraffin-embedded tissue specimens from all 134 patients were deparaffinized, dehydrated, and predigested with proteinase K. The sections were then hybridized with a digoxigenin-labeled EBV encoding RNA (EBER) 1 oligonucleotide probe (EBV detection kit; Nichirei, Tokyo, Japan) for 2 hours at 37°C. Anti-digoxigenin antibody-alkaline phosphatase was used with a nitro blue tetrazolium/5-bromo-4-chloro-3-indolyl phosphate substrate to detect the EBER1 signal. Positive control specimens were provided by the manufacture. As a negative control, the EBER1 probe was omitted from the reaction sequence.

### Statistics

Correlations between the incidence of DNMT1 immunoreactivity and variables including clinicopathological parameters, the DNA methylation status of CpG islands, *E-cadherin* expression, and *H. pylori* and EBV infection were analyzed using the chi-square test. Correlations between the PCNA-labeling index and clinicopathological parameters or DNMT1 immunoreactivity were analyzed using the Mann-Whitney *U*-test or the Kruskal-Wallis test. Differences with *P* values <0.05 were considered significant.

## Results

### Clinicopathological Significance of DNMT1 Protein Overexpression in Gastric Cancers

Immunoreactivity for DNMT1 was detected in the nuclei, but not in the cytoplasm or cell membranes, of cells in the proliferating zones of foveolar epithelia, lymphocytes, and cancer cells (Figure 1). To discriminate definitely positive cases from cases with leaky background level signal, if more than 30% of the cells in a tissue sample exhibited nuclear staining the sample was considered to show positive immunoreactivity. None (0%) of the 134 noncancerous epithelia exhibited DNMT1 immunoreactivity (except in the proliferative zones, which acted as the positive internal control for the analysis), whereas 97 (72%) of the 134 gastric cancers were DNMT1-positive.

Correlations between the incidence of nuclear immunoreactivity for DNMT1 and the clinicopathological features of the gastric cancers are shown in Table 1. DNMT1 protein overexpression was significantly associated with the degree of histological differentiation (*P* < 0.001).


Next, we evaluated cellular phenotypes based on immunohistochemistry for Muc2 and human gastric mucin, as shown in Figure 2. Fifty (37%) of the cancers showed a gastric phenotype, 34 (26%) showed an intestinal phenotype, and a further 50 (37%) showed a mixed cellular phenotype. There was no significant correlation between DNMT1 protein overexpression and the cellular phenotypes.

We then focused on the histological features of the noncancerous mucosae, as intestinal metaplasia is considered to be a precancerous lesion for adenocarcinomas with an intestinal phenotype. There was no significant correlation between DNMT1 protein overexpression in the gastric cancers and the presence or absence (or, if present, the degree) of intestinal metaplasia in the corresponding noncancerous mucosae (data not shown).

DNMT1 protein overexpression was not significantly associated with other parameters relating to cancer aggressiveness, such as the depth of invasion, vascular involvement, or lymph node metastasis.

### Correlation between DNMT1 Protein Overexpression and the DNA Methylation Status of Multiple CpG Islands

Figure 3 shows examples of the PCR products from MSP and COBRA. The incidence of DNA methylation of the C-type CpG islands of each of the genes and MINT clones tested in the noncancerous mucosae and gastric cancers is summarized in Table 2. DNA methylation of at least one C-type CpG island was seen in 50 (48%) of the 105 noncancerous mucosae and 83 (80%) of the 105 gastric cancers examined. For all patients showing DNA methylation of a certain CpG island in both their



noncancerous mucosa and cancer, the signal intensity of the reaction products reflecting the presence of methylated DNA was increased in the cancer compared with the corresponding noncancerous mucosa (Figure 3A). The DNA methylation status of each C-type CpG island in the gastric cancers is shown in Figure 4. When DNA hypermethylation was seen on three or more C-type CpG islands, we regarded the patient as being CpG island methylator phenotype

**Figure 1.** Immunohistochemical examination using anti-human DNMT1 goat polyclonal antibody. In the noncancerous mucosal sample from patient S121 (A), only the proliferative zones of the foveolar epithelia showed nuclear immunoreactivity for DNMT1, whereas the other epithelial cells did not. Although most cancer cells in a well-differentiated adenocarcinoma from patient S91 lacked nuclear immunoreactivity for DNMT1 (B), the moderately differentiated adenocarcinoma from patient S115 (C) and the poorly differentiated adenocarcinoma from patient S130 (D) showed strong nuclear immunoreactivity. Original magnifications:  $\times 180$  (A);  $\times 360$  (B-D).


(CIMP)-positive, based on previously described criteria.<sup>17</sup> Twenty-five (24%) of the 105 gastric cancers were considered CIMP-positive. Furthermore, there were significant correlations between DNMT1 protein overexpression and DNA hypermethylation of each CpG island of the *hMLH1* ( $P = 0.024$ ) and *THBS-1* ( $P = 0.043$ ) genes in the gastric cancers (Table 3). There was also a significant correlation between DNMT1 protein overexpression and CIMP for the gastric cancer tissues ( $P = 0.007$ , Table 3).

**Table 1.** DNMT1 Protein Expression and the PCNA-Labeling Index in the Gastric Cancers


Variables	Analyzed	DNMT-1 positive [number of cases (%)]	P <sup>†</sup>	PCNA-labeling index [mean ± SD (%)]	P
Tumor differentiation					
Well differentiated	23	7 (30%)	<.001	62 ± 25	0.165 <sup>‡</sup>
Moderately differentiated	31	18 (58%)			
Poorly differentiated	80	72 (90%)			
Phenotype*					
Gastric type	50	38 (76%)	0.137	60 ± 29	0.793 <sup>‡</sup>
Intestinal type	34	21 (62%)			
Mixed type	50	34 (68%)			
Depth of invasion					
Mucosa/submucosa	38	28 (74%)	0.423	56 ± 32	0.125 <sup>‡</sup>
Muscularis propria/subserosa	11	6 (55%)			
Serosa	85	63 (74%)			
Vascular involvement					
Negative	55	41 (75%)	0.494	56 ± 30	0.533 <sup>§</sup>
Positive	79	56 (71%)			
Lymphnode metastasis					
Negative	81	60 (74%)	0.931	59 ± 27	0.465 <sup>§</sup>
Positive	53	37 (70%)			

\*Cellular phenotypes are defined as described in the Materials and Methods section.

<sup>†</sup>Chi-square test, <sup>‡</sup>Kruskal-Wallis test. <sup>§</sup>Mann-Whitney U-test.



**Figure 2.** Immunohistochemical examination using anti-human gastric mucin (A) and Muc2 (B) mouse monoclonal antibodies. A: An adenocarcinoma with a gastric phenotype and noncancerous foveolar epithelia (\*) from patient S19. Both showed strong cytoplasmic immunoreactivity for human gastric mucin. B: An adenocarcinoma with an intestinal phenotype from patient S78, showing strong cytoplasmic immunoreactivity for Muc2. Original magnifications, ×360.



**Figure 3.** Examples of PCR products from DNA methylation analyses of multiple CpG islands in patients with gastric cancer (S). N, noncancerous mucosae; T, cancer tissue. The DNA methylation status of the CpG islands of the *p16* (A), *hMLH1* (B), and *E-cadherin* (C) genes was evaluated by MSP. In this analysis, the PCR products generated by primer sets M and U reflect the presence of methylated and unmethylated genes, respectively. Primer set W was used to confirm the completeness of the bisulfite modification. A: Methylated gene was detected in S24N, but the signal intensity of the M fragment was markedly increased in S24T compared with S24N. This indicates that a greater number of cells had undergone DNA hypermethylation in T than in N, although N can contain precursor cells for cancers and/or precancerous lesions. DNA methylation of the CpG islands of the *THBS-1* (D) gene and the *MINT-1*, -2, -12, and -31 clones (E, F, G, and H, respectively) was evaluated by COBRA. In this analysis, only methylated genes (arrows) were digested by the restriction enzymes.

**Table 2.** The DNA Methylation Status of the C-Type CpG Islands of Various Genes and Clones in Noncancerous Mucosae and Gastric Cancers

CpG islands	Number of tissue samples (%)			
	Noncancerous mucosae		Gastric cancers	
	Analyzed	DNA hypermethylation detected	Analyzed	DNA hypermethylation detected
p16	104	18 (17)	103	23 (22)
hMLH1	105	15 (14)	105	18 (17)
THBS-1	105	2 (2)	104	25 (24)
MINT1	105	23 (22)	105	39 (37)
MINT2	105	1 (1)	105	26 (25)
MINT12	104	6 (6)	104	18 (17)
MINT31	105	0 (0)	105	10 (10)

### *Correlation between DNMT1 Protein Overexpression and Reduced E-Cadherin Expression*

E-cadherin was detected in the cell membranes of epithelia from all (100%) of the 134 noncancerous mucosae. However, E-cadherin protein expression was considered to be reduced when more than 50% of the gastric cancer cells in a particular patient's sample lacked or showed only slight membranous E-cadherin immunoreactivity. Reduced E-cadherin expression was observed in 59 (44%) of the 134 gastric cancers (Figure 5). The incidence of reduced E-cadherin expression was 22% in well, 23% in moderately, and 59% in poorly differentiated adenocarcinomas, respectively, and reduced E-cadherin expression was significantly associated with poorer tumor differentiation ( $P = 0.002$ ). DNA methylation of CpG island of the *E-cadherin* gene was seen in 20 (19%) of the 105 gastric cancers examined (Figure 3C) and there was a significant correlation between DNA hypermethylation of CpG island of the *E-cadherin* gene and reduced E-cadherin expression in the gastric cancers ( $P < 0.001$ ). Furthermore, there was a significant correlation between DNMT1 protein overexpression and reduced E-cadherin expression in gastric cancers ( $P = 0.014$ , Table 3). In fact, coincidence of nuclear immunoreactivity of DNMT1 and lack of membrane immunoreactivity of E-cadherin was frequently observed in individual cancer cells.

### *DNMT1 Protein Expression and the PCNA-Labeling Index*

Examples of the results of immunohistochemistry for PCNA are shown in Figure 6. To evaluate the PCNA-labeling index, ~300 cells per specimen were examined at a magnification of  $\times 400$  under a microscope and the cells that did and did not show nuclear immunoreactivity for PCNA were counted. The PCNA-labeling index was expressed as the percentage of the total cells that showed nuclear immunoreactivity. The PCNA-labeling index was increased even in well-differentiated adenocarcinomas, in which the incidence of overexpression of DNMT1 protein was still low (Table 1), and coincidence of nuclear immunoreactivity of PCNA and lack of nuclear immunoreactivity of DNMT1 was frequently observed in individual well-differentiated cancer cells. Thus, DNMT1

protein overexpression was not significantly associated with the PCNA-labeling index in the gastric cancers ( $P = 0.309$ , Figure 7).

### *Correlation between DNMT1 Protein Overexpression and Etiological Factors*

To understand the background behind DNMT1 protein overexpression, etiological factors considered to be involved in gastric carcinogenesis were examined. Although 44 (46%) of the 96 patients examined showed *H. pylori* infection in their noncancerous mucosae, there was no significant correlation between *H. pylori* infection and DNMT1 protein overexpression ( $P = 0.113$ ). Similarly, there was no correlation between *H. pylori* infection and CIMP ( $P = 0.146$ ). The incidence of EBV infection was examined by *in situ* hybridization in the same cohort (Figure 8). Four patients (4%) had EBV infection in their cancer cells, and all four of these cancers showed DNMT1 protein overexpression. Furthermore, EBV infection was significantly associated with DNA hypermethylation of five or more C-type CpG islands ( $P < 0.001$ ).

### *Discussion*

We believe that this is the first report describing immunohistochemical examination for DNMT1 protein in gastric cancers. We have previously reported an increase in DNMT1 mRNA expression levels in gastric cancers compared with corresponding noncancerous mucosae.<sup>16</sup> In the present study, we showed that DNMT1 expression was also increased at the protein level in gastric cancers, suggesting that DNMT1 overexpression has some significance during gastric carcinogenesis. DNMT1 protein overexpression showed no significant correlations with either the cellular phenotype (gastric type versus intestinal type) or the presence, absence, or degree of intestinal metaplasia (a precancerous lesion for adenocarcinomas with an intestinal phenotype) in corresponding noncancerous mucosae, suggesting that DNMT1 protein overexpression is associated with gastric carcinogenesis regardless of the cellular origin or phenotype. Our results also suggest that DNMT1 may particularly affect the stage of development at which cancers begin to show

Case	DNMT1	p16	MLH1	THBS-1	MINT1	MINT2	MINT12	MINT31	Number of methylated CpG islands
S1	+								≥ 5 CIMP- positive
S2	+								
S3	+								
S4	ND								
S5	ND								
S6	+								
S7	+								
S8	+								
S9	+								
S10	+								
S11	+							≥ 3 CIMP- positive	
S12	+								
S13	+								
S14	+								
S15	+								
S16	+								
S17	+								
S18	+								
S19	+								
S20	+								
S21	ND							< 3 CIMP- negative	
S22	+								
S23	+								
S24	+								
S25	+								
S26	+								
S27	+								
S28	+								
S29	+								
S30	+								
S31	ND								
S32	+								
S33	+								
S34	+								
S35	+								
S36	+								
S37	+								
S38	+								
S39	+								
S40	+								
S41	ND								
S42	-		ND						
S43	+								
S44	+								
S45	+								
S46	+								
S47	ND								
S48	+								
S49	+								
S50	+								
S51	+								
S52	+								
S53	+								
S54	+								
S55	+								
S56	+								
S57	+								
S58	+								
S59	+								
S60	+								
S61	+								
S62	+								
S63	+								
S64	+								
S65	+								
S66	+								
S67	+								
S68	+								
S69	+								
S70	+								
S71	+								
S72	+								
S73	+								
S74	+								
S75	+								
S76	+								
S77	+			ND					
S78	+								
S79	+								
S80	+								
S81	+								
S82	+						ND		
S83	+								
S84	+			ND					
S85	+								
S86	+								
S87	+								
S88	+								
S89	+								
S90	+								
S91	+								
S92	+								
S93	+								
S94	+								
S95	+								
S96	+								
S97	+								
S98	+								
S99	+								
S100	+								
S101	+								
S102	+								
S103	ND								
S104	ND								
S105	ND								

**Figure 4.** Protein expression of DNMT1 and DNA methylation profiles for seven C-type CpG islands in 105 gastric cancers. The DNA methylation status was examined using MSP or COBRA (see Figure 3). Patient numbers are indicated on the **vertical columns** and the seven CpG islands are indicated on the **top row**. +, DNMT1 protein overexpression-positive; -, DNMT1 protein overexpression-negative; **solid box**, methylated; **open box**, unmethylated; ND, not done. When DNA hypermethylation was seen in three or more CpG islands, the patient was regarded as being CIMP-positive.

poorer differentiation. However, in a previous study, overexpression was detected even in precancerous conditions when we used quantitative reverse transcriptase-PCR analysis to examine DNMT1 mRNA expression levels in a cohort with hepatocellular carcinomas (HCCs).<sup>15</sup> Although there appears to be a discrepancy between the previous and present findings, when the same cohort was examined by immunohistochemistry protein overexpression was not detected in precancerous


**Table 3.** Correlations between DNMT1 Protein Overexpression and the DNA Methylation Status of Each C-Type CpG Island, CIMP and E-Cadherin Protein Expression in Gastric Cancers

CpG islands	DNMT1 expression (number of tissue samples)		P*
	Positive	Negative	
p16			
Methylated	13	3	0.361
Unmethylated	56	24	
hMLH1			
Methylated	20	2	0.024
Unmethylated	49	25	
THBS-1			
Methylated	18	2	0.043
Unmethylated	51	25	
MINT1			
Methylated	28	6	0.091
Unmethylated	41	21	
MINT2			
Methylated	18	4	0.237
Unmethylated	51	23	
MINT12			
Methylated	13	3	0.361
Unmethylated	56	24	
MINT31			
Methylated	10	0	0.503
Unmethylated	59	27	
CIMP			
Positive	10	1	0.007
Negative	59	26	
E-cadherin expression			
Maintained	48	27	0.014
Reduced	49	10	

\* Chi-square test. Reduced: over 50% of the cancer cells in a particular patient's sample lacked or showed only slight E-cadherin immunoreactivity.


ous conditions but only in moderately or poorly differentiated HCCs.<sup>21</sup> This may be attributable to methodological differences: quantitative reverse transcriptase-PCR is so sensitive that it can detect small elevations in DNMT1 mRNA levels in precancerous conditions, whereas immunohistochemistry cannot detect such elevations until protein expression reaches a certain level in more malignant HCCs.<sup>21</sup> By analogy with hepatocarcinogenesis, we cannot rule out the possibility that a small elevation in DNMT1 expression had already occurred in the earlier stages of gastric carcinogenesis, before the DNMT1 expression level reached the threshold of detection for the immunohistochemical methods used.

Regional DNA hypermethylation of CpG islands was detected even in noncancerous mucosae, which can contain precursor cells for cancers and/or precancerous lesions, such as intestinal metaplasia. However, the incidence and degree of DNA hypermethylation was increased in the gastric cancers compared with the noncancerous mucosae. These data are consistent with previous findings in precancerous conditions and cancers of various organs.<sup>3-8</sup> Twenty-four percent of the gastric cancers were CIMP-positive, confirming that aberrant DNA methylation is associated with the multistage development of certain subgroups of gastric cancers.




**Figure 5.** Immunohistochemical examination using the anti-human E-cadherin mouse monoclonal antibody HECD-1. Although the well-differentiated adenocarcinoma from patient S31 maintained strong E-cadherin immunoreactivity at the cell-cell borders (A), E-cadherin expression was reduced in the poorly differentiated adenocarcinoma from patient S64 (B). Original magnifications,  $\times 360$ .


Targeting of the substrate DNA by DNMT1 may be disrupted by mechanisms such as dysfunction of p21WAF1,<sup>26</sup> which competes with DNMT1 for binding to PCNA, in cancer cells.<sup>12</sup> Moreover, it has recently been suggested that DNMT1 is capable of *de novo* methylating activity as well as having a maintenance function.<sup>13</sup> Therefore, it is feasible that, in cancers, DNMT1 participates in the DNA hypermethylation of CpG islands that are not methylated in normal cells. We have previously reported that DNMT1 mRNA overexpression correlates significantly with CIMP in gastric and colorectal cancers.<sup>16</sup> In the present study, we demonstrated a significant correlation between DNMT1 expression and CIMP in gastric cancers, even at the protein level. Moreover, among the C-type CpG islands examined, those of the *hMLH1* and *THBS-1* genes may be particularly targeted by overexpressed DNMT1. This is compatible with previ-



**Figure 6.** Immunohistochemical examination using anti-human PCNA mouse monoclonal antibody. A: A well-differentiated adenocarcinoma from patient S125. B: A poorly differentiated adenocarcinoma from patient S30. Original magnifications,  $\times 360$ .



**Figure 7.** Average PCNA-labeling indices in DNMT1 protein overexpression-positive ( $n = 97$ ) and -negative ( $n = 37$ ) gastric cancers. Error bar, SD. DNMT1 protein overexpression was not significantly associated with the PCNA-labeling index in the gastric cancers ( $P = 0.309$ , Mann-Whitney *U*-test).



**Figure 8.** *In situ* hybridization investigation for EBV infection using the EBER1 oligonucleotide probe. A poorly differentiated adenocarcinoma from patient 545, showing diffuse positivity in the nuclei. Original magnification,  $\times 180$ .

ous reports that silencing of the *hMLH1* gene results in frequent microsatellite instability in gastric cancers.<sup>27,28</sup>

E-cadherin is one of the most important molecules involved in intracellular adhesion and cancer morphogenesis.<sup>29</sup> In addition to the development of mutations,<sup>30–32</sup> it has become apparent that the *E-cadherin* gene can be silenced by DNA hypermethylation around its promoter regions in gastric cancers.<sup>33,34</sup> Indeed, reduced E-cadherin protein expression associated with DNA hypermethylation around the promoter region of the *E-cadherin* gene was significantly correlated with poorer glandular differentiation of the cancers examined in this study. We showed a significant correlation between DNMT1 protein overexpression and reduced E-cadherin protein expression in gastric cancers, indicating that the *E-cadherin* gene may also be a target of overexpressed DNMT1.

Although DNMT1 is a major DNMT in humans, so far two other enzymes, DNMT3a and DNMT3b, have also been shown to possess DNMT activity.<sup>1</sup> Genomic methylation patterns may be established through cooperation among these three DNMTs, even in cancer cells.<sup>13</sup> Moreover, it may be that unknown co-factors potentially target these DNMTs to unmethylated substrates in a sequence-specific manner in cancers. Thus, DNMT1 overexpression alone may not explain regional DNA hypermethylation in cancers. Further studies on the cooperation between DNMT1 and other components of the DNA methylation machinery in tissue specimens may further our understanding of the basis of regional DNA hypermethylation during gastric carcinogenesis. Although the degree of DNA hypermethylation in noncancerous mucosae seemed slight, DNA methylation of CpG islands actually occurred in a considerable number of noncancerous mucosae that can contain precursor cells for cancer. However, DNMT1 overexpression was not detected immunohistochemically in noncancerous mucosae except for proliferative zones, in which DNMT1 performs maintenance methylation at the replication foci. The protein

expression levels of DNMT1 in noncancerous mucosae may not reach the threshold that can be detected by the present immunohistochemical method, even though it promotes slight DNA hypermethylation. Otherwise, slight DNA hypermethylation in noncancerous mucosae may be attributable to alterations in components of the DNA methylation machinery other than DNMT1. Although frequent regional DNA hypermethylation has been reported in gastric type tumors,<sup>35</sup> DNMT1 protein expression was not different between gastric and intestinal phenotypes in the present study. We assume again that unknown components of the DNA methylation machinery may potentially target DNMT1 to substrate DNA, or that DNMTs other than DNMT1 may also participate in regional DNA hypermethylation in cancers showing gastric phenotype.

On the other hand, the overall 5-methylcytosine level is lower in cancer tissues than in normal tissues.<sup>1</sup> In common with the previously reported immunohistochemical findings for DNMT1 in colorectal cancers<sup>36</sup> and HCCs,<sup>21</sup> some cancer cells exhibited very weak or no DNMT1 immunoreactivity, even in a gastric cancer that was considered DNMT1-positive according to the criteria described in the Materials and Methods section. Cancer cells with low DNMT1 levels may be prone to overall DNA hypomethylation.

An initial increase in DNMT1 mRNA expression in colon cancers became far more modest when the expression level was normalized according to that of a cell proliferation marker.<sup>37</sup> DNMT1 mRNA is expressed mainly during the S-phase and, because tumor tissue is presumed to contain a greater proportion of dividing cells than normal tissue, some debate has arisen as to whether increased DNMT1 expression is because of an increase in the proportion of dividing cells or to an acute increase in DNMT1 expression per individual cell. This continuing discussion prompted us to compare DNMT1 immunoreactivity and the PCNA-labeling index in gastric cancers. In the present study, DNMT1 protein overexpression was not significantly associated with increased proliferative activity of gastric cancer cells; we have previously observed a similar discrepancy between DNMT1 expression and cell proliferative activity in precancerous conditions for urinary bladder carcinomas<sup>38</sup> and in certain subgroups of HCCs.<sup>21</sup> These findings suggest that DNMT1 protein overexpression does not result entirely from increased numbers of dividing cells during carcinogenesis.

Finally, we focused on etiological factors believed to be involved in gastric carcinogenesis to clarify the background behind DNMT1 protein overexpression. Although DNMT1 protein overexpression was not significantly associated with the incidence of *H. pylori* infection in corresponding noncancerous mucosae, all four of the EBV-positive cancers showed DNMT1 protein overexpression. EBV infection was significantly associated with the DNA hypermethylation of five or more C-type CpG islands, in accordance with a previous report that the average number of methylated CpG islands was higher in EBV-positive gastric cancers than in EBV-negative ones.<sup>39</sup> These observations indicate that DNMT1 may play an important

role in EBV-related gastric carcinogenesis via aberrant DNA methylation. Although it has previously been reported that transfection of latent membrane protein-1 of EBV induces the expression and activity of DNMT1,<sup>40</sup> latent membrane protein-1 is not typically expressed in EBV-associated gastric cancers.<sup>41</sup> Although the molecular mechanism governing how EBV infection results in overexpression of DNMT1 protein requires further elucidation, and our present findings must be interpreted with caution because of the small number of EBV-positive gastric cancers studied, EBV infection and other etiological factors may be associated with DNMT1 overexpression and contribute toward gastric carcinogenesis by inducing frequent DNA hypermethylation of multiple CpG islands.

## References

1. Jones PA, Baylin SB: The fundamental role of epigenetic events in cancer. *Nat Rev Genet* 2002, 3:415–428
2. Vachtenheim J, Horakova I, Novotna H: Hypomethylation of CCGG sites in the 3' region of H-ras protooncogene is frequent and is associated with H-ras allele loss in non-small cell lung cancer. *Cancer Res* 1994, 54:1145–1148
3. Kanai Y, Ushijima S, Tsuda H, Sakamoto M, Sugimura T, Hirohashi S: Aberrant DNA methylation on chromosome 16 is an early event in hepatocarcinogenesis. *Jpn J Cancer Res* 1996, 87:1210–1217
4. Kanai Y, Ushijima S, Hui AM, Ochiai A, Tsuda H, Sakamoto M, Hirohashi S: The E-cadherin gene is silenced by CpG methylation in human hepatocellular carcinomas. *Int J Cancer* 1997, 71:355–359
5. Eguchi K, Kanai Y, Kobayashi K, Hirohashi S: DNA hypermethylation at the D17S5 locus in non-small cell lung cancers: its association with smoking history. *Cancer Res* 1997, 57:4913–4915
6. Kanai Y, Ushijima S, Ochiai A, Eguchi K, Hui AM, Hirohashi S: DNA hypermethylation at the D17S5 locus is associated with gastric carcinogenesis. *Cancer Lett* 1998, 122:135–141
7. Kondo Y, Kanai Y, Sakamoto M, Mizokami M, Ueda R, Hirohashi S: Genetic instability and aberrant DNA methylation in chronic hepatitis and cirrhosis—a comprehensive study of loss of heterozygosity and microsatellite instability at 39 loci and DNA hypermethylation on 8 CpG islands in microdissected specimens from patients with hepatocellular carcinoma. *Hepatology* 2000, 32:970–979
8. Kanai Y, Ushijima S, Tsuda H, Sakamoto M, Hirohashi S: Aberrant DNA methylation precedes loss of heterozygosity on chromosome 16 in chronic hepatitis and liver cirrhosis. *Cancer Lett* 2000, 148:73–80
9. Bestor T, Laudano A, Mattaliano R, Ingram V: Cloning and sequencing of a cDNA encoding DNA methyltransferase of mouse cells. The carboxyl-terminal domain of the mammalian enzymes is related to bacterial restriction methyltransferases. *J Mol Biol* 1988, 203:971–983
10. Okano M, Xie S, Li E: Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat Genet* 1998, 19:219–220
11. Bestor TH: Activation of mammalian DNA methyltransferase by cleavage of a Zn binding regulatory domain. *EMBO J* 1992, 11:2611–2617
12. Chuang LS, Ian HI, Koh TW, Ng HH, Xu G, Li BF: Human DNA-(cytosine-5) methyltransferase-PCNA complex as a target for p21WAF1. *Science* 1997, 277:1996–2000
13. Rhee I, Bachman KE, Park BH, Jair KW, Yen RW, Schuebel KE, Cui H, Feinberg AP, Lengauer C, Kinzler KW, Baylin SB, Vogelstein B: DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* 2002, 416:552–556
14. Sun L, Hui AM, Kanai Y, Sakamoto M, Hirohashi S: Increased DNA methyltransferase expression is associated with an early stage of human hepatocarcinogenesis. *Jpn J Cancer Res* 1997, 88:1165–1170
15. Saito Y, Kanai Y, Sakamoto M, Saito H, Ishii H, Hirohashi S: Expression of mRNA for DNA methyltransferases and methyl-CpG-binding proteins and DNA methylation status on CpG islands and pericentromeric satellite regions during human hepatocarcinogenesis. *Hepatology* 2001, 33:561–568
16. Kanai Y, Ushijima S, Kondo Y, Nakanishi Y, Hirohashi S: DNA methyltransferase expression and DNA methylation of CpG islands and peri-centromeric satellite regions in human colorectal and stomach cancers. *Int J Cancer* 2001, 91:205–212
17. Toyota M, Ahuja N, Ohe-Toyota M, Herman JG, Baylin SB, Issa JP: CpG island methylator phenotype in colorectal cancer. *Proc Natl Acad Sci USA* 1999, 96:8681–8686
18. Sipponen P: *Helicobacter pylori*: a cohort phenomenon. *Am J Surg Pathol* 1995, 19(Suppl 1):S30–S36
19. Shibata D, Weiss LM: Epstein-Barr virus-associated gastric adenocarcinoma. *Am J Pathol* 1992, 140:769–774
20. Shimoyama Y, Hirohashi S, Hirano S, Noguchi M, Shimozato Y, Takeichi M, Abe O: Cadherin cell-adhesion molecules in human epithelial tissues and carcinomas. *Cancer Res* 1989, 49:2128–2133
21. Saito Y, Kanai Y, Nakagawa T, Sakamoto M, Saito H, Ishii H, Hirohashi S: Increased protein expression of DNA methyltransferase (DNMT) 1 is significantly correlated with the malignant potential and poor prognosis of human hepatocellular carcinomas. *Int J Cancer* 2003, 105:527–532
22. Egashira Y, Shirmoda T, Ikegami M: Mucin histochemical analysis of minute gastric differentiated adenocarcinoma. *Pathol Int* 1999, 49:55–61
23. Herman JG, Graff JR, Myohanen S, Nelkin BD, Baylin SB: Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci USA* 1996, 93:9821–9826
24. Herman JG, Umar A, Polyak K, Graff JR, Ahuja N, Issa JP, Markowitz S, Willson JK, Hamilton SR, Kinzler KW, Kane MF, Kolodner RD, Vogelstein B, Kunkel TA, Baylin SB: Incidence and functional consequences of hMLH1 promoter hypermethylation in colorectal carcinoma. *Proc Natl Acad Sci USA* 1998, 95:6870–6875
25. Xiong Z, Laird PW: COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res* 1997, 25:2532–2534
26. Baylin SB: Tying it all together: epigenetics, genetics, cell cycle, and cancer. *Science* 1997, 277:1948–1949
27. Yanagisawa Y, Akiyama Y, Iida S, Ito E, Nomizu T, Sugihara K, Yuasa Y, Maruyama K: Methylation of the hMLH1 promoter in familial gastric cancer with microsatellite instability. *Int J Cancer* 2000, 85:50–53
28. Fleisher AS, Esteller M, Tamura G, Rashid A, Stine OC, Yin J, Zou TT, Abraham JM, Kong D, Nishizuka S, James SP, Wilson KT, Herman JG, Meltzer SJ: Hypermethylation of the hMLH1 gene promoter is associated with microsatellite instability in early human gastric neoplasia. *Oncogene* 2001, 20:329–335
29. Hirohashi S: Inactivation of the E-cadherin-mediated cell adhesion system in human cancers. *Am J Pathol* 1998, 153:333–339
30. Oda T, Kanai Y, Oyama T, Yoshiura K, Shimoyama Y, Birchmeier W, Sugimura T, Hirohashi S: E-cadherin gene mutations in human gastric carcinoma cell lines. *Proc Natl Acad Sci USA* 1994, 91:1858–1862
31. Becker KF, Atkinson MJ, Reich U, Becker I, Nekarada H, Siewert JR, Hoffer H: E-cadherin gene mutations provide clues to diffuse type gastric carcinomas. *Cancer Res* 1994, 54:3845–3852
32. Muta H, Noguchi M, Kanai Y, Ochiai A, Nawata H, Hirohashi S: E-cadherin gene mutations in signet ring cell carcinoma of the stomach. *Jpn J Cancer Res* 1996, 87:843–848
33. Yoshiura K, Kanai Y, Ochiai A, Shimoyama Y, Sugimura T, Hirohashi S: Silencing of the E-cadherin invasion-suppressor gene by CpG methylation in human carcinomas. *Proc Natl Acad Sci USA* 1995, 92:7416–7419
34. Tamura G, Yin J, Wang S, Fleisher AS, Zou T, Abraham JM, Kong D, Smolinski KN, Wilson KT, James SP, Silverberg SG, Nishizuka S, Terashima M, Motoyama T, Meltzer SJ: E-cadherin gene promoter hypermethylation in primary human gastric carcinomas. *J Natl Cancer Inst* 2000, 92:569–573
35. Endoh Y, Tamura G, Ajioka Y, Watanabe H, Motoyama T: Frequent hypermethylation of the hMLH1 gene promoter in differentiated-type tumors of the stomach with the gastric foveolar phenotype. *Am J Pathol* 2000, 157:717–722
36. De Marzo AM, Marchi VL, Yang ES, Veeraswamy R, Lin X, Nelson WG: Abnormal regulation of DNA methyltransferase expression during colorectal carcinogenesis. *Cancer Res* 1999, 59:3855–3860

37. Lee PJ, Washer LL, Law DJ, Boland CR, Horon IL, Feinberg AP: Limited up-regulation of DNA methyltransferase in human colon cancer reflecting increased cell proliferation. *Proc Natl Acad Sci USA* 1996, 93:10366-10370
38. Nakagawa T, Kanai Y, Saito Y, Kitamura T, Kakizoe T, Hirohashi S: Increased DNA methyltransferase (DNMT) 1 protein expression in human transitional cell carcinomas of the urinary bladder. *J Urol* (in press)
39. Kang GH, Lee S, Kim WH, Lee HW, Kim JC, Rhyu MG, Ro JY: Epstein-Barr virus-positive gastric carcinoma demonstrates frequent aberrant methylation of multiple genes and constitutes CpG island methylator phenotype-positive gastric carcinoma. *Am J Pathol* 2002, 160:787-794
40. Tsai CN, Tsai CL, Tse KP, Chang HY, Chang YS: The Epstein-Barr virus oncogene product, latent membrane protein 1, induces the downregulation of E-cadherin gene expression via activation of DNA methyltransferases. *Proc Natl Acad Sci USA* 2002, 99:10084-10089
41. Rowlands DC, Ito M, Mangham DC, Reynolds G, Herbst H, Hallissey MT, Fielding JW, Newbold KM, Jones EL, Young LS, Niedobitek G: Epstein-Barr virus and carcinomas: rare association of the virus with gastric adenocarcinomas. *Br J Cancer* 1993, 68:1014-1019

## Focus on gastric cancer

Toshikazu Ushijima<sup>1</sup> and Mitsuru Sasako<sup>2,\*</sup>

<sup>1</sup>Carcinogenesis Division, National Cancer Center Research Institute

<sup>2</sup>Department of Surgery

National Cancer Center Central Hospital, Tokyo, Japan 104-0045

\*Correspondence: msasako@gan2.ncc.go.jp

### Epidemiology and incidence statistics

Gastric cancer is the second most common cancer in the world (Ferlay et al., 2001). It is unique in that its time trend and geographical distribution are very informative in estimating its risk factors. In the US, the crude mortality rate in Caucasian males was 33/100,000 in the early 20<sup>th</sup> century, and this declined to 5/100,000 in the late 20<sup>th</sup> century. The declining trend is worldwide, and the decline began earlier in developed countries. However, even among them, mortality is still high in Korea (43/100,000), Russia (35/100,000), Japan (31/100,000), and Portugal (22/100,000). The age-adjusted incidence reaches as high as 70/100,000 in Korean and Japanese males. The male to female ratio is consistently two to one in many geographical regions.

The decline took place following the popularization of refrigerators, which resulted in a decreased intake of salt and an increased intake of fruit and vegetables (Palli, 2000; Potter et al., 1997). The preventive effects of fruit and vegetables are consistently confirmed by many epidemiological studies. Most epidemiological studies have shown the promoting effect of salt and the preventive effect of vitamin C. The effects of salt were also shown by animal experiments. Some epidemiological studies suggest that consumption of grilled meat/fish increases the risk, and that the consumption of carotenoids and green tea reduce the risk. Epidemiological data linking *N*-nitrosamines to gastric cancers have so far been inconclusive, although their carcinogenicity at high doses is proven.

Infection by *Helicobacter pylori* is prevalent in areas with high incidences of gastric cancers, and increases the risk of gastric cancer. However, in some Asian countries, such as India and Thailand, incidences of gastric cancers are not high in spite of the high *H. pylori* infection rates, a phenomenon known as the "Asian Enigma" (Miwa et al., 2002). Possible explanations for this include host genetic factors, different virulence among strains of *H. pylori*, and dietary factors. Polymorphisms of proinflammatory cytokine genes have been shown to associate with risk of *H. pylori*-related gastric cancers (El-Omar et al., 2000).

### Animal models

A rat model for gastric cancers induced by a chemical carcinogen, *N*-methyl-*N*'-nitro-*N*-nitrosoguanidine, has been widely used for a variety of purposes, such as evaluation of various promoting and preventing factors and clarification of genes involved in genetic susceptibility. (Yamashita et al., 2002). A model in which *H. pylori* could infect an animal was established using Mongolian gerbils, which contributed to clarification of the strong promoting effect of *H. pylori* (Shimizu et al., 1999).

In addition, there have been more than 10 lines of genetically modified mice that show hyperplasia of the gastric epithelium and/or intestinal metaplasia (Gut et al., 2002). These mouse models were created by targeting genes involved in ion trans-

port, signal transduction, transcriptional regulation, and cell adhesion. Development of gastric cancers was observed in mice lacking the pS2 trefoil protein, those lacking *Smad4/Dpc4*, those lacking the SHP2 binding site on the Ii-6 family corepressor gp130, and those lacking *RUNX3* (Judd et al., 2004; Lefebvre et al., 1996; Xu et al., 2000; Li et al., 2002).

### Disease mechanism and molecular targets

#### **Histological classification and gastric/intestinal phenotypes**

Histological classification of gastric cancers is different between Japan and Western countries. Generally, "differentiated" and "undifferentiated" types in Japanese classification correspond to "intestinal" and "diffuse" types, respectively, in the Western classification established by Lauren. It has been considered that intestinal-type gastric cancers are associated with intestinal metaplasia, whereas diffuse-type gastric cancers are originated from gastric mucosa proper. Recent analysis of gastric and intestinal phenotypes in early gastric cancers has shown that cancer cells with gastric phenotypes were present in both intestinal and diffuse types of gastric cancer. Furthermore, phenotypic expression in gastric cancer cells was shown to be independent of phenotypic changes in the surrounding gastric mucosa (Tatematsu et al., 2003).

#### **Gastric cancer predisposition**

Germline mutations of *E-cadherin* were first found in a large family from New Zealand in which diffuse-type gastric cancers took place at an early age (Guilford et al., 1998). Although *E-cadherin* germline mutations are very rare, the finding provided to be useful information for clinicians to manage high-risk patients. Gastric cancers, mainly of the intestinal type, can be associated with hereditary nonpolyposis colorectal cancer (HNPCC) syndrome, most cases of which are caused by germline mutations of mismatch repair genes *hMLH1* or *hMSH2*, and are more prominently manifested in older generations of HNPCC patients. Patients with familial adenomatous polyposis, which is caused by germline mutations of *APC*, and Peutz-Jeghers syndrome also have increased risk for gastric cancer (Oberhuber and Stolte, 2000).

#### **Molecular alterations in gastric cancer**

Many genes have been analyzed in attempts to understand the molecular bases for human gastric cancers, but only a few with frequent alterations have been identified (Table 1). Oncogenic activations of  $\beta$ -*catenin* (17%–27% in intestinal type) and *K-ras* (0%–18% in both histological types) have been found in human gastric cancers (Lee et al., 2002; Park et al., 1999). In addition, amplifications of the *c-erbB2* and the *c-met* genes have each been found in approximately 10% of both histological types.

As for tumor-suppressor genes, *p53* mutations are repeatedly reported in gastric cancers of the diffuse type (0%–21%) and intestinal type (36%–43%) (Maesawa et al., 1995).

**Table 1.** Histology and genetic alterations of gastric cancers

	Diffuse type (%)	Intestinal type (%)
Oncogene activation		
<i>β-catenin</i>	0	17–27
<i>K-ras</i>	0–6	0–18
<i>c-erbB2</i>	12–13	12–13
Inactivation of tumor suppressor genes		
<i>p53</i>	0–21	36–43
<i>APC</i>	0–5	0
<i>E-cadherin</i>		
Mutation	33–50	0
Methylation	79	55
<i>p16</i>		
Mutation	0	0
Methylation	11*	50*
Microsatellite instability	5–32	23–41

\*Incidences are overestimated due to analysis of CpG islands in exons.

Mutations of the *APC* tumor suppressor gene are found frequently in gastric adenomas, but only rarely in gastric cancers; this is clearly different from the similar frequencies of *APC* mutations in colorectal adenomas and carcinomas (Lee et al., 2002; Maesawa et al., 1995). Somatic mutations of *E-cadherin* are observed specifically in sporadic diffuse-type gastric cancers (33%–50%) (Becker et al., 1994). *RUNX3* was recently shown to be a tumor-suppressor gene of gastric cancers, although its mutations were rare (Li et al., 2002).

Microsatellite instability (MSI) is observed in 5%–10% of diffuse-type gastric cancers and in 15%–40% of intestinal-type gastric cancers. The major mechanism for the MSI in gastric cancer is inactivation of the mismatch repair gene *hMLH1* resulting from hypermethylation of its promoter (Fang et al., 2003). Similarly, mutation of the *p16* gene is infrequent, but hypermethylation of *p16* is common (25%–42% overall) in gastric cancer, with the intestinal type having higher incidence (Ding et al., 2003; Oue et al., 2002).

#### Factors that induce molecular alterations

Although *hMLH1* and *p16* can be inactivated in gastric cancers by mutations or by promoter hypermethylation, inactivation by methylation is much more frequent than mutation in sporadic gastric cancers. The second hit in *E-cadherin* germline mutation carriers is also generally due to methylation (Machado et al., 2001). A genome-wide scan for aberrant methylations revealed silencing of nine genes in gastric cancers (Kaneda et al., 2002). Even in noncancerous gastric mucosae (Waki et al., 2002), aberrant methylation can be present. These findings suggest that aberrant methylation is deeply involved in gastric carcinogenesis, and aberrant methylation seems to be useful as a new target for diagnostics and prevention of gastric cancers.

The presence of Epstein-Barr virus (EBV) is observed in 7%–20% of gastric cancers, being slightly more frequent in diffuse-type gastric cancers. EBV is clonal in cancer tissue, and is maintained as a plasmid. EBV has been shown to extend cell generations of gastric epithelial cells in in vitro cell culture, but it cannot immortalize them (Takada, 2000). Recently, EBV-associated gastric cancers were shown to be more frequently associated with promoter methylation of *p16* (Kang et al., 2002).

There has been discussion about whether intestinal metaplasia (IM) is a precancerous lesion for gastric cancers. Although gastric cancers are frequently accompanied by IM, no molecular alterations that cause both IM and gastric cancers

have been identified. It is thus more likely that factors that induce molecular alterations for IM, such as *H. pylori* infection (Uemura et al., 2001), also induce molecular alterations for gastric cancers.

#### Diagnosis of gastric cancers

Most patients with gastric cancer are diagnosed when they undergo endoscopy and biopsy after exhibiting symptoms. In Japan, about 25% of patients are diagnosed by mass screening or a personal health check (Japanese National Gastric Cancer Registry). In high-risk areas of this disease, the most important issue is the education of general practitioners and the public to make them aware of the risk of this cancer. Early diagnosis used to be made by a barium meal study, especially in mass screening in Japan (Oshima, 1997). Endoscopy is being used more and more for secondary prevention in combination with a serum test of pepsinogen subtypes. However, there is a consensus that the efficacy of mass screening itself should be reevaluated (Tsubono and Hisamichi, 2000).

At an early phase of development, a well-differentiated carcinoma (WDC) replaces the mucosa of atrophic gastritis or IM without showing any invasion. As tumors progress, they start to invade the lamina propria mucosae or the muscularis mucosae, then the submucosal layer. As these invasive parts are often missed by biopsy, the lesions are often diagnosed as dysplasia. Thus, many lesions initially diagnosed as severe dysplasia turned out to be an invasive cancer, sometimes invading even the muscularis propria, after histological evaluation of resected materials (Fertitta et al., 1993).

#### Diagnostic criteria for early gastric cancers and endoscopic mucosal resection

Diagnostic criteria of WDC differs to some extent between the West and the East (Schlemper et al., 1997). In Western countries, the diagnosis of adenocarcinoma is made only when pathologists can recognize the evidence of invasion, while the term cancer is used in the East when cellular or structural atypia is evident, even without evidence of invasion. WHO classification now clearly states that the lesions called severe dysplasia/adenoma in the West are the same as noninvasive mucosal carcinoma in the East, and this is a result of pathologists' mutual communication and cooperation (Fenoglio-Preiser et al., 1997). The Western policy runs the risk of overlooking true cancers, but the Eastern policy may induce overtreatment. However, as the result of the development of the technique of endoscopic mucosal resection (EMR), the majority of such lesions are now treated endoscopically in Japan (Ono et al., 2001). Thus, paradoxically, "severe dysplasia" is often treated by surgery in the West, and "noninvasive mucosal carcinoma" is treated by EMR. This treatment can be applied exclusively to mucosal cancer, for which endoscopic ultrasound (EUS) is sometimes helpful. Because the histology of the entire specimen resected using EMR can be examined in detail, additional surgery can be applied without much delay if a patient's tumor is found to have submucosal invasion. Because of these potential advantages, distribution of the EMR technique to the West is urgently needed.

#### Metastases and their diagnosis

Gastric cancer remains a localized disease for a long time and metastasizes slowly. Table 2 shows the incidence of metastasis to lymph nodes, the liver, and the peritoneum according to tumor depth. Metastasis to sites other than these three sites is rare. Systemic metastasis seldom occurs until the late phase of

**Table 2.** Incidence of metastasis and five-year survival rates by tumor size and depth

Depth		Number of cases	Incidence (%)			Five-year survival rate (%)
			LN metastasis	Liver metastasis	Peritoneal metastasis	
pT1	M	1063	3.3	0.0	0.0	93.3
	SM	881	17.4	0.1	0.0	88.9
pT2	MP	436	46.4	1.1	0.5	81.3
	SS	325	63.7	3.4	2.2	65.8
pT3	SE	1232	78.9	6.3	17.8	35.5
pT4	SI	724	89.8	15.5	41.6	10.1
Overall		4683	47.8	4.5	11.5	60.3

Patients operated on between 1972–1991 at National Cancer Center Hospital, including exploratory laparotomy. pT1: pathologically confirmed tumor invasion of mucosa and/or muscularis mucosa (M) or submucosa (SM). pT2: pathologically confirmed tumor invasion of muscularis propria (MP) or subserosa (SS). pT3: pathologically confirmed tumor penetration of serosa (SE). pT4: pathologically confirmed tumor invasion of adjacent structures (SI).

local invasion (T3/4). By deeper invasion, nodal metastasis occurs more massively and to more distant areas. Nearly 20% of T2 tumors have metastasis at the second tier nodes. Systemic and local recurrences of T1/T2 lesions are rare when treated by proper lymph node dissection, while local recurrence is frequent after limited surgery (Sasako, 2003).

Conventional CT scanning is useful in detecting enlarged nodes, which are often irresectable. However, 25% of metastatic nodes are 5 mm or less and undetectable by any imaging diagnostic tool, including MRI, PET scan, or EUS (Noda et al., 1998).

#### Treatment of gastric cancers and its recent advances

Tumors without distant metastasis are potentially curable, and treatment comprises resection of the primary tumor and control of lymph node metastasis. For differentiated-type T1 mucosal cancers, EMR is often successful, as metastasis does not generally occur (Gotoda et al., 2000). The Japanese Gastric Cancer Treatment Guideline indicates the criteria for EMR as follows: mucosal cancer of intestinal type, no ulcer nor ulcer scar in the lesion, and size smaller than 21 mm (Nakajima, 2002). For more advanced lesions, gastrectomy of over 2/3 of the stomach with proper lymph node dissection is regarded as standard treatment even in the West (Sasako, 2003; Allum et al., 2002; NCCN Guideline [[http://www.nccn.com/physician\\_gls/f\\_guidelines.html](http://www.nccn.com/physician_gls/f_guidelines.html)]), in spite of the negative results of two large randomized trials (Bonenkamp et al., 1999; Cuschieri et al., 1999).

Tumors with distant metastasis are mostly incurable at present, with the rare exceptions of those with solitary liver metastasis or peritoneal nodules. For these advanced or recurrent tumors, chemotherapy shows a modest effect, and cure by medical treatment is rare, even in combination with radiotherapy. Combination chemotherapy using 5-fluorouracil with other agents remains the most popular regimen.

#### Chemoradiotherapy and D2 surgery

Recently, chemoradiotherapy (CRT) after a potentially curative operation was shown to improve the results of surgery without lymph node dissection (MacDonald et al., 2001). As adjuvant chemotherapy has not proven its efficacy over surgery alone, these results strongly suggest the efficacy of radiotherapy to achieve good local control. However, the results achieved by limited surgery followed by CRT are still worse than those of extended surgery, so-called D2 nodal dissection. Currently, questions regarding whether CRT in combination with limited surgery can replace D2 surgery and whether CRT can improve the results of D2 surgery alone remain to be answered. The for-

mer should be evaluated in the Western specialized centers, where D2 surgery can be carried out safely with sufficient quality. If this proves the efficacy of CRT, both questions should be investigated in Japan. Meta-analysis evaluating the effect of adjuvant chemotherapy without irradiation after curative surgery for gastric cancer suggested strongly the effect of the treatment. As none of the large sized trials has proven the effect of adjuvant chemotherapy, it is urgent to establish standard adjuvant treatment. At the moment, a large randomized trial is going on using TS-1, which showed the highest response rate as a single agent in the past. In Western countries, neoadjuvant chemotherapy for advanced gastric cancer is now being tested in a few large phase III trials. Neoadjuvant CRT is just now under investigation as a phase II trial in some American institutions.

#### New chemotherapeutic agent

Some new chemotherapeutic agents, such as Irinotecan, TS-1, and Docetaxel, show promise as being more effective than conventional drugs. A combination chemotherapy including TS-1 has shown a response rate of over 70% (Koizumi et al., 2003). Further studies may change the chemotherapy for gastric cancer.

#### Future challenges

Severe dysplasia/noninvasive mucosal carcinoma could contain different entities that have different abilities to invade the lamina propria mucosae. However, key molecular alterations that determine this progression are unknown. The presence of lymph node or distant metastasis is a very important factor in deciding a treatment strategy, but accurate diagnosis is still difficult. Clarification of molecular alterations that are closely linked with these characteristics will be beneficial to decide on a treatment strategy for individual cases. Popular use of EMR raises a new question, whether or not a secondary cancer will arise from the remnant stomach, and prediction of risk for developing gastric cancer is becoming more important. Recent genomic approaches demonstrate great potential for addressing these issues (Hasegawa et al., 2002). The more important and appropriate questions we ask, the more useful these new approaches will be.

#### References

- Allum, W.H., Griffin, S.M., Watson, A., and Colin-Jones, D. (2002). Guidelines for the management of oesophageal and gastric cancer. *Gut* 50, 1–23.
- Becker, K.F., Atkinson, M.J., Reich, U., Becker, I., Nekarda, H., Siewert, J.R., and Hofler, H. (1994). E-cadherin gene mutations provide clues to diffuse