

Table 1
Clinicopathological characteristics of the 105 patients treated by gefitinib

Category	n	%
Age		
Median (range)	61.9 (37–86)	
Gender		
Male	61	58.1
Female	44	41.9
Clinical stage		
IIIA	2	1.9
IIIB	7	6.7
IV	96	91.4
Performance status		
0	31	29.5
1	51	48.6
2	18	17.1
3–4	5	4.8
Histologic type		
Adenocarcinoma	88	83.8
Squamous	6	5.7
Large	3	2.9
Adenosquamous	3	2.9
Undefined	5	4.8
Smoking history		
None	62	59.0
Current + former	43	41.0
Serum CEA level		
<5 ng/ml	39	37.1
≥5 ng/ml	66	62.9
No. of prior chemotherapy regimens		
0	11	10.5
1	38	36.2
2	26	24.8
3 or more	30	28.6
Response to gefitinib		
Complete response	2	1.9
Partial response	26	24.8
Stable disease	36	34.2
Progressive disease	36	34.3
Not evaluable	5	4.8

9.3 months. We analysed the effect of pretreatment serum CEA level on the survival of patients treated by gefitinib. Table 3 shows a comparison of the pretreatment clinicopathological characteristics between the patients with an elevated CEA level and normal CEA level. Although the adenocarcinoma patients tended to have a more elevated CEA level than other cell types, there was no significant difference between the two groups with respect to the analysed categories. The survival curve of the two levels of serum CEA showed that survival of patients with higher pretreatment CEA level to be significantly better (Fig. 1). A univariate analysis of several prognostic factors using Cox proportional hazards model indicated that younger age, presence of adenocarcinoma, good PS and elevated serum CEA levels to be positive prognostic factors for gefitinib treat-

Table 2
Comparison of pretreatment clinicopathological characteristics among patients with response, stable disease and progressive disease by gefitinib treatment

Category	CR + PR (n = 28)	SD (n = 36)	PD (n = 36)	P value
	n	n	n	
Age				
<65	17	22	16	0.17
≥65	11	14	20	
Gender				
Male	13	19	24	0.10
Female	15	17	12	
Histologic type				
Adenocarcinoma	26	30	28	0.11
Non-adenocarcinoma	2	6	8	
Clinical stage				
III	1	2	4	0.23
IV	27	34	32	
Performance status				
≤1	24	29	26	0.18
≥2	4	7	10	
Smoking history				
None	18	14	11	0.0092
Current + former	10	22	25	
No. of prior regimens				
≤1	14	11	21	0.39
≥2	14	25	15	
Serum CEA level				
<5 ng/ml	6	12	19	0.009
≥5 ng/ml	22	24	17	
Total (%)	28 (28%)	36 (36%)	36 (36%)	

ment (Table 4). A multivariate analysis using a stepwise method also confirmed that a good PS and elevated serum CEA levels to be independent prognostic factors (Table 5).

4. Discussion

Gefitinib is a tyrosine kinase inhibitor of EGFR, which has the potential to reduce tumour volume in NSCLC patients. Two large phase II studies conducted in pretreated non-small lung cancer patients have demonstrated a response rate of 18% and 11% [3,4]. In the analysis of the former trial, Japanese patients observed higher response rate than non-Japanese patients (27.5% vs. 10.4%, odds ratio = 3.27; $P = 0.0023$). In a multivariate analysis employed at 10% significance level, a good PS, female, adenocarcinoma and a history of receiving prior immuno/hormonal treatment were all found to be independent predictable factors for response. Also a retrospective study demonstrated that patients with adenocarcinoma of the bronchioloalveolar subtype and no history of smoking were more likely to

Table 3
Comparison of pretreatment clinicopathological characteristics between patients with elevated serum CEA level and those without

Category	CEA < 5 ng/ml	CEA ≥ 5 ng/ml	P value
	n	n	
Age (<65, ≥65)	20, 19	37, 29	0.78
Gender (Male, female)	23, 16	38, 28	>0.99
Histologic type (Adeno, non-adeno)	29, 10	59, 7	0.080
Clinical stage (III, IV)	6, 33	3, 63	0.12
Performance status (≤1, ≥2)	33, 6	49, 17	0.32
Smoking history (None, current + former)	14, 25	29, 37	0.54
No. of prior regimens (≤1, ≥2)	18, 21	31, 35	>0.99
Total (%)	39 (37.1%)	66 (62.9%)	

have an objective response to gefitinib treatment (odds ratio = 13.5 and 4.2) [5]. However, no report has so far investigated the relationship between the serum CEA concentration and responses to. In this study, a similar overall response rate (26.7% of all patients treated) as that reported in the former study of Japanese patients and association of smoking history to gefitinib sensitivity was also observed. We also demonstrate for the first time, that patients with a serum CEA concentration of over 5 ng/ml were more sensitive to gefitinib treatment than those with a concentration of below 5 ng/ml. Moreover, those with an elevated serum CEA level showed a significantly better prognosis for gefitinib treatment than those with no such increased levels based




Fig. 1. The survival curves of gefitinib treated patients with elevated serum CEA (≥5 ng/ml) and normal CEA levels (<5 ng/ml).

Table 4
Analysis of various pretreatment prognostic factors influencing survival of patients treated with gefitinib (Cox proportional hazards model)

Variables	Hazard ratio	95% CI	P value
Age (<65 vs. ≥65)	1.72	1.02–2.90	0.042
Gender (Male vs. female)	0.69	0.40–1.17	0.17
Histologic type (Adeno vs. non-adeno)	1.88	1.01–3.51	0.046
Clinical stage (III vs. IV)	0.78	0.33–1.83	0.57
Performance status (≤1 vs. ≥2)	5.41	3.07–9.62	<0.0001
Smoking history (None vs. current + former)	1.56	0.90–2.68	0.11
No. of prior regimens (≤1 vs. ≥2)	1.35	0.79–2.29	0.28
CEA (<5 vs. ≥5 ng/ml)	0.53	0.32–0.90	0.018

Table 5
Multivariate analysis of various pretreatment prognostic factors influencing survival of patients treated with gefitinib

Variable	Category	Hazard ratio	P value
Performance status	(≤1 vs. ≥2)	6.10	<0.0001
CEA	(<5 vs. ≥5 ng/ml)	0.44	0.0027

on both univariate and the multivariate analyses. These data were very surprising since an elevated serum CEA level is generally considered to be a negative prognostic factor for NSCLC [13].

CEA was first described as a specific antigen that was present in both the fetal colon and colon adenocarcinoma [14]. It is a member of the immunoglobulin supergene family, which is a cell surface adhesion protein, and it is thought to play a role in cell-to-cell adhesion [15]. The overexpression of CEA has been found in many other types of carcinomas and is thought to play a role in tumorigenesis [16]. Screaton and colleagues have recently discovered that CEA has a dominant effect in blocking differentiation and it also cooperates with Myc and Bcl-2 in cellular transformation [17]. In addition, it can also inhibit cell death induced by a loss of anchorage to the extra cellular matrix (anoikis) [18]. Our data reported here suggests that NSCLC cells which produce an abundant amount of CEA protein tend to be more sensitive to the EGFR tyrosine kinase inhibitor gefitinib, and indicates that CEA proteins may play an important role in EGFR signaling in cancer cells. If this is true, then the serum CEA may be an important surrogate marker of the gefitinib treatment. In clinical practice, in the course of treating patients with gefitinib, we believe that a change in serum CEA levels seem to closely represent the burden of a CEA positive tumour.

Recently, mutations in the tyrosine kinase domain of EGFR have been found to be strongly associated with the sensitivity of NSCLC to gefitinib [6,7]. Mutations were detected within an ATP binding pocket of the catalytic domain, and the EGFR mutants also had an enhanced tyrosine kinase activity in response to the ligand. Moreover, current studies have shown that such downstream molecules as Akt and STAT3/5 play a crucial role in the anti-apoptotic pathways of the mutant EGFR in tumour cells [8,9]. Since CEA protein has been demonstrated to have an anti-apoptotic effect in cancer cells, it is possible that an anti-apoptotic signal of the mutant EGFR may elevate the expression level of CEA protein.

In conclusion, this study indicates that the serum CEA level may be a useful predictive factor for the efficacy of gefitinib treatment, while also being a prognostic factor for advanced NSCLC patients undergoing this treatment. Further basic research and clinical studies are needed to elucidate the relationship between sensitivity to gefitinib and the CEA protein.

Conflict of interest statement


None declared.

Acknowledgements

We thank Dr. Brian T. Quinn for critical comments on the manuscript, and Ms. Yumiko Oshima for her help in preparing the manuscript.

References

1. The Non-small Cell Lung Cancer Collaborative Group. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. *Br Med J* 1995, **311**, 899–909.
2. Wakeling AE, Guy SP, Woodburn JR, et al. ZD1839 (Iressa): an orally active inhibitor of epidermal growth factor signaling with potential for cancer therapy. *Cancer Res* 2002, **62**, 5749–5754.
3. Fukuoka M, Yano S, Giaccone G, et al. Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2003, **21**, 2237–2246.
4. Kris MG, Natale RB, Herbst RS, et al. Efficacy of gefitinib, an inhibitor of the epidermal growth factor receptor tyrosine kinase, in symptomatic patients with non-small cell lung cancer: a randomized trial. *JAMA* 2003, **290**, 2149–2158.
5. Miller VA, Kris MG, Shah N, et al. Bronchioloalveolar pathologic subtype and smoking history predict sensitivity to gefitinib in advanced non-small-cell lung cancer. *J Clin Oncol* 2004, **22**, 1103–1109.
6. Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004, **350**, 2129–2139.
7. Paez JG, Janne PA, Lee JC, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 2004, **304**, 1497–1500.
8. Sordella R, Bell DW, Haber DA, et al. Gefitinib-sensitizing EGFR mutations in lung cancer activate anti-apoptotic pathways. *Science* 2004, **305**, 1163–1167.
9. Cappuzzo F, Magrini E, Ceresoli GL, et al. Akt phosphorylation and gefitinib efficacy in patients with advanced non-small-cell lung cancer. *J Natl Cancer Inst* 2004, **96**, 1133–1141.
10. Travis WD, Colby TV, Corrin B, et al. Histological typing of lung and pleural tumors. *World Health Organization international histological classification of tumors*. 3rd ed. Berlin, Springer, 1999.
11. International-Union-against-cancer. TNM classification of malignant tumors. 5th ed. New York, Wiley; 1997.
12. Green S, Weiss GR. Southwest Oncology Group standard response criteria, endpoint definitions and toxicity criteria. *Invest New Drugs* 1992, **10**, 239–253.
13. Concannon JP, Dalbow MH, Hodgson SE, et al. Prognostic value of preoperative carcinoembryonic antigen (CEA) plasma levels in patients with bronchogenic carcinoma. *Cancer* 1978, **42**, 1477–1483.
14. Gold P, Freedman SO. Demonstration of tumor-specific antigens in human colonic carcinomata by immunological tolerance and absorption techniques. *J Exp Med* 1965, **121**, 439–462.
15. Benchimol S, Fuks A, Jothy S, et al. Carcinoembryonic antigen, a human tumor marker, functions as an intercellular adhesion molecule. *Cell* 1989, **57**, 327–334.
16. Hammarstrom S. The carcinoembryonic antigen (CEA) family: structures, suggested functions and expression in normal and malignant tissues. *Semin Cancer Biol* 1999, **9**, 67–81.
17. Screaton RA, Penn LZ, Stanners CP. Carcinoembryonic antigen, a human tumor marker, cooperates with Myc and Bcl-2 in cellular transformation. *J Cell Biol* 1997, **137**, 939–952.
18. Ordonez C, Screaton RA, Ilantzis C, et al. Human carcinoembryonic antigen functions as a general inhibitor of anoikis. *Cancer Res* 2000, **60**, 3419–3424.



Long-term survivors in stage IV non-small cell lung cancer

Tatsuro Okamoto*, Riichiroh Maruyama, Fumihiko Shoji, Hiroshi Asoh, Jiro Ikeda, Tetsuya Miyamoto, Tomomi Nakamura, Tetsuro Miyake, Yukito Ichinose

Department of Thoracic Oncology, National Kyushu Cancer Center, 3-1-1, Notame, Minami-ku, Fukuoka 811-1395, Japan

Received 6 April 2004; received in revised form 18 June 2004; accepted 21 June 2004

KEYWORDS

Non-small cell lung cancer;
Stage IV disease;
Long-term survivor;
Surgery;
Chemotherapy;
Radiotherapy;
Multimodality treatment;
Prognostic factor

Summary

Background and Objectives: To determine the prognostic factors for long-term survivors (LTS) with stage IV non-small cell lung cancer (NSCLC) who had undergone various treatments.

Patients and methods: From 1990 to 1999, 222 NSCLC patients with stage IV disease, who had been treated in our department, were reviewed. As the initial treatment, 135 patients (48%) were treated with chemotherapy alone, 52 patients with a combination of chemotherapy and radiotherapy, 19 patients underwent an operation with or without any other therapeutic modalities and 16 were received radiotherapy alone.

Results: Seventeen (7.7%) patients survived for more than 2 years, and all but one had adenocarcinoma. Among these LTS, eight patients received surgery as the initial therapy, and 16 (94.1%) received some type of local-control therapy, including surgery or radiotherapy, during the course of their disease. Regarding the clinical characteristics between LTS and others (non-LTS), an early N status, a single metastatic site, a good performance status, and surgery for initial therapy were all found to be significantly important factors for LTS. A multivariate analysis using a logistic regression model also showed an early N status and surgical treatment to be significantly associated with LTS.

Conclusions: Selected patients with an early N status may be appropriate candidates for aggressive multimodality treatment including surgery, in order to provide a long-term survival for stage IV NSCLC.

© 2004 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Approximately 80% of all lung cancer patients have non-small cell lung cancer (NSCLC) [1]. Although

* Corresponding author. Tel.: +81 92 541 3231;
fax: +81 92 551 4585.
E-mail address: tokamoto@nk-cc.go.jp (T. Okamoto).

a surgical resection is the most effective therapy for an early stage NSCLC; most patients with NSCLC have advanced disease which cannot be resected at the initial presentation [2]. Many recent studies have demonstrated that platinum-based chemotherapy may have a better potential for prolonging survival than the best supportive care for these patients [3,4]. A greater than 20% reduction in the risk of death and an improvement in the median survival time of at least 6 weeks, have been reported.

Since 1990, several new agents, referred to as third generation agents, have been developed including: paclitaxel; docetaxel; vinorelbine; gemcitabine; and irinotecan. Several recent phase III trials have clearly demonstrated platinum-based combination chemotherapy with these new generation agents to be preferable to single agent chemotherapy or a combination of platinum and old generation agents, including vindesine and etoposide [5,6]. In spite of the recent advances in chemotherapy, the prognosis of patients with stage IV NSCLC is still poor and most large phase III trials have shown a median survival time of from 8 to 10 months and a 1-year survival rate of 30–35% [2].

Among many patients with stage IV NSCLC, there must be various types of patients whose clinical characteristics or clinical course differ, and we also sometimes experience a long-term survival in clinical practice. There are several reports on long-term survivors (LTS) among NSCLC patients with stage IV disease NSCLC [7–11], however, only 4 to 6% of patients with stage IV have been reported to survive for more than 2 years. In these studies, almost all the patients were initially treated with chemotherapy; however, there is still little detailed information regarding the clinical course of LTS. In the present paper, we therefore investigated the clinical characteristics of LTS among all the patients with stage IV NSCLC who were treated at our institution, and analyzed the clinical prognostic factors regarding such patients, in order to obtain a better understanding for the treatment of this metastatic disease.

2. Methods and materials

Between January 1990 and December 1999, 1251 patients with primary lung cancer were treated at the Department of Thoracic Oncology of National Kyushu Cancer Center. One thousand and one hundred fifteen patients (89.1%) had NSCLC. Among them, 284 of NSCLC (25.5%) patients had stage IV disease and 222 received some kind of

Table 1 Details of the initial treatment for patients with stage IV NSCLC

	Number of patients	Percentage
Chemotherapy alone	135	60.8
Chemotherapy + radiotherapy	52	23.4
Radiotherapy alone	16	7.2
Surgery alone	9	4.1
Surgery + chemotherapy	6	2.7
Surgery + radiotherapy	1	0.5
Surgery + chemotherapy + radiotherapy	3	1.4
Total	222	100.0

treatment. As the initial treatment, chemotherapy was performed in 135 patients (48%), a combination of chemotherapy and radiotherapy in 52, an operation with or without other therapeutic modalities in 19 and radiotherapy alone in 16 (Table 1).

The histological analysis of the tumor was based on the WHO classification for cell types [12]. A panel of pathologists reviewed the histopathologic materials. The clinical stage of these patients was determined based on the TNM classification of the Union Internationale Contre Cancer (UICC) [13]. For TNM staging, all patients underwent a computed tomography (CT) scan of the thorax and the upper part of abdomen, a bone scintigram, and a brain CT or MRI. Statistical significance for the various clinicopathological factors between LTS and non-LTS was evaluated using the χ^2 test, Fisher's exact probability test, or the Mann–Whitney test. A multivariate analysis of the clinical factors for LTS was performed by a logistic regression model. The survival curves for all patients and surgically treated patients were estimated according to the Kaplan–Meier method [14]. The terminal event was death due to any cause. Differences were considered to be statistically significant if the *P* value was below 0.05.

3. Results

3.1. Long-term survivors

The median survival time (MST) for all 222 patients who received any kind of treatment was 6.8 months, and the 1-year survival rates was 27.5%.

Among the 222 patients, 17 (7.7%) patients survived for more than 2 years. On the other hand, the MST, 1- and 2-year survival rates for patients who did not receive any treatment was 3.5 months, 8 and 0%. The details of the individual characteristics and the clinical course of LTS are presented in Table 2. Fifteen patients received chemotherapy during the course of their disease. Eight patients received surgery as the initial therapy, and 16 (94.1%) received some type of local-control therapy, including surgery or radiotherapy, during the course of their disease. All but one had adenocarcinoma and 16 were classified as having an ECOG performance status 0 or 1. Eight patients survived for more than 3 years, and three of them have been alive for more than 8 years. These three had single metastasis at the initial treatment and underwent a resection of the primary site.

3.2. Three cases who survived for more than 8 years

Patient 1 had squamous cell carcinoma in segment 3 (S3) of the left upper lobe, which was diagnosed based on a transbronchial biopsy. Two small metastases in segment 6 (S6) of the left lower lobe were demonstrated by a chest CT scan. A left upper lobectomy and a segmentectomy of S6 were performed and pulmonary metastases were also confirmed by a pathological examination. Patient 2 had adenocarcinoma in the left upper lobe and bone metastasis in the left sacrum. A pathological diagnosis of the metastasis was performed by a bore needle biopsy. He underwent concurrent chemoradiotherapy for preoperative therapy, with uracil-tegafur and cisplatin plus radiotherapy (45 gray) for bone metastasis. A left upper lobectomy and an intra-operative radiation (25 gray) were performed. Jejunal metastasis was detected 7 months later, which was again removed by operation. These two patients have so far shown no recurrence. Patient 3 had a huge mass shadow in the right middle and lower lobe with a nodular shadow in segment 6 of the left lower lobe. Brushing cytology under bronchoscopy was performed for each lesion and adenocarcinoma was detected from both specimens. A bilobectomy of the right middle and lower was performed. After surgery, five cycles of trans-bronchial interferon-gamma therapy using nebulizar was given in accordance with his request for more than a year. In addition, 2 years later, a total of 10 cycles of chemotherapy with four regimens were given over a 4-year period. He is alive with multiple pulmonary metastases.

3.3. Factors associated with LTS

A comparison of clinical characteristics between LTS and the others (non-LTS) are shown in Table 3. An early N status, a single metastatic site, a good performance status, and surgery as the initial therapy were significantly predominant in the LTS. The multivariate analysis using a logistic regression model showed an early N status and surgical treatment to be significantly associated with LTS (Table 4).

3.4. Surgery as the initial treatment

Among 19 patients who underwent surgical treatment, 17 patients had a single metastatic site; four brain, four bone, four lung, two cervical lymph nodes (LN), one axillary LN, one adrenal, and one subcutaneous metastasis. Two patients had two metastatic sites; one had adrenal gland and bone metastasis, while the other had adrenal gland and brain metastasis. All of the surgically treated patients had a controlled metastatic site and a good performance status (0 or 1). Fifteen patients were males and four were females, and the average age was 57.6 years (46–75). Thirteen had adenocarcinoma, two large cell carcinoma, two adenosquamous cell carcinoma, and two squamous cell carcinoma.

A lobectomy was performed for 10 patients, a bilobectomy for three patients, a pneumonectomy for five patients, and a wedge resection for one patient. The site where metastasis exists had no statistical impact on survival among surgical patients. When these surgical patients were divided into two categories according to the pathological N status, the 2- and 5-year survivals of the nine patients with N0 and N1 disease were 70 and 47%, respectively, while only one out of nine with N2 and N3 disease survived more than 2 years.

4. Discussion

Many randomized trials and several recent meta-analyses have demonstrated platinum-based chemotherapy to have a greater potential of prolonging survival than the best supportive care, and chemotherapy for patients with a good performance status is thus recommended [3,4]. However, the prognosis of patients with stage IV NSCLC is still poor; most large phase III trials have shown a median survival time of from 8 to 10 months and a 1-year survival rate of 30–35% [2]. We have hardly experienced that patient with metastatic NSCLC

Table 2 Individual characteristics of the 17 patients with >2 years survival

Patients	Age	Gender	PS	Survival (months)	Cell type	Alive/dead	Metastatic sites	Clinical stage	Initial therapy	CT regimen (initial treatment)	Response to CT	Further treatment
1	60	Male	0	120.3	sq	Alive	Lung (s)	T2N1	S (primary + pulmonary meta)			–
2	47	Male	1	108.9	ad	Alive	Bone (s)	T3N0	CT + RT (bone) → S (primary)	CDDP + UFT	NC	S (small intestine)
3	46	Male	1	102.8	ad	Alive	Lung (s)	T2N0	S (primary)			CT
4	58	Male	0	56.2	ad	Dead	Adrenal gland (d)	T2N3	CT + RT (primary + adrenal)	CDDP + UFT	PR	CT, RT (abdominal LN)
5	49	Male	0	53.9	ad	Alive	Axillary LN (m)	T2N3	CT → S (primary)	CDDP + DOC	PR	–
6	71	Female	1	41.3	ad	Dead	Lung (m)	T2N2	CT	S-1	NC	CT, RT (bone)
7	53	Male	1	38.0	ad	Dead	Lung (m), cervical LN	T4N0	CT	CDDP + VP16	NC	CT, RT (whole brain)
8	73	Male	1	36.5	ad	Dead	Adrenal gland (s)	T2N0	CT	CDDP + VDS	NC	–
9	54	Male	0	34.3	ad	Dead	Bone (s)	T2N1	S (bone) → CT → S (primary)	CDDP + UFT	NC	RT (bone), RT (radiosurgery), S (inguinal LN)
10	55	Female	0	32.1	ad	Dead	Lung (m)	T2N0	CT	DOC	PR	CT, RT (whole brain)
11	53	Male	1	31.9	ad	Dead	Brain (d)	T1N2	CT + RT (whole brain)	CDDP + DOC	NC	CT + RT (primary)
12	73	Female	0	30.0	ad	Alive	Bone (m)	T2N0	S (primary) + RT (bone)			CT
13	65	Male	0	28.2	ad	Dead	Lung (m)	T2N0	CT	CDDP + VDS + MMC	PR	CT, RT (bone), RT (whole brain)
14	67	Male	0	26.8	ad	Dead	Brain (m), lung (m)	T4N3	CT	CDDP + DOC	PR	CT, RT (whole brain)
15	68	Female	2	26.3	ad	Dead	Bone (d)	T4N0	CT + RT (primary)	CDDP + UFT	PR	–
16	50	Female	0	25.1	ad	Dead	Brain (s)	T2N0	S (brain) → S (primary) → RT (whole brain)			CT, CT + RT (adrenal)
17	70	Male	1	24.9	ad	Alive	Brain (s)	T2N0	S (brain) → S (primary) → RT (whole brain)			–

PS: performance status; s: single; d: double; m: multiple; LN: lymph node; CT: chemotherapy; RT: radiotherapy; S: surgery; CDDP: cisplatin; DOC: docetaxel; VDS: vindesine; MMC: mitomycin.

Table 3 Differences between the clinical characteristics of long-term survivors (LTS) and non-LTS

Characteristics		Non-LTS (n)	LTS (n)	P value
Age	<65/≥65	84/121	10/7	>0.999
Gender	Male/Female	152/53	12/5	0.973
Histologic type	Adenocarcinoma	156	16	—
	Squamous cell	35	1	
	Large cell	10	0	
	Adeno-squamous cell	4	0	
T stage	T1/T2/T3/T4/Tx	20/82/21/79/3	1/12/1/3/0	0.108
N stage	N0/N1/N2/N3/Nx	26/20/74/82/3	10/2/2/3/0	0.0003
Performance status	0/1/2/3/4	56/91/36/15/7	9/7/1/0/0	0.010
Number of metastatic sites	1/≥2	120/85	15/2	0.031
Metastatic sites				
Brain	Yes/no	61/144	4/13	0.791
Lung	Yes/no	82/123	7/10	>0.999
Bone	Yes/no	96/109	4/13	0.109
Liver	Yes/no	24/181	0/17	0.277
Adrenal gland	Yes/no	28/176	2/15	>0.999
Initial therapy	Surgery +/-	11/194	8/9	<0.0001

Table 4 Multivariate analysis using a logistic regression model for LTS in all patients with stage IV NSCLC

Variable	Category	n	OR	95%CI	P value
T stage	T1,2/T3,4	115/104	1.49	0.38–5.43	0.560
N stage	N0,1/N2,3	58/161	4.31	1.25–14.9	0.021
Performance status	0,1/2,3	162/60	3.75	0.39–30.30	0.266
Number of metastatic sites	1/≥2	135/87	3.21	0.63–16.13	0.159
Initial therapy	Surgery +/-	19/203	6.54	1.71–25.0	0.006

OR: odds ratio; CI: confidence interval.

could be successfully cured by chemotherapy alone in clinical practice.

Some studies in the literature have reported LTS among unresectable cases of NSCLC [7–11]. Patients with stages III and IV disease were included in these studies, and they were mainly treated with initial chemotherapy followed by other treatments. Only 4–6% of stage IV patients have been reported to have survived for more than two years. A similar number of patients (7.7%) survived more than two years in the present series. There are two different groups among LTS regarding the clinical course; first are those who have chemotherapy-sensitive tumors; and thus, underwent long-term chemotherapy alone, while the second group received multimodality therapy including systemic chemotherapy plus any local-control therapy as surgery or radiotherapy during the clinical course. The majority of LTS with NSCLC belong to the latter group. In the study of European Lung Cancer Working Party, 79% of the patients with unresectable NSCLC who survived more than 4 years were treated

after chemotherapy by other therapeutic modalities [9]. Satoh and coworkers [10] reported that nine out of the 14 survivors of more than 2 years underwent chest irradiation or surgery after initial chemotherapy, and other studies also reported similar data. In the present study, all but one LTS of more than 2 years received multimodality therapies, and two potential cured patients, who survived for more than 8 years, received surgery for the primary site.

There have been many studies about local treatment for metastatic NSCLC, and among them, treatment for brain metastasis have been most frequently reported. Local-control therapy using surgery or stereotactic radiosurgery (SRS) for the patients with solitary brain metastasis have a positive impact on prolonging survival [15,16]. The ASCO treatment guidelines state that in patients with controlled disease outside of the brain who have an isolated cerebral metastasis in a resectable area, a resection followed by whole brain radiotherapy is superior to whole brain radiotherapy alone

[17]. Concerning other metastatic sites, while no sufficient evidence to support a routine resection of the metastasis exists, there have been small series indicating an improvement in the long-term survival rates of patients after a surgical resection of solitary metastases of the adrenal gland and other sites [18,19]. In the present paper, the specific site of the metastasis had no significant influence on the LTS; however, an early N status, good performance status, single site of distant metastasis, and surgical treatment as initial treatment were all found to be significant factors in the long-term survival. When surgery is considered for stage IV patients, those with a good performance status and a controlled metastatic site are selected in clinical practice. Therefore, it is compatible that, in a multivariate analysis, surgical treatment, and an early N status proved to be significant prognostic factors for LTS.

Among the surgically treated patients in our series, patients with $N \leq 1$ disease had a better survival than those with $N \geq 2$ diseases. Two studies demonstrated that an N0 status was an important prognostic factor for the surgical treatment of primary lung cancer with synchronous brain metastases [20,21]. These data suggest that patients with an early N status may have an advantage for surgical treatment of stage IV NSCLC.

5. Conclusion

Adequate treatment including multimodality strategies, therefore, plays an important role in the long-term survival of stage IV NSCLC, and an aggressive treatment strategy including surgery might thus be considered for selected patients with an early N status.

Acknowledgement

We thank Dr. Brian T. Quinn for critical comments on the manuscript and Ms. Yumiko Oshima for her helps in preparing the manuscript.

References

- [1] Pass H, Mitchell J, Johnson D, Turrisi A, Minna J. Lung cancer; principles and practice. Philadelphia: Lipponcott Williams & Wilkins; 2000.
- [2] Socinski MA, Morris DE, Masters GA, Lilenbaum R. Chemotherapeutic management of stage IV non-small cell lung cancer. *Chest* 2003;123:2265–435.
- [3] The Non-small Cell Lung Cancer Collaborative Group. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. *Br Med J* 1995;311:899–909.
- [4] Souquet PJ, Chauvin F, Boissel JP, Cellerino R, Cormier Y, Ganz PA, et al. Polychemotherapy in advanced non small cell lung cancer: a meta-analysis. *Lancet* 1993;342:19–21.
- [5] Wozniak AJ, Crowley JJ, Balcerzak SP, Weiss GR, Spiridonidis CH, Baker LH, et al. Randomized trial comparing cisplatin with cisplatin plus vinorelbine in the treatment of advanced non-small-cell lung cancer: a Southwest Oncology Group study. *J Clin Oncol* 1998;16:2459–65.
- [6] Bonomi P, Kim K, Fairclough D, Cella D, Kugler J, Rowinsky E, et al. Comparison of survival and quality of life in advanced non-small-cell lung cancer patients treated with two dose levels of paclitaxel combined with cisplatin versus etoposide with cisplatin: results of an Eastern Cooperative Oncology Group trial. *J Clin Oncol* 2000;18:623–31.
- [7] Finkelstein DM, Ettinger DS, Ruckdeschel JC. Long-term survivors in metastatic non-small-cell lung cancer: an Eastern Cooperative Oncology Group Study. *J Clin Oncol* 1986;4:702–9.
- [8] Albain KS, Crowley JJ, LeBlanc M, Livingston RB. Survival determinants in extensive-stage non-small-cell lung cancer: the Southwest Oncology Group experience. *J Clin Oncol* 1991;9:1618–26.
- [9] Sculier JP, Paesmans M, Libert P, Bureau G, Dabouis G, Thiriaux J, et al. European Lung Cancer Working Party. Long-term survival after chemotherapy containing platinum derivatives in patients with advanced unresectable non-small cell lung cancer. *Eur J Cancer* 1994;30A:1342–7.
- [10] Satoh H, Ishikawa H, Yamashita YT, Naito T, Takahashi H, Kamma H, et al. Analysis of long-term survivors after platinum containing chemotherapy in advanced non-small cell lung cancer. *Anticancer Res* 1998;18:1295–8.
- [11] Chevalier T, Brisgand D, Soria JC, Douillard JY, Pujol JL, Ruffie P, et al. Long term analysis of survival in the European randomized trial comparing vinorelbine/cisplatin to vindesine/cisplatin and vinorelbine alone in advanced non-small cell lung cancer. *Oncologist* 2001;6(Suppl 1):8–11.
- [12] Travis WD, Colby TV, Corrin B, Shimosato Y, Brambilla E. Histological typing of lung and pleural tumors, 3rd ed. World Health Organization international histological classification of tumors. Berlin: Springer Verlag; 1999.
- [13] Cancer IUA. TNM classification of malignant tumors. 5th ed. New York: Wiley-Liss; 1997.
- [14] Kaplan E, Meier P. Nonparametric estimation of incomplete observations. *J Am Stat Assoc* 1958;53:457–81.
- [15] Patchell RA, Tibbs PA, Walsh JW, Dempsey RJ, Maruyama Y, Kryscio RJ, et al. A randomized trial of surgery in the treatment of single metastases to the brain. *N Engl J Med* 1990;322:494–500.
- [16] Kondziolka D, Patel A, Lunsford LD, Kassam A, Flickinger JC. Stereotactic radiosurgery plus whole brain radiotherapy versus radiotherapy alone for patients with multiple brain metastases. *Int J Radiat Oncol Biol Phys* 1999;45:427–34.
- [17] Pfister DG, Johnson DH, Azzoli CG, Sause W, Smith TJ, Baker Jr S, et al. American Society of Clinical Oncology treatment of unresectable non-small-cell lung cancer guideline: update 2003. *J Clin Oncol* 2004;22:330–53.
- [18] Luketich JD, Burt ME. Does resection of adrenal metastases from non-small cell lung cancer improve survival? *Ann Thorac Surg* 1996;62:1614–6.
- [19] Schuchert MJ, Luketich JD. Solitary sites of metastatic disease in non-small cell lung cancer. *Curr Treat Options Oncol* 2003;4:65–79.

[20] Mussi A, Pistoiesi M, Lucchi M, Janni A, Chella A, Parenti G, et al. Resection of single brain metastasis in non-small-cell lung cancer: prognostic factors. *J Thorac Cardiovasc Surg* 1996;112:146–53.

[21] Billing PS, Miller DL, Allen MS, Deschamps C, Trastek VF, Pairolero PC. Surgical treatment of primary lung cancer with synchronous brain metastases. *J Thorac Cardiovasc Surg* 2001;122:548–53.

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Triplet Chemotherapy with Cisplatin, Gemcitabine and Vinorelbine for Malignant Pleural Mesothelioma

Riichiroh Maruyama, Fumihiro Shoji, Tatsuro Okamoto, Tetsuya Miyamoto, Tetsuro Miyake, Tomomi Nakamura, Jiro Ikeda, Yoshiro Aoki, Hiroshi Wataya, Hiroshi Asoh and Yukito Ichinose

Department of Thoracic Oncology, Kyushu Cancer Center, Fukuoka, Japan

Received February 22, 2005; accepted June 1, 2005; published online July 8, 2005

Background: The incidence of malignant pleural mesothelioma (MPM) is expected to increase due to delayed control of occupational exposure to asbestos in Japan. We investigated the use of triplet combination chemotherapy with cisplatin (CDDP), gemcitabine (GEM) and vinorelbine (VNR) for the treatment of Japanese patients with MPM.

Methods: From December 2000 to August 2003, 12 patients received the following regimen: CDDP 40 mg/m², GEM 800 mg/m² and VNR 20 mg/m² on days 1 and 8 every 4 weeks. Among the 12 patients, six selected patients underwent an extrapleural pneumonectomy (EP) after a median of three cycles of triplet chemotherapy.

Results: The overall response rate for all patients and the response rate for chemotherapy-naïve cases were 58 and 67%, respectively. The median survival time and survival rate at 2 years for all patients were 11 months and 50%, respectively. The 2-year survival rates for the patients with and without EP were 83.3 and 16.7%, respectively.

Conclusions: Triplet chemotherapy with CDDP, GEM and VNR was thus found to be highly effective for patients with MPM and its toxicity was manageable. A multi-institutional phase II trial is now being planned to establish the effectiveness of this new regimen in chemotherapy-naïve patients with MPM.

Key words: malignant pleural mesothelioma – triplet chemotherapy – extrapleural pneumonectomy

INTRODUCTION

Malignant pleural mesothelioma (MPM) is a relatively rare neoplasm in Japan. The number of deaths and the proportional mortality rates from this disease in 2001 were 722 and 0.26%, respectively (1). Epidemiological studies have shown that the incidence of malignant mesothelioma is increasing worldwide, due to the occupational exposure to asbestos (2,3). While this disease may have already peaked in the USA because of the earlier control of asbestos use (4), a continuing increase of malignant mesothelioma has been shown in Japan based on the analysis of registered autopsy cases (5). The natural history is characterized by a median survival of 9–14 months, with <5% 5-year survivors. The objective response rates of 16–48% and median survivals of 9.4–11.2 months have been reported with the combination of cisplatin (CDDP) and gemcitabine (GEM) in malignant mesothelioma (6–8). GEM has shown a synergistic effect with such drugs as CDDP, vinorelbine (VNR), ifosfamide and mitomycin (9). VNR is a semi-synthetic derivative of vinblastine which is structurally modified in the catharanthine nucleus. Recently, an objective response rate of

24% has been reported with the single agent VNR in a single institution study (10). Therefore, the triplet combination chemotherapy using CDDP, GEM and VNR seems to have a potential anti-tumor activity against MPM. The Southern Italy Cooperative Oncology Group has already investigated the combination of CDDP, GEM and VNR in patients with advanced non-small cell lung carcinoma (NSCLC) as a three-armed randomized phase III trial comparing this triplet regimen with either CDDP and VNR or CDDP and GEM (11). The CDDP–GEM–VNR triplet combination produced a highly significant survival gain when compared with the CDDP–VNR doublet treatment. According to their schedule, the patients received CDDP 50 mg/m², GEM 1000 mg/m² and VNR 25 mg/m² intravenously on days 1 and 8 every 3 weeks. Under this regimen, however, Hesketh and associates observed a high rate of severe hematological toxicity, especially febrile neutropenia, despite a low delivered dose intensity of CDDP–GEM–VNR (12). Therefore, the initial doses of each drug were reduced to 80% of their planned dose in this study.

PATIENTS AND METHODS

PATIENTS

From December 2000 to August 2003, 12 patients with histologically proven MPM were enrolled into a single-center

For reprints and all correspondence: Dr Riichiroh Maruyama, Department of Thoracic Oncology, Kyushu Cancer Center, 3-1-1, Notame, Minami-ku, Fukuoka 811-1395, Japan. E-mail: rmaruyama@nk-cc.go.jp

study of CDDP–GEM–VNR triplet combination. The study had appropriate institutional ethical review board approval, and all patients provided their written, informed consent according to institutional guidelines. All patients were required to be age ≤ 75 years old, with an Eastern Cooperative Oncology Group (ECOG) performance status (PS) of 0–1 and with no uncontrolled cardiac or hepatic diseases. All patients had an adequate bone marrow function [defined as total leukocyte count $\geq 4 \times 10^9/l$, absolute neutrophil count (ANC) $\geq 2 \times 10^9/l$, platelet count $\geq 100 \times 10^9/l$ and hemoglobin level ≥ 9.5 g/dl]; adequate renal function (defined as serum creatinine level \leq upper normal limit for the laboratory, or creatinine clearance ≥ 60 ml/min); and an adequate hepatic function (defined as total bilirubin level ≤ 1.5 times the upper limit of normal and serum AST and/or ALT and alkaline phosphatase levels ≤ 2 times the upper normal limit for the laboratory). The clinical or pathological stage of the disease was based on the International Mesothelioma Interest Group (IMIG) staging system (13). The histological analysis of the tumor was based on the WHO classification for cell types (14). The toxicity criteria were based on the National Cancer Institute–Common Toxicity Criteria (NCI-CTC), version 2.0 (<http://ctep.info.nih.gov/>). The clinico-pathological characteristics of the patients are shown in Table 1. The doublet regimens (CDDP plus GEM or carboplatin plus etoposide) were performed in two patients as first-line chemotherapy.

TREATMENT SCHEDULE

The patients received the following regimen: CDDP 40 mg/m², GEM 800 mg/m² and VNR 20 mg/m² on days 1 and 8 every 4 weeks. In the absence of either disease progression or unacceptable toxicity levels, the patients were scheduled to receive the treatment for three cycles (maximum of six cycles). If grade 3 or more leukopenia and neutropenia, grade 2 or more thrombocytopenia, or grade 2 or more non-hematological toxicities occurred on day 8, the treatment on that day was skipped. If the total leukocyte count and ANC were $\geq 3 \times 10^9/l$ and $\geq 1.5 \times 10^9/l$, respectively and if the other eligibility criteria were satisfied, then the patients could receive the next cycle. If these toxicities persisted after 6 weeks from day 1 of the previous cycle, then the treatment regimen was discontinued. The blood counts and chemistries were examined at least once a week. Patients should not receive prophylactic granulocyte colony-stimulating factor (G-CSF) during any cycle. G-CSF may be used only for patients who have ANC $< 0.5 \times 10^9/l$, neutropenic fever or documented infections while neutropenic.

DOSE ADJUSTMENTS

As shown in Fig. 1, the doses of each drug were reduced by 25% in patients in whom grade 4 hematological toxicities or grade 3 or greater non-hematological toxicities (excluding nausea and vomiting) occurred, or in whom the scheduled treatment was skipped on day 8 in the previous cycle.

Table 1. Clinico-pathological characteristics of the patients

Parameter	No.	%
Median age, years (range)	53 (34–67)	
Gender		
Male	11	92
Female	1	8
Performance status (ECOG)		
0	8	67
1	4	33
Histological type		
Epithelioid	7	58
Sarcomatoid	1	8
Biphasic	3	25
Unknown	1	8
IMIG stage		
I	6	50
II	1	8
III	3	25
IV	2	17
Prior treatments		
Chemotherapy	2	17
Radiotherapy	1	8
None	9	75
Exposure to asbestos		
Yes	7	58
No	5	42

ECOG, Eastern Cooperative Oncology Group.
IMIG, International Mesothelioma Interest Group.




Figure 1. A schematic drawing of the dose modification in this study.

TUMOR ASSESSMENT DURING AND AFTER TREATMENT

The change in disease was assessed by measuring the thickness of the circumferential pleural tumor at three separate levels on transverse sections on the computed tomography (CT) findings of the chest, at baseline and at every other cycle (6). The sum of the measurements of tumor thickness at the three levels defined the unidimensional size. The measurability of target lesions at

baseline and the response criteria were based on the Response Evaluation Criteria in Solid Tumours (RECIST) (15). In brief, lesions that can be accurately measured in at least one dimension ≥ 20 mm with conventional techniques or as ≥ 10 mm with a spiral CT scan were defined as measurable lesions. The response criteria will be categorized as follows: complete response (CR), the disappearance of all target lesions; a partial response (PR), at least a 30% decrease in the sum of the pleural thickness at three separate levels; progressive disease (PD), at least a 20% increase in the sum of the pleural thickness at three separate levels or the appearance of one or more new lesions; and stable disease (SD), neither sufficient shrinkage to qualify for PR nor a sufficient increase to qualify for PD.

EXTRAPLEURAL PNEUMONECTOMY AFTER CHEMOTHERAPY

In addition to the chemotherapy inclusion criteria as mentioned above, patients were also candidates to undergo an extrapleural pneumonectomy (EP) if the following conditions were all satisfied: (i) age of 65 years or younger; (ii) tumor confined to stage I or II or with part of the tumor at stage III according to the IMIG staging system (13); (iii) EP would allow a complete resection of all gross disease; (iv) the room air oxygen partial pressure was >70 mmHg; and (v) a post-resection forced expiratory volume of 1 s ($FEV_{1.0}$) >600 ml per body surface area was predicted.

END-POINTS AND SAMPLE SIZE CONSIDERATION

The primary end-point of this study was the response rate of MPM of those who underwent this treatment. The secondary end-points were overall survival, toxicity, morbidity and mortality after surgery in the patients who underwent EP. The sample size was calculated based on an expected response rate of 54% and an acceptable lowest rate of 20%, with α and β errors of 0.05 and 0.2, respectively; a total of 11 patients were required using the one-sample multiple testing procedure of Fleming (16). In this design, when the number of responses exceeds four of 11 cases, it leads to the rejection of the hypothesis that the true response rate is $<20\%$. The accrual period and follow-up after accrual closure were 3 and 2 years, respectively.

STATISTICAL ANALYSIS

The duration of response was measured from the first day of a PR to the first date of progression or death due to any causes. The survival was calculated from the date of first chemotherapy dose until death due to any cause or the last follow-up (censored). The survival curve was produced using the Kaplan-Meier method (17). All data were analyzed using Abacus Concepts, Survival Tools for StatView (Abacus Concepts, Inc., Berkeley, CA).

RESULTS

TREATMENT TOXICITY AND RESPONSE

A total of 35 cycles of treatment (28 cycles with full dose and seven cycles with one level dose reduction) were delivered to the 12 patients. All cycles with a dose reduction were due to myelosuppression. The median number of cycles per patient was three. Out of 35 cycles, the frequency of treatment skipping on day 8 was only three cycles (myelotoxicity in one cycle and patient's refusal in two). As a consequence of delay and/or dose reductions, the median relative dose intensity (actually delivered $\text{mg}/\text{m}^2/\text{week}$ divided by the planned $\text{mg}/\text{m}^2/\text{week}$) of all drugs was 91%. Except for six candidates who were indicated to undergo EP, the reasons for discontinuing the chemotherapy were progression after SD in two patients, progression after PR, PD, patient's refusal and transition to other treatments in one case each. This regimen was associated with a manageable toxicity. No toxic deaths occurred. Grade 3-4 leukopenia, neutropenia, anemia and thrombocytopenia occurred in 50, 92, 33 and 17%, respectively (Table 2). According to the G-CSF dose criteria, eight patients received G-CSF out of 12 patients during 10 cycles out of 35. The median duration of grade 4 neutropenia was 3 days (range: 2-7 days). Severe non-hematological toxicity was uncommon. The overall response rate for all patients and that only for chemotherapy-naïve cases were 58 and 67%, respectively (Table 3). The median duration of response was 6 months (range, 2-19 months). Among the 12 patients, six patients were eligible for EP after a median of three cycles of triplet chemotherapy (one CR, four PR and one SD). All six were

Table 2. Hematological and non-hematological toxicity

NCI-CTC grade	0 (%)	1 (%)	2 (%)	3 (%)	4 (%)	3/4 (%)
Leukopenia	0	1 (8)	5 (42)	3 (25)	3 (25)	6 (50)
Neutropenia	0	1 (8)	0	5 (42)	6 (50)	11 (92)
Hb	0	2 (17)	6 (50)	3 (25)	1 (8)	4 (33)
PLT	4 (33)	4 (33)	2 (17)	2 (17)	0	2 (17)
AST or/and ALT	9 (75)	1 (8)	2 (17)	0	0	0

NCI-CTC, National Cancer Institute Common Toxicity Criteria; Hb, hemoglobin; PLT, platelets; AST, aspartate aminotransferase; ALT, alanine aminotransferase.

Table 3. Objective tumor response

	All cases (n = 12)	Chemotherapy-naïve cases (n = 9)
Complete response	1	1
Partial response	6	5
Stable disease	4	2
Progressive disease	1	1
Response rate (%)	58	67
95% confidence interval	30-86	36-98




Figure 2. Overall survival in all patients.

males with a median age of 50 years (range 34–54). Three patients had stage I, one stage II and two stage III disease. The histological types were epithelioid type in four and biphasic type in two. The median duration of hospitalization after EP was 33 days (range 19–63). There were neither deaths nor major morbidity during the perioperative period except thoracic empyema in one patient. Although there were no pathological CRs (Ef3: no viable tumor cells in resected specimens), some pathological response was recognized in all six patients (Ef1a, $\geq 2/3$ viable tumor cells, in two patients; Ef1b, $1/3 \leq$ viable tumor cells $< 2/3$, in two; and Ef2, $< 1/3$ viable tumor cells, in two).

SURVIVAL

The median follow-up duration of the patients was 13 months (range 4–43). The median survival time was 11 months for all patients and 6 months for the patients who did not undergo EP. The survival of the patients who underwent EP has not yet achieved a median value. The overall survival rate at 2 years was 50% (Fig. 2). The 2-year survival rates for the patients with and without EP were 83.3 and 16.7%, respectively. Among the six patients who underwent EP, the sites of first recurrence were local in three and liver metastasis in one.

DISCUSSION

Triplet chemotherapy with CDDP, GEM and VNR used in this trial produced a 58% objective response rate in patients with MPM. Selected patients underwent EP after a median of three cycles of chemotherapy without any perioperative deaths; this triplet chemotherapy was thus found to be feasible for the induction setting. Ichinose et al. presented a multicenter phase II trial using 80% of the dose of the previous CDDP–GEM–VNR triplet regimen in 80 patients with advanced NSCLC (18). The predominant toxicity was hematological: grade 3–4 neutropenia and thrombocytopenia occurred in 84 and 44% of the patients, respectively. The frequency of grade 3–4 non-hematological toxicity was low ($< 5\%$). Sixty-three percent of the enrolled patients completed > 4 cycles in this

trial. As a consequence of delay and/or dose reductions, the median relative dose intensity (actually delivered $\text{mg}/\text{m}^2/\text{week}$ divided by the planned $\text{mg}/\text{m}^2/\text{week}$) of all drugs was $> 82\%$. The triplet chemotherapy administered at a lower dose demonstrated a sufficiently effective activity with feasibility and tolerable toxicity in Japanese patients. In this study, the response rate only for chemotherapy-naïve cases was 67%. The median survival time and survival rate at 2 years for all patients were 11 months and 50%, respectively. Our results indicate that a CDDP–GEM–VNR triplet regimen is a promising treatment for MPM.

EP involves the removal of the complete pleural envelope and all of its contents, including the ipsilateral lung, diaphragm and a portion of the pericardium. The peri-operative mortality and morbidity rates of this procedure were high because of the operative invasiveness, but with more experience and better pre-operative management, the peri-operative mortality is decreasing. In the largest series of EP for MPM reported by Sugarbaker et al., the peri-operative mortality rate and some morbidity, including atrial and ventricular arrhythmias, the most common minor morbidity, were 3.8 and 50%, respectively (19). A recent publication from a Swiss group demonstrated that neoadjuvant chemotherapy consisting of CDDP and GEM followed by EP in MPM had no peri-operative mortality (20). EP can therefore be safely performed after neoadjuvant chemotherapy in an experienced center. Rusch and associates reported that hemithoracic radiation after a complete surgical resection dramatically reduced local recurrence and was associated with a prolonged survival for early-stage MPM (21). Adjuvant radiation administered to 57 patients (54 undergoing EP and three undergoing pleurectomy/decortication) at a median dose of 54 Gy was well tolerated (grade 0–2 fatigue, esophagitis), except for one late esophageal fistula. In our trial, local recurrence was the most common form of relapse in the six patients who underwent EP. Therefore, hemithoracic radiation after EP may be considered for local control with acceptable toxicities. In their report, the initial site of relapse in the patients, especially in those with stage III disease, who underwent EP with post-operative radiation was mainly distant metastases (21). However, previous experience with post-operative chemotherapy after EP does not suggest a marked survival benefit in the patients with stage III tumors (19). Combination chemotherapies in series with > 15 patients since 1990 are shown in Table 4. Reported combinations do not consistently appear to provide satisfactory results. A treatment regimen with pemetrexed (Alimta) plus CDDP and vitamin supplementation was recently reported to result in a superior survival time, time to progression and response rate in comparison with treatment with CDDP alone in patients with MPM in a randomized phase III trial (22). The response rates were 41.3% in the pemetrexed plus CDDP arm versus 16.7% in the CDDP alone arm. The median survival time for patients treated with pemetrexed plus CDDP was longer than that for patients receiving CDDP alone: 12.1 versus 9.3 months, thus indicating a highly statistically significant difference ($P = 0.020$, two-sided log-rank test).

Table 4. Reported combinations in series with more than 15 patients

Regimen	n	RR (%)	MST (months)
Doxorubicin/CDDP (23,24)	24–35	14–25	9–10
Doxorubicin/ifosfamide (25)	22	32	7
Cyclophosphamide/doxorubicin/CDDP (26)	23	30	14
CDDP/mitomycin (24)	35	26	8
CDDP/methotrexate/vinblastine (27)	17	53	14
CDDP/irinotecan (28)	15	27	7
CDDP/GEM (6–8)	21–52	16–48	9–11
CBDCA/pemetrexed (29)	25	32	15
CBDCA/GEM (30)	50	26	15
Oxaliplatin/GEM (31)	25	40	13
Mitomycin/methotrexate/mitoxantrone (32)	22	32	14

RR, response rate; MST, median survival time; CDDP, cisplatin; GEM, gemcitabine; CBDCA, carboplatin.

A phase I/II trial is now underway using this doublet regimen in Japan; therefore, the effectiveness and toxicity for Japanese patients with MPM is uncertain. In addition to local therapy, induction systemic chemotherapy using so-called new drugs, including CDDP–GEM–VNR triplet combination or CDDP plus the pemetrexed doublet, might produce a prolonged survival in patients with MPM, a disease for which standard treatment remains to be established.

In conclusion, CDDP–GEM–VNR combination is feasible at doses which, for the respective drugs, have a proven therapeutic effect in MPM patients. This combination is selectively manageable for induction chemotherapy followed by EP. A multi-institutional phase II trial is now being planned to establish the effectiveness of this new triplet regimen in chemotherapy-naïve patients with MPM as an intergroup study of the Japan Clinical Oncology Group and the West Japan Thoracic Oncology Group.

Acknowledgments

We thank Dr Brian T. Quinn for critical comments on the manuscript, and Miss Yumiko Oshima for reviewing the patient charts.

References

1. Statistics and Information Department MsS, Ministry of Health, Labour and Welfare of Japan. Vital Statistics of Japan 2001. Tokyo: Health and Welfare Statistics Association;2001.
2. Peto J, Hodgson JT, Matthews FE, Jones JR. Continuing increase in mesothelioma mortality in Britain. *Lancet* 1995;345:535–9.
3. Peto J, Decarli A, La Vecchia C, Levi F, Negri E. The European mesothelioma epidemic. *Br J Cancer* 1999;79:666–72.
4. Price B. Analysis of current trends in United States mesothelioma incidence. *Am J Epidemiol* 1997;145:211–8.
5. Murai Y. Malignant mesothelioma in Japan: analysis of registered autopsy cases. *Arch Environ Health* 2001;56:84–8.
6. Byrne MJ, Davidson JA, Musk AW, Dewar J, van Hazel G, Buck M, et al. Cisplatin and gemcitabine treatment for malignant mesothelioma: a phase II study. *J Clin Oncol* 1999;17:25–30.
7. Nowak AK, Byrne MJ, Williamson R, Ryan G, Segal A, Fielding D, et al. A multicentre phase II study of cisplatin and gemcitabine for malignant mesothelioma. *Br J Cancer* 2002;87:491–6.
8. van Haarst JM, Baas P, Manegold C, Schouwink JH, Burgers JA, de Bruin HG, et al. Multicentre phase II study of gemcitabine and cisplatin in malignant pleural mesothelioma. *Br J Cancer* 2002;86:342–5.
9. Peters GJ, Bergman AM, Ruiz van Haperen VW, Veerman G, Kuiper CM, Braakhuys BJ, et al. Interaction between cisplatin and gemcitabine *in vitro* and *in vivo*. *Semin Oncol* 1995;22:72–9.
10. Steele JP, Shamash J, Evans MT, Gowar NH, Tischkowitz MD, Rudd RM, et al. Phase II study of vinorelbine in patients with malignant pleural mesothelioma. *J Clin Oncol* 2000;18:3912–7.
11. Comella P, Frasci G, Panza N, Manzione L, De Cataldis G, Cioffi R, et al. Randomized trial comparing cisplatin, gemcitabine, and vinorelbine with either cisplatin and gemcitabine or cisplatin and vinorelbine in advanced non-small-cell lung cancer: interim analysis of a phase III trial of the Southern Italy Cooperative Oncology Group. *J Clin Oncol* 2000;18:1451–7.
12. Hesketh PJ, Nauman CJ, Hesketh AM, LaPointe J, Fogarty K, Oo TH, et al. Unfavorable therapeutic index of cisplatin/gemcitabine/vinorelbine in advanced non-small-cell lung cancer. *Clin Lung Cancer* 2002;4:47–51.
13. Rusch VW. A proposed new international TNM staging system for malignant pleural mesothelioma. From the International Mesothelioma Interest Group. *Chest* 1995;108:1122–8.
14. Travis WD CT, Corrin B, Shimosato Y, Brambilla E, and Collaborators from 14 Countries. World Health Organization. International Histological Classification of Tumors. *Histological Typing of Lung and Pleural Tumors*, 3rd edn. Berlin: Springer Verlag;1999.
15. Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000;92:205–16.
16. Fleming TR. One-sample multiple testing procedure for phase II clinical trials. *Biometrics* 1982;38:143–51.
17. Kaplan E, Meier P. Nonparametric estimation from incomplete observation. *J Am Stat Assoc* 1958;53:457–81.
18. Ichinose Y, Itoh K, Imamura F, Araki J, Matsumoto M, Yamamoto H, et al. Triplet combination chemotherapy of cisplatin (P), gemcitabine (G) and vinorelbine (V) with lower doses maintains a sufficient effect while avoiding toxicity in non-small cell lung cancer (NSCLC). *Proc Am Soc Clin Oncol* 2004;23.
19. Sugarbaker DJ, Flores RM, Jaklitsch MT, Richards WG, Strauss GM, Corson JM, et al. Resection margins, extrapleural nodal status, and cell type determine postoperative long-term survival in trimodality therapy of malignant pleural mesothelioma: results in 183 patients. *J Thorac Cardiovasc Surg* 1999;117:54–63;discussion 63–5.
20. Weder W, Kestenholz P, Taverna C, Bodis S, Lardinois D, Jerman M, et al. Neoadjuvant chemotherapy followed by extrapleural pneumonectomy in malignant pleural mesothelioma. *J Clin Oncol* 2004;22:3451–7.
21. Rusch VW, Rosenzweig K, Venkatraman E, Leon L, Raben A, Harrison L, et al. A phase II trial of surgical resection and adjuvant high-dose hemithoracic radiation for malignant pleural mesothelioma. *J Thorac Cardiovasc Surg* 2001;122:788–95.
22. Vogelzang NJ, Rusthoven JJ, Symanowski J, Denham C, Kaukel E, Ruffie P, et al. Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. *J Clin Oncol* 2003;21:2636–44.
23. Ardizzone A, Rosso R, Salvati F, Fusco V, Cinquegrana A, De Palma M, et al. Activity of doxorubicin and cisplatin combination chemotherapy in patients with diffuse malignant pleural mesothelioma. An Italian Lung Cancer Task Force (FONICAP) phase II study. *Cancer* 1991;67:2984–7.
24. Chahinian AP, Antman K, Goutsou M, Corson JM, Suzuki Y, Modeas C, et al. Randomized phase II trial of cisplatin with mitomycin or doxorubicin for malignant mesothelioma by the Cancer and Leukemia Group B. *J Clin Oncol* 1993;11:1559–65.
25. Dirix LY, van Meerbeeck J, Schrijvers D, Corthouts B, Prove A, van Marck E, et al. A phase II trial of dose-escalated doxorubicin and ifosfamide/mesna in patients with malignant mesothelioma. *Ann Oncol* 1994;5:653–5.

26. Shin DM, Fossella FV, Umsawasdi T, Murphy WK, Chasen MH, Walsh G, et al. Prospective study of combination chemotherapy with cyclophosphamide, doxorubicin, and cisplatin for unresectable or metastatic malignant pleural mesothelioma. *Cancer* 1995;76:2230–6.
27. Hunt KJ, Longton G, Williams MA, Livingston RB. Treatment of malignant mesothelioma with methotrexate and vinblastine, with or without platinum chemotherapy. *Chest* 1996;109:1239–42.
28. Nakano T, Chahinian AP, Shinjo M, Togawa N, Tonomura A, Miyake M, et al. Cisplatin in combination with irinotecan in the treatment of patients with malignant pleural mesothelioma: a pilot phase II clinical trial and pharmacokinetic profile. *Cancer* 1999;85:2375–84.
29. Hughes A, Calvert P, Azzabi A, Plummer R, Johnson R, Rusthoven J, et al. Phase I clinical and pharmacokinetic study of pemetrexed and carboplatin in patients with malignant pleural mesothelioma. *J Clin Oncol* 2002;20:3533–44.
30. Favaretto AG, Aversa SM, Paccagnella A, Manzini Vde P, Palmisano V, Oniga F, et al. Gemcitabine combined with carboplatin in patients with malignant pleural mesothelioma: a multicentric phase II study. *Cancer* 2003;97:2791–7.
31. Schutte W, Blankenburg T, Lauerwald K, Schreiber J, Bork I, Wollscgkaeger B, et al. A multicenter phase II study of gemcitabine and oxaliplatin for malignant pleural mesothelioma. *Clin Lung Cancer* 2003;4:294–7.
32. Pinto C, Marino A, Guaraldi M, Melotti B, Piana E, Martoni A, et al. Combination chemotherapy with mitoxantrone, methotrexate, and mitomycin (MMM regimen) in malignant pleural mesothelioma: a phase II study. *Am J Clin Oncol* 2001;24:143–7.

CASE REPORT

Intrathoracic Omental Herniation through the Esophageal Hiatus in a Young Patient

We herein present a case of intrathoracic omental herniation through the esophageal hiatus in a young patient. A 21-year-old obese man was asymptomatic, and his chest X-ray demonstrated a large, sharply defined mass. A computed tomography scan of the thorax indicated a large retrocardial mediastinal mass in which the density indicated the presence of fatty tissue judging from the Hounsfield unit range. A thoracotomy was performed under a diagnosis of either mediastinal lipoma or liposarcoma with an encapsulated fatty mass, measuring 17×12×8 cm in size. The mass, however, proved to be an omental herniation through the esophageal hiatus. It is generally assumed that the major contributing factors leading an individual to develop an omental herniation through the esophageal hiatus include aging and obesity. This is the first report of omental herniation through the esophageal hiatus in a patient still in his twenties. (*Jpn J Thorac Cardiovasc Surg* 2005; 53: 452–454)

Key words: omentum, hernia, esophageal hiatus, young patient

Riichiro Maruyama, MD, Tetsuya Miyamoto, MD, Fumihiko Shoji, MD,
Tatsuro Okamoto, MD, Tetsuro Miyake, MD,
and Yukito Ichinose, MD.

Omental herniation through the esophageal hiatus, without stomach involvement, is rare. Omental herniation has been previously misdiagnosed as a lipomatous tumor.¹ To the best of our knowledge, there have been only eight reports on omental herniation through the esophageal hiatus in the Japanese and English literatures within last three decades.^{1–8} In addition, there have also been no reports on young patients with this problem up to now.

Case

A 21-year-old obese man (height: 174 cm, weight: 90 kg, body mass index: 29.7) was admitted to our department for an investigation of a chest roentgenographic abnormality. He was asymptomatic, and his physical examination and laboratory data were all within the normal limits. A postero-anterior view of his chest X-ray demonstrated a large, sharply defined mass. The mass

measured over 15 cm in diameter with a smooth outline, it did not demonstrate either calcification or cavitation, and it extended from the retrocardial mediastinum to the diaphragm level (Fig. 1A). A computed tomography (CT) scan of the thorax showed the presence of a mass, in which the density indicated the presence of fatty tissue according to the Hounsfield unit (HU) range,⁹ in the lower thorax, located behind the heart, and beside the thoracic vertebrae (Fig. 1B). The HU of water and air in the lung parenchyma were 0 and –1,000, respectively. The HU of the fatty tissue are usually in the range of –50 to –200. Both coronal and sagittal sections of T1-weighted magnetic resonance imaging (MRI) clearly depicted a fatty mass on both sides of the diaphragm (Fig. 2). A thoracotomy was performed under an initial diagnosis of either mediastinal lipoma or liposarcoma which demonstrated an encapsulated fatty mass, measuring 17×12×8 cm in size, but which proved to be an omental herniation through the esophageal hiatus. The esophageal hiatus was repaired after the omental mass and hernia sac had first been resected. The patient had an uneventful postoperative recovery without any complications.

Discussion

The omentum sometimes herniates as a result of

From Department of Thoracic Oncology, Kyushu Cancer Center, Fukuoka, Japan.

Received for publication November 15, 2004.

Accepted for publication May 6, 2005.

Address for reprints: Riichiro Maruyama, MD, Department of Thoracic Oncology, Kyushu Cancer Center, 3-1-1 Notame, Minami-ku, Fukuoka 811-1395, Japan.

Eliska suggested that the phreno-esophageal membrane can act as one of the main mechanisms of competence in the cardio-esophageal junction, and thus preventing reflux and hiatal hernia.¹⁰ This membrane has two limbs, one which passes through the esophageal hiatus, and then attaches itself to the esophagus in the chest, while the other limb extends downward and attaches itself to the esophagus in the abdomen. The membrane is normally rich in elastic fibers, but in people over 50 years of age, the amount of elastic tissue tends to decrease and the esophagus becomes much more mobile within the hiatus. In obese patients, the omentum may also be more prominent than usual, thus predisposing it to herniation without any significant gastroesophageal hernia. It is generally assumed that the major contributing factors leading to an omental herniation through the esophageal hiatus include aging and obesity. Therefore, previous reports about omental herniation all described either middle-aged or elderly patients.¹⁻⁸ This is the first report of intrathoracic omental herniation through the esophageal hiatus in a patient still in his twenties. The weakness of the phreno-esophageal membrane was not recognized based on an intraoperative view. In addition to obesity, these findings indicate the possibility that some congenital anomaly may thus have existed in the esophageal hiatus in our case and an omental mass thereafter formed as the patient became older.

We thank Dr. Brian Quinn for critical comments on the manuscript and Miss Yumiko Oshima for help in preparing the manuscript.

REFERENCES

1. Anderson TM, Gibbs JF, Kollmorgen DR, Urschel JD. Paraesophageal omental hernia mimics pleural lipomatous tumor. *J Cardiovasc Surg (Torino)* 1999; 40: 757-9.
2. Rohlfing BM, Korobkin M, Hall AD. Computed tomography of intrathoracic omental herniation and other mediastinal fatty masses. *J Comput Assist Tomogr* 1977; 1: 181-3.
3. Tamura A, Murakami K, Sato K, Komatsu H, Yoneda R, Takahashi K. A case of intrathoracic omental herniation through the esophageal hiatus (Eng abstr). *Nihon Kyobu Shikkan Gakkai Zasshi* 1988; 26: 1010-4.
4. Lee MJ, Breathnach E. CT and MRI findings in paraesophageal omental herniation. *Clin Radiol* 1990; 42: 207-9.
5. Rockoff SD, Aaron BL, Black C, Kathuria R, Biben L. Diagnosis of paraesophageal omental hiatal hernia by magnetic resonance imaging. *Chest* 1993; 103: 285-7.
6. Saijo Y, Honda H, Nishigaki Y, Noro T. A case of paraesophageal omental herniation (Eng abstr). *Nihon Kokyuki Gakkai Zasshi* 1998; 36: 67-70.
7. Kato N, Iwasaki H, Rino Y, Imada T, Amano T, Kondo J. Intrathoracic omental herniation through the esophageal hiatus: Report of a case. *Surg Today* 1999; 29: 347-50.
8. Kubota K, Ohara S, Yoshida S, Nonami Y, Takahashi T. Intrathoracic omental herniation through the esophageal hiatus: A case report. *Radiat Med* 2001; 19: 307-11.
9. Hounsfield GN. Computed medical imaging. *Science* 1980; 210: 22-8.
10. Eliska O. Phreno-oesophageal membrane and its role in the development of hiatal hernia. *Acta Anat (Basel)* 1973; 86: 137-50.

UFT plus gemcitabine combination chemotherapy in patients with advanced non-small-cell lung cancer: a multi-institutional phase II trial

Y Ichinose^{*,1}, T Seto², H Semba², K Itoh³, Y Inoue⁴, F Tanaka⁵, J Araki⁶, M Tamanoi⁷, H Yamamoto⁸ and N Iwamoto⁹

¹Department of Thoracic Oncology, National Kyushu Cancer Center, 3-1-1, Notame, Minami-ku, Fukuoka 811-1395, Japan; ²Division of Respiratory Diseases, Kumamoto Regional Medical Center, Kumamoto, Japan; ³Department of Respiratory Medicine, Shin Beppu Hospital, Oita, Japan; ⁴Department of Respiratory Medicine, Isahaya Insurance General Hospital, Nagasaki, Japan; ⁵Department of Respiratory Medicine, Kumamoto City Hospital, Kumamoto, Japan; ⁶Department of Respiratory Medicine, Yamaguchi Central Hospital, Yamaguchi, Japan; ⁷Department of Respiratory Medicine, Minamata General Medical Center, Kumamoto, Japan; ⁸Department of Respiratory Medicine, Asou Iizuka Hospital, Fukuoka, Japan; ⁹Respiratory Organ and Diabetes Center, Saiseikai Kumamoto Hospital, Kumamoto, Japan

A multi-institutional phase II trial was conducted to evaluate the efficacy and toxicity of combination chemotherapy consisting of gemcitabine and UFT, which is composed of tegafur and uracil, for non-small-cell lung cancer (NSCLC) patients. Patients with advanced NSCLC received an oral administration of UFT (tegafur 200 mg m⁻²) b.i.d. from days 1 to 14 and intravenous injection of gemcitabine 900 mg m⁻² on days 8 and 15. This treatment was repeated every 4 weeks. A total of 44 patients were enrolled into this trial. The median age of all patients was 74 years, with 23 patients younger than 75 years and 21 patients with 75 years of age or older. A total of 18 patients (41%) achieved a partial response. The median survival time was 13.2 months and the 1-year survival rate was 59%. The most common grade 3–4 toxicity was neutropenia (57%). The frequency of grade 3 nonhaematologic toxicities was less than 5%. In addition, no significant difference in the response, survival or toxicities was observed between the patients younger than and those older than 75 years of age. This combination chemotherapy demonstrated a promising effectiveness and acceptable toxicity in patients with advanced NSCLC, even in patients older than 75 years.

British Journal of Cancer (2005) 93, 770–773. doi:10.1038/sj.bjc.6602781 www.bjcancer.com

Published online 20 September 2005

© 2005 Cancer Research UK

Keywords: uracil–tegafur; UFT; gemcitabine; NSCLC; elderly

UFT is an oral anticancer agent composed of tegafur and uracil at a 1:4 fixed molar ratio (Fujii *et al*, 1979). Although the clinical effectiveness of the single agent against advanced non-small-cell lung cancer (NSCLC) has not been evaluated in adequate sample size (Keicho *et al*, 1986), a recent randomised phase III trial in 984 patients with completely resected stage I adenocarcinoma demonstrated that postoperative adjuvant chemotherapy with UFT significantly prolonged the survival of patients in comparison to observation alone (Kato *et al*, 2004). The combination chemotherapy of UFT plus cisplatin has also been reported to be an effective treatment for advanced NSCLC. The combination chemotherapy consisting of a daily administration of UFT for 2 or 3 weeks and a bolus injection of cisplatin at the mid-cycle of the administration of UFT yields a response rate of 29–38%, and a median survival time of 10–13 months (Ichinose *et al*, 1995, 2000; Saito *et al*, 2001).

Gemcitabine is an active anticancer agent for the treatment of NSCLC. The objective response rates of patients with advanced NSCLC treated with gemcitabine alone and the combination chemotherapy of platinum plus gemcitabine range from 20 to 26%

and from 25 to 61%, respectively (Harper, 2003). A median survival time of 8–16 months in patients treated with the combination chemotherapy has been reported (Harper, 2003).

Both 5-FU generated from tegafur in UFT and gemcitabine are antimetabolites but inhibit DNA synthesis via a different pathway. The main mechanism of inhibiting DNA synthesis by 5-FU is due to 5-FU-derived fluorodeoxy monophosphate binding to thymidylate synthase (Madajewicz *et al*, 1984). The incorporation of gemcitabine triphosphate, which is generated from the phosphorylation of gemcitabine by deoxycytidine kinase, into DNA is most likely the major mechanism by which gemcitabine exerts its cytotoxic action (Huang *et al*, 1991). Such different antitumour mechanisms suggest a potential synergism between 5-FU and gemcitabine. This potential synergism has been indeed observed in *in vitro* studies using various cancer cell lines (Schulz *et al*, 1998; Peters *et al*, 2000).

In our prior phase I trial, the combination chemotherapy of UFT plus gemcitabine was found to be feasible (Seto *et al*, 2002). The most appropriate schedule and dosing was 200 mg m⁻² of UFT b.i.d. for 14 consecutive days with 900 mg m⁻² gemcitabine on days 8 and 15. The main toxicity was haematologic. The overall response rate was 33%, while the rate was 45% in the 13 patients without any prior chemotherapy. With these backgrounds, we conducted a phase II trial of combination chemotherapy using

*Correspondence: Dr Y Ichinose; E-mail: yichinos@nk-cc.go.jp

Received 22 March 2005; revised 10 August 2005; accepted 10 August 2005; published online 20 September 2005