

図 11 マウスエイズ進展に対するビタミンEの抑制メカニズム

ビタミンEはTNF- α の産生を抑制するだけでなく、ウイルス感染細胞内でTNF- α 刺激によって産生される活性酸素産生を直接抑制することによって、NF- κ Bを介した核内での転写を抑制し、ウイルス増殖を阻止する。

モデル動物のマウスやネコ等を用いて、抗酸化ビタミンの摂取効果が検討されており、エイズ進展に対する抗酸化ビタミン摂取の有効性が報告されている。ビタミンE補足効果に関しては、マウスのエイズモデルを用いた多くの研究成果があり、前述の転写因子NF- κ Bの発現をビタミンEが抑制することが見出されている⁹⁾(図10)。その他、ビタミンEの免疫賦活作用を加味して、エイズ進展に対するビタミンEの抑制作用が考えられている(図11)。

5. アレルギーとビタミンE

アレルギーとは免疫系が過剰に働き、結果として生体に危害を与えてしまう状態である。近年、ビタミンEが抗アレルギー効果を有することが見出され、その治療効果が期待されている。最近のアレルギーとビタミンEに関する報告を表3にまとめた^{10~16)}。ヒトおよびアレルギーモデルマウスにおいてビタミン

表3 最近のアレルギーとビタミンEに関するエビデンス

対 象	結 論	文献番号
鼻アレルギー モデルマウス (BALB/c マウス, 雌)	アレルギー症状 改善 (くしゃみ, 鼻汁など) 血清 IgE 濃度 低下 IL-2, IFN- γ 産生 変化なし IL-4, IL-5 産生 低下 高ビタミンE 摂取は鼻アレルギー反応を抑制する。	10
ヒト 2,633 名 18 ~ 70 歳	ビタミンE 摂取量と血清総 IgE 濃度には負の相関がある。 ビタミンE 摂取量とアトピーのリスク発生には負の相関がある。	11
ヒト アトピー性 皮膚炎患者 96 名 10 ~ 60 歳	血清 IgE 濃度 低下 ビタミンE 摂取群の 80%の者の症状が改善 ビタミンE 摂取はアトピー性皮膚炎の治療に対して優れた効果を有する。	12
ヒト 末梢血 T 細胞	ビタミンE はヒト末梢血 T 細胞の IL-4 タンパクレベルと mRNA における IL-4 発現を抑制する。	13
イヌの肥満細胞腫 (C2)	ビタミンE 処理によりヒスタミンおよび PGD ₂ 放出が抑制され, キマーゼ活性が低下した。	14
オボアルブミン 感作マウス (BALB/c マウス, 雌)	低ビタミンE 食群および高ビタミンE 食群において IgE 産生が抑制された。ビタミンE と β -カロテンをあわせて投与することにより, より強く IgE 産生が抑制された。	15
ヒト アレルギー性鼻炎 (花粉症) 患者 112 名 18 ~ 70 歳	ビタミンE 摂取群において症状のスコアが低下した。ビタミンE 摂取は花粉症に対して有効である可能性が示唆された。	16

E 補足によりアレルギー症状の軽減, 血清免疫グロブリン E (IgE) 濃度の低下, インターロイキン 4 (IL-4) 産生の低下などが見出されている。筆者らのオボアルブミン誘発食物アレルギーモデルマウスを用いた研究においても, 高ビタミ




図 12 食物アレルギーに対するビタミンEの発症予防効果

ビタミンEはB細胞の幼若化能を抑制するだけでなく、ヘルパーT細胞の幼若化能を抑制し、B細胞の形質細胞への分化を妨げる。これらの結果、IgE産生は抑制され、マスト細胞からのヒスタミン、LTB₄の分泌が抑制される。ビタミンEはこれらの種々の過程で抗アレルギー作用を発揮する。

ンE食投与により IgE 産生が抑制され、それがヘルパーT細胞割合の低下やヘルパーT細胞およびB細胞幼若化能の低下と関連することを認めている。さらに、アレルギー症状を直接引き起こす化学伝達物質であるヒスタミンやロイコトリエン B₄ (LTB₄) の血清濃度もまた低下することを認めている (図 12)。ビタミンEのアレルギー抑制メカニズムについては、ビタミンEがIgE産生を直接的に抑制することが見出されている。また、ヒスタミン遊離過程において活性酸素種(ROS)が必要であり、それがマスト細胞の脱顆粒を引き起こすシグナル伝達を調節していることが知られている。このことから、ビタミンEが抗酸化作用を介してROS発生を抑制することにより、ヒスタミン遊離を抑制している可能性が示唆されている。

6. ビタミンE同族体の免疫賦活作用

以上、これまで述べたビタミンEと免疫に関する研究成果のほとんど全てが α -トコフェロールを用いたものである。自然界にはこの他に β , γ および δ -トコフェロールが存在し、側鎖に2重結合を有するトコトリエノールを加えると、 α -トコフェロール以外に7種の同族体が存在する。これら同族体は食品を通じて摂取され、腸管で吸収されるが、血液および組織中には圧倒的に α -トコフェロールの存在が優位である。しかし、最近、その他の同族体が微量でより大きな生理活性を有することが証明されており、免疫分野においてもそれら同族体の免疫賦活作用が注目されている。筆者らの行ったヒト末梢血リンパ球を用いての *in vitro* 実験では、クロマン環のメチル基数が少なくなるほど濃度依存




図13 ビタミンE同族体との *in vitro* 培養に伴うヒト末梢血リンパ球幼若能の変化

ビタミンE同族体は100%エタノールに可溶化後、終濃度0.2%となるようにして種々の濃度に調整した。これらビタミンE同族体とヒト末梢血リンパ球をCon A存在下で *in vitro*, 72時間培養することによりT細胞幼若化能を測定した。

的にTリンパ球幼若化能の著しい低下を認めた(図13)。これはWuらの老齡マウス脾臓細胞を用いた実験結果¹⁷⁾と類似している。しかし、 γ -、 δ -トコフェロール、あるいはクロマン環に全くメチル基のないトコールを各々kg当たり500mg含む食餌で8週齡のラットを4週間飼育した場合には、*in vitro*実験で見られたような細胞障害作用は全くみられず、トコール群の脾臓細胞では α -トコフェロール投与群よりもさらに高いT細胞幼若化能を認めた。さらに、脾臓細胞からのIL-2産生、および脾臓細胞のIL-2に対する感受性から、これらビタミンE同族体の免疫賦活作用を見た場合、 α -トコフェロール群ではIL-2産生亢進を介してT細胞幼若化能の上昇が誘導されているのに対し、トコール群ではIL-2産生に対してはほとんど影響はみられず、むしろIL-2に対する感受性の亢進によりT細胞幼若化能が高まっていることを認めた。これらの結果は、ビタミンE同族体が同じようにT細胞機能を亢進しても、その作用機序が異なることを意味しており、これら同族体を適当な比率で混合することにより、より高い免疫賦活作用が誘導される可能性が考えられ、この分野での今後の研究が期待される。

(森口 覚, 兼安真弓)

文 献

- 1) Meydani S N, et al : Vitamin E supplementation enhances cell-mediated immunity in healthy elderly. *Am J Clin Nutr* 52 : 557-563, 1990
- 2) Sakai S, Moriguchi S : Long-term feeding of high vitamin E diet improves the decreased mitogen response in rat splenic lymphocytes with aging. *J Nutr Sci Vitaminol* 43 : 113-122, 1997
- 3) Moriguchi S, Kobayashi N, Kishino Y : Effects of vitamin E deficiency on the functions of splenic lymphocytes and alveolar macrophages. *J Nutr Sci Vitaminol* 35 : 419-430, 1989
- 4) Sharp A K, Colston M J : Elevated macrophage activity in nude mice. *Exp Cell Biol* 52 : 44-47, 1984
- 5) 渡邊陽子ほか : 加齢に伴う細胞性免疫能低下とNOに関する疫学的研究. 必須アミノ酸研究 158 : 39-44, 2000

- 6) Moriguchi S, et al : Vitamin E is an important factor in T cell differentiation in thymus of F344 rats. *J Nutr Sci Vitaminol* 39 : 451-463, 1993
- 7) Moriguchi S : The role of vitamin E in T-cell differentiation and the decrease of cellular immunity with aging. *Biofactors* 7 : 77-86, 1998
- 8) Moriguchi S, et al : Vitamin E supplementation induces an early recovery of cellular immunity decreased following x-ray irradiation. *Nutr Res* 16:645-656, 1996
- 9) Hamada M, et al : Vitamin E suppresses the development of murine AIDS through the inhibition of nuclear factor-kappa B expression. *Nutr Res* 20 : 1163-1171, 2000
- 10) Zheng K, et al : Effect of dietary vitamin E supplementation on murine nasal allergy. *Am J Med Sci* 318 : 49-54, 1999
- 11) Fogarty A, et al : Dietary vitamin E, IgE concentrations, and atopy. *Lancet*. 356 : 1573-1574, 2000
- 12) Tsourelis-Nikita E, et al : Evaluation of dietary intake of vitamin E in the treatment of atopic dermatitis: a study of the clinical course and evaluation of the immunoglobulin E serum levels. *Int Dermatol*. 41 : 146-150, 2002
- 13) Li-Weber M, et al : Vitamin E inhibits IL-4 gene expression in peripheral blood T cells. *Eur J Immunol* 32 : 2401-2408, 2002
- 14) Gueck T, et al : Influence of vitamin E on mast cell mediator release. *Vet Dermatol* 13 : 301- 305, 2002
- 15) Bando N, et al : Inhibition of immunoglobulin E production in allergic model mice by supplementation with vitamin E and beta-carotene. *Biosci. Biotechnol Biochem*. 67 : 2176-2182, 2003
- 16) Shahar E, et al : Effect of vitamin E supplementation on the regular treatment of seasonal allergic rhinitis. *Ann Allergy Asthma Immunol* 92 : 654-658, 2004
- 17) Wu D, et al : In vitro supplementation with different tocopherol homologues can affect the function of immune cells in old mice. *Free Radical Biol. Med* 28 : 643-651, 2000

Vitamin E

Guest Editor: Etsuo Niki

Role of Vitamin E in Immune System

Satoru Moriguchi* and Mayumi Kaneyasu

Department of Nutrition, Faculty of Human Life Science, Yamaguchi Prefectural University, Yamaguchi 753–8502, Japan

Received 29 August, 2003; Accepted 26 September, 2003

Summary Vitamin E is a potent antioxidant and has an ability to modulate host immune functions. This manuscript consists of six parts: (1) vitamin E deficiency and immunity, (2) vitamin E supplementation and immunity, (3) vitamin E and the decreased cellular immunity with aging, (4) vitamin E and T-cell differentiation in the thymus, (5) vitamin E and acquired immune deficiency syndrome (AIDS), and (6) vitamin E and the abnormal increase of host immune functions such as allergy and autoimmune diseases. In vitamin E deficiency most of the immune parameters show a downward trend, which is associated with increased infectious diseases and the incidence of tumors. In contrast, vitamin E supplementation has various beneficial effects on the host immune system. The decreased cellular immunity with aging or during the development of AIDS is markedly improved by the intake of a high vitamin E diet. In addition, vitamin E plays an important role in the differentiation of immature T cells in thymus. Vitamin E deficiency induces the decreased differentiation of immature T cells, which results in the early decrease of cellular immunity with aging in spontaneously hypertensive rats. Conversely, vitamin E supplementation induces a higher differentiation of immature T cells *via* increased positive selection by thymic epithelial cells, which results in the improvement of decreased cellular immunity in the aged. Furthermore, some reports have shown that vitamin E has an ability to modulate the development of allergy or autoimmune diseases such as rheumatoid arthritis and systemic lupus erythematosus. Taken together, all of the evidence suggest that vitamin E is a potent vitamin for modulating not only decreased immunity shown in the aged but also abnormally increased host immune system in patients with allergy or autoimmune diseases.

Key Words: vitamin E, aging, T-cell differentiation, AIDS, allergy

Introduction

Vitamin E, α -tocopherol, acts as an antioxidant in cellular membranes and as a free radical scavenger by blocking the peroxidation of polyunsaturated fatty acids (PUFA). The damage of cellular membrane

induced by free radicals results in changes of membrane-related cellular functions, inducing in some cases, the destruction of cells. In immune cells, α -tocopherol content is known to be higher than in other cells, because the cellular membrane plays an important role in the immune response. This suggests that vitamin E is an important nutrient in maintaining normal immune functions. Furthermore, it is estimated that one-fourth of Japan's popu-

*To whom correspondence should be addressed.

lation will be older than 65 years of age at the beginning of the twenty-first century. Accompanying this increased percentage of elderly people will be an increased incidence of geriatric disorders and infectious diseases associated with the decrease of cellular immunity that occurs during aging. If this decrease of cellular immunity can be diminished, it may improve the health state of elderly people and prolong their life expectancy. Meydani *et al.* have found that a higher intake of vitamin E is needed to maintain immune functions in the aged people [1] and improves the decreased cellular immune functions that occur with aging [2]. This finding is related to the decreases in the serum vitamin E level during aging and the actions of vitamin E as an antioxidant and immunostimulator. This review focuses on the effect of both vitamin E deficiency and supplementation on the immune responses in animals and humans, and the immunomodulating effects of vitamin E in aging. In addition, as recent topics in the research on vitamin E and the immune response, the actions of vitamin E on both T cell differentiation in the thymus and abnormal increases of immune functions such as allergy and autoimmune disease are discussed.

I. Effects of Vitamin E Deficiency and Supplementation on Immunity

Vitamin E Deficiency and Immunity

As summarized in Table 1, most studies have shown that vitamin E deficiency induces the impairment of both humoral and cellular immunity. These immunosuppressive effects of vitamin E deficiency are associated with increased production of free radicals, which results in increased lipid peroxidation in cell membranes. In fact, macrophages and neutrophils from vitamin E deficient rats had higher O_2 consumption, H_2O_2 release, and peroxidized lipids in their membranes [3, 4]. Although vitamin E deficiency appears to induce the decreased cellular immunity as described above, the phagocytic function of alveolar macrophages (AM) from rats fed a vitamin E deficient diet for 4 months was 2 times higher than that of control rats (Fig. 1) [5]. In addition, those AM did not respond to the *in vitro* treatment with macrophage-activating factor (MAF) prepared from Con A-activated rat splenic lymphocytes. Since MAF production from splenocytes activated *in vitro* with Con A-Sepharose beads decreased in vitamin E deficient rats, the higher phagocytic activity

Table 1. Vitamin E deficiency and the immune response.

Immune response	Result	Species
T-cell mitogenesis	Decreased	Humans, mice, rats, pigs, dogs
Interleukin-2 production	Decreased	Humans, rats
B-cell mitogenesis	Decreased	Mice, rats
Natural killer cell activity	Decreased	Mice, rats
Plaque-forming cell	Decreased	Mice
Antibody titer	Decreased	Mice, chickens
Macrophage phagocytosis	Decreased	Rats
	Increased	Rats
Polymorphonuclear leucocyte phagocytosis	Decreased	Humans, rats
Polymorphonuclear leucocyte chemotaxis	Decreased	Humans, rats




Fig. 1. Phagocytosis of ^{51}Cr -labeled opsonized SRBC by AM of rats fed a control diet or vitamin E-deficient diet. AM (2×10^5) were preincubated for 4 h in a medium consisting of RPMI 1640 with 5% fetal bovine serum (Med) or in a medium with macrophage-activating factor (MAF, 1/162 dilution). They were then incubated with ^{51}Cr -labeled opsonized SRBC for 2 h. Each value is the mean \pm SD for triplicate cultures. * Significantly different from cultures with medium ($p < 0.05$); ** significantly different from cultures with medium in control rat ($p < 0.01$). (Moriguchi *et al.*: *J. Nutr. Sci. Vitaminol.*, 35, 419-430, 1989)

of AM from rats fed the vitamin E deficient diet does not appear to be induced by MAF. Sharp and Colston demonstrated the presence of highly activated macrophages in nude mice and considered that hyperactivity seen in the macrophages of nude mice resulted from a lack of T cell-mediated suppression [6]. As described above, vitamin E deficiency causes immunodepression, including T and B cell responses, which may result in inducing the higher phagocytic activity of AM.




Fig. 2. Phagocytic activity of fractionated AM (A) and the proportions of fractionated AM (B) in rats fed a control or vitamin E-deficient diet. After AM were collected by tracheopulmonary lavage with warm saline, separation of the AM fraction was performed by using discontinuous Percoll gradient centrifugation. Discontinuous Percoll solutions were prepared by diluting a stock of Percoll with sterile saline to specific gravities of 1.050, 1.060, 1.070, and 1.080 and subsequently layered into centrifuge tubes. AM suspensions were carefully layered on the top of the diluted Percoll solutions and centrifuged at 400 g for 30 min. The cells localized at each interface area were then carefully collected with a Pasteur pipette. The fractions were designated I to IV in order of increasing density. In this experiment, the phagocytic activity of AM was assessed by measuring the absorbance of SRBC phagocytosed within AM. * Significantly different from other fractions ($p < 0.05$).

In a recent study, we separated AM into four fractions (I to IV) by discontinuous Percoll density-gradient centrifugation. The degree of AM maturation is high from Fraction IV to Fraction I. As shown in Fig. 2A, the phagocytic activity of AM is the highest in Fraction III. We have found that vitamin E deficiency decreased the number of AM and increased the percentage of AM in Fraction III having the

Table 2. Vitamin E supplementation and the immune response.

Immune response	Result	Species
T-cell mitogenesis	Increased	Humans, mice, rats, pigs
	No change	Humans
Interleukin-2 production	Increased	Humans, mice, rats
T-cell helper activity	Increased	Mice
B-cell mitogenesis	Increased	Mice, rats
Natural killer cell activity	Increased	Mice, rats
Plaque-forming cell	Increased	Mice, rats
Antibody titer	Increased	Mice, rats, sheep
Serum antibodies	No change	Humans
Macrophage phagocytosis	Increased	Rats
Polymorphonuclear leucocyte phagocytosis	Increased	Humans
Delayed type hypersensitivity	Increased	Humans
	No change	Humans

higher phagocytic ability against opsonized sheep red blood cells (SRBC) (Fig. 2B) [7]. These results suggest that vitamin E deficiency appears to induce the destruction of highly matured AM (Fraction I), which results in a decreased number of AM and a higher phagocytic activity of AM.

Vitamin E supplementation and immunity

Contrary to the effect of vitamin E deficiency on the immune response, most studies have shown that vitamin E supplementation enhances the immune response as summarized in Table 2. As its mechanism, Meydani *et al.* found that vitamin E supplementation suppresses Prostaglandin E₂ (PGE₂) synthesis in spleen homogenate from old mice [2] and Baehner *et al.* showed that the *in vitro* administration of vitamin E prevents autoxidative damage to the membrane of polymorphonuclear leukocytes by scavenging H₂O₂ [8]. Because both PGE₂ and H₂O₂ are potent inhibitors of several lymphocyte functions, including mitogenesis, cytolysis and antibody production [9–11], vitamin E supplementation decreases the formation of PGE₂ and H₂O₂, which may result in the enhancement of mitogenic response and natural killer cell activity in splenic lymphocytes.

Studies on the effect of vitamin E supplementation on macrophage function are exceedingly few compared with those on lymphocyte functions. In particular, studies concerning AM, which play an important role in the defense against bacterias and neoplasms in the lung, are very limited. As shown in




Fig. 3. Phagocytosis of ⁵¹Cr-labeled opsonized SRBC by AM of rats fed high vitamin E diets. AM (2×10^5) were preincubated for 4 h in medium or in medium with macrophage-activating factor (MAF, 1/162 dilution). They were then incubated with ⁵¹Cr-labeled opsonized SRBC for 2 h. (A) Phagocytic activity of AM incubated with medium only. Each value represents the radioactivity of ⁵¹Cr-labeled opsonized SRBC within AM and is the mean \pm SD for 10 rats. (B) The enhancement of phagocytic activity following *in vitro* treatment with MAF. Each value represents the stimulation index and is the mean \pm SD for 10 rats. The stimulation index was calculated by assigning a value of 1 to the phagocytic activity of control AM incubated with medium only and by comparing this to the phagocytic activity of AM from each group treated with MAF. * Significantly different from the controls (50 mg of vitamin E/kg of diet) ($p < 0.05$). (Moriguchi *et al.*: *J. Nutr.*, **120**, 1096–1102, 1990)

Fig. 3A, the ability of AM to phagocytose opsonized SRBC was significantly and dose-dependently enhanced by feeding high vitamin E diets [12]. In addition, those AM from rats fed high vitamin E diets exhibited decreasing responses to MAF, prepared from splenic lymphocytes *in vitro* activated with Con A for 48 h, with increasing contents of




Fig. 4. *In vitro* effect of vitamin E (2 μ g/ml) on the proliferation of whole and M ϕ -depleted splenocytes with Con A (5 μ g/ml) for 72 h. Unshaded, medium; shaded, vitamin E. ** Significantly different from whole splenocytes incubated with medium alone ($p < 0.01$). (Oonishi *et al.*: *J. Nutr. Sci. Vitaminol.*, **41**, 445–453, 1995)

vitamin E (Fig. 3B). These results suggest that AM from rats fed high vitamin E diets had already been activated by MAF.

Macrophages are known to be major prostaglandin-producing cells and also play a critical role in the regulation of immune responses by releasing cytokines that affect lymphocyte functions [13, 14]. As described previously, it has been reported that a high vitamin E diet decreases the production of PGE₂ from macrophages and enhances cellular immune functions [12, 15]. However, the precise mechanism by which vitamin E stimulates lymphocyte proliferation remains unclear. Specifically, it is not known whether vitamin E first stimulates macrophage function or lymphocyte functions. Oonishi *et al.* investigated the *in vitro* effect of vitamin E on the proliferation of both whole splenocytes and macrophage-depleted splenocytes [16]. They found that the proliferation of whole splenocytes was significantly higher than that of macrophage-depleted splenocytes at all concentration of Con A (0.5–10 μ g/ml). In addition, when whole and macrophage-depleted splenocytes were preincubated with vitamin E (2 μ g/ml) for 24 h, the proliferation of whole splenocytes was significantly enhanced compared with that of whole splenocytes preincubated with medium alone. In contrast, macrophage-depleted splenocytes did not show any increase of splenic lymphocyte proliferation following *in vitro* pretreatment with vita-




Fig. 5. Scheme on the activation of macrophages or natural killer (NK) cells by vitamin E.

min E (Fig. 4). This result suggests that vitamin E has the ability to stimulate splenic macrophages directly. In these *in vitro* experiment, the culture medium contained 2-mercaptoethanol (2-ME; 50 μ M), which is also a potent antioxidant. The addition of 2-ME to culture media has been found to enhance the DNA synthetic response of lymphocytes to mitogens [17]. Oonishi *et al.* also found that the addition of 2-ME to the culture media induces a higher proliferation of splenic lymphocytes, comparable to that of vitamin E (2 μ g/ml) [16]. This suggests that the action of vitamin E to enhance lymphocyte proliferation may be masked by the addition of 2-ME to the culture media. Furthermore, they investigated whether or not macrophages pretreated with both vitamin E (2 μ g/ml) and Con A (5 μ g/ml) *in vitro* for 24 h had the ability to enhance the proliferation of splenic lymphocytes. As its result, it was found that the addition of macrophages pretreated with both vitamin E and Con A to splenic lymphocyte cultures could induce even higher proliferation of splenic lymphocytes compared with the addition of splenic macrophages pretreated with Con A alone. Taken together, vitamin E may first activate macrophage function as an immunostimulator rather than an antioxidant, and then activate lymphocytes, including T cells and NK cells, as summarized in Fig. 5.

II. Vitamin E and Decreased Cellular Immunity with Aging

Animal Studies

Even though the life expectancy of experimental animals such as mice and rats is very short, it needs at least 1 to 2 years to investigate the effect of vitamin E on immune functions with aging. A good model for studying aging has been established in mice, which are called as the senescence-accelerated mouse (SAM) [18]. It is known that this mouse strain shows a marked decrease of splenic lymphocyte proliferation with mitogens from 6 months of age [19]. Muraga *et al.* have tried to investigate whether the decreased mitogen response of splenic lymphocytes in SAM is associated with the nutritional status of vitamin E and whether a high vitamin E diet can restore the decreased mitogen response of splenic lymphocytes in SAM. They found that there was no significant difference in serum vitamin E concentration between old SAM-P1 and SAM-R1 as control. Since a high vitamin E diet induced a similar increase of serum vitamin E concentration in both old SAM-P1 and SAM-R1, the vitamin E status in SAM-P1 appears not to be impaired and maintain a normal level [20]. In addition, a high vitamin E diet did not restore the decreased mitogen response of splenic lymphocytes in SAM-P1. Because Muraga *et al.* found that the percentage of adherent cells, mainly macrophages, in splenocytes was remarkably decreased in old SAM-P1 and their macrophages did not have the enhancing effect on proliferation of splenic lymphocytes from young SAM-R1, they came to conclusion that the decreased mitogen response of splenic lymphocytes in SAM-P1 is not associated with vitamin E status and is due to the decreases of both number and function of macrophages.

In rats, since spontaneously hypertensive rats (SHR) were derived from Wistar Kyoto rats (WKY) by Okamoto and Aoki [21], they are now widely used as an animal model for the study of human essential hypertension. Because SHR exhibit an accelerated decrease in, and abnormalities of, cellular immune functions with aging, Takeichi *et al.* proposed that SHR are a good model for the study of not only human essential hypertension but also the mechanism of aging [22]. It has been found that SHR show the remarkable decreases in mitogenesis and NK activity of splenic lymphocytes in the early stage of life (3 months old), and vitamin E levels in





Fig. 6. Proliferation of thymocytes with Con A in WKY and SHR fed a control or high vitamin E diet for 40 weeks. (○) WKY; (●) WKY+vitamin E; (□) SHR; (■) SHR+vitamin E. +Significantly different from thymocyte proliferation with Con A in WKY or SHR fed a control diet at 4 weeks old ($p < 0.05$). * Significantly different from WKY fed a control diet at the same age ($p < 0.05$). # Significantly different from SHR fed a control diet at the same age ($p < 0.05$). (Moriguchi *et al.*: *Nutr. Res.*, 15, 401–414, 1995)

serum and thymus of SHR decreased significantly more during aging in SHR than in WKY [23]. It was considered that there are several explanations for decreased vitamin E levels in serum and thymus of SHR. First, because levels of vitamin E in the serum and thymus of SHR were increased when the rats consumed a high vitamin E diet, the decreased vitamin E level of SHR does not appear to be due to malabsorption of vitamin E. Second, the corn oil used in this experiment contained a considerable quantity of PUFA, which may also further promote the wastage of vitamin E in SHR. In addition, vitamin E supplementation (500 IU/kg of diet) prevented the decrease in proliferation of thymocytes observed in SHR after Con A stimulation (Fig. 6). However, NK activity of splenocytes from SHR was not affected by the high vitamin E diet and was significantly lower than that of WKY. In contrast, phagocytic activity of AM against opsonized SRBC was significantly higher in SHR than in WKY. As it is known that macrophages are nonspecifically activated in athymic, nude mice [24], AM in SHR, which have decreased functions of T cells, also may be nonspecifically activated. In short, this activation of AM may represent compensation for the defect in T cell-mediated immune system of SHR with aging.

The decrease in T cell function in SHR is also associated with increased production of natural thymocytotoxic autoantibody (NTA) with aging [25]. We have also found that NTA titer in the serum of SHR steadily increased with aging and that vitamin E supplementation suppressed this increase [26]. The increase of NTA in SHR is relevant to the impairment of T cell differentiation in the thymus. In other words, the decrease of NTA titer in the serum of SHR may be a mechanism in which the high vitamin E diet restores cellular immune responses decreased with aging. This is related to normalize T cell differentiation in the thymus of SHR. In fact, we have found that vitamin E supplementation improves the decreased expressions of CD4 and CD8 antigens on the membrane of thymic lymphocytes of SHR [26]. These data support that SHR is a good model for aging, especially for studying the effect of vitamin E supplementation and the possibility exists that high vitamin E diet has a potent effect on T cell differentiation in thymus. The effect of vitamin E on T cell differentiation in thymus will be described in the following head.

Human studies

Recent reviews on the beneficial effect of vitamin E for the immune system in the elderly have been summarized by Landmark [27], and Meydani and Beharka [28]. The effects of vitamin E supplementation on immune responses have been epidemiologically investigated on free-living elderly people. Goodwin and Garry failed to find any correlation between vitamin E intake and immune responses such as mitogen response, delayed cutaneous hypersensitivity, and serum antibodies in a population of healthy adults (65–94 years old) [29]. As its reason, they proposed the possibility that some of the previously reported immunoenhancing properties of megadose vitamins may be due to a nonspecific adjuvant effect disappearing with time. On the other hand, Chavance *et al.* found that plasma vitamin E levels were positively correlated with delayed type hypersensitivity response to diphtheria toxoid, candida, and trichophyton in 100 healthy subjects (>60 years old) [30]. They have also found that blood vitamin E concentrations were negatively correlated with the incidence of infectious disease episodes. Payette *et al.* also found that a negative correlation between dietary vitamin E level and IL-2 production in free-living elderly people [31]. However, because this study was performed by using dietary vitamin E



(-) : suppressive effect

Fig. 8. A mechanism of vitamin E action on the improvement of cellular immunity decreased with aging.

blood lymphocytes with Con A ($p < 0.05$) (Fig. 7B) [35]. This evidence suggests that the correction using plasma lipids needs to assess the effect of vitamin E on immune functions in the aged. In addition, although Meydani *et al.* found that PGE₂ level in blood was significantly increased in the aged, which associated with decreased cellular immunity in the aged [36], nitric oxide (NO) produced from macrophage and other cells has also an inhibitory effect on T cell proliferation with mitogen [37]. NO activates cyclooxygenase (Cox), leading to the production of PGE₂. Age-related increase in the production of PGE₂ may be due to increased NO production with aging, which appears to cause the decreased cellular immunity in the aged as shown in Fig. 7A. In other words, the decreased cellular immunity in the aged with the increase level of blood PGE₂ is closely associated with the increased NO production and the action of vitamin E improving the decreased cellular immunity in the aged is due to inhibit NO production from macrophages and other cells (Fig. 8). In fact, a negative correlation between plasma NO concentration and α -tocopherol/VLDL-cholesterol was seen as shown in Fig. 7C. These data support that it is important for elderly to take an enough intake of vitamin E for preventing the decrease of cellular immunity.





Fig. 9. Proportions of CD4⁺CD8⁻ (unshaded) and CD4⁻CD8⁺ (shaded) T cells in thymocytes of rats fed control, vitamin E free, and high vitamin E diets for 7 weeks. Significantly different from the control group (** $p < 0.01$, *** $p < 0.001$). (Moriguchi *et al.*: *J. Nutr. Sci. Vitaminol.*, **39**, 451-463, 1993)

III. Vitamin E and T-Cell Differentiation in the Thymus

From the experimental results in SHR described previously, Moriguchi *et al.* speculated that vitamin is an important nutrient in T-cell differentiation in thymus. To test this hypothesis, 6-week-old Fisher rats were separated into three groups and fed diets containing various levels of α -tocopherol acetate: 0 (vitamin E free), 50 (regular), and 500 (high vitamin E)/kg of diet for 7 weeks. In this experiment rats fed the vitamin E free diet showed not only a decreased number of thymocytes but also a significant decrease of the percentage of CD4⁺CD8⁻ (helper/inducer) T cells in thymocytes compared with rats fed the regular diet (Fig. 9) [38]. In rats fed a high vitamin E diet, the number of thymocytes was not altered, but the percentages of both CD4⁺CD8⁻ and CD4⁻CD8⁺ T cells in their thymocytes increased significantly. Furthermore, IL-2 production by thymocytes was significantly greater than that in rats fed the regular diet. As Moriguchi *et al.* [12] have previously noted that AM can be greatly activated by high vitamin E diets, it is conceivable that an increase of macrophage function as antigen-presenting cells (APC) in the thymus may bring about the increased production of IL-2. In addition, the marked increases of CD4⁺CD8⁻ and CD4⁻CD8⁺ T cells in thymocytes may be directly related to the increased production of



(-) : suppressive effect

Fig. 8. A mechanism of vitamin E action on the improvement of cellular immunity decreased with aging.

blood lymphocytes with Con A ($p < 0.05$) (Fig. 7B) [35]. This evidence suggests that the correction using plasma lipids needs to assess the effect of vitamin E on immune functions in the aged. In addition, although Meydani *et al.* found that PGE₂ level in blood was significantly increased in the aged, which associated with decreased cellular immunity in the aged [36], nitric oxide (NO) produced from macrophage and other cells has also an inhibitory effect on T cell proliferation with mitogen [37]. NO activates cyclooxygenase (Cox), leading to the production of PGE₂. Age-related increase in the production of PGE₂ may be due to increased NO production with aging, which appears to cause the decreased cellular immunity in the aged as shown in Fig. 7A. In other words, the decreased cellular immunity in the aged with the increase level of blood PGE₂ is closely associated with the increased NO production and the action of vitamin E improving the decreased cellular immunity in the aged is due to inhibit NO production from macrophages and other cells (Fig. 8). In fact, a negative correlation between plasma NO concentration and α -tocopherol/VLDL-cholesterol was seen as shown in Fig. 7C. These data support that it is important for elderly to take an enough intake of vitamin E for preventing the decrease of cellular immunity.




Fig. 9. Proportions of CD4⁺CD8⁻ (unshaded) and CD4⁻CD8⁺ (shaded) T cells in thymocytes of rats fed control, vitamin E free, and high vitamin E diets for 7 weeks. Significantly different from the control group (** $p < 0.01$, *** $p < 0.001$). (Moriguchi *et al.*: *J. Nutr. Sci. Vitaminol.*, **39**, 451–463, 1993)

III. Vitamin E and T-Cell Differentiation in the Thymus

From the experimental results in SHR described previously, Moriguchi *et al.* speculated that vitamin is an important nutrient in T-cell differentiation in thymus. To test this hypothesis, 6-week-old Fisher rats were separated into three groups and fed diets containing various levels of α -tocopherol acetate: 0 (vitamin E free), 50 (regular), and 500 (high vitamin E)/kg of diet for 7 weeks. In this experiment rats fed the vitamin E free diet showed not only a decreased number of thymocytes but also a significant decrease of the percentage of CD4⁺CD8⁻ (helper/inducer) T cells in thymocytes compared with rats fed the regular diet (Fig. 9) [38]. In rats fed a high vitamin E diet, the number of thymocytes was not altered, but the percentages of both CD4⁺CD8⁻ and CD4⁻CD8⁺ T cells in their thymocytes increased significantly. Furthermore, IL-2 production by thymocytes was significantly greater than that in rats fed the regular diet. As Moriguchi *et al.* [12] have previously noted that AM can be greatly activated by high vitamin E diets, it is conceivable that an increase of macrophage function as antigen-presenting cells (APC) in the thymus may bring about the increased production of IL-2. In addition, the marked increases of CD4⁺CD8⁻ and CD4⁻CD8⁺ T cells in thymocytes may be directly related to the increased production of

IL-2 by thymocytes in the thymic medulla of rats fed the high vitamin E diet.

It is known that PGE₂ has an inhibitory effect on IL-2 production by activated T cells [39]. A marked decrease of IL-2 production following consumption of vitamin E free diet and an increase after consumption of a high vitamin E diet may be closely related to the degree of PGE₂ synthesis. In fact, Moriguchi *et al.* have reported an increase in PGE₂ production in vitamin E deficiency and a decrease after high vitamin E supplementation [38]. These changes in PGE₂ production by thymocytes may affect production of IL-2 by T cells in thymocytes and result in a decreased percentage of CD4⁺CD8⁻ T cells in thymocytes of the vitamin E free group and increased percentage in those of the high vitamin E group. In fact, since addition of indomethacin, an inhibitor of PGE₂ synthesis, to the thymocyte cultures enhanced IL-2 production by thymocytes of rats fed a vitamin E free diet, the increased production of PGE₂ in thymocytes appears to be closely related to the decreased production of IL-2 in the vitamin E free group.

Thymic epithelial cells (TEC) play an important role in the differentiation of T cells [40]. Moriguchi and Itoh have found that the ability of TEC isolated from thymocytes of rats fed a high vitamin E diet to bind to immature T cells was greater than that of TEC from rats fed a regular diet [41]. Anderson *et al.* showed that the contact of immature T cells to TEC is more important than soluble factors from TEC in positive selection of CD4⁺CD8⁻ T cells [42]. Moriguchi and Itoh have also found that the supernatant of a TEC culture does not affect the differentiation of T cells *in vitro*. These results suggest that vitamin E enhances T cell differentiation through increased binding of immature T cells to TEC in the thymic cortex. Although it is not known how vitamin E increases the binding of immature T cells to TEC, vitamin E may enhance the expression of adhesion molecules. Moriguchi and Itoh have found that vitamin E supplementation or *in vitro* addition of vitamin E to the culture medium enhances the expression of intracellular adhesion molecule-1 (ICAM-1) in TEC. However, they have failed to find that *in vitro* incubation with macrophages isolated from rats fed a high vitamin E diet induces a significant change in T cell subsets in immature T cells. These results suggest that vitamin E enhances T-cell differentiation not through macrophage function (negative selection) but through an

increased binding capacity of TEC to immature T cells *via* increased expression of ICAM-1 (positive selection).

IV. Vitamin E and Immune Diseases

Vitamin E and acquired immune deficiency syndrome (AIDS)

Acquired immune deficiency syndrome (AIDS) is a clinical disorder caused by the human immunodeficiency virus (HIV). It represents the end point in a progressive sequence of immunosuppressive changes that render the body highly susceptible to tumors and opportunistic infections. Currently, AIDS is one of the most serious public health problems in the world. Although many studies have been done from the viewpoints of prevention and treatment for AIDS, there are still few studies on the effect of nutrition against the development of AIDS.

Since reactive oxygen species promote HIV replication [43], previous studies have found that antioxidants such as vitamin C and *N*-acetyl-L-cysteine (NAC), the precursor of glutathione (GSH), are effective in inhibiting HIV replication [44]. Odeleye and Watson have reviewed the potential role of vitamin E in the treatment of immunologic abnormalities during AIDS [45]. They described that vitamin E is nontoxic over a wide range of intakes and a moderately high dose may be used to target and stimulate some specific immune cells destroyed by HIV infection. Moseson *et al.* have also reviewed the potential role of nutritional factors in the induction of immunologic abnormalities in HIV-positive homosexual men [46]. Since malnutrition and intestinal nutrient malabsorption have been found in AIDS patients, they have described that dietary manipulations might diminish the immune defects in HIV infection and enhance resistance to opportunistic infections.

Murine AIDS has often been used as an AIDS model instead of human AIDS. Although the etiology between human and murine AIDS, induced by LP-BM5 retrovirus infection, is different in some points, including the changes of the T cell subsets, both are closely similar in the functional changes following development of AIDS. Murine AIDS is characterized by lymphadenopathy, splenomegaly, hypergammaglobulinemia, deficient B-cell response to mitogen, functional deficiency of T cells, and cytokine dysregulation as shown in human AIDS [47]. In addition, it is also known that azidodideoxythymi-




Fig. 10. Effect of vitamin E supplementation on the expression of NF- κ B in cytoplasm and nucleus of splenic lymphocytes of rats infected with LP-BM5 retrovirus. Twenty microgram of each protein was analyzed by Western blotting. (a) Control diet; (b) high vitamin E diet before LP-BM5 retrovirus infection; (c) high vitamin E diet after LP-BM5 retrovirus infection. (Hamada *et al.*: *Nutr. Res.*, **20**, 1163–1171, 2000)




Fig. 11. Mechanism of the improvement of immune dysfunction in murine AIDS following vitamin E supplementation.

dine (AZT), a drug for treatment of human AIDS, is effective for treatment of murine AIDS [48]. These reports support that murine AIDS is a useful model for studying the treatment and prevention of human AIDS.

It is known that NF- κ B enhances HIV replication through activation of the HIV long-terminal repeat (LTR) [49]. In addition, reactive oxygen species promote the dissociation of NF- κ B from I- κ B and subsequently increase the production of NF- κ B [43, 44]. As shown in Fig. 10, Hamada *et al.* have found that vitamin E supplementation suppresses the expression of NF- κ B in splenic lymphocytes from mice infected with LP-BM5 retrovirus [50]. Especially, the suppressive effect of vitamin E is strongly shown in the nucleus rather than in the cytoplasm. This fact




Fig. 12. Effect of vitamin E supplementation on serum IgE production from TDI-sensitized mice. Mice were sensitized by dropping 2 μ l of 5% toluene diisocyanate (TDI) dissolved in ethyl acetate from an autopipette into the nostrils under slight ether anesthesia for consecutive days. This was repeated after a week of rest. No sensitization groups were similarly treated with the vehicle, ethyl acetate. The mice were again allowed 1 week of rest, after which all groups were provoked with 5 μ l of 2.5% TDI without anesthesia to induce nasal allergy-like symptoms. One week after provocation, the mice were anesthetized with sodium pentobarbital and were exsanguinated by cardiac puncture. Each value is the mean \pm SE. * Significantly different from no sensitization mice fed a control or vitamin E-supplemented diet ($p < 0.01$). # Significantly different from TDI sensitized mice fed a control diet ($p < 0.01$). (Zheng *et al.*: *Am. J. Med. Sci.*, **318**, 49–54, 1999)

suggests that the suppressive action of vitamin E against NF- κ B expression is not only the dissociation of NF- κ B from I- κ B but also the translocation of NF- κ B to the nucleus. Figure 11 shows that there are two systems of vitamin E actions improving immune dysfunction in murine AIDS. First, vitamin E increases the production of IFN- γ , an antiviral cytokine produced from activated T cells. Vitamin E may inhibit the development of murine AIDS through the enhancement of host antiviral activity. Second, since vitamin E is a potent antioxidant and decreases the production of TNF- α , which enhances the production of reactive oxygen species and subsequently increases the expression of NF- κ B, vitamin E may directly and indirectly inhibit the production of reactive oxygen species and NF- κ B expression,

and result in inhibiting the development of murine AIDS.

Vitamin E and abnormal increase of immune system

In this part we describe effects of vitamin E supplementation on two types of diseases with immunological abnormality such as allergy and autoimmune disease. Zheng *et al.* have found that high doses of vitamin E supplementation suppress nasal allergy [51]. As shown in Fig. 12, groups B [control+toluene diisocyanate(TDI) sensitization] and D (vitamin E+TDI sensitization) had higher ($p<0.01$) serum IgE concentrations compared with groups A (control) and C (vitamin E). However, serum IgE concentrations in group D were significantly lower than those in group B ($p<0.05$). As its mechanism it has been shown that vitamin E has a direct inhibitory effect on serum IgE formation in mice [52]. In addition, vitamin E acts as an antioxidant in cellular membranes and scavenges free radicals by blocking the peroxidation of PUFA, modifying prostaglandin formation, and thereby enhancing production of PGI₂ [53]. It is known that PGI₂ has an inhibitory effect on the formation of histamine [54], which is a major chemical mediator in the events of TDI-induced allergy, and causes itching, sneezing, and rhinorrhea. From these reports it appears that vitamin E may have reduced nasal allergic symptoms by inhibition of the production of histamine. Fogarty *et al.* have also investigated the relation between dietary vitamin E intake and serum IgE concentrations and atopy, measured as allergen skin sensitization, in a random sample of 2,633 adults [55]. As its result they found that higher concentrations of vitamin E intake were associated with lower serum IgE concentrations and a lower frequency of allergen sensitization.

By using MRL/MP-lpr/lpr (MRL/lpr) mice Weimann and Hermann have found that vitamin E beneficially affects the development of the SLE-like (systemic lupus erythematosus) autoimmune disease [56]. In addition, Kuemmerie *et al.* investigated the effects of fish oil, with and without α -tocopherol, on the course of IgA nephropathy [57] and found that vitamin E, more so than fish oil, mitigates the injury and promotes repair in experimental IgA nephropathy. Bae *et al.* investigated on plasma antioxidant/oxidant status and the dietary nutrient intake of 97 consecutive patients with SLE and 97 age- and sex-matched healthy controls [58]. They have found that plasma antioxidant status is impaired and dietary antioxidant intake is decreased in patients with SLE.

In particular, they found the plasma α -tocopherol concentration was lower in those patients. As described above, vitamin E supplementation may be effective for treatment of those patients having vitamin E deficient status. Further studies are necessary to define the exact mechanism of vitamin E modulating the immunological abnormalities in allergy and autoimmune diseases.

References

- [1] Meydani, S.N., Barklund, M.P., Liu, S., Meydani, M., Miller, R., Cannon, J., Morrow, F., Rocklin, R., and Blumberg, J.: Vitamin E supplementation enhances cell-mediated immunity in healthy elderly. *Am. J. Clin. Nutr.*, **52**, 557-563, 1990.
- [2] Meydani, S.N., Meydani, M., Verdon, C.P., Shapiro, A.C., Blumberg, J.B., and Hayes, K.C.: Vitamin E supplementation suppresses prostaglandin E₂ synthesis and enhances the immune response in aged mice. *Mech. Age. Dev.*, **34**, 191-201, 1986.
- [3] Machlin, L.: Vitamin E and prostaglandins, in Tocopherol, Oxygen, and Biomembranes, ed. by de Duve, C. and Hayaishi, O., Elsevier/North-Holland Biomedical Press, Amsterdam, pp. 208-215, 1978.
- [4] Sharmanov, A.T., Aidarkhanov, B.B., and Kurmangaliev, S.M.: Effect of vitamin E on oxidative metabolism of macrophages. *Bull. Exp. Biol. Med.*, **101**, 723-725, 1986.
- [5] Moriguchi, S., Kobayashi, N., and Kishino, Y.: Effects of vitamin E deficiency on the functions of splenic lymphocytes and alveolar macrophages. *J. Nutr. Sci. Vitaminol.*, **35**, 419-430, 1989.
- [6] Sharp, A.K. and Colston, M.J.: Elevated macrophage activity in nude mice. *Exp. Cell Biol.*, **52**, 44-47, 1984.
- [7] Moriguchi, S. and Muraga, M.: Vitamin E and immunity, in Vitamins and Hormones, Vol. 59, ed. by Litwack, G., Academic Press, New York, pp.305-336, 2000.
- [8] Baehner, R.L., Boxer, L.A., Allen, L.M., and Davis, J.: Autoxidation as a basis for altered function by polymorphonuclear leukocytes. *Blood*, **50**, 327-335, 1977.
- [9] Goodwin, J.S., Messner, R.P., and Peake, G.T.: Prostaglandin suppression of mitogen-stimulated lymphocytes *in vitro*. Changes with mitogen dose and preincubation. *J. Clin. Invest.*, **72**, 753-760, 1978.
- [10] Segman, W.E., Gindhart, T.D., Blackman, M.A., Dalal, B., Talal, N., and Werb, Z.: Suppression of natural killing *in vitro* by monocytes and polymorphonuclear leukocytes. *J. Clin. Invest.*, **69**, 876-888, 1982.
- [11] El-Hag, A., Lipsky, P.E., Bennett, M., and Clark, R.A.: Immunomodulation by neutrophil myeloperoxidase and hydrogen peroxide: Differential susceptibility of human lymphocyte function. *J. Immunol.*, **9**, 3420-3426, 1986.
- [12] Moriguchi, S., Kobayashi, N., and Kishino, Y.: High

- dietary intakes of vitamin E and cellular immune functions in rats. *J. Nutr.*, **120**, 1096–1102, 1990.
- [13] Fidler, I.J.: Macrophages and metastasis—A biological approach to cancer therapy. *Cancer Res.*, **45**, 4714–4726, 1985.
- [14] Nathan, C.F.: Secretory products of macrophages. *Clin. Invest.*, **79**, 319–326, 1987.
- [15] Romach, E.H., Kidao, S., Sanders, B.G., and Kline, K.: Effects of RRR- α -tocopheryl succinate on IL-1 and PGE₂ production by macrophages. *Nutr. Cancer*, **20**, 205–214, 1993.
- [16] Oonishi, K., Moriguchi, S., and Kishino, Y.: The role of macrophages in increased mitogen response of rat splenic lymphocytes following *in vitro* incubation with vitamin E. *J. Nutr. Sci. Vitaminol.*, **41**, 445–453, 1995.
- [17] Lemke, H. and Opitz, H.G.: Function of 2-mercaptoethanol as a macrophage substitute in the primary immune response *in vitro*. *J. Immunol.*, **117**, 388–395, 1976.
- [18] Takeda, T., Hosokawa, M., Takeshita, S., Irino, M., Matsushita, T., Tomita, Y., Yasuhira, K., Hamamoto, H., Shimizu, K., Ishii, M., and Yamamuro, T.: A new murine model of accelerated senescence. *Mech. Ageing Dev.*, **17**, 183–194, 1981.
- [19] Umezawa, M., Hamada, K., Naiki, H., Chen, W.H., Hosokawa, M., Hosono, M., Hosokawa, T., and Takeda, T.: Effects of dietary restriction on age-related immune dysfunction in the senescence accelerated mouse (SAM). *J. Nutr.*, **120**, 1393–1400, 1990.
- [20] Muraga, M., Umegaki, K., Yamamoto, S., and Moriguchi, S.: Decreased mitogen response of splenocytes in senescence accelerated mice (SAM-P1) is not restored by high vitamin E diet. *Nutr. Res.*, **19**, 1393–1400, 1999.
- [21] Okamoto, K. and Aoki, K.: Development of a strain of spontaneously hypertensive rats. *Jpn. Circ. J.*, **27**, 287–293, 1963.
- [22] Takeichi, N., Suzuki, K., and Kobayashi, H.: Characterization of immunological depression in spontaneously hypertensive rats. *Eur. J. Immunol.*, **11**, 483–487, 1981.
- [23] Moriguchi, S., Maekawa, K., Miwa, H., and Kishino, Y.: Effect of vitamin E supplementation on cellular immune functions decreased with aging in spontaneously hypertensive rats. *Nutr. Res.*, **13**, 1039–1051, 1993.
- [24] Sharp, A.K. and Colston, M.J.: Elevated macrophage activity in nude mice. *Exp. Cell Biol.*, **52**, 44–47, 1984.
- [25] Takeichi, N., Ba, D., and Kobayashi, H.: Natural cytotoxic autoantibody against thymocytes in spontaneously hypertensive rats. *Cell Immunol.*, **60**, 181–190, 1981.
- [26] Moriguchi, S., Maekawa, K., Okamoto, M., Oonishi, K., and Kishino, Y.: Vitamin E prevents the decrease of cellular immune function with aging in spontaneously hypertensive rats. *Nutr. Res.*, **15**, 401–414, 1995.
- [27] Landmark, K.: Vitamin E is beneficial for the immune system in the elderly. *Tidsskr. Nor. Laegeforen.*, **118**, 4403–4404, 1998.
- [28] Meydani, S.N. and Beharka, A.A.: Recent developments in vitamin E and immune response. *Nutr. Rev.*, **56**, S49–58, 1998.
- [29] Goodwin, J.S. and Garry, P.J.: Relationship between megadose vitamin supplementation and immunological function in a healthy elderly population. *Clin. Exp. Immunol.*, **51**, 647–653, 1983.
- [30] Chavance, M., Brubacher, G., Herbeth, B., Vernhes, G., Mikstachi, T., Dete, F., Fournier, C., and Janot, C.: Immunological and nutritional status among the elderly, in *Nutrition, Immunity and Illness in the Elderly*, ed. by Chandra, R.K., Pergamon Press, New York, pp.137–142, 1985.
- [31] Payette, H., Rpla-Pleszczynski, M., and Bhadrian, P.: Nutrition factors in relation to cellular and regulatory immune variables in a free-living elderly population. *Am. J. Clin. Nutr.*, **52**, 927–932, 1990.
- [32] Nagel, J.E., Chopra, R.K., Chrest, F.J., McCoy, M.T., Schneider, E.L., Holbrook, N.J., and Adler, W.H.: Decreased proliferation, interleukin 2 synthesis, and interleukin 2 receptor expression are accompanied by decreased mRNA expression in phytohemagglutinin-stimulated cells from elderly donors. *J. Clin. Invest.*, **81**, 1096–1102, 1988.
- [33] De Waart, F.G., Portengen, L., Doekes, G., Verwaal, C.J., and Kok, F.J.: Effect of 3 months' vitamin E supplementation on indices of the cellular and humoral immune response in elderly subjects. *Br. J. Nutr.*, **78**, 761–774, 1997.
- [34] De la Fuente, M., Fernandez, M.D., Burgus, M.S., Soler, A., Prieto, A., and Miquel, J.: Immune function in aged women is improved by ingestion of vitamins C and E. *Can. J. Physiol. Pharmacol.*, **76**, 373–380, 1998.
- [35] Watanabe, Y., Muraga, M., and Moriguchi, S.: Epidemiological study on the relationship between decreased cellular immunity and nitric oxide (NO) in the aged. *Rep. Res. Comm. Essent. Amino Acids*, **158**, 39–44, 2000 (in Japanese).
- [36] Beharka, A.A., Wu, D., Ham, S.N., and Meydani, S.N.: Macrophage prostaglandin production contributes to the age-associated decrease in T cell function which is reversed by the dietary antioxidant vitamin E. *Mech. Ageing Dev.*, **93**, 59–77, 1997.
- [37] Fu, Y. and Blankenhorn, E.P.: Nitric oxide-induced anti-mitogenic effects in high and low responder rat strains. *J. Immunol.*, **148**, 2217–2222, 1992.
- [38] Moriguchi, S., Miwa, H., Okamura, M., Maekawa, K., Kishino, Y., and Maeda, K.: Vitamin E is an important factor in T cell differentiation in thymus of F344 rats. *J. Nutr. Sci. Vitaminol.*, **39**, 451–463, 1993.
- [39] Rappaport, R.S. and Dodge, G.R.: Prostaglandin E inhibits the production of human interleukin 2. *J. Exp. Med.*, **155**, 943–948, 1982.
- [40] Eshel, I., Savion, N., and Shoham, J.: Analysis of thymic stromal cell subpopulation grown *in vitro* on extra-

- cellular matrix in defined medium. II. Cytokine activities in murine thymic epithelial and mesenchymal culture supernatants. *J. Immunol.*, **144**, 1563–1570, 1990.
- [41] Moriguchi, S. and Itoh, T.: Vitamin E enhances T cell differentiation through increased epithelial cell function in rats thymus. *Nutr. Res.*, **17**, 873–883, 1997.
- [42] Anderson, G., Owen, J.J.T., Moore, N.C., and Jenkinson, E.J.: Thymic epithelial cells provide unique signals for positive selection of CD4⁺CD8⁻ thymocytes *in vitro*. *J. Exp. Med.*, **179**, 2027–2031, 1994.
- [43] Schreck, R., Rieber, P., and Baeuerle, P.A.: Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF- κ B transcription factor and HIV-1. *EMBO J.*, **10**, 2247–2258, 1991.
- [44] Staal, F.J., Roederer, M., Herzenberg, L.A., and Herzenberg, L.A.: Intracellular thiols regulate activation of nuclear factor κ B and transcription of human immunodeficiency virus. *Proc. Natl. Acad. Sci. USA*, **87**, 9943–9947, 1990.
- [45] Odeleye, O.E. and Watson, R.R.: The potential role of vitamin E in the treatment of immunologic abnormalities during acquired immune deficiency syndrome. *Prog. Food Nutr. Sci.*, **15**, 1–19, 1991.
- [46] Moseson, M., Zeleniuch-Jacquotte, A., Belsito, D.V., Shore, R.E., Marmor, M., and Pastermack, B.: The potential role of nutritional factors in the induction of immunologic abnormalities in HIV-positive homosexual men. *J. Acquir. Immune Defic. Syndr.*, **2**, 235–247, 1989.
- [47] Morse, III, H.C., Chattopadhyay, S.K., Makino, M., Fredrickson, T.N., Hugin, W., and Hartley, J.W.: Retrovirus-induced immunodeficiency in the mouse: MAIDS as a model for AIDS. *AIDS*, **6**, 607–621, 1992.
- [48] Portnoi, D., Staal, A.M., Schwartz, D., Merigan, T.C., Herzenberg, L.A., and Basham, T.: Zidovudine (azido dideoxythymidine) inhibits characteristic early alterations of lymphoid cell population in retrovirus-induced murine AIDS. *J. Immunol.*, **144**, 1705–1710, 1990.
- [49] Lenardo, M.J. and Baltimore, D.: NF- κ B: A pleiotropic mediator of inducible and tissue-specific gene control. *Cell*, **58**, 227–229, 1989.
- [50] Hamada, M., Yamamoto, S., Kishino, Y., and Moriguchi, S.: Vitamin E suppresses the development of murine AIDS through the inhibition of nuclear- κ B expression. *Nutr. Res.*, **20**, 1163–1171, 2000.
- [51] Zheng, K.-C., Adjei, A., Shinjo, M., Shinjo, S., Todoriki, H., and Arizumi, M.: Effect of dietary vitamin E supplementation on murine nasal allergy. *Am. J. Med. Sci.*, **318**, 49–54, 1999.
- [52] Inagaki, N., Nagai, H., and Koda, A.: Effect of vitamin E on IgE antibody formation in mice. *J. Pharm. Dyn.*, **7**, 70–74, 1984.
- [53] Douillet, C., Bost, M., Accominotti, M., Borson-Chazot, F., and Ciavatti, M.: Effect of selenium and vitamin E supplementation on lipid abnormalities in plasma aorta, and adipose tissue of Zucker rats. *Biol. Trace Elem. Res.*, **65**, 221–236, 1998.
- [54] Tanaka, Y., Imai, T., Horikoshi, S., Kurokawa, M., Kimura, T., Matsuura, T., Idaira, K., Adachi, M., and Takahashi, T.: Effect of thromboxane A₂ and prostaglandin I₂ on the release of chemical mediators from human peripheral leukocytes. *Aterugi*, **42**, 1677–1682, 1993 (in Japanese).
- [55] Fogarty, A., Lewis, S., Weiss, S., and Britton, J.: Dietary vitamin E, IgE concentrations, and atopy. *Lancet*, **356**, 1573–1574, 2000.
- [56] Weimann, B.J. and Hermann, D.: Inhibition of autoimmune deterioration in MRL/lpr mice by vitamin E. *Int. J. Vitam. Nutr. Res.*, **69**, 255–261, 1999.
- [57] Kuemmerie, N.B., Chan, W., Krieg, R.J., Jr., Norkus, E.P., Trachtman, H., and Chan, J.C.: Effects of fish oil and α -tocopherol in immunoglobulin A nephropathy in the rat. *Pediatr. Res.*, **43**, 791–797, 1998.
- [58] Bae, S.C., Kin, S.J., and Sung, M.K.: Impaired antioxidant status and decreased dietary intake of antioxidants in patients with systemic lupus erythematosus. *Rheumatol. Int.*, **22**, 238–243, 2002.

Beneficial effects of the vegetable juice Aojiru on cellular immunity in Japanese young women

Kanako Ogawa^{a,*}, Kaoru Nakada^b, Satoe Yamashita^b, Tsuneo Hasegawa^a,
Satoru Moriguchi^b

^a*Q'sai Company, Ltd., 1-7-16 Kusagae, Chuoku, Fukuoka, Japan 810-8606*

^b*Department of Nutrition, Faculty of Human Life Science, Yamaguchi Prefectural University, 3-2-1 Sakurabatake, Yamaguchi, Japan 753-8502*

Received 23 January 2003; received in revised form 29 October 2003; accepted 9 December 2003

Abstract

This study was performed to investigate the effect of vegetable juice, Aojiru, on cellular immune functions of healthy young women ($n = 19$, aged 18–22 yr). They drank 180 mL of Aojiru daily for 4 weeks, and then their blood samples were collected at 2 and 4 weeks after study onset. Aojiru intake had no effect on the proportion of either natural killer cells (NK) or T-lymphocytes in their peripheral blood. There was a positive correlation between NK activity and the proportion of CD16⁺ CD56⁺ (NK) cells in their peripheral blood lymphocytes (PBL). In addition, NK activity per cell increased with prolonged Aojiru intake. Serum interleukin-2 (IL-2) concentration in subjects was significantly higher after 4 week-feeding of Aojiru compared to levels at the onset of this experiment. This result suggests that Aojiru has an ability to enhance NK activity, possibly through the increased production of IL-2. Therefore, it may be a useful supplement to promote health. © 2004 Elsevier Inc. All rights reserved.

Keywords: Aojiru; Kale; NK activity; Interleukin-2

1. Introduction

In Japan, lifestyle-related diseases such as cancer, heart disease, cerebrovascular disease, diabetes, hyperlipidemia, and hypertension have increased in recent years. Among these

* Corresponding author. Tel.: (0940) 37-3991; fax: (0940) 37-3988.

E-mail address: ogawa@kyusai.co.jp (K. Ogawa).