

表 *Clinostomum* 人体咽喉頭寄生報告例(本邦)

	年齢	性別	住所	主訴	著者
1	22	F	大阪		Yamashita(1938)
2	38	F	富山	咽頭痛, 咽喉頭異常感	堀(1942)
3	30	M	島根	咽喉頭異常感(疼痛感)	Kamo et al.(1962)
4	34	M	長崎	咽喉頭異常感, 嚥下痛	坂口, 他(1966)
5	53	M	岐阜	右側口腔違和感	佐野, 他(1980)
6	31	F	愛知	発咳、血痰、疼痛	熊田, 他(1983)
7	35	M	熊本	咽喉頭異常感(疼痛感)	Hirai et al.(1987)
8	57	F	滋賀	咽喉頭異常感, 軽い嚥下痛	古川・宮里(1987)
9	29	F	福岡	咽頭部違和感	村田・高尾(1988)
10	15	F	島根	咽頭異物感	山根, 他(1990)
11	54	F	佐賀	咽頭異物感, 軽い嚥下痛	梅崎, 他(1990)
12	70	F	秋田	咽頭痛	Yoshimura et al(1991)
13	68	M	島根	咽喉頭異常感	Isobe et al(1994)
14	37	F	佐賀	咽喉頭異常感	木船・上坂(1994)
15	40	F	佐賀	咽頭痛	木船, 他(1994)
16	27	F	島根	咽頭異物感	前嶋, 他(1996)
17	29	F	広島	咽頭違和感	白井, 他(1998)
18	20	F	山口	咽喉頭異常感	木船, 他(2000)
19	27	F	愛知	咽喉頭異常感	山本, 他(2002)
20	55	F	群馬	咽頭部違和感	椋, 他(2003)
21	27	M	佐賀	咽頭痛, 咽喉頭異常感	柿添, 他(2004)
22	46	F	熊本	咽頭痛, 咽頭部違和感	自験例(2005)

Altered expression of goblet cell- and mucin glycosylation-related genes in the intestinal epithelium during infection with the nematode *Nippostrongylus brasiliensis* in rat.

JUNKO YAMAUCHI,^{1,2} YUICHI KAWAI¹, MINORU YAMADA¹ RYUICHI UCHIKAWA¹, TATSUYA TEGOSHI¹ and NAOKI ARIZONO¹

¹Department of Medical Zoology and ²Department of Molecular Gastroenterology and Hepatology, Kyoto Prefectural University of Medicine, Kyoto, Japan

Running head: Mucin response in nematode infection

Address correspondence to:

Naoki Arizono

Telephone: +81 75 251 5325

Fax: +81 75 251 5328

E-mail: arizonon@koto.kpu-m.ac.jp

Summary

Intestinal nematode infection induces marked goblet cell hyperplasia and mucus secretion, but the mechanisms of regulation of the changes still remain to be elucidated. In the present study, epithelial cells were isolated from the rat small intestine at various times after *Nippostrongylus brasiliensis* infection, and the levels of expression of goblet cell- and mucin glycosylation-related genes were estimated by semi-quantitative reverse transcription (RT)-PCR. Among the genes investigated, mucin core peptide (MUC) 2, sialyltransferase (Siat) 4c and trefoil factor family (TFF) 3 were upregulated as early as 2-4 days post-infection, suggesting that they are associated with an early innate protective response. Seven days post-infection and thereafter, when the nematodes reached maturity, significant upregulation of MUC3, MUC4, resistin-like molecule β (Relm β) and 3O-sulfotransferase (3ST)1 was observed, while 3ST2 expression levels increased after the majority of the worms were expelled from the intestine. Similar alterations of glycosylation-related gene expression were also observed in mast-cell-deficient *Ws/Ws* rats, suggesting that mast cells in the epithelium are not relevant to the upregulation of these genes. The present finding that the expression level of each goblet cell- or glycosylation-related gene was altered differently during the time course of infection indicates the progression of sequential qualitative changes in the mucus layer after infection.

Key words: mucin; glycosylation; goblet cell; intestine; *Nippostrongylus*

Naoki Arizono, Department of Medical Zoology, Kyoto Prefectural University of Medicine, Kawaramachi-Hirokoji, Kyoto 602-8566, Japan.

e-mail: arizonon@koto.kpu-m.ac.jp

Introduction

Intestinal nematode parasites such as ascaris and hookworms infect more than 1.3 billion people in the world. Although they cause relatively little mortality, infections result in high levels of morbidity that can result in developmental retardation in infected children (1). Although it has been clarified that Th2 cytokines and/or Th2 cytokine-dependent responses such as mastocytosis and eosinophilia might have crucial roles in protective immunity against certain nematode infections (2), further clarification of factors or mechanisms that are responsible for colonization of worms in and/or rejection of worms from the intestine is an issue of importance for developing more effective measures to combat intestinal parasites. Among factors of potential importance, particularly interesting factors are produced in and secreted from goblet cells, including mucins and other secretory peptides. In fact, intestinal nematode infection induces marked goblet cell hyperplasia and mucus secretion, which may be induced directly via the local release of bioactive factors or indirectly via the activation of host immune cells (3,4).

The nematode *Nippostrongylus brasiliensis*, a lumen-dwelling parasite, is a suitable model for studying human and other clinically relevant hookworms because of the similarities of habitat and life cycle (5). Interestingly, the mucus response which is induced by infection with *N. brasiliensis* has been suggested to be responsible for the rejection of nematodes from the intestine, whereas mast cells are not essential for expulsion of this parasite (3,4). It has been shown that alterations in the terminal sugars of goblet cell mucins are associated with *N. brasiliensis* worm expulsion from the intestine by studies of lectin-binding as well as by biochemical analyses (6-10). Recently, Knight *et al.* (11) and Pemberton *et al.* (12) reported that the expression levels of a variety of goblet cell- and/or mucin-related genes were altered during infection with *Trichinella spiralis* in mice. The results indicated that molecular changes that are

induced in the epithelial and/or mucus layer during nematode infection are a complicated series of changes whose mechanisms of regulation are virtually unknown, as are the roles of each molecule in mucosal protection.

We therefore attempted to determine by semi-quantitative reverse transcription (RT)-PCR whether there were alterations of the expression of some goblet cell-related genes in the small intestine during the time-course of *N. brasiliensis* infection in rats. The genes examined included those of mucin core peptides MUC2, MUC3 and MUC4 (13-16) and goblet cell-specific secretory peptides such as Relm β (17) and intestinal trefoil factor (TFF3)(18,19). Because of the possible importance of the terminal structure of oligosaccharide chains of mucins and/or membrane glycoproteins for colonization and/or rejection of worms, we also examined the gene expression of sialyltransferase, sulfotransferase, fucosyltransferase and some histo-blood group transferases, which might modulate the terminal sugar chains of mucin and/or membrane glycoproteins (20-26).

MATERIALS AND METHODS

Animals, nematode infection and autopsy

Specific-pathogen-free male Brown Norway/Sea (BN) rats and male Fischer (F)-344 rats were purchased from SLC Inc. (Shizuoka, Japan). SPF male and female mast cell-deficient *Ws/Ws* rats were produced in our laboratory as described previously (27). Animals at 8 weeks of age were injected subcutaneously with 2,000 *N. brasiliensis* infective-stage (L3) larvae as described elsewhere (27). The animals were allowed to feed *ad libitum* throughout the experiment.

Preparation of intestinal epithelial cells

The animals were sacrificed with an overdose of ether after overnight fasting with free access to water. The separation of intestinal epithelial cells was carried out at 4°C in EDTA-Hanks' solution (Ca²⁺, Mg²⁺-free Hanks' balanced salt solution supplemented with 10 mM HEPES, pH 7.3, 1 mM DTT and 1.0 mM EDTA) as described elsewhere (28) with slight modifications. In brief, a piece of jejunum 18-22 cm from the pyloric ring and a piece of ileum 10-14 cm from the ileocecal junction, or a 4-cm-long segment of the proximal colon, were removed, opened longitudinally and cut into segments 1 cm in length. After a brief wash in PBS, 4 pieces of tissue were put into a 15-ml tube containing 4 ml of EDTA-Hanks' solution, and debris attached to the mucosal surface was removed by vigorously shaking the tubes 15 times by hand. The tissues were then transferred into another tube containing EDTA-Hanks' solution. After 75 min on ice with occasional agitation of the tissues by inverting the tubes, the epithelial cells were separated by 60 strokes of vigorous shaking of the tube by hand. After discarding the tissue, detached epithelial cells were collected by centrifugation at 600 xg for 3 min at 4°C, washed once with EDTA-Hanks' solution, and the cell pellets were stored at -80°C until use. Giemsa staining of the separated epithelial fractions showed not only epithelial cells, but also a small number of mononuclear cells, the majority of which we considered might have been intraepithelial lymphocytes. Histological examination of the tissue after collection of the epithelia showed that villus epithelium was separated completely, whereas the epithelial lining cells in the lower part of crypts were still attached to the tissue in approximately half of the crypts. In uninfected rats, the basal lamina of the epithelium was intact and lamina propria cells were retained in the tissue, but in animals after 10 days of infection, the basal lamina was partly obscured, indicating that some propria mucosal cells contaminated the epithelial fractions. Concerning the stomach, a mucosal scrape specimen of the glandular stomach was

prepared by scraping the mucosa with the edge of a glass slide and stored at -80°C until use.

Extraction of total RNA, cDNA synthesis, RT-PCR and relative quantification

Total RNA was extracted using TRIZOL Reagent (Life Technologies, Rockville, Maryland). Two-microgram aliquots of RNA were reverse transcribed in 20 µl of reverse transcription buffer containing 5 mM MgCl₂, 1 mM dNTP mixture, 1U/µl RNase inhibitor, 0.25 U/µl AMV reverse transcriptase and 0.125 µM oligo dT-adaptor primer (Takara RNA LA PCR kit, Takara Biomedicals, Osaka, Japan) at 42°C for 50 min. One-microliter aliquots of the synthesized cDNA were mixed with Sybr Green PCR master mix (Applied Biosystems, Foster City, CA) with appropriate primers and amplified using a real-time PCR system 7300 (Applied Biosystems, Foster, CA, U.S.A.).

The sense and antisense primers used were:

5'-CGGATCCAATGGAACAGTGG-3' and 5'-TGCCACTGGTAGGATGATTG-3' for MUC2; 5'-GTTTCAACTCGACTGCCACC-3' and 5'-ATAGCTGCAGTTCTTGGAGG-3' for MUC3; 5'-GCGGAAGAGGAGTGGAGAAG-3' and 5'-AGATGGCCAGTAGCAAGAGG-3' for MUC4; 5'-TTCCTTCTCTCGCTGATGGT-3' and 5'-GCAGTGGCAAGTAGTTCCAT-3' for Relmβ; 5'-ATGGAGACCAGAGCCTTCTG-3' and 5'-TGGGATGCTGGAGTCAAAACA-3' for Tff3; 5'-CTACACCTCTGCGACTTGGT-3' and 5'-GGTTCTTGACAGCTCCCATC-3' for Siat4c; 5'-CCCTTCCCTGAGATCCAGA-3' and 5'-CCGGCCTTTGGACTCATGTA-3' for 3ST1; 5'-CCCAGATCCACTTCGTCAGT-3' and 5'-AAAATTCCCGGAGCTGGTCT-3' for 3ST2; 5'-AGCAATGGCATGAGATGGTG-3' and 5'-TCTGGAAGGGTGAAGTTAGC-3' for FUT1; 5'-GGTGCCGGGAGAACATTAAT-3' and 5'-GAGAATCCGGAAGGGTGTAG-3' for FUT2; 5'-GATTTCCCTAGTGCTGCCTC-3' and 5'-GTTGTGGATACTCTTGGGCT-3' for FUT4; 5'-ATGTACAAGTGGCCAGCCTA-3' and 5'-GAATCTTCCCTTCCCCAGAG-3' for Lew 1.

Abbreviated terms for each gene are listed in Table 1. The specificity of each amplified product was confirmed by dissociation analyses giving a single sharp

dissociation peak, the absence of the amplified product without reverse transcription, and the appearance of a band of the expected size on electrophoresis of the amplified product. For the amplification of β -actin, Actb primers (Rn00667869, Applied Biosystems) and TaqMan PCR master mix (Applied Biosystems) were used. For relative quantification, standard curves of the threshold cycle (Ct) of amplification of each target against log ng total RNA were created using cDNA samples which showed the lowest Ct value in preliminary runs, and relative quantification was performed for each sample. All quantified values were normalized to those of β -actin (quantified value for a certain target/quantified value for β -actin).

Tissue preparation for histology and goblet cell count

A segment of the jejunum 22-26 cm distal to the pyloric ring and a segment of ileum 6 -10 cm from the ileocecal junction were removed, opened longitudinally, fixed in 4% buffered formalin overnight and embedded in paraffin in such a position that histological sections could be cut perpendicular to the luminal surface. Five-micrometer sections were cut and the periodic acid-Schiff (PAS) reaction with hematoxylin nuclear staining was carried out. Ten villi, which were cut as nearly perpendicularly as possible, were selected per animal, and the numbers of goblet cells and numbers of epithelial nuclei in each villus were counted under a microscope. Goblet cell number/100 epithelial cells was calculated as [Number of goblet cells/number of epithelial nuclei] x 100. The average number of goblet cells/100 epithelial cells in 10 villi was used as the representative value in a given animal, and means and SE of 4 animals were calculated.

Worm counts

After removing the jejunal and ileal segments for separation of the epithelium and tissue section preparation, the numbers of worms in other parts of the small intestine were determined by the saline incubation method.

Statistical analysis

Student's *t*-test (2-tailed) was employed for statistical analysis; a *P* value of less than 0.05 was considered significant.

RESULTS AND DISCUSSION

Expression of goblet cell- and glycosylation-related genes along the gastrointestinal tract

The relative expression levels of goblet cell- and mucin glycosylation-related genes in the normal BN rat gastrointestinal tract were examined by semi-quantitative RT-PCR (Table 1). Except for Lew1 and 3ST2, the goblet cell- and mucin synthesis-related genes examined were expressed at significantly higher levels in the colon than in the jejunum. This may partly reflect the greater abundance of goblet cells in the colon than in the small intestine, in which absorptive cells are the major epithelial constituent, and is consistent with previous reports that MUC2 and MUC4 gene/protein expression levels were higher in the colon than in the small intestine (14,16). In the stomach, MUC2, MUC3 and MUC4 expression was undetectable, while abundant expression of glycosylation-related genes was observed. This is consistent with the fact that MUC5AC and MUC6 are the major mucin core peptides in the stomach (29).

Goblet cell hyperplasia during the course of Nippostrongylus brasiliensis infection in the small intestine

N. brasiliensis larvae reach the small intestine as early as 2-3 days after cutaneous infection, develop to sexual maturity, and begin to lay eggs by 7 days post-infection (PI). However, infection does not continue for a long time: in the normal rat, the majority of adult worms are rejected from the small intestine around 14 days PI by a T-cell dependent mechanism, leaving only a small number of residual worms in the intestine (5, 30). To examine the kinetics of goblet cell response during the time course of infection, BN rats were infected with 2,000 L3 larvae of *N. brasiliensis* and autopsied 7, 14 and 21 days PI. The numbers of worms recovered from the intestine excluding the intestinal segments used for tissue preparation and epithelial separation were 345.8 ± 180.4 , 40.0 ± 17.1 and 42.2 ± 32.9 (average \pm SD) after 7, 14 and 21 days of infection, respectively, showing that the majority of worms were rejected from the intestine by 14 days PI, but a small number of worms escaped the rejection and continued to parasitize the rat at least until 21 days PI. PAS staining of the jejunal and ileal tissue sections revealed goblet-cell hyperplasia 7 and 14 days PI, while goblet cell numbers decreased to preinfection levels by 21 days PI (Table 2).

Alterations of goblet cell- and glycosylation-related gene expression during the course of Nippostrongylus brasiliensis infection in the small intestine

At least 3 types of mucins are expressed in the small intestine: MUC2, which is restricted to goblet cells, MUC3, which is expressed in both columnar and goblet cells, and MUC4, which is expressed in columnar cells (13-16). RT-PCR analyses of gene expression in isolated epithelial cells of BN rats showed upregulation of MUC2, MUC3 and MUC4 in the jejunum, but not in the ileum after infection (Fig. 1), indicating that upregulation of mucin core peptide genes occurred mainly in the local mucosa, which

was parasitized by large numbers of worms. In F-344 rats, similar kinetics of gene expression were observed after nematode infection (data not shown). It has been reported that MUC2 and MUC3 mRNA expression was increased in the small intestinal epithelium after infection with the epithelium-invading nematode *T. spiralis* in mice (31). The present findings that not only MUC2 and MUC3, but also MUC4 mRNA expression was upregulated after infection suggest that the mucin response to nematode infection might have occurred not only in goblet cells, but also in columnar cells.

The gene expression of goblet cell-specific non-mucin peptide Relm β , whose expression is dependent on Th2 cytokines and has a potentially important role in protective immunity (11,17), was also upregulated after infection (Fig. 1), consistent with previous reports showing that Relm β expression was induced after infection of mice with *Trichuris muris*, *T. spiralis* and *N. brasiliensis* (11,17). The finding that strong upregulation of Relm β occurred not only in the jejunum, but also in the ileum, where scarcely any parasites were found, may reflect the systemic effect of Th2 cytokines. TFF3, which might protect mucous epithelia from a range of insults and is known to contribute to mucosal repair (18,19), was also upregulated, consistent with its upregulation in *T. spiralis*-infected mice (32). In a previous report, TFF3 gene expression was not altered in mice infected with *N. brasiliensis* (33); the discrepancy may possibly be due to the difference of host species.

Not only the production and secretion of mucins and goblet cell-specific peptides, but also the glycosylation status of the terminal sugar chains of mucins and/or membrane glycoproteins might have an important role in the colonization of nematodes in and/or rejection of nematodes from the intestine. Although a variety of glycosyltransferases may be involved in the modulation of the terminal sugar chains of mucins and membrane glycoproteins, little is known about the dynamic changes of these genes after infection, except that the upregulation of Siat 4c, FUT2 or A-type

transferase has been reported in mice or rats after infection with *T. spiralis* or *N. brasiliensis* (9-11). The present results showed that the gene expression of Siat 4c, 3ST1, 3ST2, FUT2 and Lew1 was upregulated after infection (Fig. 1), showing for the first time that not only sialyltransferase and fucosyltransferase but also sulfotransferase gene expression changed during nematode infection. The slight upregulation of the FUT2 and Lew1 genes after infection suggests that some changes also occurred in the histo-blood antigens after infection. Although the upregulation of these genes was transient and the levels of the mRNAs returned to steady state levels by 21 days PI, 3ST2 expression was instead upregulated 21 days PI, suggesting that the expression of the two sulfotransferases is regulated by different mechanisms.

Several types of glycosylation-related genes, especially O-sulfotransferase genes, are considered to be specifically expressed in mast cells, which synthesize sulfated glycosaminoglycans such as heparin and/or chondroitin sulfate (34). It is well established that the number of mast cells increases not only in the propria mucosa, but also in the epithelium, after *N. brasiliensis* infection (35). Thus, intraepithelial mast cells contaminating the epithelial fraction might be responsible for the upregulation of 3ST1 and/or 3ST2 after infection. To clarify this point, we examined the gene expression in the epithelial fraction of *Ws/Ws* rats, which have a small deletion of the *c-kit* gene and consequently lack mast cells (35). As shown in Fig. 2, the expression of Relm β , Siat4c and 3ST1 was upregulated 7 days PI, and that of 3ST2 was upregulated 21 days PI, indicating that mast cells are not relevant to the expression of these genes. These results indicate that the glycosylation status of mucins and/or membrane glycoproteins was modified markedly during the course of infection. Although the precise roles of these changes in inflammatory conditions still remain to be elucidated, some reports have indicated a crucial role of the sialylation and sulfation levels of mucins in *Strongyloides venezuelensis*-infected mice and rats, with the sulfation levels

in goblet cells affecting the establishment and distribution of the nematodes in the intestine (36, 37).

Since *N. brasiliensis* larvae reach the small intestine as early as 2-3 days PI, early responses of goblet cell- and mucin glycosylation-related genes after infection were examined. As shown in Fig. 3, MUC2, TFF3 and Siat 4c expression in the jejunal epithelium was upregulated as early as 2-4 days after infection.

Taken together, the present findings show that the expression of each goblet cell- and glycosylation-related gene was altered differently during the course of infection (Fig. 4). The early onset of Siat4c, MUC2 and TFF3 gene upregulation might be associated with innate immunity to pathogens or may be related to an acute phase reaction. MUC3, MUC4, Relm β and 3ST1 gene upregulation, which occurred in the adult-worm colonization and rejection stages, might be regulated by different mechanisms from those of early activated genes, possibly through T-cell- or Th2 cytokine-dependent acquired immunity. Moreover, the upregulation of sialyl- and sulfo-transferase genes suggests that the sialylation and sulfation status of the mucins, membrane glycoproteins and/or histo-blood group antigens might be altered dramatically during nematode infections. The results imply that goblet cell and/or mucin responses to infection are a far more complicated series of changes than previously thought, involving the sequential expression of many factors that can individually or coordinately affect the colonization or rejection of pathogens. Further elucidation of the roles and mechanisms regulating the expression of each mucin and non-mucin secretory peptide and regulating the glycosylation status would lead to development of new strategies for the treatment or prevention of not only intestinal nematode infection but also infection with other pathogens.

Acknowledgements

This work was supported by a grant-in-aid for scientific research (B16390127) from the Japan Society for the Promotion of Science, and research grants from the Ministry of Health, Labour and Welfare, Japan.

REFERENCES

1. WHO Expert Committee. Prevention and control of schistosomiasis and soil-transmitted helminthiasis. World Health Organization Technical Report Series 2002;912:1-57.
2. Lawrence CE. Is there a common mechanism of gastrointestinal nematode expulsion?. *Parasite Immunol* 2003;25:271-81.
3. Miller HR. Gastrointestinal mucus, a medium for survival and for elimination of parasitic nematodes and protozoa. *Parasitology* 1987;94:S77-S100.
4. Nawa Y, Ishikawa N, Tsuchiya K, Horii Y, Abe T, Khan AI, et al. Selective effector mechanisms for the expulsion of intestinal helminths. *Parasite Immunol* 1994;16:333-8.
5. Ogilvie BM, Jones VE. *Nippostrongylus brasiliensis*: a review of immunity and host-parasite relationship in the rat. *Exp Parasitol* 1971;29:138-77.
6. Ishikawa N, Horii Y, Nawa Y. Immune-mediated alteration of the terminal sugars of goblet cell mucins in the small intestine of *Nippostrongylus brasiliensis*-infected rats. *Immunology* 1993;78:303-7.
7. Ishikawa N, Horii Y, Oinuma T, Suganuma T, Nawa Y. Goblet cell mucins as the selective barrier for the intestinal helminths: T-cell-independent alteration of goblet cell mucins by immunologically 'damaged' *Nippostrongylus brasiliensis* worms and its significance on the challenge infection with homologous and heterologous parasites. *Immunology* 1994;81:480-6.
8. Karlsson NG, Olson FJ, Jovall PA, Andersch Y, Enerback L, Hansson GC. Identification of transient glycosylation alterations of sialylated mucin oligosaccharides during infection by the rat intestinal parasite *Nippostrongylus brasiliensis*. *Biochem J* 2000;350:805-14.

9. Olson FJ, Johansson ME, Klinga-Levan K, Bouhours D, Enerback L, Hansson GC, et al. Blood group A glycosyltransferase occurring as alleles with high sequence difference is transiently induced during a *Nippostrongylus brasiliensis* parasite infection. *J Biol Chem* 2002;277:15044-52.
10. Holmen JM, Olson FJ, Karlsson H, Hansson GC. Two glycosylation alterations of mouse intestinal mucins due to infection caused by the parasite *Nippostrongylus brasiliensis*. *Glycoconj J* 2002;19:67-75.
11. Knight PA, Pemberton AD, Robertson KA, Roy DJ, Wright SH, Miller HR. Expression profiling reveals novel innate and inflammatory responses in the jejunal epithelial compartment during infection with *Trichinella spiralis*. *Infect Immun* 2004;72:6076-86.
12. Pemberton AD, Knight PA, Gamble J, Colledge WH, Lee JK, Pierce M, et al. Innate BALB/c enteric epithelial responses to *Trichinella spiralis*: inducible expression of a novel goblet cell lectin, intelectin-2, and its natural deletion in C57BL/10 mice. *J Immunol* 2004;173:1894-901.
13. Audie JP, Janin A, Porchet N, Copin MC, Gosselin B, Aubert JP. Expression of human mucin genes in respiratory, digestive, and reproductive tracts ascertained by in situ hybridization. *J Histochem Cytochem* 1993;41:1479-85.
14. Matsuoka Y, Pascall JC, Brown KD. Quantitative analysis reveals differential expression of mucin (MUC2) and intestinal trefoil factor mRNAs along the longitudinal axis of rat intestine. *Biochim Biophys Acta* 1999;1489:336-44.
15. Sheng Z, Wu K, Carraway KL, Fregien N. Molecular cloning of the transmembrane component of the 13762 mammary adenocarcinoma sialomucin complex. A new member of the epidermal growth factor superfamily. *J Biol Chem* 1992;267:16341-6.

16. Rong M, Rossi EA, Zhang J, McNeer RR, van den Brande JM, Yasin M, et al. Expression and localization of Muc4/sialomucin complex (SMC) in the adult and developing rat intestine: implications for Muc4/SMC function. *J Cell Physiol* 2005;202:275-84.
17. Artis D, Wang ML, Keilbaugh SA, He W, Brenes M, Swain GP, et al. RELMbeta/FIZZ2 is a goblet cell-specific immune-effector molecule in the gastrointestinal tract. *Proc Natl Acad Sci U S A*. 2004;101:13596-600.
18. Suemori S, Lynch-Devaney K, Podolsky DK. Identification and characterization of rat intestinal trefoil factor: tissue- and cell-specific member of the trefoil protein family. *Proc Natl Acad Sci U S A*. 1991;88:11017-21.
19. Taupin D, Podolsky DK. Trefoil factors: initiators of mucosal healing. *Nat Rev Mol Cell Biol* 2003;4:721-32.
20. Hounsell EF, Feizi T. Gastrointestinal mucins. Structures and antigenicities of their carbohydrate chains in health and disease. *Med Biol* 1982;60:227-36.
21. Delmotte P, Degroote S, Merten MD, Van Seuning I, Bernigaud A, Figarella C, et al. Influence of TNFalpha on the sialylation of mucins produced by a transformed cell line MM-39 derived from human tracheal gland cells. *Glycoconj J* 2001;18:487-97.
22. Delmotte P, Degroote S, Lafitte JJ, Lamblin G, Perini JM, Roussel P. Tumor necrosis factor alpha increases the expression of glycosyltransferases and sulfotransferases responsible for the biosynthesis of sialylated and/or sulfated Lewis x epitopes in the human bronchial mucosa. *J Biol Chem* 2002;277:424-31.
23. Hull SR, Laine RA, Kaizu T, Rodriguez I, Carraway KL. Structures of the O-linked oligosaccharides of the major cell surface sialoglycoprotein of MAT-B1 and MAT-C1 ascites sublines of the 13762 rat mammary adenocarcinoma. *J Biol Chem* 1984;259:4866-77.

24. Brockhausen I. Sulphotransferases acting on mucin-type oligosaccharides. *Biochem Soc Trans* 2003;31:318-25.
25. Becker DJ, Lowe JB. Fucose: biosynthesis and biological function in mammals. *Glycobiology* 2003;13:41R-53R.
26. Isshiki S, Kudo T, Nishihara S, Ikehara Y, Togayachi A, Furuya A, et al. Lewis type 1 antigen synthase (beta3Gal-T5) is transcriptionally regulated by homeoproteins. *J Biol Chem* 2003;278:36611-20.
27. Nishida M, Uchikawa R, Tegoshi T, Yamada M, Matsuda S, Sasabe M, et al. Lack of active lung anaphylaxis in congenitally mast cell-deficient *Ws/Ws* rats sensitized with the nematode *Nippostrongylus brasiliensis*. *APMIS* 1998;106:709-16.
28. Hyoh Y, Ishizaka S, Horii T, Fujiwara A, Tegoshi T, Yamada M, et al. Activation of caspases in intestinal villus epithelial cells of normal and nematode infected rats. *Gut* 2002;50:71-7.
29. Buisine MP, Devisme L, Savidge TC, Gespach C, Gosselin B, Porchet N, et al. Mucin gene expression in human embryonic and fetal intestine. *Gut* 1998;43:519-24.
30. Uchikawa R, Yamada M, Matsuda S, Tegoshi T, Nishida M, Kamata I, et al. Dissociation of early and late protective immunity to the nematode *Nippostrongylus brasiliensis* in Brown Norway and Fischer-344 rats. *Parasitology* 1996;112:339-45.
31. Shekels LL, Anway RE, Lin J, Kennedy MW, Garside P, Lawrence CE, et al. Coordinated Muc2 and Muc3 mucin gene expression in *Trichinella spiralis* infection in wild-type and cytokine-deficient mice. *Dig Dis Sci* 2001;46:1757-64.
32. Kamal M, Wakelin D, Ouellette AJ, Smith A, Podolsky DK, Mahida YR. Mucosal T cells regulate Paneth and intermediate cell numbers in the small intestine of *T. spiralis*-infected mice. *Clin Exp Immunol* 2001;126:117-25.

33. Tomita M, Itoh H, Ishikawa N, Higa A, Ide H, Murakumo Y, et al. Molecular cloning of mouse intestinal trefoil factor and its expression during goblet cell changes. *Biochem J* 1995;311:293-7.
34. Wlad H, Maccarana M, Eriksson I, Kjellen L, Lindahl U. Biosynthesis of heparin. Different molecular forms of O-sulfotransferases. *J Biol Chem* 1994;269:24538-41.
35. Arizono N, Kasugai T, Yamada M, Okada M, Morimoto M, Tei H, et al. Infection of *Nippostrongylus brasiliensis* induces development of mucosal-type but not connective tissue-type mast cells in genetically mast cell-deficient *Ws/Ws* rats. *Blood* 1993;81:2572-8.
36. Ishikawa N, Shi BB, Khan AI, Nawa Y. Reserpine-induced sulphomucin production by goblet cells in the jejunum of rats and its significance in the establishment of intestinal helminths. *Parasite Immunol* 1995;17:581-6.
37. Maruyama H, Hirabayashi Y, el-Malky M, Okamura S, Aoki M, Itagaki T, et al. *Strongyloides venezuelensis*: longitudinal distribution of adult worms in the host intestine is influenced by mucosal sulfated carbohydrates. *Exp Parasitol* 2002;100:179-85.

TABLE 1. *Relative expression levels of goblet cell- and mucin glycosylation-related genes in the gut mucosa of normal BN rats. Total RNA was extracted from epithelium separated from the jejunum, ileum or proximal colon, or was extracted from the glandular stomach mucosal scrape, and RT-PCR was performed. Levels of each gene expression were normalized to that of β -actin. In the Table levels in the jejunum were arbitrary expressed as 1.00.*

	Stomach	Jejunum	Ileum	Colon
MUC2 (mucin core peptide 2)	UD	1.0 \pm 0.2	4.5 \pm 1.8	153.2 \pm 49.8*
MUC3 (mucin core peptide 3)	UD	1.0 \pm 0.4	1.5 \pm 0.3	7.5 \pm 3.0*
MUC4 (mucin core peptide 4)	UD	1.0 \pm 0.3	0.7 \pm 0.1	41.0 \pm 13.1*
Relm β (resistin-like molecule β)	2.8 \pm 1.7	1.0 \pm 0.4	0.4 \pm 0.1	47.3 \pm 21.1*
TFF3 (intestinal trefoil factor)	0.2 \pm 0.1	1.0 \pm 0.4	2.7 \pm 0.5*	3.9 \pm 0.4*
Siat 4c (α -2,3-sialyltransferase IV)	3.6 \pm 1.0*	1.0 \pm 0.2	11.7 \pm 4.1*	7.0 \pm 3.0*
3ST1 (3-O sulfotransferase-1)	12.7 \pm 2.2*	1.0 \pm 0.2	1.3 \pm 0.2	15.4 \pm 3.0*
3ST2 (3-O sulfotransferase-2)	1.9 \pm 0.7	1.0 \pm 0.4	1.1 \pm 0.3	3.3 \pm 1.6
FUT1 (α -1,2-fucosyltransferase 1)	1.0 \pm 0.2	1.0 \pm 0.2	1.5 \pm 0.1	13.8 \pm 0.5*
FUT2 (α -1,2-fucosyltransferase 2)	3.6 \pm 0.6*	1.0 \pm 0.1	1.5 \pm 0.3	14.3 \pm 2.8*
FUT4 (α -1,3-fucosyltransferase 4)	4.7 \pm 0.9*	1.0 \pm 0.1	0.7 \pm 0.1*	3.0 \pm 0.2*
Lew 1 (Lewis type 1 antigen synthase: β 1,3-N-acetylglucosaminyltransferase 5)	0.1 \pm 0.0*	1.0 \pm 0.3	0.3 \pm 0.0	2.9 \pm 0.9

Data shown are mean \pm SE of 4 rats. *Significantly different from the levels in the jejunum ($p < 0.05$). UD: undetectable.